GLUON DENSITY IN NUCLEI

A.L.A yala $F^{\circ a}$, M.B.G ay Ducati^{a)} and E.M.Levin^{c)d)y}

^{a)} Instituto de F sica, Univ. Federal do Rio Grande do Sul Caixa Postal 15051, 91501–970 Porto A legre, RS, BRAZIL

^{b)} Instituto de F sica e Matematica, Univ. Federal de Pelotas Campus Universitario, Caixa Postal 354, 96010–900, Pelotas, RS, BRAZIL

> ^{c)} Theory Division, Ferm iNationalAccelerator Laboratory Batavia, IL 60510 - 0500, USA

^{d)} Theory D epartment, P etersburg N uclear P hysics Institute 188350, G atchina, St. P etersburg, RUSSIA

Tak, given by EM. Levin at RHIC '96 Summer Study, BNL, LI, July 7-19, 1996

A bstract: In this talk we present our detail study (theory and numbers) [1] on the shadowing corrections to the gluon structure functions for nuclei. Starting from rather contraversial information on the nucleon structure function which is originated by the recent HERA data, we develop the G lauber approach for the gluon density in a nucleus based on M ueller form ula β] and estimate the value of the shadowing corrections in this case. Than we calculate the rst corrections to the G lauber approach and show that these corrections are big. Based on this practical observation we suggest the new evolution equation which takes into account the shadowing corrections and solve it. We hope to convince you that the new evolution equation gives a good theoretical tool to treat the shadowing corrections for the gluons density in a nucleus and, therefore, it is able to provide the theoretically reliable initial conditions for the tim e evolution of the nucleus – nucleus cascade. The initial conditions should be xed both theoretically and phenom enologically before to attack such m ore complicated problem s as the m ixture of hard and soft processes in nucleus nucleus interactions at high energy or the theoretically reliable approach to hadron or/and parton cascades for high energy nucleus-nucleus interaction.

E-m ail: ayala@ if.ufrgs.br

E-m ailgay@ ifufrgsbr

^y E-m ail: levin@ hep anlgov; leving@ ccsg.tau ac.il

1 Introduction.

The main goal of this talk is to share with you our experience and results that we got during the last two years reconsidering the whole issue of the shadow ing corrections (SC) to the gluon density in nuclei [1]. The title which rejects the key problem s that we are going to discuss is: $\$ All (theory and numbers) about the SC to gluon density in nuclei"

It is well known that the gluon density is the most important physical observable that governs the physics at high energy (low B jorken x) in deep inelastic processes [2]. D ealing with nucleus we have to take into account the shadow ing correction (SC) due to rescattering of the gluon inside the nucleus, which is the main point of interest in this paper. We show that SC can be treated theoretically in the fram ework of perturbative QCD (pQCD) and can be calculated using the inform ation on the behavior of the gluon structure function for the nucleon.

The outline of the talk books as follows. We start with ourm otivation answering the question why we got interested in the SC for nucleus gluon density. In section 3 we will discuss the theory and numerics of the G lauber (M ueller) approach emphasizing it's theory status and the estimates for the SC that came out of it. A fler short discussion in section 4 the rst corrections to the G lauber approach we will present what we consider as a right way of doing, namely, the new evolution equation that sum sall SC (section 5). In section 6 we are going to discuss our next steps that we plan to do in a nearest future, while in section 7 we will give our answer to the hot question: and what ?, trying to collect all problem s of RHIC physics that we will be able to answer using our approach.

2 M otivation.

Let us start with a brief summary of the HERA results for the nucleon structure functions (parton densities in a nucleon). The experiment [4] shows that the deep inelastic structure function $F_2(x;Q^2)$ increases in the region of small x (at high energies):

$$F_2(x;Q^2) / \frac{1}{x^{0.2}}$$
 for $10^2 > x > 10^5$

at large and sm all (Q^2 1 2 GeV^2) values of the photon virtualities Q^2 .

At rst sight we can conclude from the analysis based on the D G LA P evolution equations undertaken through all the world R [6] [7] [8] that :

1. The DGLAP evolution equations work quite well and no other ingredients are needed to describe all the HERA data.

2. The parton cascade is rather deluted system of partons with small parton – parton interaction which can be neglected in a rst approximation. In other words we do not need any SC to describe the experimental data.

3. The phenom enological input, namely, the quark and gluon distribution at initial virtuality $Q^2 = Q_0^2$ can be chosen at su ciently low values of Q^2 using the backward evolution of the experimental data in the region of Q^2 4 5G eV². Even more, the craziest parameterization that we have seen in our life – the GRV one [6] does it's job perfectly well, starting with $Q_0^2 = 0.3$ G eV²?!.

W hat we have discussed is moreless common opinion of all experts in D IS and one can ind it in many plenary and review talks during the last two years.

However we would like to draw your attention to several facts which do not t to this common scheme:

1. The best parameterization of the HERA data is not the solution of the DGLAP equations but a simple form ula [9]:

$$F_2(x;Q^2) = a + m \log \frac{Q^2}{Q_0^2} \log \frac{x_0}{x}$$

with a = 0.078; m = 0.364; $x_0 = 0.074$; $Q_0^2 = 0.5$ GeV². It is clear that this simple formula cannot be a solution of the DGLAP evolution equations. To make obvious this remark it is enough to recalculate the gluon structure function from the above expression as it has been done in Ref.[9]. Indeed, xG (x;Q²) turns out to be equal to

$$xG(x;Q^2) = 3 \log \frac{x_0}{x}$$

without any Q^2 -dependence within the direct contradiction with the DGLAP evolution.

2. Using the HERA data we can evaluate the parameter which characterizes the value of the SC, namely [2]

$$= \frac{3}{Q^2 R^2} \times G (x; Q^2) ; \qquad (1)$$

where xG $(x;Q^2)$ is the gluon structure function and R^2 is the radius of area populated by gluons in a nucleon. The physical meaning of becomes clear if we rewrite it in the form

= xG (x;Q²)
$$\frac{(GG)}{R^2}$$
;

where (GG) is the cross section of two gluon interaction in our parton cascade calculated by M ueller and Q in [10], namely, (GG) = $\frac{3^2 s}{Q^2}$. The physical meaning of this form what is the probability of the gluon – gluon interaction inside the parton cascade. It looks very natural if we compare eq. (1) with the small parameter for proton – nucleus interaction. Indeed, the



Figure 1: The J= production without a) and with b) dissociation of the proton.

parameter which governs the value of the G lauber corrections for proton – nucleus interaction $_{pA} = A \frac{(pp)}{R_A^2}$, where A is the number of the constituents (nucleons), (pp) is the cross section of the interaction of our constituents and R_A^2 is the area populated by nucleons. The question arises what is the value of R in eq. (1)? Using the new HERA data on photoproduction of J/ m eson [11] we are able to estimate the value of R² in the de nition of (see eq. (1)). To illustrate the point we picture in Fig.1 the process of J/ photoproduction in the additive quark model (AQM). We see that we have two processes with di erent slopes (B) in t (or in b_2^2): the J/ production without (Fig.1a) (B $_{el} = 5 \text{ GeV}^2$) and with (Fig.1b) (B $_{in} = 1.66 \text{ GeV}^2$) dissociation of the proton. The AQM gives us the simplest estimates for the resulting slope (R²) in eq. (1) if we neglect any slope from the Pomeron – J/ vertex in Fig.1, nam ely

$$\frac{1}{R^2} = \frac{1}{4} f \frac{3}{2B_{el}} + \frac{1}{2B_{in}} g = \frac{1}{5} G eV^2 :$$
 (2)

F ig 2 shows the contour plot for using the GRV parameterization [6] for the gluon structure function and the value of $R^2 = 5G eV^2$. One can see that reaches = 1 at HERA kinematic region, meaning shadowing corrections take place.

3. The situation boks even m ore contraversparam eterizationial if we plot the average value of the anomalous dimension $< > = 0 \ln (xG (x;Q^2))=0 \ln Q^2$ in the GRV parameterization. Fig.3 shows two remarkable lines: < > = 1, where the deep inelastic cross section reaches the value compatible with the geometrical size of the proton, and < > = 1=2, which is the characteristic line in whose vicinity both the BFKL Pomeron (see Ref.[12]) and the GLR equation [2] should take over the DGLAP evolution equations.

W ewill discuss below the de nition of the anom alous dimension and why this ratio is the average anom alous dimension.



Figure 2: Contour plot for for $R^2 = 5G ev^2$.

the BFKL and the GLR equations, what we need to know right now, is only the fact that both equations give the signal of the new physics. The HERA data passed over the second line and even for su ciently small values of Q² they crossed the rst one without any indication of a strange behaviour near these lines.

Concluding this brief sum mary of the HERA data and physics behind them we would like to repeat that to our taste the situation at HERA looks very controversial and the statement that the DGLAP evolution works is institut not the last outcome of the HERA data. On the other hand we have to develop the new approach to the SC, more general than the GLR one, which will allow us to give reliable estimates for the SC in the kinematic region to the left of the line < > = 1=2. This is why we decided to reconsider everything that has been known about the SC, trying to forget everything that we knew about them, and to start our analisys of the SC from the very beginning. We also decide to choose the gluon density in a nucleus as a laboratory or training ground for the new approach to the problem of the SC.

We have three reasons for such a choice: (i) the nucleus D IS is easier to handle theoretically, as we will show in the main body of our talk; (ii) the previous analysis of the SC shows that this is mainly density e ect in the parton cascade (see review [13] for example) and we anticipate larger gluon density for D IS with a nucleus; (iii) the R H IC is coming and the gluon density in nuclei will provide the initial condition for any phenom enological cascades for nucleus – nucleus interaction at high energies.

3 The Glauber approach in QCD

3.1 The Mueller formula.



Figure 3: Contours for < > = 1 and < = 1/2.



Figure 4: The structure of the parton cascade in the G lauber (M ueller) form u.a. A denotes the nucleus, N – the nucleon, G (Q²) – the virtual gluon and $_{\rm N}$ (r_t^2) is the nucleon cross section.

The idea how to write the G lauber form ula in QCD was originally form ulated in two papers Ref.[14] and Ref. [3]. However, the key paper for our problem is the second paper of A. M ueller who considered the G lauber approach for the gluon structure function. Nevertheless, it is easier to explain the main idea considering the penetration of quark – antiquark pair, produced by the virtual photon, through the target. W hile the boson projectile is traversing the target, the distance r_2 between the quark and anti-quark can vary by amount $r_2 / R_A \frac{k_t}{E}$ where E denotes the energy of the pair in the target rest fram e and R_A is the size of the target (see Fig.4). The quark transverse momentum is $k_t / 1=r_2$. Therefore

$$r_{?} / R \frac{k_{t}}{E} r_{?};$$
 (3)

and is valid if

$$r_{2}^{2} s 2m R;$$
 (4)

where s = 2m E. In term s of B jorken x, the above condition looks as follows

$$x \qquad \frac{1}{2m R}: \tag{5}$$

Therefore the transverse distance between quark and antiquark is a good degree of freedom [14][3][17]. A shas been shown by A M ueller, not only quark – antiquark pairs can be considered in a such way. The propagation of a gluon through the target can be treated in a sim ilar way as the interaction of gluon – gluon pair with de nite transverse separation r_t with the target. It is easy to understand if we remem ber that virtual colorless graviton or H iggs boson is a probe of the gluon density.

The total cross section of the absorption of gluon (G $\,$) with virtuality Q 2 and B jorken x can be written in the form :

$${}^{A}_{tot}(G) = {}^{Z_{1}}_{0} dz {}^{Z} \frac{d^{2}r_{t}}{2} {}^{Z} \frac{d^{2}b_{t}}{2} {}^{G}_{2} (Q^{2};r_{t};x;z) {}_{A}(x;r_{t}^{2}) [{}^{G}_{2} (Q^{2};r_{t};x;z)]; (6)$$

where z is the fraction of energy which is carried by the gluon, $\frac{G}{2}$ is the wave function of the transverse polarized gluon and $_{A}(x;r_{t}^{2})$ is the cross section of the interaction of the GG - pair with transverse separation r_{t} with the nucleus. This cross section can be written in the form :

$$_{A}(x;r_{?}^{2}) = 2 d^{2}b_{L} \operatorname{Im} a(x;r_{?};b_{L});$$
 (7)

where a is the elastic amplitude for which we have the s-channel unitarity constraint:

$$2 \operatorname{Im} a(x; r_{2}; b_{1}) = \frac{1}{2} (x; r_{2}; b_{1}) + G_{in} (x; r_{2}; b_{1}); \qquad (8)$$

where G_{in} is the contribution of all the inelastic processes. Let us recall that two terms in eq. (8) have di erent physical meaning: the left hand side and the statem in the right hand

side describe the interference between the incoming plane wave and outgoing spherical wave which amplitude is the elastic scattering amplitude (a). These two terms cannot be calculated using a classical approach or simple M onte Carlo – like model. The Quantum M echanics of the interaction is mostly absorbed in these two terms while the last term has a simple probabilistic meaning, namely, the probability of any inelastic interactions, and can be treated almost classically and, for certain, in the probabilistic way, for example in M onte Carlo-like models. The unitarity establishes the correlation between two unknowns a and G in and has the general solution:

$$a(x;r_{?};b_{t}) = if 1 \qquad e^{\frac{1}{2}(x;r_{?};b_{t})} g;$$
(9)

$$G_{in}(x;r_{?};b_{t}) = 1 \qquad e^{(x;r_{?};b_{t})} :$$

One can see that has a simple physical meaning, namely e is the probability that GG-pair has no inelastic interaction during the passage through the target. The opacity is an arbitrary real function, which can be specified only in more detail theory or model approach than the unitarity constraint. One of such specific model is G lauber approach or E ikonal model.

However, before we will discuss thism odel let us make one in portant remark on the strategy of the approach to the SC . We are trying to built a model or theory for the total cross section (or for the gluon structure function) not because the SC should be the strongest one in this particular observable, but because if we will be able to calculate opacity we will have the theory or model for all inelastic processes. Indeed, using AGK cutting rules [15] we can calculated any inelastic process, if we know , in accordance with the s-channel unitarity. It is worthwhile mentioning that the inverse procedure does not work. If we know the SC in all details for a particular inelastic process, say for the inclusive production, we cannot reconstruct all other process and the total cross section in particular.

Now, let us built the G lauber approach. First, let us assume that is small (1) and it's b dependence can be factorised as $= {}^{e}(x;r_{?})S(b)$ with the normalization: ${}^{R}d^{2}b S(b_{?}) = 1$. Expanding eq. (9) and substituting it in eq. (7), one can obtain:

$$_{A}(x;r_{?}) = e(x;r_{?})$$
 (10)

At small the cross section of the deep inelastic process with a nucleus is proportional to the number of nucleons in a nucleus (A), namely,

$$A(x;r_{?}) = A_{N}(x;r_{?})$$
:

To calculate ^e we need to substitute everything in eq. (6) and use the formula for $_{A}(G) = \frac{4}{Q^{2}} x G_{A}(x;Q^{2})$ as well as the expression for the wave function of the GG-pair in the virtual gluon probe. Such calculations has been done in Ref.[3] and we recapture here the result (see for example Ref. [1] form ore details):

$$e = A_{N}(x;r_{?}) = \frac{3^{2}}{4}r_{?}^{2}xG(x;\frac{4}{r_{?}^{2}}):$$
 (11)

The G lauber (eikonal) approach is the assumption that $= {}^{e}S(b_{t}) \text{ with } {}^{e}$ of eq. (11) not only in the kinematic region where is small but everywhere. From the point of view of the structure of the nal state this assumption means that the rich typical inelastic event was modeled as a sum of the diraction dissociation of GG – pair plus uniform in rapidity distribution of produced gluons. For example, we neglected in the G lauber approach all rich structure of the large rapidity gap events including the diractive dissociation in the region of large mass.

Substituting everything in eq. (11) and eq. (6) and using the wave function calculated by M ueller in $\operatorname{Ref}[\beta]$ we obtain the G lauber (M ueller) form ula for the gluon structure function:

$$xG_{A}(x;Q^{2}) = \frac{4}{2} \frac{x^{2}}{x} \frac{dx^{0}}{x^{0}} \frac{d^{2}r_{t}}{r_{t}^{4}} \frac{d^{2}r_{t}}{r_{t}^{4}} \frac{d^{2}r_{t}}{r_{t}^{4}} \frac{d^{2}r_{t}}{r_{t}^{4}} 2^{n} \frac{d^{2}r_{t}}{r_{t}^{2}} \frac{d^{2}r_{t}}{r_{t}^{2}} (x^{0}r_{t}^{2}) \frac{d^{2}r_{$$

It is easy to see that the rst term in the expansion of eq. (12) with respect to gives the DGLAP equation in the region of sm all x.

To calculate the pro le function S (b_t) we make the usual assumption that $\langle b_t^2 \rangle_N$ in the interaction of G G - pair with the nucleon is much smaller than the nucleus radius ($\langle b_t^2 \rangle_N$ R_A. Therefore, S (b_t) can be expressed through the nucleon wave function in a nucleus, namely

$$S_{A}(q_{z};b_{t}) = dz_{1} e^{iq_{z}z_{1}} a_{A}(z_{1};b_{t};r_{2};:::r_{1};r_{A}) (z_{1};b_{t};r_{2};:::r_{1};r_{A})$$

$$(z_{1};b_{t};r_{2};:::r_{1};r_{A}) a^{A} r_{1};$$
(13)

where the wave function is norm alized as

^Z
_A
$$(z_1; b_c; r_2; ::: r_i; r_A)$$
 $(z_1; b_c; r_2; ::: r_i; r_A) \int_{i=1}^{Y^A} d^3 r_i = A$: (14)

A sum ing that there is no correlation between nucleons in a nucleus and the simple G aussian form of a single nucleon wave wave function we derive the G aussian parameterization for $S(b_{c})$, namely

$$S_{A}(q_{z}; b_{1}^{2}) = \frac{A}{R_{A}^{2}} e^{\frac{b_{1}^{2}}{R_{A}^{2}}} e^{\frac{R_{A}^{2}}{R_{A}^{2}}} ;$$
 (15)

where the mean radius $R_{\rm A}^{\,2}\,$ is equal to

$$R_A^2 = \frac{2}{5} R_W^2 S$$

and R_{WS} is the size of the nucleus in the W ood-Saxon parameterization. We choose $R_{WS} = r_0 A^{\frac{1}{3}}$ with $r_0 = 1.3$ fm in all our calculation. We are doing all calculation in the rest frame of the nucleus where we can neglect the change of energy for the recoil nucleon in the nonrelativistic

theory for the nucleus. Indeed, its energy is $E_{p^0} = m + \frac{q^2}{2m}$ and $\frac{q^2}{2m} = q_z$. At high energy (sm all x) we can neglect also the q_z -dependance (see Ref.[1] for details).

U sing G aussian parameterization for S (b) (see eq. (15)) we can take the integral over b_t and obtain the answer (N $_c$ = N $_f$ = 3)

$$xG_{A}(x;Q^{2}) = \frac{2R_{A}^{2}}{2} x^{0} \frac{1}{x} \frac{dx^{0}}{x^{0}} \frac{z}{\frac{1}{Q^{2}}} \frac{dr_{t}^{2}}{r_{t}^{4}} C + \ln(g(x^{0};r_{t}^{2}))E_{1}(g(x^{0};r_{t}^{2}))^{0}$$
(16)

where C is the Euler constant and E_1 is the exponential integral (see Ref.[23] Eq. 5.7.11) and

$$_{G} (\mathbf{x}^{0}; \mathbf{r}_{t}^{2}) = \frac{3 _{S} A r_{t}^{2}}{2 R_{A}^{2}} \mathbf{x}^{0} G_{N}^{D G LAP} (\mathbf{x}^{0}; \frac{1}{r_{t}^{2}})$$
(17)

The eq. (12) is the master equation of this section and it gives a way to estimate the value of the SC.We would like to stress that we have only adjusted the approach of Ref. [3] to the rescattering in a nucleus. It means that we did nothing except that we share the responsibility with A.Mueller for eq. (12).

O ne can see that the M ueller form ula of eq. (12) depends only on . If is small (1), we can expand eq. (16) and obtain the DGLAP evolution equation for the gluon structure function. If 1, we can use the asymptotic form ula for E₁ and obtain:

$$xG_{A}(x;Q^{2}) = \frac{2R_{A}^{2}}{2} \int_{x}^{2} \frac{1}{x^{0}} \frac{dx^{0}}{x^{0}} \int_{\frac{1}{Q_{0}^{2}(x^{0})}}^{\frac{1}{Q_{0}^{2}}} \frac{dr_{2}^{2}}{r_{2}^{4}} C + \ln(G(x^{0};r_{2}^{2})^{0};$$

where Q_0^2 (x⁰) is the solution of the equation:

$$_{\rm G} ({\rm x}^0; {\rm r}_{\rm t}^2 = \frac{1}{Q_0^2 ({\rm x}^0)}) = 1 :$$
 (18)

In Fig.5 are plotted the contours of for a nucleon target that give an idea in which kinem atic region we expect big SC.

3.2 Theory status of the M ueller form ula.

In this section we shall recall the main assumptions that have been made to obtain the Mueller formula.

1. The gluon energy (x) should be high (sm all) enough to satisfy eq. (1) and $_{\rm S} \ln (1=x)$ 1. The last condition means that we have to assume the leading $\ln (1=x)$ approximation of perturbative QCD for the nucleon gluon structure function.

2. The DGLAP evolution equations hold in the region of small x or, in other words, $_{\rm S}\ln(1=r_{\rm P}^2)$ 1.0 ne of the lessons from HERA data is the fact that the GLAP evolution can describe the experimental data.



Figure 5: The contours of for Nucleon, Ca and Au.

These two assumptions mean that we describe the gluon emission in so called D ouble Log Approximation (DLA) of perturbative QCD, or in other words, we extract from each Feynman diagram of the order $\binom{n}{S}$ the contribution of the order ($_{S}$ ln 1=x ln Q²=Q₀²)ⁿ, neglecting all other contributions of the same diagram. In term softhe DGLAP evolution, we have to assume that the DGLAP evolution equations describe the gluon emission in the region of small x. However, the rst assumption is very important for the whole picture, since it allows us to treat successive rescatterings as independent and simplifies all formulae reducing the problem to an eikonal picture of the classical propagation of a relativistic particle with high energy (E 1 , where is the scattering radius in the nuclear matter) through the nucleus. The second one simplifies calculations but we can consider the BFKL evolution [12] instead of the DGLAP one.

3. Only the fastest partons (GG pairs) interact with the target. This assumption is an artifact of the G lauber approach, which looks strange in the parton picture of the interaction. Indeed, in the parton m odel we rather expect that all partons not only the fastest ones should interact with the target. In the next section we will show that corrections to the G lauber approach due to the interaction of slower partons are essential in QCD too.

4. There are no correlations (interaction) between partons from the di erent parton cascades (see Fig.4). This assumption means that even the interaction of the fastest GG-pair was taken into account in the M ueller form ula only approximately and we have to assume that we are dealing with large number of colours to trust the M ueller form ula. Indeed, it has been proven that correlations between partons from di erent parton cascades lead to corrections to the M ueller form ula of the order of $1=N_c^2$, where N_c is the number of colours (see R ef.[1] and references therein for detail discussions on this subject).

5. There are no correlations between di erent nucleons in a nucleus.

6. The average b for GG pair-nucleon interaction is much smaller than R_A .

The last two are usual assumptions to treat nucleus scattering. We have used the specic G aussian parameterization for b_t dependence. A loo, one can easily generalize our formula in more general case, as W ood-Saxon parameterization [18].

3.4 The modi ed Mueller form ula.

The next step of our approach is to give an estim ate of the SC using the M ueller form ula. However, before doing so, we have to study how well works the DLA of perturbative QCD which was heavily used in the derivation of the M ueller form ula. Let us recall that the solution of the DGLAP evolution equations can be easily found in the moments space. For any function f(x) we denote the moment f(!) as

$$f(!) = \int_{0}^{Z_{1}} dx x^{!} f(x) :$$

Note that the moment variable ! is chosen such that the ! = 0 moment m easures the number of partons, and the moment ! = 1 m easures their momentum. An alternative moment variable N = ! 1 is often found in the literature. The x-distribution can be reconstructed by considering the inverse M ellin transform, which for the gluon distribution reads:

$$xG(x;Q^{2}) = \frac{1}{2 i} \int_{C}^{Z} d! g_{in}(!;Q^{2}_{0}) e^{! \ln(1=x) + (!) \ln(Q^{2}=Q^{2}_{0})};$$
(19)

where the contour of integration C is taken to the right of all singularities and function g_{in} is defined by the initial gluon distribution at $Q^2 = Q_0^2$. The anomalous dimension (!) has to be calculated in perturbative QCD and can be written in the form :

$$(!) = \frac{SN_{c}}{!} + \frac{2 \frac{4}{S}N_{c}^{4}}{4} \frac{1}{!^{4}} + O(\frac{5}{S}) + O(S) :$$
(20)

In the DLA we take only the 1st term of this series, namely,

$$^{\text{DLA}} = \frac{\text{SN}_{\text{C}}}{!} \frac{1}{!};$$

In the BFKL evolution equation all terms of the order $\left(\frac{s}{!}\right)^n$ have to be taken into account. They generate the BFKL anom alous dimension of the form :

$${}^{BFKL}(!) = \frac{{}^{SN_{c}}}{!} \frac{1}{!} + \frac{2 {}^{4}_{S}N_{c}^{4}}{4} \frac{3}{!^{4}} + \frac{X^{1}}{{}^{1}_{n=5}}C_{n} \left(\frac{s}{!}\right)^{n} j_{!!L}! \frac{1}{2} + \frac{1}{2} + \frac{1}{!} \frac{1}{!$$

where ${}^{BFKL}(! = !_L) = 1=2$. The main qualitative property of the BFKL anom abus dim ension is the fact that it cannot exceed the value 1/2.

The momentum conservation means that (! = 1) = 0. None of the DLA or the BFKL anom alous dimension satisfies this equation, because they give the good approximation to the fullanom alous dimension only in the region of smallvalues of ! or, in other words, in the region of small x.

The DLA anom alous dimension leads to the simple evolution equation:

$$\frac{(e^2 \times G (x; Q^2))}{(e \ln (1=x))(e \ln Q^2)} = -\frac{s N_c}{s} \times G (x; Q^2) :$$
(21)

S

Now let us estimate how well works the DLA. In all our numerical estimates we use the GRV parameterization [6] for the nucleon gluon distribution, which describes all available experimental data quite well, including recent HERA data at low x. Moreover, GRV is suited for our purpose because (i) the initial virtuality for the GLAP evolution is small ($Q_0^2 = 0.25 \text{GeV}^2$) and we can discuss the contribution of the large distances in MF having some support from experimental data; (ii) in this parameterization the most essential contribution comes from the region where

 $_{\rm s}\ln Q^2$ 1 and $_{\rm s}\ln 1=x$ 1. This allows the use of the double leading log approximation of pQCD, where the MF is proven [3]. It should be also stressed here, that we look at the GRV parameterization as a solution of the DGLAP evolution equations, disregarding how much of the SC has been taken into account in this parameterization in the form of the initial gluon distribution.

However, in spite of the fact that the GLAP evolution in the GRV parameterization starts from very low virtuality (Q_0^2 0.25G eV²) it turns out that the DLA still does not work quite well in the accessible kinematic region ($Q^2 > 1 \text{GeV}^2$; x > 10⁵). To illustrate this statement we plot in Fig.5 the ratio:

$$\frac{\langle \underline{SN_{c}} \rangle}{\underline{SN_{c}}} = \frac{\frac{\ell^{2} \times G^{GRV} (x \mathcal{Q}^{2})}{\ell \ln (1=x) \ell \ln \mathcal{Q}^{2}}}{x G^{GRV} (x; \mathcal{Q}^{2})}$$

This ratio is equal to 1 if the DLA holds. From Fig.5 one can see that this ratio is rather around 1/2 even at large values of Q^2 .



Figure 6: The ration $\frac{\langle s \rangle}{s}$ for di erent values of Q² in the GRV parameterization

We can understand why the corrections to the DLA is so big modeling the complicated expression for of eq. (20) by simple formula [19] $^{\text{y}}$:

$$(!) = \frac{SN_c}{!} \quad f = 1g:$$
 (22)

Eq. (22) has correct the DLA lim it at small! and it satisfies the momentum conservation ((! = 1) = 0). The typical values of ! in all available parametrizations, even in the GRV

 $^{^{\}rm y}$ W e are very grateful to Yu. Dokshitzer for enlighting discussions on this problem during the RHIC '96 W orkshop

, which is the closest to the DLA, is < ! > 0.5. Therefore, we have about 50% correction to the DLA. Therefore, the DLA cannot provide a reliable estimates for the gluon structure function.

On the other hand, our master equation (see eq. (12)) is proven in DLA.W illing to develop a realistic approach in the region of not ultra sm all x ($x > 10^4$) we have to change our master equation (eq. (12)). We suggest to substitute the full DGLAP kernel (the full expression of eq. (20)) in the rst term of the rh.s. This procedure gives

$$xG_{A}(x;Q^{2}) = xG_{A}(x;Q^{2})(eq: (12)) + AxG_{N}^{GRV}(x;Q^{2})$$
$$A - \frac{sN_{c}}{x} \frac{Z_{1}Z_{Q^{2}}}{xQ_{0}^{2}} \frac{dx^{0}}{x^{0}} \frac{dQ^{02}}{Q^{02}} x^{0}G_{N}^{GRV}(x^{0};Q^{02}):$$
(23)

The above equation includes also $A \times G_N^{GRV} (x; Q_0^2)$ as the initial condition for the gluon distribution and gives $A \times G_N^{GRV} (x; Q^2)$ as the rst term of the expansion with respect to $_G$. Therefore, this equation is an attempt to include the full expression for the anom alous dimension for the scattering o each nucleon, while we use the DLA to take into account all SC. Our hope, which we will con rm by numerical calculation, is that the SC are small enough for $x > 10^{-3}$ and we can be not so careful in the accuracy of their calculation in this kinematic region. Going to smaller x, the DLA becomes better and eq. (23) tends to our master equation (12).

The gluon structure function for nucleon (A = 1).

In this subsection we are going to check how eq. (23) describes the gluon structure function for a nucleon, which is our main ingredient in the M ueller form ula. We calculate st the ratio

$$R_{1}^{N} = \frac{xG^{A}(x;Q^{2})(eq:(23))}{xG_{N}^{GRV}(x;Q^{2})};$$
(24)

for A = 1, which is shown in Fig.7. From this ratio we can see the general behavior of the SC as a function of $\ln(1=x)$ and Q^2 . In the region of the HERA data, $3 < \ln(1=x) < 10$, and $Q^2 > 2 \text{ G eV}^2$ [4], the SC are not bigger than 15%. The SC give a contribution bigger than 20% only at very sm all value of x, where we have no experimental data.

In the sem iclassical approach (see [2]), the nucleon structure function is supposed to have Q^2 and x dependence as

$$xG^{N}(x;Q^{2}) / fQ^{2}g^{<} f \frac{1}{x}g^{<} :$$
 (25)

W e can calculate both exponents using the de nitions

<
$$! > = \frac{(! \ln (xG^{N} (x;Q^{2})))}{(! \ln (1=x))}$$
 : (26)



Figure 7: The SC for nucleon (A = 1) as a function of ln (1=x) and Q^2 , where ratio R_1 compares xG^A with xG (GRV) distribution.

$$< > = \frac{(\ln (xG^{N} (x;Q^{2})))}{(\ln (Q^{2}=Q_{0}^{2}))};$$
 (27)

The eq.(26) gives the average value of the elicitie power < ! > of the gluon distribution, $xG(x;Q^2) / x^{<!>}$, which is suitable to study the small x behavior of the gluon distributions. Fig.8 shows the calculation of < ! > the nucleon distribution for eq. (23) and for GRV gluon distribution, both as functions of ln (1=x) for di erent values of Q^2 . From the gure, we can see that the elicitie powers of xG^A (A = 1) and xG (GRV) have the same general behavior in the small x limit but the nucleon distribution is slightly suppressed. We calculate also, in the same kinemical region, the exponent < >, given by eq (27). This is the average value of the anom alous dimension, which describes the elicitie dependence of the distribution in Q^2 variable. Figs.9 shows < > for the nucleon and GRV distributions, indicating that the Q^2 dependence is slightly soften by the SC.

Comparing gures 8 and 9, we can conclude that even these more detailed characteristic of the gluon structure function have not been seriously a ected by the SC in the nucleon case.

W e also use the DGLAP evolution equations to predict the value of the deep inelastic structure function F_2 from the xG^A gluon distribution. Summing the DGLAP evolution equations



Figure 8: The e ective power < ! > calculated for xG A (A = 1) and the GRV distribution.



Figure 9: The elective power of Q 2 dependence calculated for xG $^{\rm A}$ (A = 1) and the GRV distribution.

for each quark avor, the function F_2 may be written [20]

$$F_{2} = \frac{s(Q^{2})}{q} \frac{x}{q} e_{q}^{2} \frac{Q^{2}}{Q_{0}^{2}} \frac{dQ^{2}}{Q^{2}} \frac{z}{1} \frac{x}{z} [z^{2} + (1 - z)^{2}] \frac{x}{1 - z} G^{N} (\frac{x}{1 - z}; Q^{0^{2}})$$
(28)

where the sea quark distributions have been neglected in comparison with the gluon distribution. Fig.10 shows the prediction for F_2 from xG^A and from the GRV distribution, compared with experimental data. A swe can see, the magnitude of the suppression due to the SC is less than 10% in the region of the HERA data and this suppression is smaller than the experimental error.



Figure 10: F_2 from xG^A and the GRV distribution, compared with experimental data [4].

From the above results we can conclude that eq. (23) gives a good description for the gluon structure function for nucleon and describes the available experimental data. Therefore, it can be taken as a correct rst approximation in the approach to the nucleus case.

3.6 The gluon structure function for nucleus.

In the fram ework of perturbative approach it is only possible to calculate the behavior of the gluon distribution at small distances. The initial gluon distribution should be taken from the experiment. A ctually the initial virtuality Q_0^2 should be big enough to guarantee that we are dealing with the leading twist contribution. Our main assumption is that we start the QCD evolution with a small value of Q_0^2 considering that the MF is a good model for high twist contributions in D IS o nucleus.

	$Q^2 = 1 \text{G eV}^2$		$Q^2 = 10G eV^2$	
х	R _{1N}		R _{1N}	
10 ²	0.94	0.0416	0.98	0.014
10 ³	0.92	0.0616	0.94	0.034
10 4	0.88	0.094	0.92	0.0563
10 5	8.0	0.145	0.86	0.093

Table 1: Values of R_{1N} and for parameterization $R_1 = R_{1N} A$

The scale of the SC governs by the value of $_{\rm A}$, namely they are big for $_{\rm A} > 1$ and sm all for $_{\rm A} < 1$. Fig.4 shows the plot of $_{\rm A} = 1$ for di erent nuclei. O ne can see that the SC should be essential for heavy nuclei starting from C a at the accessible experimentally kinematic region.

Now we extend the de nition of R_1 for the nucleus case

$$R_{1} = \frac{xG^{A}(x;Q^{2})}{AxG_{N}^{GRV}(x;Q^{2})};$$
(29)

where the num erator is calculated using eq.(23). Figure 11 shows the results for the calculations of R₁ as a function of the variables $\ln (1=x)$, $\ln Q^2$ and $A^{1=3}$. Fig.11a presents the ratio R₁ for two di erent values of Q^2 and for di erent nuclei. The suppression due to the SC increases with $\ln (1=x)$ and is much bigger than for the nucleon case. For A = 40 (Ca) and $Q^2 = 10 \text{ GeV}^2$, the suppression varies from 4% for $\ln (1=x) = 3 \text{ to } 25\%$ for $\ln (1=x) = 10$. For A = 197 (Au) the suppression is still bigger, going from 6% to 35% in the same kinematic region. Fig. 11b shows the same ratio for di erent values of Q^2 for the gold. The suppression decreases with Q^2 . Figs. 11c and d show the R₁ ratio as a function of $A^{1=3}$ and x for a xed value of Q^2 . A sexpected, the SC increases with A. An interesting feature of this gure is the fact that the curves tend to straight lines as x increases. It occurs because, as x grows, the structure function xG (G R V) become es maller, and the correction term of (23) proportional to dom inates. Since is proportional to A ¹⁼³, the curves behave as straight lines. The decrease of suppression with Q^2 is illustrated in m one detail in Figs. 11e and fwhich presents R₁ as a function of $\ln Q^2$ for di erent values of x for C a and Au, respectively. The e ect is pronounced for small Q^2 and x and dim inishes as $\ln Q^2$ increases.

This picture (Fig.11) shows also that the gluon structure function is far away from the asymptotic one. The asymptotic behavior R_1 ! 1 (see Figs.11e and f) occurs only at very high value of Q² as well as in the GLR approach (see ref. [25]). The asymptotic A-dependence $(R_1 / A^{\frac{1}{3}})$ has not been seen in the accessible kinematic range of Q² and x (see Figs.11c and d and Table 1). This result also has been predicted in the GLR approach [24]. We want also to mention that parameterization $R_1 = R_{1N} A$ does not t the result of calculations quite



Figure 11: R_1 as a function of $\ln (1=x)$, $\ln Q^2$ and $A^{1=3}$: a) R_1 as a function of $\ln (1=x)$ for di erent nucleus and di erent values of A; b) R_1 as a function of $\ln Q^2$ for di erent values of x_B for Au; c) and d) R_1 as a function of $A^{1=3}$ for di erent Q^2 ; e) and f) R_1 dependence on Q^2 for C a and Au.

well for 1Gev^2 Q^2 20GeV^2 and 10^2 x 10^5 . For x 10^2 the parameterization $R_1 = R_{1N}$ $R^0 A^{\frac{1}{3}}$ with parameters R_{1N} and R^0 for each value of Q^2 , works much better relation that only the rest correction to the Born term is essential in the Mueller formula.

We extend also the calculation of the exponents < ! > and < > of the sem iclassical approach for the nuclear case. We calculate the elective power of the nuclear gluon distribution <math>< ! > using the expression

$$< ! > = \frac{(! \ln (xG^{A} (x;Q^{2})))}{(! \ln (1=x))}$$
 : (30)

Fig.12 shows the results as functions of $\ln(1=x)$ for di erent values of Q² and di erent nucleus. The SC decreases the elective power of the nuclear distribution, giving rise to a lattening of the distribution in the small x region.



Figure 12: < ! > for di erent values of Q² and A.

It is also interesting to notice that at small values of Q², the elective power tends to be rather small, even in the nucleon case, at very small x. However it should be stressed that the elective power remains bigger than the intercept of the so called "soft" Pomeron [21], even in the case of a su ciently heavy nucleus (Au), for Q² > 1G eV². Now adays, many parameterizations [22] with matching of "soft" and "hard" Pomeron have appeared triggered by new HERA data on di raction dissociation [26]. These parameterization used Pomeron-like behavior namely, xG (x;Q²) / x^{! (Q²)}. However, if the Pomeron is a Regge pole, ! cannot depend on Q², and the only reasonable explanation is to describe ! (Q²) as the result of the SC. Looking at Fig.12 we can claim the SC from the MF cannot provide su ciently strong SC to reduce the value of ! to 0:08, a typical value for the soft Pomeron [21], at least for Q²

The calculation of the elective value of the anom alous dimension may help us to estimate what distances work in the SC corrections. This elective exponent is given by

< > =
$$\frac{(2 \ln (xG^{A} (x;Q^{2})))}{(2 \ln (Q^{2}=Q_{0}^{2}))}$$
: (31)



Figure 13: < > for di erent Q² and A.

Fig.13 shows the results as functions of ln (1=x) for di erent values of Q² and for two nuclei. We see that the values of at ln (1=x) 5, for both Ca and Au, is very close to the results for GRV and for nucleon case. At sm aller values of x, the anom alous dimension presents a sizeable reduction, which increases with A.For ln (1=x) > 15, < > tends to zero unlike in the DGLAP evolution equations (see Fig.10 for the GRV parameterization). A nalysing the Q² dependence, we see that < > is bigger than 1 only for Q² = 0.5 G eV². For Q² = 1.0 G eV², the anom alous dimension is close to 1=2, and for Q² > 5.0 G eV² it is always sm aller than 1=2.

U sing sem iclassical approach, we see that

$$/ \frac{1}{Q^2} (Q^2)$$
; (32)

and if 1, the integral over r_t in the master equation (23) becomes divergent, concentrating at small distances.

If 1 > 1=2, only the rst SC term, namely, the second term in expansion of the master equation, is concentrated at small distances, while higher order SC are still sensitive to small r_2 behavior. Fig.13 shows that this situation occurs for $Q^2 > 1 \text{ G eV}^2$, and even for $Q^2 = 1 \text{ G eV}^2$ at very small values of x. W e will return to discussion of these properties of the anom alous dimension behavior in the next section.

3.7 The gluon life time cuto .

In the D IS the incident electron penetrates the nucleus and radiates the virtual photon whose lifetime $/\frac{1}{m_x}$ [27]. We can recover three di erent kinem atic regions:

1. $=\frac{1}{m_X} < R_{NN}$, where R_{NN} is the characteristic distances between the nucleons of the nucleus. This virtual photon can be absorbed only by one nucleon and the total cross section is (A) = A (p).

 $2.R_A > = \frac{1}{m_x} > R_{NN}$, where R_A is the nucleus radius. In this kinem atic region the virtual photon can interact with the group of nucleons. However, (A) is still proportional to A since the number of nucleons in a group is much less than A.

3. $=\frac{1}{m_X} > R_A$. Here, before reaching the front surface of the nucleus, the virtual photon \decom poses" in the developed parton cascade which then interacts with the nucleus. It can be shown [28] that the absorption cross section of the virtual photon will now be proportional to the surface area of the nucleus (A) / $A^{\frac{2}{3}}$, because the wee partons of the parton cascade are absorbed at the surface and do not penetrate into the centre of the nucleus.

Everything that we have discussed have been calculated in the third kinematic region. For the RHIC energies we have to develop some technique how to penetrate into the second one. To do this we have to remember that the opacity (or _G) actually depends on the longitudinal part of the momentum transfer (q_z) which could be calculated in terms of x and x^{0} of our master equation (12), namely, $q_{z} = (x + x^{0})m$ (see Ref.[1]). Recalling that opacity / $r_{2}^{2} xG(x;q_{z};\frac{1}{r_{2}^{2}}) S(b_{t};q_{z})$ we see that q_{z} - dependence enters to two factors: to gluon structure function and to the nucleon pro le function. We know how to take into account the q_{z} - dependence of the gluon structure function (see Ref. [2] where the DGLAP equation for $q_{z} \in 0$ is written). However we neglected this e ect in our present estimates, hoping that this dependence occurs on the hadron scale and cannot change too much the dependence of the SC on the number of collisions during the passage through the nucleus.

The dependence of the pro le function S $(b_t;q_z)$ on q_z have been discussed and in the Gaussian parametrization it can be factor out in the form :

$$S(b_{1};q_{2}) = S(b_{1}) \quad L(q_{1} = S(b_{1}) \quad e^{\frac{R_{A}^{2}}{4}q_{2}^{2}}:$$
 (33)

This q_z -dependence takes into account the fact that the virtual gluon can interact with the target only during the nite time = 1=m x undergoing < R_A collisions. Using eq. (33), we can obtain:

$$x G_{A} (x;Q^{2}) = A x G_{N} (x;Q^{2})$$

$$A \frac{S N_{c}}{x} \frac{Z^{-1}}{x} \frac{Z}{x^{0}} \frac{dQ^{0}}{Q^{0}} L (q_{z}) x^{0} G_{N} (x^{0};Q^{0})$$

$$+ \frac{2R_{A}^{2}}{x} \frac{Z^{-1}}{x} \frac{dx^{0}}{x^{0}} \frac{dr_{t}^{2}}{\frac{1}{2}} \frac{dr_{t}^{2}}{r_{t}^{4}} fC + \ln (L (q_{z}) - G (x^{0};r_{t}^{2})) + E_{1} (L (q_{z}) - G (x^{0};r_{t}^{2})) g:$$

$$(34)$$

Fig. 14 shows the result of our calculations. C om paring Fig.11 with this picture, one can see that the nite lifetime of the virtual gluon a ects the behavior of the gluon structure function only at su ciently large x (x 10²) diminishing the value of the SC in this kinematic region. This e ect turns out to be very important for the RHIC energies and has to be studied in more details.

4 First corrections to the G lauber (M ueller) Approach.

In this section we discuss the corrections to the G lauber approach (the M ueller form ula of eq. (12). To understand how big could be the corrections to the G lauber approach we calculate



Figure 14: R_1 for CA and Au with gluon life time cuto .

the second iteration of the M ueller form ula of eq. (12). A shas been discussed, eq. (12) describes the rescattering of the fastest gluon (gluon -gluon pair) during the passage through a nucleus (see Fig.1). In the second iteration we take into account also the rescattering of the next to the fastest gluon. This is a well de ned task due to the strong ordering in the parton fractions of energy in the parton cascade in leading $\ln (1=x)$ approximation of pQCD that we are dealing with. Namely:

$$x_{B} < x_{n} < ::: < x_{1} < 1;$$
 (35)

where 1 corresponds to the fastest parton in the cascade.

Therefore, in the second interaction we include the rescatterings of the gluons with the energy fraction 1 and x_1 (see Fig.15). Doing the rest iteration we insert in eq. (12) G_N ($x;Q^2$) = G_N^{GRV} ($x;Q^2$). For the second iteration we calculate the gluon structure function using eq. (12) substituting

$$x G_{N} = \frac{x G_{A}^{1} (x; Q^{2})}{A} \qquad x G_{N}^{GRV} (x; Q^{2}); \qquad (36)$$

where xG_A^1 is the result of the rst iteration of eq. (12) that has been discussed in details in section 3.

Fig.16 shows the need to subtract xG_N^{GRV} in eq. (36) making the second iteration. Indeed, in the second iteration we take into account the rescattering of gluon 1^0 – gluon 2^0 pair o a nucleus. We picture in Fig.16 the 1st term of such iteration in which $G_{1^0}G_{2^0}$ pair has no rescatterings. It is obvious that it has been taken into account in our 1st iteration, so we have to subtract it to avoid a double counting.



Figure 15: The interaction with nucleons that is taken into account in the second iteration of M ueller form ula.



Figure 16: The rst term of the second iteration of eq. (36).



Figure 17: Second iteration calculations for R_1 , < ! >, and < > for Ca and Au.

O ne can see in Figs.17 that the second iteration gives a big e ect and changes crucially R₁, < >, and < ! >. The most remarkable feature is the crucial change of the value of the e ective power ! (Q²) for the \Pom eron" intercept which tends to zero at HERA kinematic region, making possible the matching with \soft" high energy phenom enology. It is also very instructive to see how the second iteration makesmore pronounced all properties of the behavior of the anom alous dimension (< >) that we have discussed. The main conclusions which we can make from Figs.17 are: (i) the second iteration gives a sizable contribution in the region $x < 10^2$ and for $x = 10^3$ it becomes of the order of the mail x we have to develop a dimension teration. It means that for such small x we have to develop a dimension which we simple iteration procedure for eq. (12). However, let us rst understand why the second iteration becomes essential to establish small parameters that enter to our problem.

As has been discussed, we use the GLAP evolution equations for gluon structure function in the region of small x. It means, that we sum the Feynman diagrams in pQCD using the following set of parameters:

s 1;
$$s \ln \frac{1}{x} < 1$$
; $s \ln \frac{Q^2}{Q_0^2} < 1$; $s \ln \frac{Q^2}{Q_0^2} \ln \frac{1}{x}$ 1: (37)

The idea of the theoretical approach of rescattering that has been form ulated in the GLR paper [2] is to introduce a new parameter z:

$$= \frac{N_{c} s A}{2Q^{2}R_{A}^{2}} xG (x;Q^{2})$$
(38)

and sum all Feynm an diagram susing the set of eq. (37) and as parameters of the problem, neglecting all contributions of the order of: $_{s}$, $_{s}$, $_{s}$ ln (1=x), $_{s}$ ln (1=x), $_{s}$ ln ($_{s}$ ln (

 $(_{\rm S} \ln (Q^2=Q_0^2) \ln (1=x))^n$ contributions in xG (x;Q²). However, it is not a complete solution. To illustrate this point let us compare the value of the second term of the expansion of eq. (12) with respect to (r_t^2) with the rst correction due to the second iteration in the rst term of such an expansion. In other words we wish to compare the values of the diagram s in Fig.18 b and Fig.18a. The contribution of the diagram of Fig.18a is equal:

$$xG(x;Q^{2})(Fig:18a) = \frac{R_{A}^{2}}{2^{2}} \frac{Z}{x^{0}} \frac{dx^{0}}{x^{0}} dQ^{2} \frac{Q^{2}}{x^{0}} \frac{Q^{2}}{4}; \qquad (39)$$

where x^0 and Q^{02} are the fraction of energy and the virtuality of gluon 1 in Fig.18a

 $^{^{\}rm z}$ In the GLR paper the notation for $~w\,as\,W$, but in this paper we use to avoid a m isunderstanding since, in D IS, W is the energy of interaction.

The diagram of Fig.18b contains one more gluon and its contribution is:

$$xG(x;Q^{2})(Fig:18b) = \frac{R_{A}^{2}}{^{2}2} \frac{N_{c}}{^{2}2} \frac{X}{x^{0}} \frac{dx^{0}}{Q^{2}} \frac{dQ^{2}}{^{2}2} \frac{dx^{"}}{x^{"}} dQ^{"^{2}} \frac{dx^{"};Q^{"^{2}}}{^{2}}$$

$$/ \frac{sN_{c}}{^{2}2} \ln(1=x) \ln(Q^{2}=Q_{0}^{2}) xG(x;Q^{2})(Fig:18a));$$
(40)

where x^0 (x^{00}) and Q^{02} (Q^{02}) are the fraction of energy and the virtuality of gluon 1 (1⁰) respectively in Fig.18b. Therefore, eq. (40) gives the contribution which is of the order of eq. (39) in the kinematic region where the set of parameters of eq. (37) holds. It means also that we need to sum all diagrams of Fig.18b type to obtain the full answer. In the diagram of Fig.18b not only one but many gluons can be emitted. Such emission leads to so called \triple ladder" interaction, pictured in Fig.18c (see ref.[2]). This diagram is the rst from so called \fan" diagrams of Fig. 18d. To sum them all we can neglect the third term in eq. (12) and treat the remained terms as an equation for xG (x;Q²). It is easy to recognize that we obtain the GLR equation [2][10]. Generally speaking the GLR equation sum s the most in portant diagrams in the kinematic region where s hn(1=x) ln(Q²=Q₀²) 1 and < 1.

5 The general approach.

5.1 W hy equation?

We would like to suggest a new approach based on the new evolution equation to sum all SC. However, rst of all we want to argue why an equation is better than any iteration procedure. To illustrate this point of view let us di erentiate the M ueller form ula with respect to $y = \ln (1=x)$ and $= \ln Q^2$. It is easy to see that this derivative is equal to

$$\frac{\partial^2 x G (x; Q^2)}{\partial y \partial} = \frac{4}{2} db_t^2 f 1 \qquad e^{\frac{1}{2} (x; r_2^2 = \frac{1}{Q^2}) S (b_t^2)} g :$$
(41)

The nice property of eq. (41) is that everything enters at small distances, therefore everything is under theoretical control. Of course, we cannot get rid of our problem s changing the procedure of solution. Indeed, the nonperturbative e ects coming from the large distances are still important but they are all hidden in the boundary and initial conditions to the equation. Therefore, an equation is a good (correct) way to separate what we know (small distance contribution) from what we don't (large distance contribution).

5.2 The generalized evolution equation.

W e suggest the following way to take into account the interaction of all partons in a parton cascade with the target. Let us di erentiate the M ueller form ula over $y = \ln (1=x)$ and $= \ln (Q^2=Q_0^2)$. It gives:

$$\frac{\theta^2 x G_A (y;)}{\theta y \theta} = \frac{2 R_A^2 Q^2}{2} f C + \ln + E_1 () g:$$
(42)



Figure 18: Corrections to the G lauber approach.

Rewriting eq. (42) in terms of given by

$$= \frac{N_{c} s}{2Q^{2}R_{A}^{2}} xG_{A} (x;Q^{2})$$
(43)

we obtain:

$$\frac{\partial^2 (y;)}{\partial y \partial t} + \frac{\partial (y;)}{\partial y} = \frac{N_c s}{f C} + \ln (y;) + E_1((y;)) g F(): \quad (44)$$

Now, let us consider the expression of eq. (44) as the equation for This equation has the following nice properties:

1. It sums all contributions of the order $(_{S}y_{})^{n}$ absorbing them in $xG_{A}(y;)$, as well as all contributions of the order of n . Therefore, this equation solves the old problem, form ulated in Ref.[2] and for N_c ! 1 eq. (44) gives the complete solution to our problem, summing all SC;

2. The solution of this equation m atches with the solution of the DGLAP evolution equation in the DLA of perturbative QCD at ! 0;

3. At sm all values of (< 1) eq. (44) gives the GLR equation. Indeed, for sm all we can expand the rhs of eq. (44) keeping only the second term. Rewriting the equation through the gluon structure function we have

$$\frac{\theta^2 \mathbf{x} \mathbf{G}_{\mathbf{A}} (\mathbf{y}; \mathbf{y})}{\theta \mathbf{y} \theta} = \frac{\mathbf{s} \mathbf{N}_{\mathbf{c}}}{\mathbf{x}} \mathbf{G} (\mathbf{x}; \mathbf{Q}^2) \qquad \frac{\mathbf{s} \mathbf{A}^2}{2\mathbf{R}_{\mathbf{A}}^2} (\mathbf{x} \mathbf{G} (\mathbf{x}; \mathbf{Q}^2))^2 ; \qquad (45)$$

which is the GLR equation [2] with the coe cient in front of the second term calculated by Mueller and Q in [10].

4. For $_{\rm S}y$ 1 this equation gives the G lauber (M ueller) form u.l., that we have discussed in details.

5. This equation almost coincide with the equation that L M cleman with collaborators [29] derived from quite di erent approach and with di erent technique. W e are sure that almost will disappear when they will do m ore careful averaging over transverse distances.

Therefore, the great advantage of this equation in comparison with the GLR one is the fact that it describes the region of large and provides the correct matching both with the GLR equation and with the G lauber (Mueller) formula.

Eq. (44) is the second order di erential equation in partial derivatives and we need two initial (boundary) conditions to specify the solution. The rst one is obvious, namely, at xed y and Q^2 ! 1

$$! \quad \frac{N_{c} S A}{2Q^{2}R_{A}^{2}} xG_{N}^{GLAP} (x;Q^{2}) :$$

The second one we can x in the following way: at $x = x_0$ ($y = y_0$) which is small, namely, in the kinematic region where $_{sy}$ 1

$$! _{in} = \frac{N_{c} s}{2Q^{2}R_{A}^{2}} xG_{A} (x;Q^{2}) ; \qquad (46)$$

where xG_A is given by the M ueller form ula (see eq. (12)). Practically, we can take $x_0 = 10^2$, because corrections to the M F are sm all at this value of $x = x_0$.

5.3 The asymptotic solution.

F inst observation is the fact that eq. (44) has a solution which depends only on y. Indeed, one can check that $= a_{\text{asymp}}(y)$ is the solution of the following equation:

$$\frac{d_{asymp}}{dy} = F (_{asymp}) :$$
 (47)

The solution to the above equation is:

$$\sum_{asymp(y)=y_0}^{Z} \frac{d^{0}}{F(0)} = y \quad y :$$
(48)

It is easy to nd the behavior of the solution to eq. (48) at large value of y since F () ! $_{\rm S}$ ln at large ($_{\rm S} = \frac{\rm N_c}{\rm s}$). It gives

$$_{asymp}$$
 ! $_{sy}\ln(_{sy})$ at $_{sy}$ 1: (49)

At small value of y, F () ! s and we have:

asymp !
$$asymp (y = y_0) e^{s(y_0)}$$
: (50)

The solution is given in Fig.19 for $_{\rm S} = 1=4$ in the whole region of y for di erent nuclei in comparison with our calculations based on the MF.W e chose the value of $_{\rm asym\,p}$ (y = y₀) from eq. (46). We claim this solution is the asym ptotic solution to eq. (44) and will argue on this point a bit later.

For nuclei the SC incorporated in the asymptotic solution turn out to be much stronger than the SC in the G lauber approach for any $Q^2 > 1 \text{ GeV}^2$ at $x > 10^2$. In this kinematic region the solution of eq. (44) is drastically di erent from the G lauber one.

A general conclusion for Fig.19 is very simple: the amount of shadowing which was taken into account in the MF is not enough , at least for the gluon structure function in nuclei at $x < 10^{2}$ and we have to solve eq. (44) to obtain the correct behavior of the gluon structure function for nuclei.



Figure 19: The G lauber approach and asymptotic solution for di erent nuclei.

Now, we would like to show that solution eq. (44) is the asymptotic solution of the new evolution equation. In order to check this we need to prove that this solution is stable. It means that if add a small function and searching for the solution to the equation in the form $= a_{\text{sym p}} +$, we have to prove that turns out to be smallnamely, . The following linear equation can be written for :

$$\frac{\partial^2 (y;)}{\partial y \partial} + \frac{\partial (y;)}{\partial y} = \frac{dF()}{d} j_{=asymp}(y) \quad (y;):$$
(51)

In Ref.[1] was proven, that the solution of eq. (51) is much smaller than .

Therefore the asymptotic solution has a chance to be the solution of our equation in the region of very small x. To prove that the asymptotic solution is the solution to the equation we need to solve our equation in the wide kinematic region starting with our initial condition. We managed to do this only in sem iclassical approach.

5.4 Sem iclassical Approach.

The sem iclassical approach has been adjusted to the solution of the nonlinear equation of eq.(44)-type in Refs. [2, 30, 31] (for simplicity, we assume that s is xed).

In the sem iclassical approach we are boking for the solution of eq. (44) in the form

where S is a function with partial derivatives: $\frac{@S}{@y} = !$ and $\frac{@S}{@} = which are smooth function of y and . It means that$

$$\frac{\mathrm{e}^2 \mathrm{S}}{\mathrm{e} \mathrm{e}_{\mathrm{Y}}} = \frac{\mathrm{e}_{\mathrm{S}}}{\mathrm{e}_{\mathrm{Y}}} \frac{\mathrm{e}_{\mathrm{S}}}{\mathrm{e}_{\mathrm{Y}}} = !$$
(53)

U sing eq.(53), one can easily rewrite eq.(44) in the form

$$\frac{\mathrm{dS}}{\mathrm{dy}}\frac{\mathrm{dS}}{\mathrm{d}} + \frac{\mathrm{dS}}{\mathrm{dy}} = \mathrm{e}^{\mathrm{S}}\mathrm{F}(\mathrm{e}^{\mathrm{S}}) \qquad (\mathrm{S})$$
(54)

or

$$! (+1) = (S)$$
 (55)

W e are going to use the m ethod of characteristics (see, for example, ref.[33]). For equation in the form

$$F(;y;S;;!) = 0$$
 (56)

we can introduce the set of characteristic lines ((y); S(y); (y); (y)), which satisfy a set of well de ned equations (see, for example, Refs. [30] [31] for the method and Ref [1] for detailed calculation). Using eq.(54) and eq.(55), we obtain the following set of equations for the characteristics:

$$\frac{d}{dy} = \frac{(S)}{(+1)^2}; \quad \frac{dS}{dy} = \frac{2+1}{(+1)^2} \quad (S); \quad \frac{d}{dy} = -\frac{0}{S} - \frac{1}{1}; \quad (57)$$

where ${}_{S}^{0} = \frac{\theta}{\theta S}$. The initial condition for this set of equations we derive from eq.(46), namely

$$S_{0} = \ln_{in} (Y_{0}; 0)$$

$$= \frac{\theta \ln_{in} (Y_{0}; 0)}{\theta} = 0$$
(58)

The main properties of these equations have been considered in R ef.[1] analytically, however, here, we restrict ourselves mostly the num eric solution of these equations.

We set the initial condition $y = y_0 = 4.6$ ($x_B = 10^2$), where the shadowing correction is not big and the evolution starts from < 0. In this case d =dy > 0 and the value of increases . At the same time dS=dy < 0 and S decreases if $_0 < 1=2$. With the decrease of S, the value of $_{\rm S}^0$ becomes smaller and after short evolution the trajectories of the nonlinear equation start to approach the trajectories of the DGLAP equations. We face this situation for any trajectory with $_0$ close to -1. If the value of $_0$ is smaller than $\frac{1}{2}$ but the value of S₀ is su ciently big, the decrease of S due to evolution cannot provide a small value for 0 (S) and increases until its value becomes bigger than $\frac{1}{2}$ at some value of $y = y_c$. In this case for $y > y_c$ the trajectories behave as in the case with $_0 > \frac{1}{2}$. For $_0 > 1=2$, the picture changes crucially. In this case, dS=dy > 0, d =dy > 0 and both increase. Such trajectories go apart from the trajectories of the DGLAP equation and nonlinear e ects play more and more in portant role with increasing y. These trajectories approach the asymptotic solution very quickly. For the num erical solution we use the 4th order Runge -K utta m ethod to solve our set of equations with the initial distributions of eq. (58). The result of the solution is given in Figs 20 and 21. In these gures we plot the bunch of the trajectories with di erent initial conditions. For the nucleon (Fig 20) we show also the dependence of along these trajectories. One can notice that the trajectories behave in the way which we have discussed in our qualitative analysis. It is interesting to notice that the trajectories, which are di erent from the trajectories of the G LAP evolution equations, start at $y = y_0 = 4.6$ with the values of Q² between 0.5G eV² and 2.5G eV², we can hope that the linear evolution equations (the DG LAP equations) will describe the evolution of the deep inelastic structure function in the limited but su ciently wide range of Q².

In Figs. 20 and 21 we plot also the lines with de nite value of the ratio $R = \frac{xG (xQ^2) (generalized equation)}{xG (xQ^2) (G LAP)}$ (horizontal lines). These lines give the way to estimate how big are the SC. One can see that they are rather big.



Figure 20: The trajectories and contour plot for the solution of the generalized evolution equation for N.R = $\frac{xG(x,Q^2)(\text{generalized equation})}{xG(x,Q^2)(\text{GLAP})}$.

We have discussed only the solution with xed coupling constant which we put equal to $_{\rm S}$ = 0.25 in the num erical calculation. The problem how to solve the equation with running coupling constant is still open.

5.5 The generalized evolution equation versus the GLR equation.

In Ref.[l] we studied in detail the solution to the GLR equation in the same sem iclassical approximation. Our conclusion is that the GLR equation gives much stronger SC than the generalized evolution equation. This di erence we can see comparing the solution to the both equation in the region ultra small x.



Figure 21: The trajectories and contour plot for the solution of the generalized evolution equation for Ca and Au. R = $\frac{xG(x,Q^2)(generalized equation)}{xG(x,Q^2)(GLAP)}$.

Indeed, our asymptotic solution turns out to be quite di erent from the GLR one. The GLR solution in the region of very small x leads to saturation of the gluon density [30, 31, 32]. Saturation means that tends to a constant in the region of small x. The solutions of eq. (44) approach the asymptotic solution at x ! 0, which does not depend on Q^2 , but exhibits su ciently strong dependence of on x (see Fig.19), namely / $_{\rm S} \ln (1=x) \ln \ln (1=x)$. The absence of saturation does not contradict any physics since gluons are bosons and it is possible to have a lot of bosons in the sam e cell of the phase space. We should adm it that A . M ueller rst cam e to the sam e conclusion using his form ula in Ref.[3].

6 Next steps.

Here, we list our problem s that have to be solved to com plete our study of the SC :

1. Calculation of $F_2^A(x;Q^2)$ to compare our calculation of the SC with the available experim ental data.

2. Recalculation of the SC using more reliable W ood-Saxon parameterization for prole function S (b₁) instead of the G aussian one. The form of the prole function especially essential to obtain a reliable estimates for the SC in the region of the moderate $x = \frac{1}{2mR_{0}}$.

3. Solution of the generalized evolution equation for running $_{\rm S}$. The experience of solving the GLR equation tells us that there is a principal di erence in the solutions for xed and running $_{\rm S}$, namely, the critical line of the GRL equation appears only for running $_{\rm S}$ [2].

We think that it is very important to study the generalized equation with running $_{\rm S}$ and to compare this solution with the solution of the GRL equation.

4. We have discussed that for RHIC energies it is very important to study in more details the e ect of the nallifetime of the gluon in a nucleus. We plan to recalculate the SC replacing xG (x;Q²) in our formulae by xG (x;Q²;q_z) for which the kernels of the evolution equations have been calculated in Ref.[2].

5. In all our calculations we neglected the parton interaction inside GG + N scattering. Our estimates, which have been presented in section 2, shows that this interaction should be very important. Indeed, for example, in the M ueller formula we have to change the parameter $_{G}$ due to the parton interaction inside the nucleon. This change is simple, the only that we need to do is to replace the number of collisions $A = R_{A}^{2}$ by

$$\frac{A}{R_A^2} ! \frac{A}{R_A^2} + \frac{1}{R_N^2}$$

in the de nition of $_{\rm G}$ in eq. (12). It means that all results will be the same but nucleus with the new elective number of nucleons:

$$A_{eff}^{\frac{1}{3}} = A^{\frac{1}{3}} + \frac{R_0^2}{R_N^2};$$

where $R_A = R_0 A^{\frac{1}{3}}$. Using our estimates for $R_N^2 = 5 \text{GeV}^2$ we can see that elective A for the gold is $A_{eff}^{\frac{1}{3}} = 9.6$ instead $A^{\frac{1}{3}} = 6$. For light nuclei the change is even more essential. Therefore, we are planning to take into account the parton interaction inside a nucleon as soon as possible.

6. We have neglected all correlations between partons of the order $\frac{1}{N_c^2}$ which could be sizable in the case of the nucleus D IS.We suppose to study this problem using the technique that has been developed in Ref.[34].

7. Everywhere through the paper we used the DLA of perturbative QCD. However, the key assumption that simplify our theoretical approach was the $_{\rm S}\ln(1=x)$ 1 approximation. We plan to develop our approach in the case of the BFKL dynamic and, therefore, to get rid of our assumption that $_{\rm S}\ln(Q^2=Q_0^2)$ 1. We consider this generalization as an important step, since our result that we have no saturation of the gluon density in nuclei even at ultra small x could be an artifact our double log approximation of perturbative QCD.

7 And what?

W e presented here our approach to the SC and a natural question arises and what? W hat and how we can do for the RHIC physics. How our approach can help in creating of the reliable



Figure 22: Four stages of nucleus - nucleus collision.

M onte Carlo code for nucleus - nucleus interaction at high energies W e are going to answer these hot question in this section.

Let us consider rst the space time structure of the nucleus – nucleus interaction (see Fig.22). One can see four stages of this process:

1. For time smaller that t_0 , where t_0 is the time of the rst parton - parton interaction, we have a very coherent system of parton, con ned in our both nuclei. We know almost nothing about this system.

2. At time t_0 the st parton - parton interaction occurs and we believe that this interaction destroys the coherence of our parton system at the very instant.

3. During time from t_0 till t_h , where t_h is the hadronization time, we have a quark -gluon stage of the process. We believe that we can reach a simple and econom is understanding this stage in framework of QCD. We also believe that new collective phenomena could be created in the nucleus -nucleus interaction during this stage of the process such as the Quark -G luon P lasm a mostly because of the high density of the produced gluons. For this stage we have the M onte C arb codes based on QCD, the lattice calculation and a lot of beautiful ideas that has been discuss at this conference.

4. The last stage - hadronization is a black box. Nothing is known, but the success of the Local-Hadron-Parton Duality in the description of the LEP data allows us to hope that this stage could be not very important for our understanding of the nucleus - nucleus collisions.

Our approach can de ne the initial condition at $t = t_0$ for the third stage. W hat can we do?

1. We are able to calculate the inclusive cross section for gluons at $t = t_0$ or, in other words, de ne the gluon distribution at $t = t_0$. A ctually, it has been done by C E scola [35] and his collaborator and has been presented at this conference. We can only improve his treatment of the SC which was based on the GLR equation. However, let us discuss brie y the form ula for the inclusive gluon cross section. It can be written using the factorization theorem [36] in the form :

$$\frac{d}{dy \, dp_t^2} = / \int dx_1 dx_2 \, x_1 G_{A_1} (x_1; p_t^2) \, x_2 G_{A_2} (x_2; p_t^2) \, \frac{s}{p_t^4}$$

where the last factor is the hard gluon – gluon cross section and y and p_t are rapidity and transverse momentum of produced gluon, respectively. One can see that this cross section is infrared unstable and diverges at small values of p_t . The SC provides a natural scale that cut o this divergence. A rough estimate for this new scale can be done from equation

$$_{\rm G}$$
 (x; $r_{\rm 2}^2 = \frac{1}{Q_0^2(x)}$) = 1

(see Fig.4). For $p_t < Q_1(x)$ the gluon structure function xG (x; p_t^2) / p_t^2 and one can see that the number of gluon with transverse momenta smaller than $p_t = Q_0(x)$ turns out to be very small.

2. We can calculate also the double inclusive cross section which gives the two gluon correlation function at $t = t_0$. We would like to stress that for nucleus – nucleus collision this correlation function is big and have to be taken into account. Indeed, we have two di erent contribution to the double inclusive process, pictured in Fig.23: the production of two gluons from one parton cascade (see Fig.23a) and from two parton cascades (see Fig.23b). However, for nucleus – nucleus collisions the rst contribution is proportional to A_1A_2 (without the SC) while the second ism uch bigger and it is of the order of $A_1^2A_2^2 \frac{R_{A_1}^2 + R_{A_2}^2}{R_{A_1}^2 R_{A_2}^2}$ (without the SC and for the G aussian prole function). Using our approach we can calculate the two gluon correlation function within better acuraccy than the above simple estimates. We hope, that these two observables: gluon distribution for the QCD motivated cascade during the third stage of our process.

3. We think that these two observables: multiplicity of gluons and two gluon correlation will be enough to de ne the initial condition for current M onte C arb codes. However, we think that these codes are doing something wrong. Indeed, we learned from A. Mueller [17] that correct degrees of freedom for parton cascading boks in the simplest way and which could be used for a probabilistic interpretation and therefore, they are natural degrees of freedom for M onte C arb simulations are not quark and gluons but colourless quark – antiquark dipoles. The gluon structure function is the probability to nd a colourless dipole with the size $r_2 = \frac{1}{Q}$. Therefore, we think that the code should be written for such dipoles and their interaction. We shall answer the questions:(i) how to calculated the average multiplicity of dipoles with the size r_2 and (ii) how to calculate the correlations between such dipoles. We are going to do this in the nearest future.

4. Now we want to discuss a hot question how to mix the "soft" and "hard" Pomerons. The common way of doing such a mixture is to use the G lauber formula and replace in this formula $(r_2^2 ! _{soft} + _{hard}. W e think this is a correct procedure to obtain an estimate how important soft or/and hard processes. In section 3 we argued that this is the most econom ic way of doing which satis es the s-channel unitarity. However, all M onte C arb program s that$



Figure 23: Double inclusive cross section in QCD .



Figure 24: Total cross section in $^2_{\rm S}$ and $^3_{\rm S}$ orders of perturbative QCD .

we know use for the calculation of hard the factorization formula, namely

$$_{hard} = \frac{!}{2} \sum_{p_0^2}^{Z} dp_t^2 x_1 G (x_2; p_t^2) x_2 G (x_2; p_t^2) \frac{2}{p_t^4};$$

which describes really the inclusive production of gluons. The factor 1/2 in front does not help because to nd hard we need to calculate the real multiplicity but not the number of gluon line in the Feynman diagram. In Fig.24 we picture the $\frac{3}{5}$ corrections to the hard cross section considering the scattering of two mesons made from heavy quarks. Perturbative QCD is certainly a good tool to study such processes. From this picture one sees that including the inclusive cross section in the place of the total we missed the radiative correction to the the partial cross section with four quarks ithe nal state.

Our way of doing is the following. We will write the Mueller formula or our more so sticated approach for dipole (with size r_2 scattering with a nucleus. To nd the proton – nucleus cross

section we need to calculate the integral:

$$(pA) = \int_{0}^{Z_{1}Y^{3}} dz_{i} \frac{Z_{Y}}{2} dz_{i}^{2} \frac{d^{2}r_{?}^{ik}}{2} p(z_{i};r_{?}^{ik} 2f1) e^{\frac{1}{2}n(r_{?}^{ik} S(b_{r}))} g(r_{?}^{ik};z_{i}):$$

To nd the wave function of the nucleon we have to use a model for example the constituent quark model or instanton liquid model. The nice feature of this form ula that the typical $r_{?}^{ik}$ will be of the order 1 G eV 1 due to the SC. It means that we need to know the wave function at su ciently sm all distances where we have some control from lattice calculations and QCD sum rules. This form ula takes into account correctly hard process and give the factorization form ula for the inclusive production. We suppose to do an estimates using the model for the nucleon wave function. If they will show that we need some admixture of the soft processes we will add to N in the above form ula in an usual phenom enologic way, using the model of, so called, soft Pom eron.

8 Conclusions.

W e have two conclusions:

1. We hope that we convinced you that we are on the way from our Really Highly Ine cient Calculation to your RHIC. Much work is need to clarify the initial condition for the QCD phase of nucleus -nucleus interaction and this is the rst and the most important task which we need to attack, since it will determ ine the correct degrees of freedom for further evolution of QCD cascades.

2. Everything that we have talked about satis es the third law of theoretical physics: Any model is a theory which we apply to a kinematic region, where we cannot prove that this theory is wrong. We may believe that correct SC will provide the picture of the nucleus – nucleus interaction in which hard and sem ihard processes will play a crucial role with only small if any contam ination of the soft contribution.

A cknow ledgem ents:

O ne of us (E M L.) is very grateful to Sid K ahana for creation a stimulating atmosphere of discussion at RHIC '96 W orkshop and for enlighting discussions of the di cult theoretical problems in the modeling heavy ion collisions. E M L. thanks all participant of the hard working group and especially Yu. D okshitzer, C E skola, A. M ueller and M Strikm an for fruitful discussions on the subject of the talk and related topics. M BGD thanks A. C apella and D Schi for enlightening discussions. W ork partially nanced by CNPq, CAPES and FINEP, Brazil.

References

- [] A L A yala, M B G ay D ucati and E M . Levin: CBPF FN -020/96, hep-ph 9604383.
- [2] L.V.Gribov, E.M. Levin and M.G. Ryskin: Phys. Rep. 100 (1983) 1.
- [3] A.H.Mueller: Nucl. Phys. B 335 (1990) 115;
- [4] H1 collaboration, T. Ahm ed et al.: Nucl. Phys. B 439 (1995) 471;
 ZEUS collaboration, M. Derrick et al.: Z. Phys. C 69 (1995) 607; H1 collaboratrion, S. Aid et al.: DESY 96-039 (1996).
- [5] V N.G ribov and L N.Lipatov: Sov.J.Nucl.Phys.15 (1972) 438; L N.Lipatov: Yad.Fiz.
 20 (1974) 181; G.A ltarelliand G.ParisiNucl.Phys.B 126 (1977) 298; Yu L Dokshitzer:
 Sov.Phys.JETP 46 (1977) 641.
- [6] M.Gluck, E.Reya and A.Vogt: Z.PhysC 67 (1995) 433.
- [7] A D. Martin, R G. Roberts and W J. Stirling: Phys. Lett. B 354 (1995) 155.
- [8] H L. Laiet al, CTEQ collaboration: Phys. Rev. D 51 (1995) 4763.
- [9] W .Buchmuller and D.Haidt: DESY 96-061, M arch 1996.
- [10] A.H.Mueller and J.Qiu: Nucl. Phys. B 268 (1986) 427.
- [11] H1 collaboration, S.A id et al.: DESY 96-037, M arch 1996.
- [12] E A. Kuraev, LN. Lipatov and V S. Fadin: Sov. Phys. JETP 45 (1977) 199; Ya.Ya. Balitskii and L.V. Lipatov Sov. J. Nucl. Phys. 28 (1978) 822; LN. Lipatov: Sov. Phys. JETP 63 (1986) 904.
- [13] E.Laenen and E.Levin: Ann.Rev.Nucl.Part.44 (1994) 199.
- [14] E M . Levin and M G Ryskin: Sov. J. Nucl. Phys. 45 (1987) 150.
- [15] V A . A bram ovski, V N . G ribov and O . V . K ancheli: Sov. J. Nucl. Phys. 18 (1973) 308.
- [16] M.Abram ow itz and IA.Stegun: \Handbook of Mathematical Functions", Dover Publication, INC, NY 1970.
- [17] A H.Mueller: Nucl. Phys. B 415 (1994) 373.
- [18] H A . Enge: Introduction to nuclear physics", Addisson-W esley Publishing Company, M assachusetts, 1971.
- [19] R K. Ellis, Z. Kunst and E. M. Levin: Nucl. Phys. B 420 (19) 94)517.

- [20] A D.Martin, W.J.Stirling and R.G.Roberts: Phys. Lett. B 354 (1995) 155.
- [21] A. Donnachie and P.V. Landsho: Phys. Lett. B 185 (1987) 403, Nucl. Phys. B 311 (1989) 509; E. Gotsman, E. Levin and U. Maor: Phys. Lett. B 353 (1995) 526.
- [22] H.Abram owicz et al.: Phys. Lett. B 269 (1991) 465 and references therein; A.Capella et al.: Phys. Lett. B 343 (1995) 403, Phys. Lett. B 345 (1995) 403; K.Golec-Biemat and J. Kwiecinski: Phys. Lett. B 353 (1995) 329.
- [23] M.Abram ow itz and IA.Stegun: \Handbook of Mathematical Functions", Dover Publication, INC, NY 1970.
- [24] E.M. Levin and M.G. Ryskin: Sov. J. Nucl. Phys. 41 (1985) 300.
- [25] J.Qiu: Nucl. Phys. B 291 (1987) 746.
- [26] ZEUS collaboration, M. Derrick et al.: Phys. Lett. B 315 (1993) 481, Phys. Lett. B 332 (1994) 228, Phys. Lett. B 338 (1994) 483;
 H1 collaboration, T. Ahm ed et al.: Nucl. Phys. B 429 (1994) 477, DESY 95-036.
- [27] V N.Gribov, B L. Io e and I.Ya. Pomeranchuk: Yad. Fiz. 2 (1965) 768; B L. Io e: Phys. Rev. Lett. 30 (1970) 123.
- [28] V N.G mbov: Sov. Phys. JETP 30 (1970) 600; ZhETF 57 (1969) 1306.
- [29] Jam al Jullian-M arian, A lex K ovner, Larry M cLerran and H erbert W eigert: HEP-M INN-96-1429.
- [30] J.C. Collins and J.Kwiecinski: Nucl. Phys. B 335 (1990) 89.
- [31] J. Bartels, J. Blum lein and G. Shuler: Z. Phys. C 50 (1991) 91.
- [32] J.Bartels and E.Levin: Nucl. Phys. B 387 (1992) 617.
- [33] IN . Sneddon: E lem ents of partial di erential equations", M c.C raw -H ill, N Y 1957.
- [34] Laenen and E. Levin: Nucl. Phys. B 451 (1995) 207.
- [35] C.Eskola: Talk at RHIC '96 W orkshop.
- [36] J. Collins, D E. Soper and G. Sterm an: Nucl. Phys. B 308 (1988) 833; In \ Perturbative Quantum Chromodynamics", ed. A. Mueller, Singapure, W S 1989 and reference therein latex rhic.txt