

hep-ph/9608310
DFF 254-7-96
July 1996

k_{\perp} -factorization and perturbative invariants at small x

Stefano Catani *

I.N.F.N., Sezione di Firenze
and Dipartimento di Fisica, Università di Firenze
Largo E. Fermi 2, I-50125 Florence, Italy

Abstract

I review the basic idea of k_{\perp} -factorization and its relation to collinear factorization. Theoretical results in resummed perturbation theory are summarized and the example of the heavy-flavour structure functions is explicitly considered. Using these results one can investigate the small- x behaviour of quantities that are independent of the non-perturbative parton densities. In particular, one can introduce physical anomalous dimensions that relate the scaling violations in different hadronic observables.

DFF 254-7-96
July 1996

*Talk given at International Workshop on Deep Inelastic Scattering and Related Phenomena, *DIS 96*, Rome, Italy, 15-19 April 1996. A shorter version will appear in the Proceedings.

1 k_\perp -factorization in hadron collisions at high energy

Hadron collisions at large transferred momentum Q ($Q \gg \Lambda$, Λ being the QCD scale) can be studied in QCD perturbation theory by computing the corresponding cross sections as power-series expansions in $\alpha_S(Q^2)$. In the high-energy regime $\sqrt{S} \gg Q$ (\sqrt{S} the centre-of-mass energy) or, equivalently, at small values of the ratio $x = Q^2/S$, the coefficients of these power-series expansions contain logarithmically-enhanced contributions of the type $\ln^n x$. As soon as $\alpha_S \ln 1/x \sim 1$, the fixed-order expansion in α_S is no longer reliable. The higher-order contributions $(\alpha_S \ln x)^n$ have to be evaluated and, possibly, resummed to all orders in perturbation theory.

The basic theoretical input for the resummation is provided by the BFKL equation [1] for the gluon distribution $\mathcal{F}(x, k_\perp; Q_0)$. This distribution describes the evolution of an initial-state gluon with momentum p^μ ($p^2 = Q_0^2 \sim 1\text{GeV}^2$) into an off-shell gluon with momentum $k^\mu = xp^\mu + k_\perp^\mu$ ($-k^2 = -k_\perp^2 \gg Q_0^2$). The evolution process is obtained by radiating final-state partons with momenta $k_{\perp i}$ ($k_\perp = -\sum_i k_{\perp i}$). The BFKL equation thus resums $(\alpha_S \ln x)^n$ terms due to gluon evolution over the large rapidity gap $y = \ln 1/x$. Since these terms are produced by emission of partons with any transverse momentum $k_{\perp i}$, no k_\perp -ordering is embodied in the BFKL equation.

Gluons are not directly observable in scattering processes. Having at our disposal the BFKL equation, we still have to relate it to physical cross sections. The relation is provided by the k_\perp -factorization theorem [2, 3, 4]. In the case of processes involving a single incoming hadron, like, for instance, in deep-inelastic lepton-hadron scattering (DIS), the cross section is written as follows

$$\sigma(x, Q^2) = \int_x^1 \frac{dz}{z} \int d^2 k_\perp \hat{\sigma}(x/z, Q^2; k_\perp) \mathcal{F}(z, k_\perp) , \quad (1)$$

$$\mathcal{F}(x, k_\perp) = \int_x^1 \frac{dz}{z} \mathcal{F}(x/z, k_\perp; Q_0) \bar{f}_g(z, Q_0) . \quad (2)$$

In Eq. (2) the lower-end gluon ($p^2 = Q_0^2$) of the BFKL distribution $\mathcal{F}(x, k_\perp; Q_0)$ is first coupled to the incoming hadron via a non-perturbative (but process-independent) gluon density $\bar{f}_g(x, Q_0)$. Then, in Eq. (1) the upper-end gluon k is coupled to the scattering process via the partonic cross section $\hat{\sigma}$. Note that, since the $(\alpha_S \ln x)^n$ contributions to $\sigma(x, Q^2)$ are due to the emission of partons with any $k_{\perp i}$, in order not to spoil the consistency of small- x resummation, the partonic cross section $\hat{\sigma}(z, Q^2; k_\perp)$ has to be properly defined (and perturbatively computed) in terms of an incoming off-shell ($k^2 = k_\perp^2$) gluon. Accordingly, an unconstrained k_\perp -integration is involved in the factorization formula (1).

2 Collinear factorization and small- x resummation

Equations (1,2) have to be compared with the customary perturbative QCD formulae as obtained by applying the factorization theorem of collinear singularities [5]. According to the latter the hadronic cross section is written in the following way

$$\sigma(x, Q^2) = \sigma_0(Q^2) \sum_a \int_x^1 \frac{dz}{z} C^a(x/z, \alpha_S(Q^2)) \tilde{f}_a(z, Q^2) , \quad (3)$$

where the rescaling factor $\sigma_0(Q^2)$ has been introduced to make dimensionless the coefficient functions $C^a(x, \alpha_S(Q^2))$. The parton densities $\tilde{f}_a(x, Q^2)$ fulfil the Altarelli-Parisi equations [6]:

$$\frac{d\tilde{f}_a(x, Q^2)}{d\ln Q^2} = \sum_b \int_x^1 dz P_{ab}(z, \alpha_S(Q^2)) \tilde{f}_b(x/z, Q^2) . \quad (4)$$

Both the coefficient functions $C^a(x, \alpha_S)$ and the Altarelli-Parisi splitting functions $P_{ab}(x, \alpha_S)$ are computable in perturbation theory order by order in α_S .

The main difference between the factorization formulae (1) and (3) is due to the presence of the additional k_\perp integration in Eq. (1). Nonetheless, k_\perp -factorization can be recast in a form that is fully consistent with collinear factorization [2, 7]. One first solves the BFKL equation and factorizes the k_\perp dependence in Eq. (2) with respect to scale-dependent parton densities. Then, one inserts this k_\perp -dependent factor into Eq. (1) and explicitly carries out the integration over k_\perp . As a result, one ends up with Eqs. (3,4) but coefficient functions and splitting functions are no longer evaluated in fixed-order perturbation theory. They are indeed supplemented with the all-order resummation of the logarithmically-enhanced contributions $\alpha_S^n \ln^m x$ at small x . k_\perp -factorization is thus a powerful tool: the *infinite* perturbative summation of the logarithmic terms in $C^a(x, \alpha_S)$ and $P_{ab}(x, \alpha_S)$ is accomplished by a *fixed-order* computation of the off-shell cross section $\hat{\sigma}(x, Q^2; k_\perp)$ [2, 7] and of the kernel of the BFKL equation [8, 9, 10].

Note that the coefficient functions and the parton densities on the right-hand side of Eq. (3) are not separately physical observables. One has some freedom in redefining them (and, correspondingly, the splitting functions in Eq. (4)), provided that their convolution, i.e. the physical cross section on the left-hand side of Eq. (3), is left unchanged. This freedom is called factorization-scheme dependence. The steps leading from the k_\perp -factorization formulae (1,2) to resummed splitting and coefficient functions can be performed by having full control of the factorization-scheme dependence [11, 7]. This property is essential for the consistency between k_\perp -factorization and collinear factorization.

To the purpose of discussing the main theoretical results of the resummation programme, it is convenient to introduce the N -moments. For any function $g(x, \dots)$, I define its N -moments $g_N(\dots)$ in the usual way:

$$g_N(\dots) \equiv \int_0^1 dx x^{N-1} g(x, \dots) . \quad (5)$$

In particular, the *anomalous dimensions* $\gamma_{ab,N}(\alpha_S)$ are related to the $N+1$ -moments of the Altarelli-Parisi splitting functions, that is, $\gamma_{ab,N}(\alpha_S) \equiv P_{ab,N+1}(\alpha_S)$. Note that logarithmic contributions of the type $\ln^{n-1} x$ in x -space correspond to multiple poles $(1/N)^n$ for $N \rightarrow 0$ in N -space.

As for the parton flavours a , it is useful to consider singlet ($a = g, S$) and non-singlet ($a = \{a_{NS}\}$) terms. Thus \tilde{f}_g is the gluon density, and the singlet-quark density is related to the quark (\tilde{f}_{q_f}) and antiquark ($\tilde{f}_{\bar{q}_f}$) densities as $\tilde{f}_S = \sum_f (\tilde{f}_{q_f} + \tilde{f}_{\bar{q}_f})$.

The first general results following from the k_\perp -factorization theorem regard the classification of the small- x logarithmic contributions. One can show [7] that logarithmic terms in the non-singlet sector (coefficient functions and splitting functions) are suppressed by

(at least) a power of x with respect to similar terms in the singlet sector. Thus, non-singlet corrections are negligible at small x and are not considered in the following (they are reviewed elsewhere in these Proceedings [12]). In the singlet channel one can show that the perturbative expansions of the splitting functions and of the coefficients functions for totally inclusive cross sections contain at most one power of $\ln x$ (or $1/N$, in N space) for each power of α_S . Thus, one can set up an improved perturbative expansion at small x by systematically resumming the leading logarithmic (LL) contributions $(\alpha_S/N)^n$, the next-to-leading logarithmic (NLL) terms $\alpha_S(\alpha_S/N)^n$ and so forth.

Among the various anomalous dimensions and coefficients functions, only the gluon anomalous dimensions $\gamma_{gg,N}$ and $\gamma_{gS,N}$ have LL contributions. They are given by

$$\gamma_{gg,N}(\alpha_S) = \frac{C_A}{C_F} \gamma_{gS,N}(\alpha_S) + \mathcal{O}\left(\alpha_S \left(\frac{\alpha_S}{N}\right)^n\right) = \gamma_N(\alpha_S) + \mathcal{O}\left(\alpha_S \left(\frac{\alpha_S}{N}\right)^n\right), \quad (6)$$

where $\gamma_N(\alpha_S)$ is the celebrated BFKL anomalous dimension [13, 2, 7], whose explicit expression will not be reported here.

All the NLL calculations carried out during the last few years lead to analytic formulae given in terms of the BFKL anomalous dimension γ_N . The NLL contributions to the quark anomalous dimensions $\gamma_{Sg,N}$ and $\gamma_{SS,N}$ have been computed in Ref. [7]. Among the NNL terms in the gluon anomalous dimensions, those proportional to N_f have been evaluated recently [10]. The calculation of the remaining NLL contributions to $\gamma_{gg,N}$ and $\gamma_{gS,N}$ is in progress [8, 9, 10]. The processes whose coefficient functions are known to NLL accuracy are the following: the structure function F_2 and the longitudinal structure function F_L in DIS [7], the heavy-flavour photoproduction [2] and hadroproduction [2, 3, 4] cross sections, the heavy-flavour cross sections via intermediate-vector-boson exchange [14] and the heavy-quark structure functions in DIS [2]. Since preliminary experimental results on charm production in DIS at HERA have been presented at this Workshop [15], in the following Section I review the resummed formulae for the heavy-quark structure functions.

Note that all the NLL results summarized above are factorization-scheme dependent [11, 16]. The factorization scheme that is actually used has to be always specified.

3 Heavy-flavour structure functions in DIS

The heavy-flavour structure functions $F_2^{Q\bar{Q}}$, $F_L^{Q\bar{Q}}$ are completely analogous to the customary DIS structure functions F_2 , F_L with the only additional constraint that heavy quarks of mass M are produced in the final state. The collinear factorization formula can be written as follows ($i = 2, L$)

$$F_i^{Q\bar{Q}}(\xi, Q^2; M^2) = \sum_a \int_\xi^1 \frac{dz}{z} C_i^{Q\bar{Q},a}(\xi/z, \alpha_S(Q^2); Q^2/M^2) \tilde{f}_a(z, Q^2) . \quad (7)$$

Note that in Eq. (7) I have defined $F_i^{Q\bar{Q}}$ as function of Q^2 , M^2 and the inelasticity variable ξ , which is related to the customary Bjorken variable x by $\xi = x(1 + 4M^2/Q^2)$. Since

$0 \leq \xi \leq 1$, considering the N -moments of $F_i^{Q\bar{Q}}(\xi, Q^2; M^2)$ with respect to ξ , one obtains

$$F_{i,N}^{Q\bar{Q}}(Q^2; M^2) = \sum_a C_{i,N}^{Q\bar{Q},a}(\alpha_S(Q^2); Q^2/M^2) \tilde{f}_{a,N}(Q^2) . \quad (8)$$

The coefficient functions $C_{i,N}^{Q\bar{Q},a}$ have been computed up to one-loop order in Ref. [17]. The corresponding resummed formulae to NLL accuracy were obtained in Ref. [2]. The gluon coefficient functions are given as follows

$$C_{i,N}^{Q\bar{Q},g}\left(\alpha_S; \frac{Q^2}{M^2}\right) = R_N \frac{Q^2}{16\pi^2\alpha M^2} h(\gamma_N) \left(\frac{M^2}{Q^2}\right)^{\gamma_N} \left(1 + \frac{Q^2}{4M^2}\right)^N K_N^{(i)}\left(\frac{Q^2}{M^2}\right), \quad (9)$$

where $\gamma_N = \gamma_N(\alpha_S)$ is the BFKL anomalous dimension in Eq. (6) and $R_N = R_N(\alpha_S)$ is a factorization-scheme dependent term (it is the same in the $\overline{\text{MS}}$ and DIS schemes: see Eq. (3.17) of the second paper in Ref. [7]). The functions $h(\gamma)$ and $K_N^{(2)}(Q^2/M^2)$ are respectively given in Eqs. (3.9) and (4.18) of the second paper in Ref. [2]. The explicit expression for $K_N^{(L)}(Q^2/M^2)$ was not reported in Ref. [2]. It is:

$$\begin{aligned} K_N^{(L)}\left(\frac{Q^2}{M^2}\right) &= \left(1 + \frac{Q^2}{4M^2}\right)^{-(N+1)} \frac{3}{(7-5\gamma_N)(1+2\gamma_N)} \left\{ \left[1 - \gamma_N + \frac{6M^2}{Q^2}\right] \right. \\ &+ \left(1 + \frac{Q^2}{4M^2}\right)^{\gamma_N-1} \left[\gamma_N(1-\gamma_N) \frac{Q^2}{2M^2} - 2(1-\gamma_N) - \frac{6M^2}{Q^2} \right] \\ &\cdot \left. F\left(1-\gamma_N, 1/2; 3/2; \frac{Q^2}{Q^2+4M^2}\right) \right\} , \end{aligned} \quad (10)$$

where $F(a, b; c; z)$ is the hypergeometric function. In the $\overline{\text{MS}}$ and DIS schemes the NLL contributions to the singlet-quark coefficient functions are obtained from those in the gluon coefficient functions by the following colour-charge relation

$$C_{i,N}^{Q\bar{Q},s}\left(\alpha_S; \frac{Q^2}{M^2}\right) = \frac{C_F}{C_A} \left[C_{i,N}^{Q\bar{Q},g}\left(\alpha_S; \frac{Q^2}{M^2}\right) - \frac{\alpha_S}{2\pi} C_{i,N}^{Q\bar{Q},g(0)}\left(\frac{Q^2}{M^2}\right) \right] , \quad (11)$$

where the second term in the square bracket is the lowest-order term in the perturbative expansion of $C_{i,N}^{Q\bar{Q},g}(\alpha_S, Q^2/M^2)$.

4 Factorization-theorem invariants at small x

So far, the most important phenomenological consequence of small- x resummation is its large effect [18, 19, 20] on the scaling violation of the proton structure function F_2 in the HERA kinematic region. To asses the relevance of this result, further investigations, in particular on the impact of subleading contributions, are necessary. At the same time, however, it is worthwhile pointing out that the factorization-scheme dependence can play an important role.

The freedom of arbitrarily choosing the factorization scheme is not a particular feature of small- x dynamics. In general, its effect amounts to a redefinition of the input parton

densities that is perturbatively under control. The effect, instead, can be quite large in the small- x region because each power of α_S can be accompanied by an enhancing logarithmic factor of $\ln 1/x$.

After having fixed splitting and coefficient functions (using either fixed-order or resummed perturbation theory), the QCD analysis of the HERA data on F_2 requires to fit the x -dependence of the parton densities at some input scale Q_0 . As a result one may find the dominant small- x behaviour $\tilde{f}_g(x, Q_0^2) \sim \text{const.}$ as well as a steeper behaviour $\tilde{f}_g(x, Q_0^2) \sim x^{-\lambda}$. No matter the behaviour one finds, one would be led to conclude that it is due to non-perturbative phenomena. Actually, by simple power counting one can argue [21] that the difference between these two extreme behaviours can consistently be attributed to different factorization schemes or, equivalently, to higher-order (i.e. beyond the LL accuracy) logarithmic contributions in perturbation theory.

This argument may appear very formal (it is indeed provoking, to some extent) and with no physical content. Actually, the factorization-scheme dependence, rather than an ambiguity in higher-order perturbative coefficients, has to be regarded more physically as a parametrization of our ignorance in factorizing perturbative from non-perturbative physics. Studying the proton structure function we have at our disposal two quantities, F_2 and $dF_2/d\ln Q^2$, and, besides perturbative QCD, we have to introduce two non-perturbative distributions, \tilde{f}_S and \tilde{f}_g . Roughly speaking, the sole proton structure function does not provide sufficient information to fully disentangle perturbative from non-perturbative dynamics at small- x .

A better theoretical control on perturbative physics can be achieved by exploiting the very physical content of the collinear-factorization theorem, that is, the universality (process independence) of the parton densities. This means that the same parton densities and the same perturbative approach have to be used to study the small- x behaviour of different physical observables. Considering more observables, besides F_2 , one can (over-)constrain the definition of the parton densities thus eliminating the factorization-scheme dependence and emphasizing the perturbative QCD component.

The k_\perp -factorization approach shares these universality features and provides resummed perturbative calculations for many different processes. Thus, one can introduce quantities that are invariant with respect to factorization-scheme transformations. As discussed in Refs. [2, 14], one can consider ratios of hadronic cross sections and properly defined K -factors (the functions $K_N^{(i)}$ in Eq. (9) are an example for that) in which the dependence on the parton densities cancels. These K -factors are factorization-scheme independent by definition. Analogously, by simply using two hadronic observables one can formulate the dynamics of scaling violation entirely in terms of perturbative quantities that play the role of physical anomalous dimensions [21].

4.1 Physical anomalous dimensions

Considering two different hadronic observables F_A and F_B (for instance, F_2 and F_L or $F_2^{Q\bar{Q}}$), one can write down evolution equations in the following form

$$\frac{dF_A(x, Q^2)}{d \ln Q^2} = \int_x^1 \frac{dz}{z} \left[\Gamma_{AA}(x/z, \alpha_S(Q^2)) F_A(z, Q^2) + \Gamma_{AB}(x/z, \alpha_S(Q^2)) F_B(z, Q^2) \right] . \quad (12)$$

The parton densities have completely disappeared. Equation (12) relates the scaling violation of two physical observables to the actual value of the same observables. Thus the kernels $\Gamma_{AB}(x, \alpha_S)$ are *physical observables* as well. Owing to the formal resemblance to the Altarelli-Parisi splitting functions in Eq. (4), the kernels $\Gamma_{AB}(x, \alpha_S)$ can be considered as *physical splitting functions* and, correspondingly, their moments $\Gamma_{AB, N}(\alpha_S)$ are *physical anomalous dimensions*.

Each of the physical splitting functions is factorization-scheme invariant and thus unambiguously computable in perturbation theory. In other words, from the viewpoint of perturbative QCD, each $\Gamma_{AB}(x, \alpha_S)$ is completely analogous to the celebrated ratio $R_{e^+e^-} = \sigma(e^+e^- \rightarrow \text{hadrons})/\sigma(e^+e^- \rightarrow \mu^+\mu^-)$ in e^+e^- annihilation.

Although the kernels Γ_{AB} can be evaluated without carrying out any factorization procedure, they can be related [21] to the customary splitting and coefficient functions. Therefore, for any process whose coefficient functions are known to one-loop order, one can straightforwardly obtain the corresponding physical anomalous dimensions to the same accuracy. Analogously, using the resummed calculations listed in Sect. 2, one can derive resummed expressions for the physical anomalous dimensions. In the case of $\{F_A, F_B\} = \{F_2, F_L\}$, for instance, the anomalous dimension Γ_{LL} contains LL terms and they are simply given by the BFKL anomalous dimension:

$$\Gamma_{LL, N}(\alpha_S) = \gamma_N(\alpha_S) + \mathcal{O}(\alpha_S(\alpha_S/N)^n) , \quad (13)$$

while Γ_{2L} (which drives the scaling violation in the quark channel) enters only to NLL order and is given by

$$\frac{\alpha_S}{2\pi} \Gamma_{2L, N}(\alpha_S) = \frac{\alpha_S}{2\pi} \left[\frac{1}{1 - \gamma_N(\alpha_S)} + \frac{3}{2} \gamma_N(\alpha_S) \right] + \mathcal{O}(\alpha_S^2(\alpha_S/N)^n) . \quad (14)$$

The expression (14) is remarkably simple in comparison with those [2, 7, 10] of other factorization-scheme-dependent quantities to NLL accuracy.

In the evolution equation (12) the small- x perturbative dynamics is completely controlled by the physical anomalous dimensions. No subtle theoretical interplay between perturbative logarithms and steepness of parton densities takes place. The physical anomalous dimensions can be evaluated both in fixed-order perturbation theory and in resummed perturbation theory. For any given set of observables and kinematic region of x , one can thus compare the two approaches and study the theoretical accuracy of the perturbative expansion. Having done that, one can go back to the customary partonic picture and investigate more safely the small- x behaviour of the non-perturbative parton densities.

A more detailed discussion on physical anomalous dimensions can be found elsewhere [21] and some phenomenological studies have already been presented at this Workshop [22].

Acknowledgments. The results reviewed in this contribution are due to a long and intense collaboration with Marcello Ciafaloni and Francesco Hautmann. This research is supported in part by EEC Programme *Human Capital and Mobility*, Network *Physics at High Energy Colliders*, contract CHRX-CT93-0357 (DG 12 COMA). I am happy to thank the Local Organizing Committee, and in particular Giulio D’Agostini, for the organization of an excellent Workshop.

References

- [1] L.N. Lipatov, Sov. J. Nucl. Phys. 23 (1976) 338; E.A. Kuraev, L.N. Lipatov and V.S. Fadin, Sov. Phys. JETP 45 (1977) 199; Ya. Balitskii and L.N. Lipatov, Sov. J. Nucl. Phys. 28 (1978) 822.
- [2] S. Catani, M. Ciafaloni and F. Hautmann, Phys. Lett. B242 (1990) 97, Nucl. Phys. B366 (1991) 135, Proceedings of the HERA Workshop, eds. W. Buchmüller and G. Ingelman (DESY Hamburg, 1991), p. 690.
- [3] J.C. Collins and R.K. Ellis, Nucl. Phys. B360 (1991) 3.
- [4] E.M. Levin, M.G. Ryskin, Yu.M. Shabel’skii and A.G. Shuvaev, Sov. J. Nucl. Phys. 53 (1991) 657.
- [5] See, for instance, J.C. Collins, D.E. Soper and G. Sterman, in *Perturbative Quantum Chromodynamics*, ed. A.H. Mueller (World Scientific, Singapore, 1989), p. 1 and references therein.
- [6] V.N. Gribov and L.N. Lipatov, Sov. J. Nucl. Phys. 15 (1972) 438, 675; G. Altarelli and G. Parisi, Nucl. Phys. B126 (1977) 298; Yu.L. Dokshitzer, Sov. Phys. JETP 46 (1977) 641.
- [7] S. Catani and F. Hautmann, Phys. Lett. 315B (1993) 157, Nucl. Phys. B427 (1994) 475.
- [8] L.N. Lipatov and V.S. Fadin, Sov. J. Nucl. Phys. 50 (1989) 712; V.S. Fadin, R. Fiore and A. Quartarolo, Phys. Rev. D 53 (1996) 2729; V.S. Fadin, in these Proceedings.
- [9] V. Del Duca, Phys. Rev. D 54 (1996) 989, preprint EDINBURGH-96-3 (hep-ph/9604250) .
- [10] G. Camici and M. Ciafaloni, preprint DFF-250-6-96 (hep-ph/9606427).
- [11] S. Catani, M. Ciafaloni and F. Hautmann, Phys. Lett. 307B (1993) 147.
- [12] J. Blümlein, these Proceedings; B. Ermolaev, these Proceedings; B.R. Webber, these Proceedings.

- [13] T. Jaroszewicz, Phys. Lett. B116 (1982) 291.
- [14] G. Camici and M. Ciafaloni, Nucl. Phys. B467 (1996) 25.
- [15] K. Daum, these Proceedings.
- [16] M. Ciafaloni, Phys. Lett. 356B (1995) 74; R.D. Ball and S. Forte, Phys. Lett. 359B (1995) 362; S. Catani, Z. Phys. C70 (1996) 263.
- [17] E. Laenen, S. Riemersma, J. Smith and W.L. van Neerven, Nucl. Phys. B392 (1993) 162.
- [18] R.K. Ellis, F. Hautmann and B.R. Webber, Phys. Lett. 348B (1995) 582; F. Hautmann, preprint CAVENDISH-HEP-95-04.
- [19] R.D. Ball and S. Forte, Phys. Lett. 351B (1995) 313, Phys. Lett. 358B (1995) 365; S. Forte, these Proceedings.
- [20] J.R. Forshaw, R.G. Roberts and R.S. Thorne, Phys. Lett. 356B (1995) 79.
- [21] S. Catani, talk presented at the UK Phenomenology Workshop on HERA Physics (Durham, September 1995), preprint DFF 248/4/96 (to appear).
- [22] R.S. Thorne, these Proceedings; R.G. Roberts and R.S. Thorne, in preparation.