## TECHNICOLOR MECHANISMS FOR SINGLE TOP PRODUCTION

LESLEY L. SMITH

Butler County Community College, 901 S. Haverhill Rd., El Dorado, KS 67042

PANKAJ JAIN

Indian Institute of Technology, Kanpur, 208016, UP, India

DOUGLAS W. MCKAY

Department of Physics and Astronomy, University of Kansas, Lawrence, KS 66045

We investigate the contribution of technicolor mechanisms to the production of single top quarks at hadron colliders. We find that a promising candidate process is gluon-gluon fusion to produce a W-boson plus technipion, with subsequent decay of the technipion to a top quark plus a bottom quark. The top-plus-bottom mode is the dominant one when the technipion mass is larger than the top mass. We calculate the total cross section and the  $p_T$  distribution for the technipion production at Tevatron and LHC energies for a range of technipion masses, starting at 200 GeV. The decay chain of technipion to top plus bottom quarks. We study the backgrounds to our process and the kinematic cuts that maximize the signal to background. We report event rate estimates for the upgraded Tevatron and the LHC.

The discovery of the top quark<sup>1</sup> has opened up an exciting area of physics. Top production by non Standard Model (SM) mechanisms are of great value in probing new physics. It has been known for some time that the SM is not completely satisfying. It does not explain electroweak symmetry breaking and has many free parameters. Technicolor (TC) has emerged as one of the possible successors to the SM, and we present results of single top production by TC here. We know the low-energy region of QCD can be represented by a nonlinear sigma-type model  $^2$ . Thus we invoke a low-energy effective Langrangian for a class of technicolor theories which is motivated by the chiral symmetry of the Technicolor Lagrangian and the observation of vector-meson dominance in QCD  $^2$ . We are interested in the production of single top quarks and we look at the gluon fusion process  $gg \rightarrow W + P_8$ where  $P_8$  indicates a color octet, SU(2) doublet technipion<sup>3</sup>. The technipion predominantly decays to a top quark and a bottom quark, assuming  $M_{P_8} > m_t^4$ . The lowest order processes contributing to this gluon fusion are: the  $ggWP_8$  4-point interaction diagram, the s-channel gluon exchange diagram, and the 2 diagrams with  $P_8$  exchange in the t and u channels.

Table	1:	$p_T$	distribution
-------	----	-------	--------------

$p_T (\text{TeV})$	0.1	0.2	0.4	0.6	0.8	1.0
$\frac{d\sigma_{TC}}{dydp_T}$ (fb TeV <sup>-1</sup> )	62.0	46.6	14.4	4.3	1.4	0.5

Table 2: Technipion cross sections and events per year

	$M_{P_8} = 240 \text{ GeV}$	$300 { m GeV}$	$350 { m GeV}$	$400 \mathrm{GeV}$					
$\sigma_{TC}$									
$\sqrt{s} = 2 \text{ TeV}$	$0.059~{\rm fb}$	0.016  fb	0.0059  fb	0.0022  fb					
$\sqrt{s} = 14 \text{ TeV}$	56  fb	$31~{\rm fb}$	20  fb	14  fb					
# events per year									
$\sqrt{s} = 2 \text{ TeV}$	0.37	0.10	0.04	0.01					
$\sqrt{s} = 14 \text{ TeV}$	$1.7 \times 10^4$	$9.7 \times 10^3$	$6.3 \times 10^3$	$4.4 \times 10^3$					

In the Farhi-Susskind type model that we use for the present analysis, the vertices needed are the single pseudoscalar effective Lagrangian  $\mathcal{L}(\phi)_{eff} = \frac{iN_{TC}}{16\pi^2 F_T} \epsilon^{\mu\nu\lambda\rho} \int dx (\partial_\mu G^b_\nu + \frac{1}{2}g_3 f^{bcd} G^c_\mu G^d_\nu) g_2 g_3 (P_8^{+,b} \partial_\lambda W^-_\rho + P_8^{-,b} \partial_\lambda W^+_\rho)$ , the normal parity  $P_8$ - $P_8$ -G effective interaction Lagrangian<sup>5</sup> and the usual QCD triple gluon vertex. Using these vertices and the tree-level diagrams described above, we computed the order  $\alpha_2^2 \alpha_3 p_T$  distributions and the total cross section for technipion production at Tevatron and LHC energies. The final state that results from the  $P_8 + W$  production and subsequent decay of  $P_8$  into t + band then t into W + b is the same WWbb final state that results from  $t + \overline{t}$ production and decay. We comment below on cuts that reduce the  $t + \overline{t}$  background. The  $p_T$  distribution for  $p\overline{p} \to W + P_8$  has its peak at  $p_T \simeq 0.1$  TeV for  $M_{P_8} = 240$  GeV and  $\sqrt{s}=14$  TeV. We show several  $\frac{d\sigma}{dydp_T}$  values in the range between 0.1 TeV and 1 TeV for y = 0,  $M_{P_8} = 240$  GeV, and  $N_{TC} = 3$ in Table 1 in units of fb TeV<sup>-1</sup>. In Table 2 we give the corresponding total cross sections for technipion production for a range of technipion masses. We also show the events per year expected with the upgraded Tevatron and with the LHC at luminosities of  $2 \times 10^{32} cm^{-2} s^{-1}$  and  $10^{34} cm^{-2} s^{-1}$ , respectively. At Tevatron energies and expected luminosities, the event rate is unobservably small. At the LHC, however, the event rate is large enough to be interesting, allowing rather severe cuts to be made to reduce background and not lose the whole signal. For  $\sqrt{s}=14$  TeV, and a top mass of 180 GeV, the single top cross section is 117 pb, while the  $t\bar{t}$  cross section is 525 pb<sup>6</sup>. Monte Carlo studies indicate that the WWbb decay products of the TC signal differ from those of the SM  $t\bar{t}$  production and decay in that the TC  $p_T$  spectrum of the recoil W is harder than the W spectrum of  $t\bar{t}$  and the b and  $\bar{b}$  are not back-to-back. For example if we impose a typical  $p_T > 400$  GeV cut on W and a cos  $\theta > 0.6$  on on the  $b\bar{b}$  opening angle, the fraction of SM  $t\bar{t}$  events passing these cuts is less than  $10^{-3}$  while the fraction of TC events passing the cuts is about  $5 \times 10^{-2}$ . By tuning these cuts, one can substantially reduce the SM  $t\bar{t}$  background in comparison to the TC signal. Thus our preliminary study indicates that a search for charged, colored technipions in the WWbb final state at the LHC is feasible.

## Acknowledgements

We thank Phil Baringer for discussions and for providing Monte Carlo studies of t and  $\overline{t}$  production. This research was supported in part by DOE grant # DE - FG02 - 85ER40214. The computational facilities of the Kansas Institute for Theoretical and Computational Science were used for part of this work. D.W.M. thanks S.Ranjbar-Daemi and Faheem Hussain for the hospitality of the high energy group at ICPT, Trieste, during the course of this work.

## References

- CDF collaboration: F. Abe et al., Phys. Rev. Lett. 74, 2626 (1995); Phys. Rev. D 52, 2605 (1995); D0 Collaboration: S. Abachi et al., Phys. Rev. Lett. 74, 2632 (1995).
- See, for example, M.E. Peskin, in *Recent Advances in Field Theory and Statistical Mechanics*, proceedings of the Les Houches Summer School, Les Houches, France, 1982, edited by J.-B. Zuber and R. Stora (Les Houches Summer School Proceedings, Vol. 39)(North-Holland, Amsterdam, 1984)
- 3. For example, in comparison to the gluon fusion process, direct production of  $P_8$  from  $tb\overline{b}$  in the sea is down by more than an order of magnitude.
- E. Eichten, I.Hinchliffe, K. Lane, C. Quigg, *Rev. Mod. Phys.* 56, 579 (1984).
- 5. D.W. McKay, B.-L. Young, Phys. Lett. B 169, 79 (1986).
- 6. P. Baringer, personal communication.