

Leptoquark Mass Limits from Single Leptoquark Production at LEP and LEP 200

Michael A. Doncheski and Stephen Godfrey

Ottawa-Carleton Institute for Physics

Department of Physics, Carleton University, Ottawa CANADA, K1S 5B6

We investigate the discovery potential for first generation leptoquarks at the LEP and LEP 200 e^+e^- colliders. We consider single leptoquark production via resolved photon contributions which offers a much higher kinematic limit than the more commonly considered leptoquark pair production process. This process also has the advantage that it is independent of the leptoquark chirality, is almost insensitive to whether the leptoquark is scalar or vector, and only depends on the leptoquark charge. We estimate that from the nonobservation of energetic e^- jet events, the limits at LEP for all types of leptoquarks are $M_{LQ} > 90 \text{ GeV}$. For the just completed $P_{\bar{S}} = 161 \text{ GeV}$ run at LEP 200 we estimate potential scalar leptoquark mass limits of $M_{LQ} > 130 \text{ GeV}$ for $Q_{LQ} = 2/3; 4/3$ and $M_{LQ} > 148 \text{ GeV}$ for $Q_{LQ} = 1/3; 5/3$ and for vector LQ 's the limits are 140 and 154 GeV respectively. Ultimately, LEP 200 should be able to achieve discovery limits for leptoquarks of 188 GeV for the $P_{\bar{S}} = 190 \text{ GeV}$ runs. In all cases we assume electromagnetic strength coupling.

Present address: Department of Physics, Pennsylvania State University, Mont Alto, PA 17237
USA

PACS numbers: 12.15.Ji, 12.60.-i, 12.90.+b, 14.80.-j

There is considerable interest in the study of leptoquarks (LQs) – colour (anti-)triplet, spin 0 or 1 particles which carry both baryon and lepton quantum numbers. Such objects appear in a large number of extensions of the standard model such as grand unified theories, technicolour, and composite models. Quite generally, the signature for leptoquarks is very striking: a high p_T lepton balanced by a jet (or missing p_T balanced by a jet, for the q decay mode, if applicable). Searches for leptoquarks have been performed by the H1 [1] and ZEUS [2] collaborations at the HERA ep collider, by the D0 [3] and CDF [4] collaborations at the Tevatron pp collider, and by the ALEPH [5], DELPHI [6], L3 [7], and OPAL [8] collaborations at the LEP e^+e^- collider. The LQ limits obtained by the LEP experiments have for the most part considered LQ pair production although DELPHI considered single LQ production via the process $Z \rightarrow LQ + e + \text{jet}$. In this letter we consider single leptoquark production in e^+e^- collisions which utilizes the quark content of a Weizacker-Williams photon radiating off one of the initial leptons [9–14]. This process offers the advantage of a much higher kinematic limit than the LQ pair production process, is independent of the chirality of the LQ, and gives similar results for both scalar and vector leptoquarks. The cross section depends on the LQ charge since the photon has a larger u quark content than d quark content and hence has a larger cross section for LQ's which couple to the u quark than for LQ's which couple to the d quark.

The most general $SU(3) \times SU(2) \times U(1)$ invariant scalar leptoquarks satisfying baryon and lepton number conservation have been written down Buchmüller et al. [15]. However, only those leptoquarks which couple to electrons can be produced in e^- collisions and for real leptoquark production the chirality of the coupling is irrelevant. For this case the number of leptoquarks reduces to four which can be distinguished by their electromagnetic charge; $Q_{em} = -1/3, 2/3, 4/3$, and $5/3$. In our calculations we will follow the convention adopted in the literature [12] where the leptoquark couplings are replaced by a generic Yukawa coupling g which is scaled to electromagnetic strength $g^2/4 = e_{em}^2$ and allow to vary.

The process we are considering is shown in Fig. 1. The parton level cross section is

trivial, given by:

$$\hat{\sigma}(s) = \frac{2^2 e_m}{M_s} (M_s^2)^{P-2} \quad (1)$$

Convoluting the parton level cross section with the quark distribution in the photon one obtains the expression

$$\begin{aligned} \sigma(s) &= \int_0^1 f_{q\gamma}(z; M_s^2) \hat{\sigma}(s) dz \\ &= f_{q\gamma}(M_s^2=s; M_s^2) \frac{2^2 e_m}{s} : \end{aligned} \quad (2)$$

We note that the interaction Lagrangian used in Ref. [13] associates a factor $1=2^{P-2}$ with the leptoquark-lepton-quark coupling. Thus, one should compare our results with those in Ref. [13] with 2^2 . We give results with 2^2 chosen to be 1.

For e^+e^- colliders the cross section is obtained by convoluting the expression for the resolved photon contribution to e^- production of leptoquarks, Eqn. (2), with the Weizsacker-Williams effective photon distribution

$$(e^+e^- \rightarrow X S) = \frac{2^2 e_m}{s} \int_{M_s^2=s}^1 \frac{dx}{x} f_{=e}(x; M_s^2=2) f_{q\gamma}(M_s^2=(xs); M_s^2) \quad (3)$$

with the Weizsacker-Williams effective photon distribution given by

$$f_{=e}(x; E) = \frac{e_m}{2} \left(\frac{[1 + (1-x)^2]}{x} \ln \frac{E^2 (1-2x+x^2)}{m_e^2 (1-x+x^2=4)} + x \ln \frac{(2-x)}{x} + \frac{2(x-1)}{x} \right) : \quad (4)$$

There exist several different quark distribution functions in the literature [16-20]. The different distributions give almost identical results for the $Q_{LQ} = 1=3; 5=3$ leptoquarks and for the $Q_{LQ} = 2=3; 4=3$ leptoquarks give LQ cross sections that vary by most a factor of two, depending on the kinematic region. We obtain our results using the GRV distribution functions [19] which we take to be representative of the quark distributions in the photon.

Before proceeding to our results we consider possible backgrounds [21]. The leptoquark signal consists of a jet and electron with balanced transverse momentum and possibly activity from the hadronic remnant of the photon. The only serious background is a hard scattering of

a quark inside the photon by the incident lepton via t-channel photon exchange; eq. 1. We plot the invariant mass distribution for this background in our plots of the LQ cross sections and find that it is typically smaller than our signal by two orders of magnitude. For the LQ invariant mass distribution we chose a 5 GeV invariant mass bin so that $dM = 5 \text{ GeV}$. Related to this process is the direct production of a quark pair via two photon fusion

$$e^+ \rightarrow e^+ q \bar{q} \quad (5)$$

However, this process is dominated by the collinear divergence which is actually well described by the resolved photon process eq. 1 given above. Once this contribution is subtracted away the remainder of the cross section is too small to be a concern [21]. Another possible background consists of ν 's pair produced via various mechanisms with one decaying leptonically and the other decaying hadronically. Because of the neutrinos in the final state it is expected that the electron and jet's p_T do not in general balance which would distinguish these backgrounds from the signal. However, this background should be checked in a realistic detector Monte Carlo to be sure. The remaining backgrounds originate from heavy quark pair production with one quark decaying semileptonically and only the lepton being observed with the remaining heavy quark not being identified as such. All such backgrounds are significantly smaller than our signal in the kinematic region we are concerned with.

In Fig. 2 we show the cross sections for $\sqrt{s} = 91.17 \text{ GeV}$. In our results we assume $\text{BR}(LQ \rightarrow e + q) = 1$. We interpret the non-observation of any e jet events by any of the LEP experiments to signify that a LQ does not exist up to the leptoquark mass that would produce at least 3 e jet events with the integrated luminosities of the combined LEP experiments. Assuming $\text{BR}(LQ \rightarrow e \text{ jet}) = 1$ this leads to LQ search limits of $M_{LQ} > 90 \text{ GeV}$ for all first generation leptoquark types. If instead $\text{BR}(LQ \rightarrow e + q) = 0.5$ and $\text{BR}(LQ \rightarrow \nu + q) = 0.5$ the second LQ decay mode would have an even more dramatic signature than the one we consider; a high p_T monojet balanced against a large missing p_T . Thus, in this case the sum of the two possible decays would give limits similar to the one

given above.

In Fig. 3 we show the cross sections for $\sqrt{s} = 161 \text{ GeV}$, the energy of the most recent LEP200 run. For an integrated luminosity of 10 pb^{-1} per LEP experiment we interpret the non-observation of any e⁺jet events for the LQ mass that would result in 3 events as scalar LQ search limits of $M_{LQ} > 130 \text{ GeV}$ for $Q_{LQ} = 2=3; 4=3$ and $M_{LQ} > 148 \text{ GeV}$ for $Q_{LQ} = 1=3; 5=3$ and 140 GeV and 154 GeV respectively for the vector leptoquarks. These limits represent a significant improvement over the LEP limits given above and are quite competitive with similar published limits obtained by the Tevatron experiments [3,4].

Figures 4 and 5 give the cross sections for $\sqrt{s} = 175 \text{ GeV}$ and $\sqrt{s} = 190 \text{ GeV}$ respectively. At present we do not know for certain what the total integrated luminosities will be for each of these energies. We use the currently planned 25 pb^{-1} at $\sqrt{s} = 175 \text{ GeV}$ and 500 pb^{-1} at $\sqrt{s} = 190 \text{ GeV}$ for each of the LEP experiments. Again, using the criteria that none of the experiments see any e⁺jet events we obtain search limits up to 170 GeV for $\sqrt{s} = 175 \text{ GeV}$ and 188 GeV for $\sqrt{s} = 190 \text{ GeV}$. The search limits for the different charge and spin assumptions are summarized in Table 1.

In this letter we have used the resolved photon contributions to single leptoquark production to estimate potential limits on leptoquark masses. This production mechanism has the advantage that it can produce both scalar and vector leptoquarks with roughly the same cross section and of being insensitive to the chiral properties of the leptoquarks. It's only dependence on LQ properties is on the LQ charge because the photon has a larger u-quark than d-quark content. The potential limits we obtain from the non-observation of LQ's at $\sqrt{s} = 161 \text{ GeV}$ are competitive with published limits of the Tevatron collaborations. In general, the limits obtainable at LEP200 will be slightly lower than direct limits that can be obtained at HERA [1,2]. Nevertheless, limits obtained in searches at LEP200 should complement those obtained at HERA and the Tevatron for two reasons: First, LQ's will have very dramatic and distinct signatures at LEP200 in contrast to the Tevatron and HERA where their presence are determined by statistical enhancements over the SM backgrounds. Second, the limits obtained at LEP200 tend to be less dependent on the properties of the

LQ's than results obtained at ep and hadron colliders. Finally, we remind the reader that our results are of course only theorist's estimates which should be examined more closely and carefully than has been described here. For example, we have not included detector acceptances in our estimates, perhaps oversimplistically assuming that the LQ decay products will decay isotropically in the detector resulting in fairly high detection efficiencies. In addition, a detailed detector Monte Carlo of possible backgrounds should be performed. Nevertheless we believe our estimates to be fairly robust and are not likely to be changed substantially by a more rigorous scrutinization.

ACKNOWLEDGMENTS

This research was supported in part by the Natural Sciences and Engineering Research Council of Canada. The authors thank Bob Carnegie for helpful conversations and his encouragement and Dean Karlen for useful discussions.

REFERENCES

- [1] H1 Collaboration, I.Abt et al, Nucl.Phys.B 396, 3 (1993); H1 Collaboration, I.Ahmed et al, Z.Phys.C 64, 545 (1994).
- [2] ZEUS Collaboration, M.Derrick et al, Phys.Lett.B 306, 173 (1993).
- [3] D0 Collaboration, S.Abachi et al, Phys.Rev.Lett. 72, 965 (1994).
- [4] CDF Collaboration, F.Abe et al, Phys.Rev.D 48, 3939 (1993).
- [5] ALEPH Collaboration, DeCamp et al, Phys.Rept. 216, 253 (1992).
- [6] DELPHI Collaboration, P.Abreu et al, Phys.Lett.B 316, 620 (1993).
- [7] L3 Collaboration, Adriani et al, Phys.Rept. 236, 1 (1993).
- [8] OPAL Collaboration, G.Alexander et al, Phys.Lett.B 263, 123 (1991).
- [9] M.A.Doncheski and S.Godfrey, Phys.Rev.D 49, 6220 (1994).
- [10] M.A.Doncheski and S.Godfrey, Phys.Rev.D 51, 1040 (1995).
- [11] O.J.Eboli, E.M.Gregores, M.B.Magro, P.G.Mercadante, and S.F.Novaes, Phys.Lett.B 311, 147 (1993).
- [12] H.Nadeau and D.London, Phys.Rev.D 47, 3742 (1993); G.Belanger, D.London and H.Nadeau, Phys.Rev.D 49, 3140 (1994).
- [13] J.L.Hewett and S.Pakvasa, Phys.Lett.B 227, 178 (1989).
- [14] For related papers see also: J.E.Cieza Montalvo and O.J.P.Eboli, Phys.Rev.D 47, 837 (1993); T.M.Aliev and Kh.A.Mustafaev, Yad.Fiz. 58, 771 (1991); V.Ilyin et al, Phys.Lett.B 351, 504 (1995); erratum B 352, 500 (1995); Phys.Lett.B 356, 531 (1995).
- [15] W.Buchmuller, R.Ruckl, and D.Wyler, Phys.Lett.B 191, 442 (1987).

- [16] A . Nicolaidis, Nucl. Phys. B 163, 156 (1980).
- [17] D W . Duke and J F . Owens, Phys. Rev. D 26, 1600 (1982).
- [18] M . Drees and K . Grassie, Z. Phys. C 28, 451 (1985); M . Drees and R . Godbole, Nucl. Phys. B 339, 355 (1990).
- [19] M . Glück, E . Reya and A . Vogt, Phys. Lett. B 222, 149 (1989); Phys. Rev. D 45, 3986 (1992); Phys. Rev. D 46, 1973 (1992).
- [20] H . Abramowicz, K . Charchula, and A . Levy, Phys. Lett. B 269, 458 (1991).
- [21] Hadronic backgrounds in e^+e^- and e^-e^- collisions and associated references are given in M A . Doncheski, S . Godfrey, and K A . Peterson, Carleton Preprint O C I P /C -94-2 (hep-ph/9407348).

TABLES

TABLE I. Mass limits and discovery limits for leptoquarks at LEP and LEP200. Our results are based on the non-observation of any leptoquark candidates when 3 are expected for the given LQ mass. We take $\epsilon = 1$ and assume $\text{BR}(LQ \rightarrow e + q) = 1$. Our results are given in GeV.

P _s (GeV)	L (pb ⁻¹)	Scalar LQ					Vector LQ				
		Q = 2=3	4=3	Q = 1=3	5=3	Q = 2=3	4=3	Q = 1=3	5=3		
91.17	720	90		90		90		91			
161	40	130		148		140		154			
175	100	154		167		162		170			
190	2000	187		188		188		189			




Fig 1: The resolved photon contribution for leptoquark production in e^+e^- collisions.




Fig 2: The cross sections for scalar leptoquark production due to resolved photon contributions in e^+e^- collisions for $\sqrt{s} = 91.17 \text{ GeV}$. chosen to be 1 and resolved photon distribution functions of Glück, Reya and Vogt [19] are used. The dashed line is the $e^+e^- \rightarrow q\bar{q}$ background. For the LQ invariant mass distribution we use a 5 GeV invariant mass bin so that $dM = 5 \text{ GeV}$.




Fig 3: The cross sections for leptoquark production due to resolved photon contributions in e^+e^- collisions for $\sqrt{s} = 161$ GeV. See Fig. 2 for details.




Fig 4: The cross sections for leptoquark production due to resolved photon contributions in e^+e^- collisions for $\sqrt{s} = 175$ GeV. See Fig. 2 for details.




Fig 5: The cross sections for leptoquark production due to resolved photon contributions in e^+e^- collisions for $\sqrt{s} = 190$ GeV. See Fig. 2 for details.