

BI-TP 96/40
August 1996

ANOMALOUS ELECTROMAGNETIC PROCESSES AT HIGH TEMPERATURES[†]

R. Baier¹, M. Dirks¹ and O. Kober¹

¹*Fakultät für Physik, Universität Bielefeld, D-33501 Bielefeld, Germany*

Abstract

Chiral theories of constituent quarks interacting with bosons and photons at high temperatures are studied. In the expected chirally symmetric phase effective electromagnetic anomalous couplings for e.g. $\pi\sigma \rightarrow \gamma\gamma$, $\gamma \rightarrow \pi\pi\pi\sigma$, etc., are derived by applying functional methods.

[†] Invited talk at the Workshop on Quantum Chromodynamics: Confinement, Collisions, and Chaos, 3 - 8 June 1996, The American University of Paris, Paris

1 Introduction

In this contribution we discuss our approach [1] to confirm the interesting conjecture by Pisarski [2, 3], who pointed out that although the chiral (abelian as well as nonabelian) anomaly [4, 5] in terms of fundamental fields is temperature independent [6], the manifestation of the anomaly, however, in terms of effective fields changes with temperature. When considering e.g. $\pi^0 \rightarrow 2\gamma$ the observation [2, 3] is: "In a hot, chirally symmetric phase, π^0 doesn't go into 2γ , but $\pi^0\sigma$ does"!

In the following we restrict our discussion to these electromagnetic couplings, which have first been found [2, 3] in the framework of the constituent quark model [7] by calculating the contribution from the Feynman one-loop triangle (box) diagrams at nonzero, high temperature T .

Our approach also starts from the linear sigma model with constituent quarks interacting with bosons and photons [7]. In order to include the effect of the axial anomaly on the bosons we transform the basis of right- and left-handed quarks following Manohar et al. [8], but we take into account a chirally symmetric phase [9] at high T . At $T = 0$ and in the spontaneously broken phase this procedure gives the anomalous couplings by applying functional methods for the evaluation of the fermion determinant in connection with path integrals [10].

2 Effective Chiral Lagrangians

We consider the $SU(2)_L \otimes SU(2)_R$ lagrangian for N_C coloured right- and left-handed quarks parametrized in the linear form ("Σ - basis"),

$$\mathcal{L} = \bar{\psi}_L i \not{\partial} \psi_L + \bar{\psi}_R i \not{\partial} \psi_R - g(\bar{\psi}_L \Sigma \psi_R + \bar{\psi}_R \Sigma^\dagger \psi_L) + \mathcal{L}_{boson}, \quad (2.1)$$

with the $SU(2)$ matrix [7] $\Sigma = \sigma + i\vec{\tau} \cdot \vec{\pi}$. The electromagnetic interaction is as usual, $\partial_\mu \rightarrow \partial_\mu + ieQA_\mu$, where A_μ is the photon field and $Q = 1/2$ ($1/3 + \tau_3$). The explicit form of the boson lagrangian \mathcal{L}_{boson} is not needed. Although in the strictly symmetric phase the constituent quark mass m has to vanish, we nevertheless introduce explicitly a breaking term

$$\mathcal{L}_m = -m\bar{\psi}\psi. \quad (2.2)$$

We treat m as an explicit regularization parameter, which is finally removed in the symmetric phase by performing the limit $m \rightarrow 0$. However, we do not break the chiral symmetry spontaneously by the standard redefinition of the scalar field σ .

We change the quark fields [8] by

$$\psi_L' \equiv \xi^\dagger \psi_L, \quad \psi_R' \equiv \xi \psi_R, \quad (2.3)$$

("ξ- basis") defining a unitary matrix ξ in such a way that the quark-pion coupling (2.1) becomes replaced by a derivative coupling. In the symmetric phase we obtain for nonvanishing m , after expanding in terms of the boson fields $(\sigma, \vec{\pi})$ up to quadratic terms,

$$\xi \simeq \exp\left[\frac{i\vec{\tau} \cdot \vec{\pi}}{2(m/g + \sigma)}\right], \quad (2.4)$$

which replaces the transformation matrix in the broken phase,

$$\xi = \exp\left[\frac{i\vec{\tau} \cdot \vec{\pi}}{2F_\pi}\right], \quad (2.5)$$

where F_π is the pion decay constant. In the chirally broken phase the transformation (2.5) may be obtained from (2.4) by first shifting $\sigma \rightarrow \langle \sigma \rangle + \sigma \doteq F_\pi + \sigma$ and then performing the well defined

limit $m \rightarrow 0$, while the dynamical σ field becomes heavy and decouples. However, approaching the symmetric phase, where $\langle \sigma \rangle$ and F_π vanish, the σ field does not decouple and the mass parameter has to be kept as $m \neq 0$.

The transformation (2.3) yields

$$\mathcal{L} = \bar{\psi}'(i\not{D} - m)\psi' + \mathcal{L}_{boson} + O(g\sigma\bar{\psi}'\psi') , \quad (2.6)$$

with

$$D_\mu = \partial_\mu + ieQA_\mu + i\bar{V}_\mu + i\bar{A}_\mu\gamma_5 , \quad (2.7)$$

where e.g. the axial current \bar{A}_μ becomes, keeping $m \neq 0$ fixed and expanding in the fields,

$$\bar{A}_\mu \simeq \frac{1}{2} \left(\frac{g}{m} \right)^2 \left[\left(\frac{m}{g} - \sigma \right) \vec{\tau} \cdot \partial_\mu \vec{\pi} - \vec{\tau} \cdot \vec{\pi} \partial_\mu \sigma \right] + \dots . \quad (2.8)$$

It is important to notice that because of the factor $(g/m)^2$ the current \bar{A}_μ becomes singular for $m \rightarrow 0$, and as a consequence the symmetric limit is not immediately obtained.

Although the lagrangian remains invariant under the transformation (2.3), the jacobian is not unity, as it is well known. Thus, in order to obtain equivalent low-energy physics in both representations, i.e. in the " Σ -" as well as in the " ξ - basis", we have to calculate the fermion determinant for the transformation (2.3) at high temperature, and finally we have to perform the chirally symmetric limit for $m \rightarrow 0$.

3 Zeta- Regularization at High Temperature

The jacobian for the transformation (2.3) of the quark fields is determined by standard steps [7]. First one defines infinitesimal transformations by introducing a continuous parameter t , $0 \leq t \leq 1$, and extending the transformation (2.4),

$$\xi \rightarrow \xi_t \equiv \exp\left[\frac{it\vec{\tau} \cdot \vec{\pi}}{2(m/g + \sigma)}\right] \equiv \exp[it\vec{\pi}] . \quad (3.1)$$

The effective lagrangian is defined by

$$\int d^4x \mathcal{L}_{odd} \equiv 2 \int_0^1 dt \operatorname{tr}(\bar{\pi}\gamma_5) , \quad (3.2)$$

where \mathcal{L}_{odd} contains the interactions in the odd-intrinsic-parity sector of the chiral model under consideration. In the following we only consider local terms as given by a Taylor expansion in t .

In order to derive the anomalous coupling in leading order for $\pi^0\sigma \rightarrow 2\gamma$ in the chirally symmetric phase, we assume that the thermal bath is constituted by the π^0 and σ fields, to which external photons are coupled, i.e. the photons are not thermalized due to their small electromagnetic coupling. In this respect we differ from Pisarski's treatment [3], where the photons are kept in thermal equilibrium with π^0 and σ .

To be explicit we have to look for the nonvanishing contributions to

$$\mathcal{L}_T^{2\gamma} = \lim_{m \rightarrow 0} 2N_C \int_0^1 dt \operatorname{tr}' \left(\frac{\tau_3 \pi^0}{2(m/g + \sigma)} \gamma_5 \zeta_s(x) |_{2\gamma} \right) , \quad (3.3)$$

where we apply the zeta function method [11] to regularize (3.3). tr' indicates the Dirac and flavour trace. This method is combined with the expansion for the heat kernel [7] (for massive fermions) $H(x, \tau)$ by the relation,

$$\zeta_s(x) \equiv \frac{1}{\Gamma(s)} \int_0^\infty d\tau \tau^{s-1} H(x, \tau) . \quad (3.4)$$

At $T = 0$, in the broken phase, the ζ -function regularization starts with a convergent expression $\zeta_s(x)$ for $s > 0$, which is then analytically continued to $s = 0$. At high T , i.e. in the chirally symmetric phase, we propose to continue in the variable s in such a way that a non-trivial limit for \mathcal{L}_{odd} , i.e. here for $\mathcal{L}_T^{2\gamma}$, is finally obtained in the $m \rightarrow 0$ limit.

A few technical steps have to be performed:

- at high T it is convenient to continue first to euclidean coordinates and momenta ($p_E \equiv (\vec{p}, p_4)$);
- introducing a complete set of energy (Matsubara) and momentum eigenstates one obtains the expansion for the heat kernel (3.4),

$$H(x, \tau) \simeq \frac{\tau^2}{2} T \Sigma \int \frac{d^3 p}{(2\pi)^3} \exp[-\tau(p_E^2 + m^2)] \hat{\sigma}^2. \quad (3.5)$$

Here only the electromagnetic couplings in (2.7) are kept, where expressed in terms of the electromagnetic field tensor we have $\hat{\sigma} \equiv eQ \sigma_{\mu\nu} F^{\mu\nu}/2$. Terms of $O(\tau^0, \tau^1)$, which either give vanishing contributions after taking the Dirac trace in (3.3), or do not contribute at $T = 0$, or lead to interactions in higher orders of the boson fields at high T , are dropped. Σ sums the fermionic frequencies $p_4 = 2\pi T(n + 1/2)$ with integer n ;

- in order to evaluate the Matsubara frequency sum we first perform the τ -integration in (3.4) inserting (3.5); it is convenient to introduce the following dimensionless functions [12] $I(y^2; s + 2, \frac{1}{2})$, where $y = m/(2\pi T)$. For $s > 0$ they have the series expansion[12],

$$\begin{aligned} T \Sigma \int \frac{d^3 p}{(2\pi)^3} \frac{1}{(p_E^2 + m^2)^{s+2}} &\equiv \frac{T^{-2s}}{(2\pi)^{2s+1}} I(y^2; s + 2, \frac{1}{2}) \\ = \frac{T^{-2s}}{(2\pi)^{2s+1}} \frac{1}{4\pi^{3/2} \Gamma(s+2)} \Sigma_0^\infty \frac{(-)^k}{k!} \Gamma(k + s + 1/2) \zeta(2k + 2s + 1, \frac{1}{2}) y^{2k}, \end{aligned} \quad (3.6)$$

where the Hurwitz zeta-function is related by $\zeta(z, \frac{1}{2}) \equiv (2^z - 1)\zeta(z)$ to the Riemann function $\zeta(z)$. It is crucial to realize that only even powers in (m/T) are present in (3.6);

- taking care of the proper dimension of τ in (3.5) we multiply by T^{2s} ; an additional normalization factor $N(s)$, with $N(s = 0) = 1$, is allowed, which for $s \leq 0$ is,

$$N(s) = \frac{(2\pi)^{2s} \Gamma(1/2 - s) \zeta(1 - 2s, \frac{1}{2})}{\Gamma(3/2) \Gamma(-s)}. \quad (3.7)$$

After performing these steps the leading term of the function $\zeta_s|_{2\gamma}(x)$ can be written as (in Minkowski metric)

$$\zeta_s|_{2\gamma}(x) \simeq \frac{\pi N(s) \Gamma(s+2)}{(2\pi)^{2s+2} \Gamma(s)} I(y^2; s + 2, \frac{1}{2}) \hat{\sigma}^2. \quad (3.8)$$

We collect the terms in $\zeta_s|_{2\gamma}(x)$ with different powers in m , i.e. in y^2 , in order to finally perform the chirally symmetric limit.

4 Anomalous Electromagnetic Couplings

We now give the anomalous coupling for $\pi^0 \sigma \rightarrow 2\gamma$ in the chirally symmetric phase. Expanding (3.3) in the fields $(\sigma, \vec{\pi})$, e.g. $1/(m/g + \sigma) \simeq g/m - (g^2/m^2)\sigma$, it is seen, that the symmetric limit $m \rightarrow 0$ is no longer trivial. However, from (3.8) and the expansion (3.6) with respect to m^2 , one finds by adjusting the value of s to $s = -1$ that the term proportional to y^2 indeed leads to a finite nonvanishing value in the $m \rightarrow 0$ limit. In addition all the other coefficients in (3.6) for $k \neq 1$ vanish due to the pole at $s = -1$ of $\Gamma(s)$ in (3.8), from which we calculate

$$\zeta_{s=-1}(x) = -\zeta(3, \frac{1}{2}) y^2 \quad \zeta_{s=0}(x) = -\frac{7\zeta(3)}{32\pi^2} y^2 \hat{\sigma}^2. \quad (4.1)$$

Keeping the term linear in the σ field the $m \rightarrow 0$ limit is well defined, and the effective lagrangian for $\pi^0 \sigma \rightarrow 2\gamma$ finally becomes

$$\mathcal{L}_T^{2\gamma} = -\frac{7\zeta(3) g^2 \alpha N_C}{96\pi^3 T^2} \epsilon^{\mu\nu\alpha\beta} F_{\mu\nu} F_{\alpha\beta} (\pi^0 \sigma). \quad (4.2)$$

The effective photon coupling to neutral particles (π^0, σ) given in (4.2) is manifestly gauge invariant. By comparing the above coupling at high T with the well known one [4, 5] at $T = 0$ we note the substitution

$$\frac{1}{F_\pi} \rightarrow \frac{7\zeta(3)g^2}{4\pi^2 T^2} \sigma, \quad (4.3)$$

which is in agreement with the derivation given by Pisarski [2].

In a corresponding fashion the anomalous couplings linear in $F_{\mu\nu}$ for e.g. $\gamma \rightarrow 3\pi \sigma$ may be derived, which turn out to be in leading order (keeping $s = -2$) proportional to $\zeta(5) g^4/T^4$ at high temperature.

Generalizing the result (4.2) to the case of $SU(3)$ allows to obtain the radiative two-photon couplings for the processes $\eta (\eta') \sigma \rightarrow 2\gamma$ in the chirally symmetric limit. There are also additional anomalous couplings between photons and mesons with more than one σ field.

5 Conclusions

Finally, one may ask about phenomenological consequences, i.e. possible observable effects, of these high temperature anomalous couplings. An interesting case for testing (4.2) - and those related to the anomalous interactions of vector mesons [2, 3] at high T - are photon- and dilepton measurements.

In this context we mention the production of low-mass lepton pairs in heavy ion collisions at high energies [13]. In order to interpret these data, especially for dilepton masses below the ρ meson mass, a detailed knowledge of the electromagnetic interactions of $\pi^0, \eta, \eta', \omega, \rho$, etc. in a high density hadronic environment (i.e. maybe in a chirally symmetric phase at high T) is important.

Anomalous couplings of interacting bosons at high temperatures, e.g. for $KK \rightarrow \pi\pi\pi\sigma$, have been derived [1, 2, 3]. However, we do not find a simple compact expression, comparable to the Wess-Zumino-Witten action [14] at $T = 0$, except that all these couplings vanish in the $T \rightarrow \infty$ limit.

Acknowledgments

We like to thank R. D. Pisarski for useful comments and discussions. R. B. thanks the organizers H. M. Fried and B. Müller, and M. Le Bellac for inviting him to this Workshop. Partial support by the EEC Programme "Human Capital and Mobility", Network "Physics at High Energy Colliders", Contract CHRX-CT93-0357 is acknowledged. M. D. is supported by Deutsche Forschungsgemeinschaft.

References

- [1] R. Baier, M. Dirks, and O. Kober, *Phys. Rev. D* **54**, 2222 (1996).
- [2] R. D. Pisarski, in *From Thermal Field Theory to Neural Networks: a Day to Remember Tanguy Altherr*, eds. P. Aurenche et al. (World Scientific Publishing, 1996), p. 41.
- [3] R. D. Pisarski, *Phys. Rev. Lett.* **76**, 3084 (1996).
- [4] S. L. Adler, *Phys. Rev.* **177**, 2426 (1969); J. S. Bell and R. Jackiw, *Nuovo Cimento* **60**, 147 (1969).
- [5] R. A. Bertlmann, *Anomalies in Quantum Field Theory* (Clarendon Press, Oxford, 1996), and references therein.
- [6] L. Dolan and R. Jackiw, *Phys. Rev. D* **9**, 3320 (1974); H. Itoyama and A. H. Mueller, *Nucl. Phys. B* **218**, 349 (1983); and references in [1-3].
- [7] J. F. Donoghue, E. Golowich, and B. R. Holstein, *Dynamics of the Standard Model* (Cambridge Univ. Press, Cambridge, 1992).
- [8] A. Manohar and G. Moore, *Nucl. Phys. B* **243**, 55 (1984).
- [9] For a recent discussion of the QCD phase diagram and applications of chiral symmetry at high temperature, see R. D. Pisarski, lectures presented at "Workshop on Finite Temperature QCD and Quark-Gluon Transport Theory", Wuhan, PRC (April 1994).
- [10] K. Fujikawa, *Phys. Rev. D* **21**, 2848 (1980); *Phys. Rev. D* **29**, 285 (1984).
- [11] N. K. Nielsen, M. T. Grisaru, H. Römer, and P. van Nieuwenhuizen, *Nucl. Phys. B* **140**, 477 (1978); and references therein.
- [12] N. P. Landsman, *Nucl. Phys. B* **322**, 498 (1989).
- [13] CERES Collaboration, G. Agakichiev et al., *Phys. Rev. Lett.* **75**, 1272 (1995).
- [14] J. Wess and B. Zumino, *Phys. Lett. B* **37**, 95 (1971); E. Witten, *Nucl. Phys. B* **223**, 422 (1983).