

OHSTPY-HEP-T-96-021
 hep-ph/9608442
 August 1996

THERMODYNAMICS OF QCD AT HIGH TEMPERATURE ^a

A. NIETO

Department of Physics, The Ohio State University, Columbus, OH 43210, USA

A hierarchy of effective field theories is used to separate the contributions from different momentum scales and to calculate the free energy of QCD at high temperature in powers of the coupling constant up to order g^5 . The behavior of the perturbative series will also be discussed.

The partition function of QCD at finite temperature is obtained from the Lagrangian $\mathcal{L}_{\text{QCD}} = (1/4)G_{\mu\nu}^a G_{\mu\nu}^a + \bar{q}\gamma_\mu D_\mu q$, with gauge coupling constant g . We consider the effective theory that results from integrating out the nonstatic modes. Such a theory, which is called Electrostatic QCD (EQCD), is made out of only the static bosonic modes and 3 dimensional. The free energy density is $F_{\text{QCD}} = T(f_E - \log \mathcal{Z}_{\text{EQCD}}/V)$, where $\mathcal{Z}_{\text{EQCD}}$ is the partition function of EQCD obtained from the Lagrangian

$$\mathcal{L}_{\text{EQCD}} = \frac{1}{4}G_{ij}^a G_{ij}^a + \frac{1}{2}(D_i A_0)^2 + \frac{1}{2}m_E^2 A_0^2 + \frac{1}{8}\lambda A_0^4 + \delta\mathcal{L}_{\text{EQCD}} \quad (1)$$

with effective gauge coupling constant g_E . $\delta\mathcal{L}_{\text{EQCD}}$ represents an infinite series of non-renormalizable terms. The effects of the fermions are incorporated into the effective parameters. By power counting, the effective parameters at leading order are: $f_E \sim T^3$, $m_E^2 \sim g^2 T^2$, $g_E^2 \sim g^2 T$, and $\lambda_E \sim g^4 T$.

The magnetostatic fields remain massless; therefore, we can go further in separating the different scales of QCD at high temperature by integrating out A_0 . We obtain an effective theory of EQCD which is called magnetostatic QCD (MQCD). The free energy density of QCD can be written as $F = T(f_E + f_M + f_G)$, where $f_G = -\log \mathcal{Z}_{\text{MQCD}}/V$, and $\mathcal{Z}_{\text{MQCD}}$ is the partition function of MQCD obtained from the Lagrangian $\mathcal{L}_{\text{MQCD}} = (1/4)G_{ij}^a G_{ij}^a + \delta\mathcal{L}_{\text{MQCD}}$. The gauge coupling constant is g_M and $\delta\mathcal{L}_{\text{MQCD}}$ represents an infinite series of non-renormalizable terms.

Again, we can use power counting to identify the order of the leading contribution to the MQCD parameters: $f_M \sim m_E^3 \sim g^3 T^3$ and $g_M^2 \sim g_E \sim g^2 T$. λ_E contributes to the free energy only at order g^6 ; so, if we are interested in the

^a Invited talk presented at DPF96 in Minneapolis, August 1996.

free energy at lower order, we can ignore λ_E . Similarly the non-renormalizable terms of EQCD can also be omitted.

The leading contribution to f_G can be obtained by realizing that the only parameter with dimensions involved in MQCD is g_M and therefore the leading contribution is of order $g_M^6 \sim g^6 T^3$. Since we are interested in computing the free energy of QCD up to order g^5 we can ignore the contribution from f_G ; it will be enough to compute f_E and f_M . We conclude that the free energy of QCD up to order g^5 is given by $F = T[f_E(T, g; \Lambda_E) + f_M(m_E^2, g_E; \Lambda_E)]$, where Λ_E is a factorization scale that separates the scales T and gT . Therefore, we have to determine f_E , m_E^2 , g_E , and f_M . Calculations for $SU(N)$ gauge theory with n_f fermions and results in analytical form are detailed in²; here, I will just state the results in numerical form for $SU(3)$ with 3 fermions.

The effective mass m_E is computed by matching the electrostatic screening mass for QCD and EQCD.

$$m_E^2 = 0.93 \frac{g^2(\mu)}{4\pi} T^2 \left[1 + \left(-0.26 + 1.43 \log \frac{\mu}{2\pi T} \right) \frac{g^2}{4\pi} \right]. \quad (2)$$

At this order in g^2 , there is no dependence on the factorization scale Λ_E .

The parameter f_E is determined by calculating the free energy in both full QCD and EQCD, and matching the two results.

$$\begin{aligned} f_E(\Lambda_E) = & -14.8 T^3 \left[1 - 0.90 \frac{g^2}{4\pi} + \right. \\ & \left. + \left(18.3 + 6.91 \log \frac{\Lambda_E}{2\pi T} - 1.30 \log \frac{\mu}{2\pi T} \right) \left(\frac{g^2}{4\pi} \right)^2 \right], \quad (3) \end{aligned}$$

where g is the coupling constant in the $\overline{\text{MS}}$ renormalization scheme at the scale μ . We have used the renormalization group equation of the coupling constant to shift the scale of the running coupling constant to an arbitrary renormalization scale μ .

Through order g^5 , f_M is proportional to the logarithm of the partition function for EQCD: $f_M = -\log \mathcal{Z}_{\text{EQCD}}/V$. Now, we have to consider the contribution to $\log \mathcal{Z}_{\text{EQCD}}$ of orders g^3 , g^4 , and g^5 which are given by the sum of 1-loop, 2-loop, and 3-loop diagrams. The details of this calculation can be found in²; the final result is

$$f_M(\Lambda_E) = -0.21 m_E^3 \left[1 - \left(0.54 + 0.72 \log \frac{\Lambda_E}{2m_E} \right) \left(\frac{g_E^2}{m_E} \right) - 0.70 \left(\frac{g_E^2}{m_E} \right)^2 \right]. \quad (4)$$

The coefficient f_M in (4) can be expanded in powers of g by setting $g_E^2 = g^2 T$ and by substituting the expression (2) for m_E^2 . The complete free energy to order g^5 is then $F = (f_E + f_M)T$; in agreement with the result obtained independently by Kastening and Zhai⁴. Note that the dependence on the arbitrary factorization scale Λ_E cancels between f_E and f_M , up to corrections that are higher order in g , leaving a logarithm of T/m_E .

We have calculated the free energy as a perturbation expansion in powers of g to order g^5 . We now ask how small $\alpha_s \equiv g^2/(4\pi)$ must be in order for the perturbation expansion to be well-behaved^{2,3}. If the series is apparently convergent, then it can plausibly be used to evaluate the free energy. For simplicity, we consider the case $n_f = 3$ and choose the renormalization scale $\mu = 2\pi T$ which is the mass of the lightest nonstatic mode. The correction to the leading order term of the free energy is a multiplicative factor $1 - 0.9\alpha_s + 3.3\alpha_s^{3/2} + (7.1 + 3.5 \log \alpha_s)\alpha_s^2 - 20.8\alpha_s^{5/2}$. If the temperature is a few times above the transition temperature (~ 200 MeV); *e.g.*, $T \simeq 350$ MeV the series has the form $1 - 0.27 + 0.54 + 0.26 - 1.03 + \dots$. It is clear that the perturbative series is not convergent at *this* temperature.

f_E and m_E contain the contributions of order T . In the case of m_E^2 , which is given by (2), the correction to the leading order result from the series $1 - 0.26\alpha_s$. The next-to-leading-order correction to m_E^2 is smaller than the leading-order term only if $T > 40$ MeV. A similar analysis of f_E gives $T > 67$ MeV. We conclude that the perturbation series for the parameters of EQCD are well-behaved provided that $T > 70$ MeV.

Expression (4) gives f_M which is the contribution from the scale T . The next-to-leading-order correction is small if Λ_E is chosen to be approximately m_E . The next-to-next-to-leading-order correction is smaller than the leading order term only if $T > 2$ GeV. Thus the perturbation series for f_M is well-behaved only for temperatures that are much higher than those required for the parameters of EQCD to have well-behaved perturbation series.

This analysis indicates that the slow convergence of the expansion for F in powers of $\sqrt{\alpha_s}$ can be attributed to the slow convergence of perturbation theory at the scale gT .

Acknowledgments

This work was supported in part by the U.S. Department of Energy, Division of High Energy Physics, under Grant DE-FG02-91-ER40690. I would like to thank E. Braaten for valuable discussions.

References

1. E. Braaten and A. Nieto, *Phys. Rev. D* **51**, 6990 (1995).
2. E. Braaten and A. Nieto, *Phys. Rev. D* **53**, 3421 (1996).
3. E. Braaten and A. Nieto, *Phys. Rev. Lett.* **76**, 1417 (1996).
4. B. Kastening and C. Zhai, *Phys. Rev. D* **52**, 7232 (1995).