

BARYONIC FORM FACTORS IN THE NONRELATIVISTIC QUARK MODEL

HAI-YANG CHENG

Institute of Physics, Academia Sinica, Taipei, Taiwan 115, Republic of China

Weak current-induced baryonic form factors at zero recoil are evaluated in the rest frame of the heavy parent baryon using the nonrelativistic quark model. Contrary to previous similar work in the literature, our quark model results do satisfy the constraints imposed by heavy quark symmetry for heavy-heavy baryon transitions. Assuming a dipole q^2 behavior, we have applied the quark model form factors to nonleptonic, semileptonic and weak radiative decays of the heavy baryons.

The general expression for the $\frac{1}{2}^+ \rightarrow \frac{1}{2}^+$ baryonic transition $B_i \rightarrow B_f$ reads

$$\begin{aligned} \langle B_f(p_f) | V_\mu - A_\mu | B_i(p_i) \rangle = & \bar{u}_f [f_1(q^2)\gamma_\mu + i f_2(q^2)\sigma_{\mu\nu}q^\nu + f_3(q^2)q_\mu \\ & - (g_1(q^2)\gamma_\mu + i g_2(q^2)\sigma_{\mu\nu}q^\nu + g_3(q^2)q_\mu)\gamma_5] u_i, \end{aligned} \quad (1)$$

where $q = p_i - p_f$. In the heavy-quark limit, the form factors f_i and g_i are related to three baryonic Isgur-Wise functions: $\zeta(\omega)$ for antitriplet-antitriplet transition, and $\xi_1(\omega)$, $\xi_2(\omega)$ for sextet-sextet transition. In general, it is difficult to estimate the $1/m_Q$ corrections to hadronic form factors in the heavy quark effective theory (HQET). However, a tremendous simplification occurs in the antitriplet-antitriplet heavy baryon transition, e.g., $\Lambda_b \rightarrow \Lambda_c$: $1/m_Q$ corrections only amount to renormalizing the function $\zeta(\omega)$ and no further new function is needed¹. This simplification stems from the fact that the chromomagnetic operator does not contribute to $\Lambda_b \rightarrow \Lambda_c$ and that the diquark of the antitriplet heavy baryon is a spin singlet.

Going beyond the antitriplet-antitriplet heavy baryon transition, the predictive power of HQET for form factors at order $1/m_Q$ is lost owing to the fact that $1/m_Q$ corrections due to wave function modifications arising from O_1 and especially O_2 are not calculable by perturbative QCD. Therefore, it is appealing to have model calculations which enable us to estimate the $1/m_Q$ corrections for other baryon form factors. To our knowledge, two different quark-model calculations^{2,3} are available in the literature. Unfortunately, none of the calculations presented in^{2,3} is in agreement with the predictions of HQET. For example, several heavy quark symmetry relations between baryon form factors are not obeyed in². While this discrepancy is resolved in³, its prediction for $\Lambda_b \rightarrow \Lambda_c$ (or $\Xi_b \rightarrow \Xi_c$) form factors at order $1/m_Q$ is still too large by a factor of 2 when compared with HQET.

In ⁴ we have shown that our prescription of quark model calculations does incorporate the features of heavy quark symmetry and hence can be applied to compute baryon form factors beyond the arena of HQET. For example, for $\Lambda_b \rightarrow \Lambda_c$ transition we obtain

$$\begin{aligned} f_1^{\Lambda_b \Lambda_c}(q_m^2) = g_1^{\Lambda_b \Lambda_c}(q_m^2) &= 1 + \frac{\Delta m \bar{\Lambda}}{4} \left(\frac{1}{m_{\Lambda_b} m_c} - \frac{1}{m_{\Lambda_c} m_b} \right), \\ f_2^{\Lambda_b \Lambda_c}(q_m^2) = g_3^{\Lambda_b \Lambda_c}(q_m^2) &= -\frac{\bar{\Lambda}}{4} \left(\frac{1}{m_{\Lambda_b} m_c} + \frac{1}{m_{\Lambda_c} m_b} \right), \\ f_3^{\Lambda_b \Lambda_c}(q_m^2) = g_2^{\Lambda_b \Lambda_c}(q_m^2) &= -\frac{\bar{\Lambda}}{4} \left(\frac{1}{m_{\Lambda_b} m_c} - \frac{1}{m_{\Lambda_c} m_b} \right), \end{aligned} \quad (2)$$

at zero recoil, where $q_m = (m_{\Lambda_b} - m_{\Lambda_c})^2$, and $\bar{\Lambda} = m_{\Lambda_c} - m_c$. The above results agree with the HQET predictions up to the zeroth order of α_s .

Two remarks are in order. First, the baryonic form factor is proportional to a flavor factor $N_{fi} = \langle B_f | b_q^\dagger b_Q | B_i \rangle_{\text{flavor}}$, which is equal to unity for heavy-to-heavy transition, but less than unity for heavy-to-light transition. For example, $N_{\Lambda_c \Lambda} = 1/\sqrt{3}$. In the literature it is customary to replace the s quark in the baryon Λ by the heavy quark Q to obtain the wave function of the Λ_Q . However, this amounts to assuming SU(4) or SU(5) flavor symmetry. Since SU(N)-flavor symmetry with $N > 3$ is badly broken, the flavor factor $N_{\Lambda_Q \Lambda}$ is no longer unity. Indeed, if $N_{\Lambda_c \Lambda}$ were equal to one, the predicted rate for $\Lambda_c \rightarrow \Lambda e^+ \nu_e$ would have been too large by at least a factor of 2 ! Second, as the conventional practice, we make the pole dominance assumption for the q^2 dependence to extrapolate the form factors from maximum q^2 to the desired q^2 point. We argued that a dipole q^2 behavior is more preferred since it is close to the baryonic Isgur-Wise function calculated recently. Nevertheless, one should bear in mind that the assumption of pole dominance for form factors is probably too simplified and this problem remains unresolved.

We have applied the quark model form factors to nonleptonic, semileptonic and weak radiative decays of the heavy baryons ⁴. In the heavy c -quark limit, there are two independent form factors in $\Lambda_c \rightarrow \Lambda$ transition

$$\langle \Lambda(p) | \bar{s} \gamma_\mu (1 - \gamma_5) c | \Lambda_c(v) \rangle = \bar{u}_\Lambda \left(F_1^{\Lambda_c \Lambda}(v \cdot p) + \not{v} F_2^{\Lambda_c \Lambda}(v \cdot p) \right) \gamma_\mu (1 - \gamma_5) u_{\Lambda_c}. \quad (3)$$

We found that

$$R \equiv F_1^{\Lambda_c \Lambda} / F_2^{\Lambda_c \Lambda} = -1 / \left(1 + 4 \frac{m_s}{\bar{\Lambda}} \right) = -0.23, \quad (4)$$

with $\bar{\Lambda} = m_{\Lambda_c} - m_s$, in accord with the CLEO result ⁵ $R = -0.25 \pm 0.14 \pm 0.08$. For the semileptonic decay $\Lambda_c \rightarrow \Lambda e^+ \nu_e$, the decay asymmetry parameter is

obtained to be $\langle\alpha\rangle = -0.84$, in good agreement with experiment ⁶ $\langle\alpha\rangle = -0.82^{+0.11}_{-0.07}$. The decay rate is predicted to be

$$\Gamma(\Lambda_c \rightarrow \Lambda e^+ \nu_e) = (N_{\Lambda_c \Lambda})^2 \times 2.11 \times 10^{11} s^{-1} = 7.1 \times 10^{10} s^{-1}, \quad (5)$$

while experimentally ⁶ $\Gamma(\Lambda_c \rightarrow \Lambda e^+ \nu_e)_{\text{expt}} = (11.2 \pm 2.4) \times 10^{10} s^{-1}$. The presence of the flavor suppression factor $N_{\Lambda_c \Lambda}$, which is missed in most literature, will of course affect the predictions on the decay rates of many decay modes involving a transition from heavy to light baryons.

Generally it is very difficult to tackle the nonleptonic weak decays of the baryons because of the presence of nonspectator W exchange effects manifested as pole contributions. Nevertheless, there are two decay modes of great interest, namely $\Lambda_c \rightarrow p\phi$ and $\Lambda_b \rightarrow \Lambda J/\psi$ which are theoretically very clean in the sense that they proceed only through the internal W -emission diagram. Noting $N_{\Lambda_c p} = \frac{1}{\sqrt{2}}$ and $N_{\Lambda_b \Lambda} = \frac{1}{\sqrt{3}}$ and using the factorization approach, we found $Br(\Lambda_c \rightarrow p\phi) = 7.1 \times 10^{-4}$, which agrees with the experimental result ⁶ $(1.06 \pm 0.33) \times 10^{-3}$. The branching ratio of $\Lambda_b \rightarrow \Lambda J/\psi$ is predicted to be 2.1×10^{-4} , which is two orders of magnitude smaller than the UA1 observation ⁷ but consistent with the most recent CDF measurement ⁸ $Br(\Lambda_b \rightarrow \Lambda J/\psi) = (4.2 \pm 1.8 \pm 0.7) \times 10^{-4}$.

Acknowledgments

This work was done in collaboration with B. Tseng and supported in part by the National Science Council of ROC under Contract No. NSC85-2112-M-001-010.

References

1. H. Georgi, B. Grinstein, and M.B. Wise, *Phys. Lett.* **B252**, 456 (1990).
2. R. Pérez-Marcial, R. Huerta, A. Garcia, and M. Avila-Aoki, *Phys. Rev.* **D40**, 2955 (1989); *ibid.* **D44**, 2203(E) (1991).
3. R. Singleton, *Phys. Rev.* **D43**, 2939 (1991).
4. H.Y. Cheng and B. Tseng, *Phys. Rev.* **D53**, 1457 (1996).
5. CLEO Collaboration, G. Crawford *et al.*, *Phys. Rev. Lett.* **75**, 624 (1995).
6. Particle Data Group, *Phys. Rev.* **D54**, 1 (1996).
7. UA1 Collaboration, C. Albarjar *et al.*, *Phys. Lett.* **B273**, 540 (1991).
8. P. Chang, private communication.