

Durham DTP/96/90
MPI-PhT/96-92
September 28th, 1996

QCD coherence and the soft limit of the energy spectrum

VALERY A. KHOZE ^{a,b,1} , SERGIO LUPIA ^{c,2} , WOLFGANG OCHS ^{c,3}

^a *Department of Physics
University of Durham, Durham DH1 3LE, UK*

^b *Institute for Nuclear Physics
St. Petersburg, Gatchina, 188350, Russia*

^c *Max-Planck-Institut für Physik
(Werner-Heisenberg-Institut)
Föhringer Ring 6, D-80805 Munich, Germany*

Abstract

The analytical perturbative approach, if taken to the limit of its applicability, allows one to predict an energy independent limit for the one-particle invariant density in QCD jets $E \frac{dn}{d^3p}$ at very small momenta p . This is a direct consequence of the colour coherence in soft gluon branching. The existing data on the charged and identified particle inclusive spectra follow this prediction surprisingly well. Further tests of the perturbatively based picture in the soft region are discussed.

¹e-mail: v.a.khoze@durham.ac.uk

²e-mail: lupia@mppmu.mpg.de

³e-mail: wwo@mppmu.mpg.de

1. Introduction

One of the most striking predictions of the perturbative approach to QCD jet physics [1,2] is the depletion of the soft particle production and the resulting approximately Gaussian shape of the inclusive distribution in the variable $\xi = \log E_{jet}/E$ for particles with energy E in a jet of energy E_{jet} (the so-called “hump-backed plateau”)[3,4]. Due to the coherence of the gluon radiation it is not the softest partons but those with intermediate energies ($E \sim E_{jet}^{0.3-0.4}$) which multiply most effectively in QCD cascades. We recall that the gluons of long wave length are emitted by the total colour current which is independent of the internal structure of the jet and is conserved when the partons split. Applying the hypothesis of the Local Parton Hadron duality (LPHD)[5], one then comes to the conclusion that the hadron spectrum at low momentum p should be nearly independent of the jet energy E_{jet} [5].

It is a very important question to understand whether there is a smooth transition between the purely perturbative regime and the soft momentum region. It looks quite intriguing that the shape of the measured charged particle energy spectra in e^+e^- annihilation appears to be surprisingly close over the whole momentum range (down to small momenta of a few hundred MeV) to the perturbative predictions based on the Modified Leading Log Approximation (MLLA)[2,6,7,8]. Moreover, the momentum spectra in the variable $\log p$, measured by the TASSO [9] and OPAL Collaborations [10] provided first evidence in favour of the scaling behaviour at small momenta: as discussed in [11] these data could be considered as a confirmation of the basic ideas of QCD coherence and LPHD. A further confirmation of this picture has been provided recently in the study of the invariant particle density $E dn/d^3p$ at low momenta and also in the low cms energy region. The spectra are found to be in a good agreement with the scaling behaviour and with the analytical perturbative expectations which become sensitive to the running of the coupling α_s at small scales [12].

In this paper we focus on the soft limit of the parton spectra, and how its scaling behaviour arises in the analytical calculations for the coherent parton cascade. We consider the invariant density $E dn/d^3p \equiv dn/dy d^2p_T$ in the limit of vanishing rapidity y and transverse momentum p_T or, equivalently, for vanishing momentum $|\vec{p}| \equiv p$, i.e.,

$$I_0 = \lim_{y \rightarrow 0, p_T \rightarrow 0} E \frac{dn}{d^3p} = \frac{1}{2} \lim_{p \rightarrow 0} E \frac{dn}{d^3p} \quad (1)$$

where the factor 1/2 takes into account that both hemisphere are added in the limit $p \rightarrow 0$. If the dual description of hadronic and partonic final states is really adequate down to very small momenta, the finite, energy independent limit of the invariant hadronic density I_0 is expected.

This will be examined for both charged and identified hadrons in the full energy range explored so far in e^+e^- annihilation. A possible growth of the spectrum in the soft region could indicate that either coherence or the local duality (or both) breaks down. Since colour coherence is a general property of QCD as a gauge theory, it is the LPHD concept that is tested in the measurements of the soft hadron distributions. Therefore, such studies could provide one with additional information on the physics of confinement.

The scaling properties of the invariant cross section or density have been discussed in the early seventies in the context of the parton model [13] and Feynman-scaling [14] (see

e.g. [15]). Note that scaling in the parton model refers to the p_T -integrated spectra. In the QCD bremsstrahlung picture of multiparticle production discussed here the p_T -integrated central rapidity plateau rises with cms energy, and scaling behaviour is retained only in the large wavelength limit (1).

It is important to test further the QCD origin of this scaling behaviour. This can be accomplished by studying the soft radiation from the emitters (“parton antennae”) corresponding to different effective colour charges. The good scaling property of the density I_0 in e^+e^- annihilation suggests taking this number as a standard scale in the comparison with other processes. Some of the results of this paper and further details have also been presented elsewhere [17].

2. Particle production in the soft limit in DLA and MLLA

DLA predictions.

We consider first the analytical predictions for the energy spectrum of partons near the soft limit. The asymptotic behaviour of the energy spectrum is obtained from the Double Log Approximation (DLA) in which energy conservation is neglected and only the leading singularities in the parton splitting functions are kept. The evolution equation of the inclusive energy distribution of partons p originating from a primary parton A is given by[3]:

$$D_A^p(\xi, Y) = \delta_A^p \delta(\xi) + \int_0^\xi d\xi' \int_0^{Y-\xi} dy' \frac{C_A}{N_C} \gamma_0^2(y') D_g^p(\xi', y'). \quad (2)$$

Here we have used the logarithmic variables $\xi = \log(1/x) = \log(Q/E)$ and $Y = \log(Q/Q_0)$ with E the particle energy and Q the jet virtuality ($Q = P\Theta$ for a jet of primary momentum P and half opening angle Θ); C_A is the respective colour factor, i.e., $C_g = N_C$ and $C_q = C_F$; γ_0 denotes the anomalous dimension of multiplicity and is related to the QCD running coupling by $\gamma_0^2 = 4N_C\alpha_s/2\pi$ or $\gamma_0^2 = \beta^2/\log(p_T/\Lambda)$ with $\beta^2 = 4N_C/b$, $b \equiv (11N_c - 2n_f)/3$; Λ is the QCD-scale and N_C and n_f are the number of colours and of flavours respectively. The shower evolution is cut off by Q_0 , such that the transverse momentum $p_T \geq Q_0$.

As can be seen from (2) the ξ -spectrum vanishes in the soft limit $\xi \rightarrow Y$ ($E \rightarrow Q_0$). The behaviour near this limit can be found by the iterative solution of (2). With two iterations one obtains

$$D_A^g(\xi, Y) = \delta_A^g \delta(\xi) + \frac{C_A}{N_C} \beta^2 \log \left(1 + \frac{Y - \xi}{\lambda} \right) \left[1 + \frac{\beta^2 \int_0^{Y-\xi} d\tau \log(1 + \frac{\tau}{\lambda}) \log(1 + \frac{\xi}{\tau+\lambda})}{\log(1 + \frac{Y-\xi}{\lambda})} \right] + \dots \quad (3)$$

where $\lambda = \log Q_0/\Lambda$. The second term of order β^2 corresponds to the emission of a single gluon and yields the leading contribution for $E \rightarrow Q_0$. It is proportional to the colour charge factor of the primary parton. Furthermore, this term does not depend on the cms energy, contrary to the higher order terms which provide the rise of the spectrum for large E with increasing \sqrt{s} . The spectrum vanishes at $E \rightarrow Q_0$ as

$$D_A^g(\xi, Y) \sim Y - \xi \sim \log E/Q_0 \sim E - Q_0. \quad (4)$$

An exact solution is known for fixed α_s [3] with similar behaviour near the boundaries as above in (3) and (4).

MLLA predictions.

A considerable improvement in accuracy for nonasymptotic energies is obtained in the application of the MLLA [6,7,2] which takes into account systematically all the corrections of relative order $\sqrt{\alpha_s}$ based on the evolution equation which includes energy conservation and the exact form of the parton splitting functions. The leading high energy MLLA contribution obeys a differential equation (see also [2,12], with $D(\xi, Y, \lambda) \equiv D_g^g$)

$$\frac{\partial^2 D(\xi, Y, \lambda)}{\partial \xi \partial Y} + \frac{\partial^2 D(\xi, Y, \lambda)}{dY^2} - \gamma_0^2 D(\xi, Y, \lambda) = -a \left(\frac{\partial}{\partial \xi} + \frac{\partial}{\partial Y} \right) \left(\frac{\alpha_s(Y + \lambda)}{2\pi} D(\xi, Y, \lambda) \right) \quad (5)$$

where $a = \frac{11}{3}N_C + \frac{2n_f}{3N_C^2}$. The limit $a = 0$ corresponds to the DLA as in (2). In case of fixed α_s the exact MLLA solution can be obtained [12] (see also [6]) by multiplying the DLA result with the factor $\exp(-a\gamma_0^2(Y - \xi)/4N_C)$. Similarly, for running α_s one can make an ansatz

$$D(\xi, Y, \lambda) = D(\xi, Y, \lambda)|_{DLA} \exp \left[-a \int_{\xi}^Y \gamma_0^2(y)/(4N_C) dy \right] \quad (6)$$

which solves Eq. (5) approximately in the region around $\xi \sim Y$.⁴ Note that the exponent in (6) can be written as $-\frac{a}{b} \log \left(\frac{Y+\lambda}{\xi+\lambda} \right)$ and behaves near the boundary like $-\frac{a}{b} \frac{Y-\xi}{Y+\lambda}$, so the slope of the distribution varies with energy like the coupling $\alpha_s \sim \gamma_0^2 = 4N_C/[b(Y + \lambda)]$ contrary to the case with fixed coupling.

Phase-space effects in the soft region.

While the limiting behaviour of the partonic energy spectrum in the soft region follows from the general principle of colour coherence the detailed form of the observable hadronic spectrum is predicted uniquely from the LPHD hypothesis only for $E \simeq p \gg Q_0$, but not near the kinematical boundary because of the sensitivity of the spectrum to the cut-off procedure. In the MLLA the partons are treated as massless with energy $E = p \geq p_T \geq Q_0$, so $\xi \leq Y$. Experimental hadronic spectra are usually presented as function of momenta p or $\xi_p = \log(1/x_p)$ which is not limited from above. The same kinematic limit for partons and hadrons is obtained if the hadronic mass m_h and the partonic p_T cut-off Q_0 are taken the same.

For the relation between parton and hadron distributions one may require that the invariant density $E dn/d^3p$ of hadrons approaches a constant limit for $p \rightarrow 0$ as is observed experimentally. For the spectra which vanish linearly as in (4) this is achieved by relating the hadron and parton spectra in a single A -jet as [3,12]

$$E_h \frac{dn(\xi_E)}{dp_h} = K_h E_p \frac{dn(\xi_E)}{dp_p} \equiv K_h D_A^g(\xi_E, Y) \quad (7)$$

with $E_h = \sqrt{p_h^2 + Q_0^2} = E_p \equiv E \geq Q_0$ and $\xi_E \equiv \xi = \log Q/E$, where K_h is a normalization parameter. If hadrons from both hemispheres are added, K_h should be replaced by $2K_h$.

⁴When eq. (6) is inserted into eq. (5) all terms cancel except for the small difference $(\partial D_{DLA}/\partial Y) \exp[-a \int_{\xi}^Y \gamma_0^2(y)/(4N_C) dy] [\frac{a\alpha_s(\xi)}{2\pi} - \frac{a\alpha_s(Y)}{2\pi}]$, which vanishes at $\xi = Y$. Exponential damping factors of this type are known to take into account single log corrections[16].

Then, indeed, for hadrons the invariant density $Edn/d^3p = K_h D_A^g(\xi, Y)/4\pi(E_h^2 - Q_0^2)$ approaches the finite limit as in (1)

$$I_0 = K_h \frac{C_A \beta^2}{8\pi N_C \lambda Q_0^2}. \quad (8)$$

In the fixed α_s limit β^2/λ is replaced by γ_0 and $I_0 \sim 1/Q_0^2$. With prescription (7) the moments of the full energy spectrum $D(\xi, Y)$ are well described by the MLLA formulae and with $Q_0 = 270$ MeV in a wide energy range [12].

The relation (7) is not unique, however. We found that the alternative prescription based on phase space arguments, $dn/d\xi_p \simeq (p/E)^3 D(\xi_E, Y)$ (see e.g. [18,20]):

$$E_h \frac{dn}{d^3p_h} = K_h \left(\frac{1}{4\pi E^2} \right) D_A^g(\xi_E, Y) \quad (9)$$

works well in e^+e^- annihilation for charged pions ($K_h = K_\pi$) in the whole energy region and for charged particles ($K_h = K_{ch}$) in the low energy region. In this application, including both hemispheres, D_A^g in eq. (9) is replaced by $2 \cdot 4/9 D_g^g$, using for D_g^g the MLLA limiting spectrum [5,18] with $Q_0 = \Lambda = 138$ MeV. The low cut-off mass is plausible in this region, which is dominated by pions.

Behaviour of analytical results

In order to illustrate the above analytical results we compare in Fig. 1 the various approximations (in case of running α_s) for low particle energies using the relation (7) between parton and hadron spectra: the DLA result in the approximation (3) with and without MLLA exponential factor (6) for $\lambda = 0.01$ and also the “limiting spectrum”, which solves the MLLA equations for $\lambda = 0$ in the full kinematic region, except very close to the boundary $\xi = Y$ where it stays finite. The normalization of the limiting spectrum is as in fits to e^+e^- annihilation; the normalizations of the DLA and MLLA curves are chosen to approach the limiting spectrum for energies $E \geq 0.5$ GeV.

These results show that in all cases considered, DLA and MLLA for both fixed and running α_s , the single particle invariant density approaches an energy independent value in the soft limit $\xi \rightarrow Y$ ($E \rightarrow Q_0$). This originates from the soft gluon emission contribution of order α_s which is determined by the total colour charge of the primary partons due to the colour coherence. In this limit the MLLA converges towards the DLA as the energy conservation constraints are unimportant and the parton splitting functions are only probed for very small fractional momenta.

3. Discussion of experimental results

Inclusive charged particle spectra

Fig. 2 shows the charged particle invariant density in e^+e^- annihilation, Edn/d^3p , as a function of the particle energy E at different *cms* energies ranging from 3 GeV up to LEP-1.5 *cms* energy (133 GeV)[9,10,22,23]. The effective mass value of $Q_0 = 270$ MeV has been used in the kinematical relations. As shown in [12] this allows a reasonably good description of the moments of the energy spectra by the MLLA limiting formulae over a large energy interval. A fit using eq. (9) is shown elsewhere (see last reference in [17]).

It is remarkable that the data from all *cms* energies tend to converge in the soft limit: we find $2I_0 \simeq 6\text{--}8 \text{ GeV}^{-2}$ (using $Q_0 = 270 \text{ MeV}$) and $2I_0 \simeq 4\text{--}6 \text{ GeV}^{-2}$ (using $Q_0 = 138 \text{ MeV}$). Inspecting the soft limit more closely, the LEP data seem to tend to a limiting value larger by about 20% as compared to the lower energy data. This may be well due to the overall systematic effect in the relative normalization of the different experiments. In this respect, it is instructive to recall that both TASSO and the LEP Collaborations have collected their data at different *cms* energies with the same accelerators and detectors, thus avoiding within their data samples any problems due to the relative normalization; in these samples the scaling works well.

A possible source of scaling violations are weak decays (for example kaons, heavy quarks). Particles from such decays (from the new “antenna”) would add incoherently to the particles produced from the primary quarks and thereby could yield a rise of the soft particle spectrum with increasing energy. It would be interesting to disentangle these possibilities experimentally and to find out the size of primary scaling violation, if any.

The data also compare well with the MLLA analytical predictions, not only concerning the energy independence of the soft limit I_0 but also the considerable energy dependence of the initial slope which is largely due to the running of α_s in the MLLA damping factor (6).

Identified particles’ spectra

Let us consider now the invariant density for π , K and p . In this case, the simplest assumption is to identify with the mass scale the particle mass itself. Fig. 3 shows the invariant cross section Edn/d^3p as a function of the particle energy E for charged pions, charged kaons and protons, as derived from the inclusive momentum spectra measured at *cms* energies from 1.6 GeV to 91 GeV[23,24]. In all cases the data tend towards a common limit for $E \rightarrow m_h$ ($p \rightarrow 0$).

Note that the scale for I_0 in eq. (8) is given not by Λ^{-2} , but by Q_0^{-2} , which can be related to an effective particle mass. Indeed, I_0 shows a clear mass dependence ($2I_0^\pi = 5\text{--}8 \text{ GeV}^{-2}$, $2I_0^K = 0.3\text{--}0.4 \text{ GeV}^{-2}$, $2I_0^p = 0.08\text{--}0.10 \text{ GeV}^{-2}$). This is consistent with $I_0^h \simeq m_h^{-2}$ (after normalization to I_0^π , one would predict $I_0^K = 0.37\text{--}0.61 \text{ GeV}^{-2}$, $I_0^p = 0.10\text{--}0.17 \text{ GeV}^{-2}$). Inclusion of the logarithmic correction $1/\lambda$ in eq. (8) would, however, require rather small values of $\Lambda \simeq 100 \text{ MeV}$. The measured values of I_0^h are also in good agreement with the expectations following from eq. (9) and the finite limit at $E \rightarrow Q_0$ of the limiting spectrum. Alternatively, one can use in the fits effective scales Q_0 which deviate from the respective hadron masses; this is also suggested by the fits to the peak region of the ξ -distribution⁵. This intriguing subject certainly deserves further detailed studies.

4. Further tests of the dual description of parton and hadron cascades in e^+e^- annihilation.

It is suggestive to relate the observed behaviour of particle spectra for vanishing momenta to the expected behaviour of soft gluon emission off the primary partons. The particle rate I_0 in this limit (see eq. (8)) cannot be directly predicted, as it depends on the normalization

⁵ For example, the maximum ξ^* of the distribution does not follow the expectations ($\xi^* \simeq 1/2 \log E_{jet}/Q_0$ in DLA) with $Q_0 = m_h$ [25].

factor and the cut-off parameter Q_0 . However, the remarkable scaling behaviour of I_0 in e^+e^- annihilation provides one with a standard scale for the comparison with other processes. It would be very interesting to study whether the soft particle production is universal, i.e., a purely hadronic quantity or whether the intensity I_0 indeed depends on the colour topology of the primary active partons in the collisions process. This could provide one with a direct support of the dual description of soft particle production in terms of the QCD bremsstrahlung.

In what follows we consider some possibilities to test this hypothesis further and discuss what one may expect from the comparison of the e^+e^- reaction with other collision processes.

gg final state

A direct test of the relevance of QCD coherence for the soft production limit is the comparison of the $q\bar{q}$ with the gg colour singlet final state. In the gg final state the same line of argument applies as above for the $q\bar{q}$ final state, but the colour charge factor is increased by $N_C/C_F = 9/4$ in eq. (8) implying roughly a doubling of the soft radiation intensity I_0 . This increased radiation has been suggested originally for the global multiplicity in gluon jets[26]; in the real life experiments various effects keep the observed increase considerably below this limit (see e.g. [8]). As the soft radiation is not strongly influenced by energy-momentum constraints (see the DLA-MLLA comparison above) the effects of the different colour charges could be more pronounced in this case.

An approximate realization of a colour octet antenna is possible in $e^+e^- \rightarrow q\bar{q}g$ with the gluon recoiling against a quasi-collinear $q\bar{q}$ pair[27] (for a recent study of such events, see [28]). The production rate of soft particles in such events is expected to be increased if it really is related to the effective colour charge of the primary partons.

3-jet events

To be more quantitative, we consider the soft radiation in 3-jet $q\bar{q}g$ events of arbitrary jet orientation into a cone perpendicular to the production plane and compare it to the radiation into the same cone in a 2-jet $q\bar{q}$ event again perpendicular to the primary $q\bar{q}$ directions. The analysis of these configurations avoids the integration over the $k_T \geq Q_0$ boundaries along the jets.

We restrict ourselves to calculations of the soft gluon bremsstrahlung of order α_s . From the experience above with two-jet events this contribution dominates in the soft limit whereas higher order contributions take over with increasing particle energy. Calculations of this type have been successfully applied to the string/drag phenomena[29,30] which refer to the particle multiplicity flows projected onto the 3-jet production plane.

Let us consider first the soft radiation into arbitrary direction \vec{n} from a $q\bar{q}$ antenna pointing in directions \vec{n}_i and \vec{n}_j [30]:

$$dN_{q\bar{q}} = \frac{dp}{p} d\Omega_{\vec{n}} \frac{\alpha_s}{(2\pi)^2} W^{q\bar{q}}(\vec{n}) \quad , \quad W^{q\bar{q}}(\vec{n}) = 2C_F(\hat{i}j) \quad (10)$$

with $(\hat{i}j) = a_{ij}/(a_i a_j)$, $a_{ij} = (1 - \vec{n}_i \vec{n}_j)$ and $a_i = (1 - \vec{n} \vec{n}_i)$. Such an antenna is realized, for example, in a $q\bar{q}\gamma$ event, and can be obtained by the appropriate Lorentz boost from the $q\bar{q}$ rest frame. The soft gluon radiation in a $q\bar{q}g$ event is given as in (10) but with the

angular factor

$$W^{q\bar{q}g}(\vec{n}) = N_C[(\widehat{1+}) + (\widehat{1-}) - \frac{1}{N_C^2}(\widehat{+-})] \quad (11)$$

where $(+, -, 1)$ refer to (q, \bar{q}, g) .

For the radiation perpendicular to the primary partons $(\widehat{ij}) = a_{ij} = 1 - \cos \Theta_{ij}$, with relative angles Θ_{ij} between the primary partons i and j . For 2-jet events of arbitrary orientation one obtains in this case

$$W_{\perp}^{q\bar{q}}(\Theta_{+-}) = 2C_F(1 - \cos \Theta_{+-}). \quad (12)$$

Correspondingly the ratio R_{\perp} of the soft particle yield in 3-jet events to that of 2-jet events in their own rest frame ($W_{\perp}^{q\bar{q}}(\pi) = 4C_F$) is given by

$$R_{\perp} \equiv \frac{dN_{\perp}^{q\bar{q}g}}{dN_{\perp}^{q\bar{q}}} = \frac{N_C}{4C_F} [2 - \cos \Theta_{1+} - \cos \Theta_{1-} - \frac{1}{N_C^2}(1 - \cos \Theta_{+-})] \quad (13)$$

It is also interesting to note the difference of this prediction to the large N_C approximation in which the $q\bar{q}g$ event is treated as a superposition of two $q\bar{q}$ dipoles (see, e.g., [29]). In this case the last term in eq. (13) drops out and $C_F = (N_C^2 - 1)/2N_C \simeq N_C/2$. This yields

$$R_{\perp} \simeq \frac{1}{2} [2 - \cos \Theta_{1+} - \cos \Theta_{1-}] \quad (\text{large } N_C) \quad (14)$$

Predictions from these formulae for R_{\perp} are presented in Table (1) for various relative angles Θ_{ij} . Note, in particular, the limiting cases $R_{\perp} = 1$ for soft or collinear primary gluon emission and the proper gg limit for the parallel $q\bar{q}$ ($\Theta_{+-} = 0$) configuration, as expected.

The role of the large- N_C limit can be investigated also by studying the production rate in 3-jet events normalized to the sum of rates from the corresponding 2-jet events (dipoles) with opening angle Θ_{1+} and Θ_{1-} respectively:

$$\begin{aligned} \tilde{R}_{\perp} &\equiv \frac{dN_{\perp}^{q\bar{q}g}}{dN_{\perp}^{q\bar{q}}(\Theta_{1+}) + dN_{\perp}^{q\bar{q}}(\Theta_{1-})} \\ &= \frac{N_C^2}{N_C^2 - 1} \left(1 - \frac{1}{N_C^2} \frac{1 - \cos(\Theta_{1+} + \Theta_{1-})}{2 - \cos \Theta_{1+} - \cos \Theta_{1-}} \right) \end{aligned} \quad (15)$$

This ratio measures directly the deviation from the large- N_C limit $\tilde{R}_{\perp} = 1$ and thereby from the $q\bar{q}$ -dipole approximation. This approximation is not necessarily limited towards soft particle production. For the simple case of Mercedes-like events ($\Theta_{1+} = \Theta_{1-} = \Theta_{+-}$) one obtains $\tilde{R}_{\perp} = 17/16 = 1.06$. The 2-jet rates for relative angle Θ_{ij} which appear in the denominator of eq. (15) could be found experimentally from the corresponding $q\bar{q}\gamma$ final states.

Let us list a few further results:

- a) A particularly simple situation is met for Mercedes-type events where no jet identification is necessary for the above measurements.
- b) The large angle radiation is independent of the mass of the quark (for $\Theta \gg m_Q/E_{jet}$).
- c) The above predictions for the ratios $R_{\perp}, \tilde{R}_{\perp}$ are derived for the soft particles according to the bremsstrahlung formula (10). One may also consider the particle flow integrated

	R_{\perp}	R_{\perp} (large N_C)
$\Theta_{1+} = \pi - \Theta_{1-}$ (collinear or soft gluons)	1	1
$\Theta_{1+} = \Theta_{+-} = \frac{5}{6}\pi$	1.21	1.18
$\Theta_{1+} = \Theta_{+-} = \frac{3}{4}\pi$	1.42	1.35
$\Theta_{1+} = \Theta_{1-} = \frac{2}{3}\pi$ (Mercedes)	1.59	1.5
$\Theta_{1+} = \Theta_{1-} = \pi$ ($q\bar{q}$ antiparallel to g)	$\frac{N_C}{C_F} = 2.25$	2

Table 1: Prediction for the ratio $R_{\perp} = dN_{\perp}^{q\bar{q}g}/dN_{\perp}^{q\bar{q}}$ from (13) and its large- N_C -approximation (14) for different configurations of the $q\bar{q}g$ events ($\Theta_{1+} \equiv \Theta_{gq}$, $\Theta_{1-} \equiv \Theta_{g\bar{q}}$, $\Theta_{+-} = 2\pi - \Theta_{1+} - \Theta_{1-}$).

over momentum, as in the discussion of the string/drag effect. In this case one has to include all higher order contributions which take into account the fact that the soft gluon is part of a jet generated from a primary parton. Then the angular flow $dN/d\Omega_{\vec{n}}$ is given by the product of the radiation factor $W(\vec{n})$ and a ‘‘cascading factor’’[30]. For the ratio of multiplicity flows one obtains the same predictions (13),(15) as the cascading factors cancel. It will be interesting to see to what extent the predicted angular dependence for both quantities – the multiplicity flows and the soft particle yields – are satisfied experimentally.

d) The similarity of particle flows in 3-jet and 2-jet events with corresponding angles Θ_{1+} , Θ_{1-} , as expected in the large- N_C approximation, should also apply to further details of the final state such as the particle ratios. Since the Lorentz transformation along the boost direction produces a larger drag for heavier particles, one may expect that after the boost the K/π and p/π ratios for soft particles decrease and then the same is true for soft particles in 3-jet events in comparison to 2-jet events in their rest frame⁶.

5. Extension to different reactions

We consider here processes with primary hadrons or photons with dominantly 2-jet final states. This includes semihard processes which are initiated by partonic 2-body scatterings. In case of quark exchange the two outgoing jets originate from colour triplet charges and I_0 should be as in e^+e^- annihilation; in case of gluon exchange I_0 should be about twice as large (N_C/C_F) as in a gg jet system. In 2-jet events the limiting density I_0 may

⁶This expectation has been verified in the JETSET Monte Carlo[31]. We thank T. Sjöstrand for providing us with this information.

be most conveniently determined as limit of $dn/dy d^2p_T$ at $y \approx 0$ for $p_T \rightarrow 0$. Final states with several well separated jets can be treated in analogy to the $e^+e^- \rightarrow 3$ -jet events discussed before.

(a) quark exchange processes

A simple example is the process $\gamma\gamma \rightarrow q\bar{q} \rightarrow 2$ -jets which dominates if either the virtuality Q^2 of the initial photon or the scattering angle photon-jet is sufficiently large (see, e.g. [32]). In this case the soft production intensity I_0 should be the same as in e^+e^- annihilation.

Another example is deep inelastic scattering at large Q^2 . In this case the current fragments in the Breit frame are expected to have the same characteristics as the quark fragments in one hemisphere of e^+e^- annihilation (for a QCD analysis, see[33]), and this is indeed observed for not too small Q^2 [34,35]. Therefore one expects again the same limit I_0 as in e^+e^- annihilation.

(b) gluon exchange processes

These are expected to dominate for virtual $\gamma_V\gamma$ or $\gamma_V p$ scattering ($Q^2 \gtrsim \text{few GeV}^2$) through the photon gluon fusion subprocess $\gamma g \rightarrow q\bar{q}$ at small Bjorken x . In this case (for a fast $q\bar{q}$ pair in one hemisphere with small opening angle) the particle jets emerging in opposite directions are generated by an effective colour octet emitter and the soft intensity I_0 should be about twice as large as in e^+e^- annihilation.

Another example is hadron-hadron collision with a particle or jet at moderate p_T ($\gtrsim 1$ – 2 GeV) at small angles so that the overall 2-jet structure is maintained. This type of scattering is dominated by gluon exchange and the same increase of I_0 is therefore expected.

(c) soft collisions (minimum bias events)

These processes (with initial hadrons or real photons) are not so well understood theoretically as the hard ones but it might be plausible to extrapolate the gluon exchange process towards small p_T [36]. Experimental data at ISR energies, however, do not follow the expectation of a doubling of I_0 in comparison to e^+e^- annihilation, rather the intensities are found to be similar[37]. On the other hand, I_0 roughly doubles when going from $\sqrt{s} \sim 20$ GeV [38,39] to $\sqrt{s} = 900$ GeV at the collider[40]. If additional incoherent sources like weak decays can be excluded, such behaviour could indicate the growing importance of one-gluon exchange expected from the perturbative picture. Then a saturation at $I_0^{hh}/I_0^{e^+e^-} \sim 2$ is expected and no additional increase for a semihard p_T trigger. A rise of I_0^{hh} could also result from incoherent multiple collisions of partons (e.g. [41]) which recently has also been postulated for γp collisions[42].

It should be noted that events from all three classes *a*, *b* and *c* can occur in the same reaction depending on the external kinematical constraints (triggers). For example, if Q^2 for fixed hadronic energy W in γp or $\gamma\gamma$ collisions is decreased from large $Q^2 \sim W^2$ towards zero one changes from (a) to (b) and finally to (c). Also, a dependence on the reference frame is expected, case (a) in Breit frame, case (b) typically in *cms* frame for small Bjorken x .

Rapidity dependence of I_0 .

The frame dependence can be studied conveniently in these exchange processes by considering $I_0(y) = \frac{dn}{dy d^2p_T} \Big|_{p_T \rightarrow 0}$ as a function of rapidity y , measured, say, in the *cms* frame.

The Breit frame is reached by a Lorentz transformation in the direction of the incoming photon, and corresponds to a different rapidity, y_{Breit} , in the *cms*. According to the above discussion, one would then expect for the processes with virtual photons at large Q^2 in general a “quark plateau” of $I_0(y)$ in the current region of length $\Delta y \approx \log(Q/m)$ near y_{Breit} and a transition to a “gluon plateau” in the complementary region of length $\Delta y \approx \log(W^2/Qm)$ where m is a typical particle mass. The existence of a plateau corresponds to the energy independence of I_0 as seen in e^+e^- annihilation. So, if gluon exchange occurs with sufficient hardness in the process, I_0 should develop a step like behaviour but never become larger than twice the I_0 value in e^+e^- annihilation. The height of the “plateau” is a direct indicator of the underlying exchange process (quark or gluon like).

6. Conclusions

The analytical perturbative approach to multiparticle production, based on the Modified Leading Log Approximation and Local Parton Hadron Duality (LPHD) has proven to be successful in the description of various inclusive characteristics of jets. It is of importance to investigate the limitations of this picture, in particular in the soft region where non-perturbative effects are expected to occur.

It is remarkable that the intensity of the soft hadron production follows to a good approximation a scaling law in a range of two orders of magnitude in the *cms* energy of e^+e^- annihilation (1.6-140 GeV). Such a scaling law is derived analytically for the soft gluons in the jet and follows directly from the coherence of the soft gluon radiation from all emitters. It appears that the production of hadrons which is known to proceed through many resonance channels nevertheless can be simulated in the average through a parton cascade down to a small scale of a few 100 MeV as suggested by LPHD. The scaling behaviour down to the very low *cms* energy of 1.6 GeV can only be explained within a parton model description if the cut-off scale Q_0 is well below 1 GeV.

It will be interesting to investigate the suggested scaling property (1) and its possible violation further in the same experiment to avoid systematic effects; this seems to be feasible at LEP and HERA for the simplest processes. In particular, the effect of weak decays on scaling violations should be clarified. The sensitivity of the soft particle production to the effective colour charge of the primary emitters can be tested through the transverse production rates in multijet events. The soft radiation in e^+e^- annihilation can be used as a standard scale in the comparison of various processes. In this way the soft particle production can be a sensitive probe of the underlying partonic process and the contributions from incoherent sources.

Acknowledgements

We would like to thank Yu. L. Dokshitzer, P. Minkowski and T. Sjöstrand for useful discussions. This work was supported in part by the UK PPARC and the EU Training and Mobility Program “Hadronic Physics with High Energy Electromagnetic Probes”, Network FMRX-CT96-0008.

References

- [1] A. Bassetto, M. Ciafaloni and G. Marchesini, Phys. Rep., **C100** (1983) 201.
- [2] Yu. L. Dokshitzer, V. A. Khoze, A. H. Mueller and S. I. Troyan, “Basics of Perturbative QCD”, ed. J. Tran Thanh Van, Editions Frontières, Gif-sur-Yvette, 1991.
- [3] Yu. L. Dokshitzer, V. S. Fadin and V. A. Khoze, Phys. Lett., **115B** (1982) 242; Z. Phys., **C15** (1982) 325.
- [4] A. Bassetto, M. Ciafaloni, G. Marchesini and A. H. Mueller, Nucl. Phys., **B207** (1982) 189.
- [5] Ya. I. Azimov, Yu. L. Dokshitzer, V. A. Khoze and S. I. Troyan, Z. Phys., **C27** (1985) 65 and **C31** (1986) 213.
- [6] Yu. L. Dokshitzer and S. I. Troyan, Proc. 19th Winter School of the LNPI, Vol. 1, p.144; Leningrad preprint LNPI-922 (1984).
- [7] A. H. Mueller, Nucl. Phys., **B213** (1983) 85; erratum quoted ibid., **B241** (1984) 141.
- [8] for a recent discussion of the phenomenological status of the analytical perturbative approach, see V. A. Khoze and W. Ochs, “Perturbative QCD approach to multiparticle production”, preprint Durham DTP/96/36, MPI-PhT-96/29 (1996).
- [9] TASSO Coll., W. Braunschweig et al., Z. Phys. **C47** (1990) 187.
- [10] OPAL Collaboration: M. Z. Akrawy et al., Phys. Lett., **B247** (1990) 617.
- [11] V. A. Khoze, in in “ Z^0 Physics”, Cargèse 1990, Eds. M. Lévy, J-L. Basdevant, M. Jacob, D. Speiser, J. Weyers and R. Gastmans, NATO ASI, Series B: Physics Vol. 261, p. 419.
- [12] S. Lupia and W. Ochs, Phys. Lett. **B365** (1996) 339; paper EPS 0803 subm. to the 1995 Int. Europhysics Conference on High Energy Physics, 27 July – 2 August 1995, Brussels, Belgium.
- [13] J. D. Bjorken and E. A. Paschos, Phys. Rev. **185** (1969) 1975.
- [14] R. P. Feynman, “Photon-Hadron Interaction”, W. A. Benjamin, NY, 1972.
- [15] G. J. Feldman and M. L. Perl, Phys. Rep. **19** (1975) 233.
- [16] Ya. I. Azimov, Yu. L. Dokshitzer, V. A. Khoze, Sov. Phys. Uspekhi **23** (1980) 732.
- [17] S. Lupia, hep-ph/9608360, to appear in the Proc. of QCD '96, (Montpellier, France, 1996), Nucl. Phys. B Proc. Suppl.;
S. Lupia, V. A. Khoze, W. Ochs, preprint MPI-PhT-96-94, to appear in the Proc. of the 7th International Workshop “Correlations and Fluctuations”, Nijmegen, Holland, June 30-July 6, 1996.
- [18] Yu. L. Dokshitzer, V. A. Khoze and S. I. Troyan, Int. J. Mod. Phys. **A7** (1992) 1875.

- [19] V. A. Khoze, S. Lupia and W. Ochs, “Perturbative description of particle spectra at LEP-1.5”, preprint Durham DTP/96/38, MPI-PhT-96/28, hep-ph/9604410 (April 1996), to appear in Phys. Lett. B
- [20] Yu. L. Dokshitzer, V. A. Khoze and S. I. Troyan, Z. Phys., **C55** (1992) 107.
- [21] Yu. L. Dokshitzer, V. A. Khoze and S. I. Troyan, in Proc. 6th Int. Conf. on Physics in Collision, ed. M. Derrick (World Scientific, Singapore, 1987), p.417;
- [22] MARK I Coll., J. L. Siegrist et al., Phys. Rev. **D26** (1982) 969.
- [23] MARK II Coll., J. F. Patrick et al., Phys. Rev. Lett. 49 (1982) 1232;
 TOPAZ Coll., R. Itoh et al., Phys. Lett. **B345** (1995) 335;
 ALEPH Coll., D. Buskulic et al., Z. Phys. **C55** (1992) 209;
 DELPHI Coll., P. Abreu et al., Phys. Lett. **B347** (1995) 447;
 L3 Coll., B. Adeva et al., Phys. Lett. **B259** (1991) 199;
 ALEPH Coll., D. Buskulic et al., CERN preprint CERN-PPE/96-43, March 1996, submitted to Z. Phys. C;
 DELPHI Coll., P. Abreu et al., Phys. Lett. **B372** (1996) 172;
 OPAL Coll., M. Z. Akrawy et al., CERN preprint CERN-PPE/96-47, March 1996, submitted to Z. Phys. C.
- [24] ADONE Coll., B. Esposito et al., Lett. Nuovo Cimento **19** (1977) 21;
 DASP Coll., R. Brandelik et al., Phys. Lett. **B67** (1977) 358;
 ARGUS Collaboration, S. Behrends et al., Phys. Rev. **D31** (1985) 2161;
 TPC/2 γ Collaboration, H. Aihara et al., Phys. Rev. Lett. **61** (1988) 1263.
- [25] DELPHI Collaboration: P. Abreu et al., Nucl. Phys. **B444** (1995) 3.
- [26] S. J. Brodsky and J. F. Gunion, Phys. Rev. Lett., **37** (1976) 402.
- [27] Yu. L. Dokshitzer, V. A. Khoze and S. I. Troyan, Sov. J. Nucl. Phys., **47** (1988) 881;
 J. W. Gary, Phys. Rev., **D49** (1994) 4503.
- [28] OPAL Collaboration, “Test of QCD analytical predictions for the mean multiplicity difference between gluon and quark jets”, paper pa02-014 submitted to the 28th International Conference on High Energy Physics, Warsaw, Poland, 25-31 July 1996.
- [29] B. Andersson, G. Gustafson, G. Ingelman and T. Sjöstrand, Phys. Rep., **97** (1983) 31;
 B. Andersson, G. Gustafson and T. Sjöstrand, Phys. Lett., **B94** (1980) 211.
- [30] Ya. I. Azimov, Yu. L. Dokshitzer, V. A. Khoze and S. I. Troyan, Phys. Lett., **B165** (1985) 147.
- [31] T. Sjöstrand, Computer Physics Commun. **82** (1994) 74.
- [32] H. Kolanoski, P. Zerwas, in “High Energy Electron-Positron Physics”, Eds. A. ALi, P. Söding, World Scientific, Singapore, 1988, p. 697.

- [33] L. V. Gribov, Yu. L. Dokshitzer, V. A. Khoze and S. I. Troyan, JETP Lett. **45** (1987) 405; Sov. Phys. JETP **68** (1988) 1303.
- [34] H1 Collaboration: S. Aid et al., Nucl. Phys. **B445** (1995) 3.
- [35] ZEUS Collaboration: M. Derrick et al., Z. Phys. **C67** (1995) 93.
- [36] F. E. Low. Phys. Rev. **D12** (1975) 163; S. Nussinov, Phys. Rev. Lett. **34** (1975) 1286.
- [37] H. Aihara et al., Phys. Lett. **B184** (1987) 114
- [38] NA22 Collaboration: M. Adams et al., Z. Phys. **C39** (1988) 311.
- [39] ABCDHW Collaboration: A. Breakstone et al., Z. Phys. **C69** (1995) 55.
- [40] UA1 Collaboration: C. Albajar et al., Nucl. Phys. **B335** (1990) 261.
- [41] T. Sjöstrand and M. van Zijl, Phys. Rev. **D36** (1987) 2019.
- [42] J. M. Butterworth, J. R. Forshaw, T. Sjöstrand and J. K. Storrow, J. Phys. **G22** (1996) 883.




Figure 1: Invariant density $E dn/d^3p$ as a function of the particle energy E for $Q_0 = 270$ MeV. Predictions at cms energies of $\sqrt{s} = 3$ GeV (lower three curves) and 91 GeV (upper three curves) using $E dn/d^3p = 2 \cdot 4/9 K_h D_g^g(\xi_E)/[4\pi(E^2 - Q_0^2)]$ with D_g^g computed in MLLA (eqs. (6,3), DLA (eq. (3)) and the Limiting Spectrum (normalization $K_h = 0.45, 0.45$ and 1.125 respectively).




Figure 2: Invariant density $E dn/d^3p$ of charged particles in e^+e^- annihilation as a function of the particle energy $E = \sqrt{p^2 + Q_0^2}$ at $Q_0 = 270$ MeV. Data points at various cms energies from SLAC, TASSO and TOPAZ Collaborations, LEP-1 and LEP-1.5[23,9,10] are compared to MLLA predictions (normalization as in Fig. 1 with $\lambda = 0.01$, $K_h = 0.45$).




Figure 3: **a)** Invariant density $E dn/d^3p$ of charged pions in e^+e^- annihilation as a function of the particle energy E at $Q_0 = 138$ MeV at various cms energies; **b)** the same as in **a)**, but for charged kaons (with $Q_0 = 494$ MeV) and protons (with $Q_0 = 938$ MeV) [24,23]. The kaon distribution is multiplied by a factor 10 for the sake of clarity.