

hep-ph/9610280

RAL-96-052

RU-96-35

Determining the Gluonic Content of Isoscalar Mesons

Frank E. Close¹

Rutherford Appleton Laboratory
Chilton, Didcot, OX11 0QX, England

Glenys R. Farrar²

Department of Physics and Astronomy
Rutgers University, Piscataway, NJ 08855, USA

Zhenping Li³

Physics Department
Peking University, Beijing, 100871, P. R. China

Abstract

We develop tools to determine the gluonic content of a resonance of known mass, width and J^{PC} from its branching fraction in radiative quarkonium decays and production cross section in collisions. We test the procedures by applying them to known qqm mesons, then analyze four leading glueball candidates. We identify inconsistencies in data for $J=0$ $f_0(1500)$ and $J=1$ $f_J(1710)$ whose resolution can quantify their glueball status. When $(f_0(1500) \rightarrow \gamma \gamma)$ and $(f_J(1710) \rightarrow \gamma \gamma)$ are known, the $nn;ss;gg$ mixing angles can be determined. The enigmatic situation in 1400-1500 MeV region of the isosinglet 0^+ sector is discussed.

¹e-mail: fec@v2.rl.ac.uk

²e-mail: farrar@farrar.rutgers.edu

³e-mail: zpli@ibm320h.phy.pku.edu.cn

1 Introduction

There has been considerable recent interest in the possible sighting of glueballs. Four states are of particular interest:

$f_0(1500)$ [1, 2, 3]

$f_J(1710)$ [4] where $J = 0$ or 2 [5]

(2230) [6]

(1440) [7], now resolved into two pseudoscalars.

In this paper we calculate the production rate of conventional mesons ($q\bar{q}$) and glueballs ($\langle G \rangle$) in the radiative decay of vector quarkonium, as a function of their mass, angular momentum, and width. If the data on the radiative production of these states are correct, we find that

(i) The $f_0(1500)$ is probably produced at a rate too high to be a $q\bar{q}$ state. The average of world data suggests it is a glueball- $q\bar{q}$ mixture.

(ii) The $f_J(1710)$ is produced at a rate which is consistent with it being $q\bar{q}$, only if $J = 2$. If $J = 0$, its production rate is too high for it to be a pure $q\bar{q}$ state but is consistent with it being a glueball or mixed $q\bar{q}$ -glueball having a large glueball component.

(iii) The (2230) , whose width is ~ 20 MeV, is produced at a rate too high to be a $q\bar{q}$ state for either $J = 0$ or 2 . If $J = 2$, it is consistent with being a glueball. The assignment $J = 0$ would require $\text{Br}(J = 0) < 3 \times 10^{-4}$, which already may be excluded.

(iv) The enhancement once called (1440) has been resolved into two states. The higher mass (1480) is dominantly ss with some glue admixture, while the lower state (1410) has strong affinity for glue.

We note what improvements in data would allow these constraints to be sharpened. We also analyze 3-state mixing in the 0^{++} sector between nn , ss and gg , showing how $(J = 0, f_0)$ and (f_0, \dots) determine mixing angles.

The interest in these states as glueball candidates is motivated on both phenomenological and theoretical grounds. Phenomenologically, these states satisfy qualitative criteria expected for glueballs[8]:

1. Glueballs should be favoured over ordinary mesons in the central region of high energy scattering processes, away from beam and target quarks. The f_J (1710) and possibly the f_0 (1500) have been seen in the central region in pp collisions[9, 10].
2. Glueballs should be produced in proton-antiproton annihilation, where the destruction of quarks creates opportunity for gluons to be manifested. This is the Crystal Barrel [11], and E 760 [12] production mechanism, in which detailed decay systematics of f_0 (1500) have been studied. The empirical situation with regard to f_J (1710) and f_0 (2230) is currently under investigation. The f_0 (1440) is clearly seen in pp annihilation [13, 14]
3. Glueballs should be enhanced compared to ordinary mesons in radiative quarkonium decay. In fact, all four of these resonances are produced in radiative J/ψ decay at a level typically of 1 part per thousand. A major purpose of this paper is to decide whether these rates indicate that these resonances are glueballs, or not.

On the theoretical side, lattice QCD predicts that the lightest "ideal" (i.e., quenched approximation) glueball be 0^{++} , with state-of-the-art mass predictions of 1.55 ± 0.05 GeV [15] and 1.74 ± 0.07 GeV [4]. That lattice QCD is now concerned with such fine details represents considerable advance in the field and raises both opportunity and enigmas. First, it encourages serious consideration of the further lattice predictions that the 2^{++} glueball lie in the 2.2 GeV region, and hence raises interest in the f_0 (2230). Secondly, it suggests that scalar mesons in the 1.5 - 1.7 GeV region merit special attention. Amsler and Close[2] have pointed out that the f_0 (1500) shares features expected for a glueball that is mixed with the nearby isoscalar members of the 3P_0 qq nonet. If the f_J (1710) proves to have $J = 2$, then it is not a candidate for the

ground state glueball and the $f_0(1500)$ will be essentially unchallenged. On the other hand, if the $f_J(1710)$ has $J = 0$ it becomes a potentially interesting glueball candidate. Indeed, Sexton, Vaccarino and Weingarten [16] argue that $f_{J=0}(1710)$ should be identified with the ground state glueball, based on its similarity in mass and decay properties to the state seen in their lattice simulation. While the consistency between theoretical mass predictions and the observed states is quite satisfactory in the 0^{++} and 2^{++} sectors, this is not the case in the 0^+ sector. Both lattice and sum rule calculations place the lightest 0^+ glueball at or above the 2^{++} glueball so that the appearance of a glueball-like pseudoscalar in the 1.4-1.5 GeV region is unexpected. It is interesting that its properties are consistent with those predicted for the gluino-gluino bound state in supersymmetry breaking scenarios with a light gluino [17]

In order to make quantitative estimates of the gluonic content of isosinglet mesons, we use their measured radiative quarkonium production rates and gamma-gamma decay widths. We apply the relationship proposed by Cakir and Farrar [18] (CF) between the branching fraction for a resonance R in radiative quarkonium decay, $b_{\text{rad}}(Q\bar{Q}_V \rightarrow R) = (Q\bar{Q}_V \rightarrow X)$ and its branching fraction to gluons, $\text{br}(R \rightarrow gg) = \text{br}(R \rightarrow \text{all})$:

$$b_{\text{rad}}(Q\bar{Q}_V \rightarrow R_J) = \frac{C_R \times H_J(x)^2 m_R}{8 (2/9) M_V^2} \text{tot br}(R_J \rightarrow gg); \quad (1)$$

where M_V and m_R are masses of the initial and final resonances, and $x = 1 - \frac{m_R^2}{M_V^2}$; C_R is a numerical factor and $H_J(x)$ a loop integral which will be discussed in section 2. For a resonance of known mass, total width (Γ_{tot}), and J^{PC} , a relationship such as eq. (1) would determine $\text{br}(R \rightarrow gg)$ if $b_{\text{rad}}(Q\bar{Q}_V \rightarrow R)$ were known. CF argued that one expects

$$\begin{aligned} \text{br}(R[qq] \rightarrow gg) &= O\left(\frac{2}{9}\right) \sim 0.1 - 0.2 \\ \text{br}(R[G] \rightarrow gg) &\sim O(1); \end{aligned} \quad (2)$$

Thus knowledge of $\text{br}(R \rightarrow gg)$ would give quantitative information on the glueball content of a particular resonance. Using $H_J(x)$ determined in the non-relativistic quark model (NRQM), CF found that known qq resonances

(such as $f_2(1270)$) satisfy the former and noted that the $f_0(1710)$ might be an example of the latter.

In the present paper we give a more general discussion of the functions $H_J(x)$ needed to employ Eq. (1), clarify some of the assumptions implicit in its derivation, and verify that application of the relation does not depend on flavor mixing. A number of experimental tests are proposed. We discuss the additional information that can be obtained when the cross section for production of the resonance in gg collisions is known.

The paper is organized as follows. We start with a section on the formalism and its model dependence (sec. 2). A general treatment of the problem requires defining form factors for the coupling of a resonance, of specified J^{PC} , to a pair of virtual gluons. The partial width $\Gamma(R \rightarrow gg)$ is a linear combination of the form factors at the on-shell-gluon point. The internal structure of the resonance determines both the relative size of the various form factors at the on-shell-gluon point, as well as their virtuality dependence, just as in the case of the nucleon electromagnetic form factors F_1 and F_2 . The $H_J(x)$'s depend on integrals of the form factors over the gluon virtualities. In sec. 2.4 we discuss higher order corrections, scale dependence, and the relationship of the $R \rightarrow gg$ form factors to the $R \rightarrow \gamma\gamma$ amplitudes which can in principle be measured in a photon-photon collider. (The phenomenology of the latter is developed in section 5). In sec. 3 we re-express eq. (1) so that its implications are more transparent and it is easier to apply to data. Our central results (eqs. 38-40) show how the spin of a resonance and its width into gluons is its production rate in radiative quarkonium decay. In section 3.2 we show that the relations do not depend on flavor mixing and that the known qq resonances $f_2(1270; 1525)$ satisfy eq.(2). In sec. 3.3 we discuss the utility of experimental study of radiative ψ decay, especially $\psi(3700)$; in sec. 3.4 we investigate 1^{++} mesons. In secs. 4.1-4.3 we apply eqs. (38-40) to the $f_0(1500)$; $f_J(1710)$, $f_2(2300)$, and $f_2(1410;1480)$ leading to the results listed at the beginning of the introduction. In section 5 we discuss how $\Gamma(R \rightarrow gg)$ in combination with J^{PC} can help to distinguish glueballs from qq states and determine

basic parameters. Section 6 considers the possibility of glueball- $q\bar{q}$ mixing involving three states, $f_0(1370)$; $f_0(1500)$ and $f_0(1710)$. In general we use the nomenclature of the 1994 edition of the Particle Data Tables throughout [5]. Readers mainly interested in the phenomenological results can proceed directly to sections 3 et seq.

2 Formalism

2.1 $\Gamma_{\text{rad}}(Q\bar{Q}_V \rightarrow \gamma + R)$

The decay width for the radiative decay of a vector heavy quarkonium state, $Q\bar{Q}_V \rightarrow \gamma + R$ is

$$\Gamma = \frac{1}{24} \frac{k^3}{M_V^2} \sum_{\text{if}} |A|^2; \quad (3)$$

where k is the photon momentum, M_V is the mass of the spin-1 $Q\bar{Q}_V$ state, and the summation is over the polarizations of the initial and final particles. If the resonance R does not contain a "valence" $Q\bar{Q}$ component, the decay occurs through a two gluon intermediate state, in leading order pQCD, and the amplitude A is given by

$$A = \frac{1}{2} \sum_{\lambda_1, \lambda_2} \int \frac{d^4k}{(2\pi)^4} \frac{1}{k_1^2} \frac{1}{k_2^2} \langle (Q\bar{Q})_V | j^a g^b \rangle \langle g^a g^b | R \rangle; \quad (4)$$

The summation is over the polarization vectors $\epsilon_{\lambda_1, \lambda_2}$ and color indices a, b of the intermediate gluons, whose momenta are denoted $k_{1,2}$. The amplitude $\langle (Q\bar{Q})_V | j^a g^b \rangle$ couples a vector $Q\bar{Q}$ state with polarization and momentum $(\epsilon; K)$ to a photon $(\epsilon; k)$ and the two virtual gluons. For heavy quarks, Q , this amplitude is reliably given by perturbative QCD. Using the non-relativistic quark model to describe the $Q\bar{Q}_V$ wavefunction (the pQCD-NRQM approximation [19, 20, 21]):

$$\langle (Q\bar{Q})_V | j^a g^b \rangle = e_Q g_s^2 \epsilon^{abc} \int \frac{d^3k}{4 M_V^3} \frac{2 i R_V(0)}{3} \frac{M_V^2}{(k+k_2)k_1 (k+k_1)k_2 (k+k)k} \epsilon_{\lambda_1}^a \epsilon_{\lambda_2}^b \epsilon_{\lambda}^c; \quad (5)$$

where

$$\begin{aligned}
a_V = & \frac{1}{2} [k_1 \cdot k_2 E_k \cdot k_2 + k_1 \cdot E_k \cdot k_1 + k_1 \cdot k_2 E_k \cdot k_1] \\
& + E_k \cdot k_1 + k_2 \cdot k_1 + k_2 \cdot k_1 + k_1 \cdot k_2 + k_1 \cdot k_2 \\
& + f_1; k_1, \quad ; kg + f_2; k_2, \quad ; kg: \quad (6)
\end{aligned}$$

$R_V(0)$ is the $(QQ)_V$ wavefunction at the origin and e_Q is the charge of the heavy quark Q .

The amplitude $\langle g^a(k_1; \lambda_1) g^b(k_2; \lambda_2) \mathcal{R} \rangle$ must be linear in λ_1 and λ_2 and Lorentz and gauge invariant. A linearly independent set of tensor structures satisfying these requirements for J^{++} states is given in [22]. Thus we can write, with the shorthand $G^i(k_1, k_2)$ and the convention that $G^{1(2)}$ refers to $k_{1(2)}^i$, and suppressing J^{PC} labels on the F_i 's:

$$\langle g^a g^b \mathcal{J}^{++} \rangle = \frac{ab A_{0^{++}}}{P} F_1(k_1^2; k_2^2) G^1 G^2 + F_2(k_1^2; k_2^2) k_1 G^1 G^2 k_2^i \quad (7)$$

$$\langle g^a g^b \mathcal{J}^+ \rangle = ab A_0 + F_1(k_1^2; k_2^2) G^1 G^2; \quad (8)$$

$$\begin{aligned}
\langle g^a g^b \mathcal{J}_1^{++} \rangle = & ab A_{1^{++}} F_1(k_1^2; k_2^2) (G^1 G^2 k_2 + G^2 G^1 k_1) \\
& + F_2(k_1^2; k_2^2) (k_1 \cdot k_2) G^1 G^2; \quad (9)
\end{aligned}$$

$$\begin{aligned}
\langle g^a g^b \mathcal{J}_2^{++} \rangle = & ab A_{2^{++}} F_1(k_1^2; k_2^2) G^1 G^2 + F_2(k_1^2; k_2^2) k_1 k_2 G^1 G^2 + \\
& F_3(k_1^2; k_2^2) k_1 G^1 G^2 k_2 + F_4(k_1^2; k_2^2) k_1 k_2 k_1 G^1 G^2 k_2^i; \quad (10)
\end{aligned}$$

where

$$P^i = g \frac{P^i P^j}{m^2}; \quad (11)$$

for a resonance with mass m and momentum P . Here λ and λ_i are the polarization vector and tensor for a vector or tensor resonance, and satisfy the relations

$$\begin{aligned}
\lambda^i \lambda^j &= g \frac{P^i P^j}{m^2} \text{ or} \\
\lambda_{00} &= \frac{1}{2} (P_0 P_0 + P_0 P_0) = \frac{1}{3} P_0 P_0; \quad (12)
\end{aligned}$$

The resonance R could be qq , glueball or mixture of both. The form factors $F_i(k_1^2; k_2^2)$ depend on the composition of R . This will be discussed

below. We adopt the normalization convention that $F_1[J^{PC}](0;0) = 1$. Also, we use the shorthand for on-shell form factors $F_i[J^{PC}](0;0) = F_i[J^{PC}]$. Note that for $k_1^2 = k_2^2 = 0$, $\langle g^a g^a j^{1++} \rangle = 0$ as it must by Furry's theorem. The constants $A_{J^{PC}}$ in Eqs. 7-10 are dependent on the coupling between gluons and the resonance constituents, the wavefunction of the resonance and its mass (specific examples for qq are given below in eq. (24)).

After summing over the color index of the initial two gluon state, the general expression for the total width of a resonance R decaying into two real gluons [5] is

$$\Gamma(R \rightarrow gg) = \frac{1}{(2J+1)} \frac{1}{2m} \sum_{1,2} \langle gg | R \rangle^2; \quad (13)$$

where $\langle gg | R \rangle$ is the coefficient of δ^{ab} in eqs. 7-10. For example in the case $R = 0^{++}$, using the matrix element $\langle gg | 0^{++} \rangle$ given by eq. 7, and summing over gluon polarizations $1,2$, gives

$$\begin{aligned} \sum_{1,2} \langle gg | 0^{++} \rangle^2 &= \frac{1}{3} \mathcal{A}_{0^{++}}^2 \int (k_1 - k_2)^2 g_{00} g_{00} + k_1^0 k_1^0 k_2^0 k_2^0 \\ &\quad + k_1^0 k_1^0 k_2^0 k_2^0 - 2k_1^0 k_1^0 (g_{01} k_2^0 + g_{01} k_1^0 k_2^0) \\ &\quad + k_1^0 k_1^0 (g_{01} k_2^0 + g_{01} k_1^0 k_2^0) - P_{11} P_{00} \\ &= \frac{3}{8} m^4 \mathcal{A}_{0^{++}}^2; \end{aligned} \quad (14)$$

which leads to

$$\Gamma(0^{++}) = \frac{3m^3}{16} \mathcal{A}_{0^{++}}^2; \quad (15)$$

The decay widths for the other states are obtained by the same procedure and read

$$\Gamma(0^{-+}) = \frac{2m^3}{3} \mathcal{A}_{0^{-+}}^2; \quad (16)$$

and

$$\Gamma(2^{++}) = \frac{m^3}{20} \mathcal{A}_{2^{++}}^2 \left[1 + \frac{m^4}{12} F_2^2[2^{++}] \right]; \quad (17)$$

While these are nominally two-gluon widths, when the scale of resolution of the gluons is taken large enough (see ref. [18] and below) they become the total

gluonic widths. Note also that since $A_{1^{++}}$ cannot be fixed in this way, we focus primarily on 0^{++} , 2^{++} , and 0^{+} states. We return to the axial mesons in sec. 3.4

Given the form factors appearing in $\langle g^a g^b \mathcal{R} \rangle$ and the NRQM pQCD formula (eq. 5) for $\langle Q Q_V j g^a g^b \rangle$, the integral $\int d^4 k \frac{1}{k_1^2 k_2^2} \langle Q Q_V j g^a g^b \rangle \langle g^a g^b \mathcal{R} \rangle$ can be carried out, thus determining the $H_J(x)$'s appearing in eq. (1). The analysis of ref.[18] assumed that the relative size of the on-shell form factors and their dependence on gluon virtualities is universal, for heavy and light qq mesons and for glueballs. There is no general reason why this should be the case. For example, we know from form factors of electromagnetic and weak currents that some aspects of form factors are universal⁴ while other aspects such as the relative magnitude of the nucleon on-shell form factors depend on detailed structure of the bound state, in particular the constituent quark magnetic moments. The next sections describe the information we presently have on the $\langle gg \mathcal{R} \rangle$ form factors.

2.2 $\langle gg \mathcal{R} \rangle$ form factors

A particular example for the $\langle gg \mathcal{R} \rangle$ form factors is the case of $\mathcal{R} = qq$, where the quantity $\langle gg \mathcal{R} \rangle$ has been modeled [19, 21, 22] as a QCD analogue [24] of the two photon coupling to positronium [25]. In the NRQM approximation [22]

$$\langle gg \mathcal{J}^{++}(qq) \rangle = c^0 \frac{1}{6} \int d^4 k \frac{1}{k_1^2 k_2^2} G^a G^a (m^2 + k_1^2) 2k_1^i G^a G^a k_2^i = (k_1^2 - k_2^2)^2; \quad (18)$$

$$\langle gg \mathcal{J}^{++}(qq) \rangle = c^0 m \frac{1}{2} \int d^4 k \frac{1}{k_1^2 k_2^2} G^1 G^2 k_2^i + G^2 G^1 k_1^i = (k_1^2 - k_2^2)^2; \quad (19)$$

and

$$\langle gg \mathcal{P}^{++}(qq) \rangle = c^0 \frac{1}{2m} \int d^4 k \frac{1}{k_1^2 k_2^2} G^a G^a e^i = (k_1^2 - k_2^2)^2; \quad (20)$$

⁴For instance, the leading Q^2 dependence of the nucleon and meson electromagnetic form factors depends only on the number of valence constituents [23].

where the constants c^0 are proportional to the derivative of the radial wavefunctions at the origin:

$$c^0 = g_s^2 \frac{1}{m^3} R^0(0); \quad (21)$$

Analogously the two gluon coupling for a 0^+ state is [22]

$$\langle gg \mathcal{J}^+ (qq) \rangle = c \quad G^a G^a = k_1 \quad k; \quad (22)$$

where

$$c = g_s^2 \frac{1}{4} \frac{1}{3m} R^0(0); \quad (23)$$

The above are particular models for Eqs. 7-10. To see the correspondence between the quantities $\langle gg \mathcal{J}^{PC} (qq) \rangle$ and Eqs. 7-10, it is convenient to note that Eq. 18 can be rewritten in the form of Eq. 7:

$$\langle gg \mathcal{J}^{++} (qq) \rangle = c m^2 \frac{2}{3} G^{1a} G^{2a} P = (k_1 \quad k)^2; \quad (24)$$

The constants A_{JPC} in our Eqs. 7-10 are thus related to the constants c and c^0 of ref. [22] (Eqs. 21 and 23 above) by

$$A_{0^{++}} = A_{2^{++}} = \frac{P}{8m} A_{1^{++}} = \frac{4}{m^2} \frac{2c^0}{2c} \\ A_{0^+} = \frac{2c}{m^2}; \quad (25)$$

and the matching implies that the only non-zero form factors for $\langle gg \mathcal{J}^{PC} \rangle$ in the NRQM are

$$F_1 [0^{++}] = F_1 [2^{++}] = F_1 [1^{++}] = \frac{m^4}{4 (k_1 \quad k)^2}; \quad (26)$$

and [32]

$$F_1 [0^+] = \frac{m^2}{2k_1 \quad k}; \quad (27)$$

Substituting Eq. 25 into Eqs. 15-17, gives

$$((qq)_{0^{++}}) = 96 \frac{2}{m^4} R^0(0)^2; \quad (28)$$

$$\langle (gg)_{2^{++}} \rangle = \frac{128}{5} \frac{s^2}{m^4} \mathcal{R}^0(0) f^2; \quad (29)$$

and

$$\langle (gg)_{0^{++}} \rangle = \frac{8}{3} \frac{s^2}{m^2} \mathcal{R}^0(0) f^2; \quad (30)$$

which agree with those in Ref. [18].

The gluon structure appearing in the form factors for the scalar and tensor NRQM mesons (e.g., $G^a G^a P$) has been widely employed also for scalar and tensor glueballs [26]. It can be considered a natural relativistic generalization of TE mode glueballs in a cavity approximation such as the MIT bag model [27]:

$$(G_{J^{++}}) = h_{1;1} j_{\mathbf{J}} + i(\gamma_{\hat{K}_1})^{(\prime)} (\gamma_{\hat{K}_2})^{(\prime)} (r) \quad (31)$$

where $r(x)$ is a radial wavefunction, and the superscripts \prime specify the projection of the angular momenta along \hat{z} . The relativistic generalization adopted in ref. [26] produces

$$(G_{0^{++}}) = \frac{1}{3} P \frac{G^{1a} G^{2a}}{8k_1 k} (x) \quad (32)$$

for the scalar and

$$(G_{2^{++}}) = \frac{G^{1a} G^{2a}}{8k_1 k} (x) \quad (33)$$

for the tensor states, where P and \mathcal{R}^0 are defined in eqs. 11 and 12. Note that in cavity approximation the same function $r(x)$ appears for both 0^{++} and 2^{++} states, so that the relative magnitudes of their form factors are fixed. The resulting matrix elements for the two gluon couplings of the glueballs are

$$\langle gg \mathcal{G}^J (J^{PC}) \rangle = A f(k_1^2; k_2^2) P^J \frac{G^{1a} G^{2a}}{k_1 k} \quad (34)$$

where $P^0 = \frac{P}{3}$ and $P^2 = \mathcal{R}^0$. The form factor $f(k_1^2; k_2^2)$ in Eq. 34 is determined by the wavefunctions $r(x)$ which appears in eqs. 32 and 33.

Comparison with eqs. (7) and (10) shows that the glueball wavefunction (34) corresponds to $A_{0^{++}} + F_1 [0^{++}](k_1^2; k_2^2) = A_{2^{++}} + F_1 [2^{++}](k_1^2; k_2^2)$; the remaining form factors vanish. Since this relation between 0^{++} and 2^{++} form factors is

the same as in the NRQM, both models give the same result for the ratio of the 2^{++} and 0^{++} widths. Thus in the limit that the masses of the scalar and tensor states are equal, independently of whether they are a pair of (NRQM) qq [24, 25, 28] or (cavity approximation) glueball states,

$$\frac{\Gamma_{2^{++}}}{\Gamma_{0^{++}}} = \frac{4}{15}; \quad (35)$$

The large mass gap between the 0^{++} glueball candidates $f_0(1500)$ and $f_0(1700)$, and the 2^{++} candidate (2230) (assuming (2230) has $J = 2$) prevents immediate application of eq. 35. However since these f_0 's have widths 100–150 MeV and the (2230) width is 20 MeV (see sec. 3.3) { either f_0 is compatible with (35). The presence of 0^{++} qq states in the vicinity of these f_0 's also complicates the situation (see sec. 6).

For a light qq system, Eq. 35 will be modified by relativistic effects [28], which increases the ratio $4=15$ to around $\frac{1}{2}$. While there are no $R \rightarrow gg$ data available, one could relate the width for $R \rightarrow gg$ to that for the $R \rightarrow gg$ at the tree level. The data for $f_0(1300) \rightarrow gg$ and $f_2(1270) \rightarrow gg$ are consistent with the result here. However, the relativistic effects on the loop integral $\int d^4x H(x)^2$ remain to be investigated.

2.3 Virtuality Dependence of Glueball Form Factors

The analysis of $J=0$ radiative decays in ref. [18] implicitly assumed that the relative size of the on-shell form factors and also their dependence on gluon virtualities are universal for qq mesons and glueballs. Our investigation of the previous section showed that the first assumption may be reasonable.

For the $\langle ggR \rangle$ form factors the situation is more complicated than for electromagnetic form factors, because the gluon virtualities can vary independently. That is, the form factors here are functions of two variables, constrained by the requirement of being even under interchange of the gluon

momenta (from Bose statistics). We can further constrain the form factors by power counting arguments. Replace the variables k_1^2 and k_2^2 by $k_1 - k_2$ and the dimensionless ratio $z = \frac{(k_1 - k_2)(k_1 + k_2)}{k_1 k_2}$. When R is an $L = 0$ bound state of two constituents, the leading large $k_1 - k_2$ behavior of $F_1(k_1^2; k_2^2)$ is $\frac{1}{(k_1 - k_2)} f(z)$. The $F_i(k_1^2; k_2^2)$ entering $\langle ggR \rangle$ with additional factors of k_i have correspondingly more rapid falloff, like F_2 compared to F_1 for the case of electromagnetic form factors. For $L = 1$ systems one expects an additional $k_1 - k_2$ suppression, where Λ is a scale reflecting the variation of the wavefunction at the origin. This scaling behavior is manifested by the NRQM results eqs. 18 - 22.

Neglect of higher twist corrections to the leading form factors and neglect of those form factors whose leading dependence falls more rapidly, can be expected to give corrections to the $H_J(x)$'s of order $m_R^2 = M_V^2$ compared to the leading terms. Since for our application this is a small quantity, we neglect these corrections. As we saw in sec. 2.2, in the NRQM $f(z) = \text{const}$. The effect of possible corrections to constancy of $f(z)$, and overall scaling behavior which differs from the NRQM, is presently under study.

2.4 Higher Order Corrections and Scale Dependence

For a resonance which couples to the two-gluon intermediate state, corrections to the above formalism involve one additional gluon loop and thus should be of order $O(\alpha_s^4)$ in the amplitude. For heavy QQ mesons the wavefunction can be treated perturbatively and it is straightforward to make a systematic expansion in the coupling constant [24, 29]. In this context it is sensible to distinguish between the components of the wavefunction in which the QQ are in a "color singlet" or "color octet" state [30]. However for the light qq mesons and glueballs of interest, defining a perturbative expansion and the relation between "constituent" and "current" partons are more subtle and we do not undertake this here. Suffice it to say that the concept of "color singlet" versus "color octet" components of the wavefunction does not have an a priori well-defined intuitive meaning as for the heavy quark system. The issue of composition is a scale-dependent question, as it is for the nucleon.

The Altarelli-Parisi evolution of the parton distribution functions is a clear illustration of this point. In principle the same is true for the heavy quark system, however the quark mass gives a natural scale in that case.

Our treatment in previous sections implicitly made use of an effective Lagrangian approach to the problem. By working with $b_{\text{rad}}(Q, Q_V, \dots + R)$, i.e., dividing $(Q, Q_V, \dots + R)$ by $(Q, Q_V, \dots + X)$ also computed in leading order perturbation theory, one removes the dependence on the effective strong coupling at the heavy quark vertices. Similarly, quoting the result in terms of $\text{br}(R \rightarrow gg)$ removes at leading order the sensitivity to the scale dependence of the definition of the R wavefunction. Of course, the concept of "gluonic width" of a qq resonance necessarily has an intrinsic scale dependence (as one goes to shorter and shorter distance scale, contributions from the parton sea invalidate simple valence intuition). Without a careful treatment of next-to-leading order corrections, we cannot specify the correct scale for α_s appearing in the estimate of eq.(1) for the gluonic branching fraction of qq mesons. For this reason, we can make only qualitative use of the gluonic branching fractions that we extract for qq mesons. However when the branching fraction of a state is found to be $> 1=2$, indicating that the state has a significant gluon component, the sensitivity on α_s is a higher order correction and we can make quantitative use of the $\text{br}(R \rightarrow gg)$ that we extract from the data.

It follows from the above discussion that we cannot expect a trivial relationship between the form factors for the amplitude $\langle gg | R \rangle$, and those for $\langle R | j \rangle$. In principle the latter can be measured as a function of photon virtualities in an e^+e^- collider. For heavy QQ resonances such as the ψ_c states, these amplitudes are identical except for the value of the overall coefficient A_{jPC} . To obtain $\langle R | j \rangle$ from $\langle gg | R \rangle$, substitute $g \rightarrow e_q$ and remove the color factor. At leading order this gives

$$\langle R | j \rangle = \frac{9e_q^4}{2} \left(\frac{1}{s} \right)^2 \langle R | gg \rangle: \quad (36)$$

We test the validity of eq. (36) for light qq resonances in section 5 by applying it to the known qq states $f_2(1270)$ and $f_2(1525)$. We then extend it to other

examples. Insisting on the naive relation (36) allows one to extract an effective value of α_s . Doing so for several $q\bar{q}$ resonances gives some idea of the sensitivity of $\text{br}(R \rightarrow g\bar{g})$ to scale.

It might be the case that the dynamics of the form factors, i.e., the functional dependence of the F_i 's on k_1^2 and k_2^2 and their relative normalization at the on-shell point, corresponds more accurately than does their overall normalization, in going from $\langle R \rightarrow j\bar{j} \rangle$ to $\langle g\bar{g}R \rangle$ ⁵. This could in principle be tested by measuring the on-shell form factors in a collider and using this dependence to predict the $H_J(x)$'s appearing in eq. (1). Once $\text{br}_{\text{rad}}(Q\bar{Q} \rightarrow V \rightarrow R)$ is measured for both $J=$ and radiative decay to a given $q\bar{q}$ meson, one can infer H_J at two values of x . With several related $q\bar{q}$ mesons of different masses, such as $f_2(1270)$ and $f_2(1520)$, this will give a number of points in $H_J(x)$.

3 Constraints From Radiative Quarkonium Decay

3.1 J^P dependence of $Q\bar{Q} \rightarrow V \rightarrow R_{J^P}$

The loop integral in eq. (4) determines the function $H_J(x)$ appearing in (1). For the NRQM wavefunctions these integrals have been evaluated [19, 32] in analytical form and are recorded in the appendix for convenience. Readers interested in the derivation of the analytic expressions are referred to those papers. The relevant functions $xH_J(x)^2$ are shown in Fig. 1 for $J^{PC} = 0^{++}; 2^{++}$ and 0^+ . The corresponding c_R 's in Eq. 1 are:

$$c_R = \begin{cases} 8 & J^{PC} = 0^+ \\ \geq 1 & J^{PC} = 0^{++} \\ \geq \frac{2}{3} & J^{PC} = 2^{++} \\ \geq \frac{5}{2} & J^{PC} = 2^{++} \end{cases} \quad (37)$$

⁵E.g., for $q\bar{q}$ states, the overall normalization of $\langle g\bar{g}R \rangle$ contains a factor α_s and is necessarily scale dependent.

In the x regime of immediate interest, $x = 0.5 - 0.75$, we note from Fig. 1 that $\frac{x \mathcal{H}_J}{30 \cdot 45} \ll 1$. This enables us to manipulate the CF expression, eq. (1), into a scaled form that exhibits the phenomenological implications immediately. Specifically, for scalar mesons

$$10^3 \text{br}(J = \text{! } 0^{++}) = \left(\frac{m}{1.5 \text{ GeV}} \right) \left(\frac{R \text{! } gg}{96 \text{ MeV}} \right) \frac{x \mathcal{H}_S(x)^2}{35}; \quad (38)$$

This is to be compared with the analogous formula for a tensor meson:

$$10^3 \text{br}(J = \text{! } 2^{++}) = \left(\frac{m}{1.5 \text{ GeV}} \right) \left(\frac{R \text{! } gg}{26 \text{ MeV}} \right) \frac{x \mathcal{H}_T(x)^2}{34}; \quad (39)$$

For pseudoscalars we find:

$$10^3 \text{br}(J = \text{! } 0^+) = \left(\frac{m}{1.5 \text{ GeV}} \right) \left(\frac{R \text{! } gg}{50 \text{ MeV}} \right) \frac{x \mathcal{H}_{PS}(x)^2}{45}; \quad (40)$$

Having scaled the expressions this way, because $\frac{x \mathcal{H}_J}{30 \cdot 45} \ll 1$ in the x range relevant for production of 1.3 - 2.2 GeV states (see Fig. 1), we see immediately that the magnitudes of the branching ratios are driven by the denominators 96 and 26 MeV for 0^{++} and 2^{++} , and 50 MeV for 0^+ . Thus if a state R_J is produced in $J = \text{! } X$ at $O(10^3)$ then $(R_J \text{! } gg)$ will typically be of the order 100 MeV for 0^{++} , $O(25 \text{ MeV})$ for 2^{++} , and $O(50 \text{ MeV})$ for 0^+ .

This immediately shows why the 2^{++} qq states are prominent: A 2^{++} state with a total width of $O(100 \text{ MeV})$ (typical for 2^{++} qq 's in this mass range [2, 31]) will be easily visible in $J = \text{! } 2^{++}$ with branching fraction $O(10^{-3})$, while remaining consistent with

$$\text{br}(R \text{! } qq) \leq 0.1 - 0.2; \quad (41)$$

Eqs. 38 - 40 not only indicate which qq states will be prominent in $J = \text{! } R$, but they also help to resolve an old paradox concerning 0^{++} production. It was recognised early on that when the gluons in the absorptive part of $J = \text{! } gg$ are classified according to their J^{PC} , the partial wave with 2^{++} was predicted to dominate. The waves with 0^+ and 0^{++} were also predicted to be significant and of comparable strength to one another [20]. When extended

to include the dispersive part [18, 19] the 0^{++} was predicted to be prominent over a considerable part of the kinematic region of interest. States with $J = 3$ were predicted to have very small rate in this process. Experimentally, all but one of these appeared to be satisfied. There are clear resonant signals in 2^{++} and 0^{++} , and no unambiguous signals have been seen with $J = 3$. However no 0^{++} signal had been isolated.

From our relations above, we see that for a 0^{++} to be produced at the 10^{-3} level in $J = 0$ radiative decay it must either have a large gluonic content and width $O(100)$ MeV or, if it is a qq meson, it must have a very large width, > 500 MeV. Taking this into account, along with the following points, the puzzle of the absence of 0^{++} signal has been resolved:

(i): The width of 3P_0 qq is predicted to be ~ 500 MeV [2, 31]. Thus production at the level $br(J = 0 \rightarrow (gg)_0 \sim 10^{-3})$ is consistent with $br(R \rightarrow gg) = O(\frac{2}{9}) \sim 0.1 - 0.2$, but the ~ 500 MeV wide signal is smeared over a large kinematic (x) range.

(ii): The ~ 100 MeV wide $f_0(1500)$ signal seen in $J = 0 \rightarrow 4$ was originally misidentified as 0^{++} , but is now understood to be 0^{++} [3].

(iii): The $f_J(1710)$ which was originally believed to be $J = 2$ may contain a contribution with $J = 0$ [3, 5].

3.2 Flavor Mixing and the $f_2(1270)$ and $f_2(1520)$ qq States

We can test this formalism by applying it first to the well known quarkonium states $f_2(1270)$ and $f_2(1525)$. The above formulae have been derived for the case that the produced meson $R(q_i q_i)$ contains a single flavour, so we begin by considering what changes occur for a state of mixed flavour. We shall see that $br(J = 0 \rightarrow R)$ and $(R \rightarrow gg)$ depend on the mass and flavour of R , but in a common way such that $br(R \rightarrow gg)$ is universal, as summarized in eqs.38-40.

For a general qq resonance

$$R = \cos \theta \psi_{ni} + \sin \theta \psi_{si}; \quad (42)$$

where $n = \frac{u\bar{u} + d\bar{d}}{2}$. Allowing for flavour symmetry breaking

$$h_{gg} \psi_{si} = r_s^2 h_{gg} \psi_{di}; \quad (43)$$

Thus

$$h_{gg} R = \left(\frac{P}{2} \cos \theta + r_s^2 \sin \theta \right) h_{gg} \psi_{di}; \quad (44)$$

so

$$(R \rightarrow gg) = \left(\frac{P}{2} \cos \theta + r_s^2 \sin \theta \right)^2 (R \rightarrow dd) \rightarrow gg \quad (45)$$

and similarly

$$(V \rightarrow R) = (V \rightarrow R \rightarrow dd) \left(\frac{P}{2} \cos \theta + r_s^2 \sin \theta \right)^2; \quad (46)$$

Evidently, the flavour factors cancel out in derivation of the expressions of the previous sections and so apply immediately to states R of arbitrary flavour mixings. We can illustrate this with the 3P_2 states $f_2(1270)$ and $f_2(1520)$, for which $\cos \theta = 1$ and 0 respectively. From eq. 45 we have

$$(f_2(n\bar{n})(1270) \rightarrow gg) = (f_2(d\bar{d})(1270) \rightarrow gg) = 2 \quad (47)$$

and

$$(f_2(s\bar{s})(1520) \rightarrow gg) = (f_2(d\bar{d})(1520) \rightarrow gg) = r_s^2; \quad (48)$$

If $qq \rightarrow gg$ is flavour blind we expect $r_s^2 = 1$.

To confront these equations with data we use the measured radiative branching ratios [5]

$$10^3 \text{ br}(J = \rightarrow f_2(1270)) = 1.4 \pm 0.14 \quad (49)$$

and

$$10^3 \text{ br}(J = \rightarrow f_2(1520)) = 0.63 \pm 0.1; \quad (50)$$

From eq. (39), we have

$$(1270 \rightarrow gg) = 41 \pm 7 \text{ M eV}; \quad (51)$$

and

$$\Gamma(f_2(1520) \rightarrow gg) = 17 \pm 2 \text{ M eV} ; \quad (52)$$

Combining these results with the measured widths,

$$\Gamma_{\text{tot}}(1270) = 185 \pm 20 \text{ M eV} ; \quad \Gamma_{\text{tot}}(1520) = 76 \pm 10 \text{ M eV} ; \quad (53)$$

we find

$$\text{br}(f_2(1270) \rightarrow gg) / \text{br}(f_2(1520) \rightarrow gg) = 0.22 \quad (54)$$

which are as expected for established $q\bar{q}$ states, see eq.(1) and ref. [18]. Inter alia this supports the idea that glueball mixing is not prominent in the 2^{++} channel at these masses[2].

If the dependence on mass is weak in going from 1270 to 1520 MeV, eqs.47 and 48 imply

$$\frac{\Gamma(f_2(1270) \rightarrow gg)}{\Gamma(f_2(1520) \rightarrow gg)} = \frac{2}{r_s^2} ; \quad (55)$$

Inserting the widths from eqs. 51 and 52 we see that $r_s \approx 0.8 < 1$, and thus $q\bar{q}$ favor can be ignored to first approximation in this analysis. We will exploit this favor independence in a later section (4.1) to probe the structure of wavefunctions for potential $q\bar{q}$ mesons.

3.3 Radiative Upsilon Decay

No peaks are seen in the photon energy spectrum in inclusive $\Upsilon \rightarrow X$ at a branching fraction sensitivity of about 10^{-4} [5]. The following analysis suggests that with only a factor of a few improvement in sensitivity, many interesting states should become evident. We could use data on $\Upsilon \rightarrow R$ in two ways. Firstly, for $R = \text{cc}$, pQCD-NRQM predictions should be reliable. Testing those predictions tests the underlying assumption of this methodology: that pQCD provides an adequate description of the $\langle Q Q_V j gg \rangle$ amplitude. Secondly, production of a given resonance in $\Upsilon \rightarrow R_J$ depends on $H_J(x)$ at x much closer to 1. This allows a more detailed examination of the form factors, as well as probing the $x \rightarrow 1$ region where resummation of perturbation theory may be required for the 0^{++} and 2^{++} cases.

Let us begin by considering $\psi \rightarrow c\bar{c} + \gamma$, where the $Q\bar{Q}$ bound states are rather well understood [19]. The branching ratio is

$$\text{br}(\psi \rightarrow c\bar{c} + \gamma) = c_J \left(\frac{4}{5}\right)_{\text{R}} \frac{x \mathcal{H}_J(x)^2}{8(2-9)M^2} m \quad (56)$$

where $c_J = 1(c); \frac{2}{3}(0)$ and $\frac{5}{2}(2)$. In all these examples $x = 1 - m_c^2/M^2 = 0.9$ so from eq. 1 we have $x \mathcal{H}_J^2 = 54(c); 32(0); 37(2)$. If we use $s(m_b) = 0.18$ and include the one loop corrections from [33], we find

$$\text{br}(\psi \rightarrow \gamma\gamma) = \frac{4}{5} (1 - 2s) = 2.8\%$$

Identifying $(c\bar{c} \rightarrow \gamma\gamma)$ with $(c\bar{c} \rightarrow \text{light hadrons})$ implies

$$\text{br}(\psi \rightarrow 2\gamma) = \text{br}(\psi \rightarrow 0) = 0.9 \cdot 10^{-5}$$

and

$$\text{br}(\psi \rightarrow c) = (2.3 - 0.9) \cdot 10^{-5}$$

Although these predicted branching ratios are small, the photons are in a region of phase space where there is little background, so a relatively short period of dedicated running at a B factory should be adequate to observe these modes. Precision data on these transitions could both validate the pQCD analysis and give insights into higher order effects including the role of colour octet components in the wavefunctions.

Data on $\psi \rightarrow f_{0,2}(1270-1700)$ may also be obtained, replacing the present upper limits 10^{-4} [5]. The kinematics here are $x = 0.97$. In this region $\mathcal{H}_S(x)^2$ and $\mathcal{H}_T(x)^2$ are dominated by $\ln(1-x)$ divergences, and the leading order pQCD predictions become unreliable. We urge that studies of pQCD resummation be made in order to analyze this process and make predictions. Data on these processes could be used both to extract $\mathcal{H}_J(x)^2$ for phenomenological use as in sec. 4, and also permit detailed testing of pQCD resummation techniques. The qualitatively different behavior of the $x \rightarrow 1$ limits of $\mathcal{H}_J(x)$ for the 0^+ and $0; 2^{++}$ cases can also be exploited to this end.

3.4 1^{++} States

Before turning to our main topic of glueball candidates, we discuss briefly the possibility of applying this formalism to 1^{++} mesons. Since for them we cannot normalize the $\langle gg\mathcal{R} \rangle$ amplitudes using the procedure of sec. 3.1 which lead to eqs. (15 - 17) it is not certain that this is possible. However in the spirit of an effective lagrangian approach it might be appropriate to consider that gluons have an effective mass, so that the amplitude $\langle gg\mathcal{R} \rangle$ need not vanish at the on-shell point when one of the gluons is longitudinal. Making the further assumption that the pQCD-NRQM approximation of ref. [29] gives an adequate description of the 1^{++} total width, with an adjustable overall normalization, one can obtain a relation of the form of eq. (1) [18]. Substituting for $H_1(x)$ leads to the scaled formula

$$10^3 \text{br}(J=1^+ \rightarrow 1^{++}) = \left(\frac{m}{1.45 \text{ GeV}}\right) \left(\frac{R \rightarrow gg}{12 \text{ MeV}}\right) \frac{x^2 H_1(x)^2}{30} \quad (57)$$

where in this application $(R \rightarrow gg)$ is the total direct coupling to gluons, not literally the coupling to two massless gluons.

It is interesting to apply the above relation to the $f_1(1285)$, $f_1(1420)$ and $f_1(1530)$ states (see also ref. [19]). Only two isoscalar mesons can be accommodated in a quarkonium nonet and there has been considerable discussion as to which of the three axial states is the odd one out (and, if it exists, what its nature is). There has been no confirmed sign of $f_1(1530)$ in $J=1$ radiative decay whereas the $f_1(1420)$ and $f_1(1285)$ are both seen. Their branching ratios are respectively [5]

$$\text{br}(J=1^+ \rightarrow f_1(1285)) = (0.65 \pm 0.10) \cdot 10^{-3} \quad (58)$$

and

$$\text{br}(J=1^+ \rightarrow f_1(1420)) \quad \text{br}(f_1(1420) \rightarrow KK) = (0.83 \pm 0.15) \cdot 10^{-3} : \quad (59)$$

Inserting these values into eq. 57 together with $\Gamma_{\text{tot}}(f_1(1285)) = 24 \pm 3 \text{ MeV}$ and $\Gamma_{\text{tot}}(f_1(1420)) = 52 \pm 4 \text{ MeV}$ give

$$\text{br}(f_1(1285) \rightarrow gg) = 0.34 \pm 0.05 \quad (60)$$

and

$$\text{br}(f_1(1420) \rightarrow gg) = (0.19 \pm 0.04) = (\text{br}(f_1(1420) \rightarrow K\bar{K})) : \quad (61)$$

If we now input $\text{br}(f_1(1420) \rightarrow K\bar{K}) = 0.67$ [34] we find

$$\text{br}(f_1(1420) \rightarrow gg) = 0.3 : \quad (62)$$

Within the uncertainties of applying these ideas to low masses (e.g. $f_1(1285)$ has a low width due to phase space suppression of $K\bar{K}$) and the ill defined branching ratio to $K\bar{K}$ for the $f_1(1420)$, these results are not inconsistent with the accepted qq interpretation of the $f_1(1285)$ and support also the quarkonium interpretation of $f_1(1420)$ (unless the $\text{br}(K\bar{K})$ should turn out to be much overestimated).

The ratio

$$\frac{\text{br}(J^= \rightarrow f_1(1420))}{\text{br}(J^= \rightarrow f_1(1285))} = 1.9 \quad (63)$$

is consistent with the quarkonium mixing arising from a quadratic mass formula for the axial nonet [34]

$$f_1(1285) = 0.94 \eta_{ni} - 0.35 \eta_{si} - 0.57 \eta_{li} + 0.83 \beta_i \quad (64)$$

$$f_1(1420) = 0.35 \eta_{ni} + 0.94 \eta_{si} - 0.83 \eta_{li} - 0.57 \beta_i \quad (65)$$

Recent data from BES [35] have large error bars but are consistent with the older data for $f_1(1420)$; they obtain $\text{br}(J^= \rightarrow f_1(1420)) = \text{br}(f_1(1420) \rightarrow K\bar{K}) = (0.76 \pm 0.46 \pm 0.18) \cdot 10^{-3}$, $m = 59 \pm 5$ MeV. They do not see any $f_1(1285)$ but this may not be surprising since they are looking in the $K\bar{K}$ mode. They also report a signal $f_1(1497)$, $m = 44 \pm 7$ MeV and $\text{br}(J^= \rightarrow f_1(1497)) = \text{br}(f_1(1497) \rightarrow K\bar{K}) = (0.52 \pm 0.23) \cdot 10^{-3}$. This state's parameters are also consistent with those expected for a quarkonium as long as $\text{br}(f_1(1497) \rightarrow K\bar{K}) = 0.5$.

The axial mesons are currently an enigma. There are three candidates where the quark model would require only two. The lattice predicts the lightest 1^{++} glueball to be at ≈ 4 GeV [15]. It is noticeable that no single experiment

sees all three and one should be cautious as to whether there are indeed three genuine states. We urge that BES, in particular, seek three f_1 signals or place limits against them in order to help clarify the above analysis. In any event, more detailed theoretical work, specifically formalizing the effective lagrangian treatment of the problem, is warranted in order to relate to the production of axial mesons in and to provide a more solid foundation to the theoretical analysis after the experimental situation becomes clear.

4 Glueball Candidates

4.1 $f_0(1500)$

We look first at the established scalar meson, $f_0(1500)$. As we shall see below, at present there are discrepancies between the values of $\text{br}(J = 0^+ \rightarrow f_0(1500))$ as determined from various experimental analyses.

The analysis of ref. [3] gives $\text{br}(J = 0^+ \rightarrow f_0(1500) \rightarrow \text{gg}) = (5.7 \text{--} 0.8) \times 10^{-4}$ with an overall 15% normalization uncertainty. The analysis of ref. [36, 37] implies that the 0^+ mode is at most 50% branching ratio and so we infer

$$\text{br}(J = 0^+ \rightarrow f_0(1500)) = (1.15 \text{--} 0.15) \times 10^{-3} \quad 15\%$$

In this case, with $\Gamma_{\text{tot}}(f_0(1500)) = 120 \text{--} 20 \text{ MeV}$, if we add errors in quadrature and use the central value, eq. 38 implies that

$$\text{br}(f_0(1500) \rightarrow \text{gg}) = 0.9 \text{--} 0.2 \quad (66)$$

This is significantly larger than the $O(\frac{2}{3})$ which would be expected for a pure $q\bar{q}$ system, and supports this state as a glueball candidate.

On the other hand BES has recently reported [38] $\text{br}(J = 0^+ \rightarrow f_0(1500) \rightarrow \pi^0 \pi^0) = 3 \text{--} 5 \times 10^{-5}$. Landua [38] combines this with the Crystal Barrel data on $\text{br}(f_0(1500) \rightarrow \text{gg})$ to get

$$\text{br}(J = 0^+ \rightarrow f_0(1500)) = (0.4 \text{--} 0.6) \times 10^{-3}$$

Thus via eq. (38),

$$\text{br}(f_0(1500) \rightarrow gg) = 0.3 \pm 0.5; \quad (67)$$

The interpretation of this state cannot be settled until the experimental situation clarifies. An order of magnitude increase in statistics for $J=0$ will enable extension of the analysis shown in Fig. 2 of ref. [3], and $J=0$ likewise needs to be improved. The neutral channel $J=0 \rightarrow \pi^0 \pi^0 \pi^0$ is particularly advantageous here as it is free from contamination and so can help to improve the quantification of $\text{br}(f_0(1500) \rightarrow \pi^0 \pi^0 \pi^0)$. These should be high priorities at a Υ -charm Factory.

We shall return to the interpretation of the $f_0(1500)$ in section 6.

4.2 $f_J(1710)$

The case of $f_J(1710)$ is particularly interesting and the conclusions depend critically on whether $J=0$ [3, 39] or $J=2$ [5, 40, 41]. It has been observed most clearly in radiative $J=0$ decay in the KK mode [40], with evidence also in the 4π mode [3]. Recently BES has reported seeing both $J=0$ and $J=2$ states in this region. We discuss the various measurements in turn.

In the KK channel, $\text{br}(J=0 \rightarrow f_J(1710) \rightarrow KK) = (0.97 \pm 0.12) \times 10^{-3}$ [5]. Assuming first that $f_J(1710)$ is a single state with $J=2$, we use eq. (39):

$$10^3 \text{br}(J=0 \rightarrow f_2(1710)) = \left(\frac{m}{1.5 \text{ GeV}}\right) \left(\frac{R \rightarrow gg}{26 \text{ MeV}}\right) \frac{\chi^2_{\text{HT}}(x) f^2}{34};$$

which implies

$$\text{br}(f_2(1710) \rightarrow gg) = \frac{(22 \pm 3) \text{ MeV}}{\text{br}(f_2(1710) \rightarrow KK)}; \quad (68)$$

No comparable signal has been seen in any other channel in $J=0 \rightarrow f_2(1710) \rightarrow X$ and it would thus appear that KK is a major mode of any $J=2$ object in $J=0$ radiative decays (the listing of decay channels for $J=0 \rightarrow f_J(1710) \rightarrow KK$ in ref. [5] suggests that this mode is greater than 50% of

the $K K + +$ channels together). With $\Gamma_{\text{tot}}(f_J(1710)) = 150 \text{ M eV}$, eq. 68 and $\text{br}(f_2(1710) \rightarrow K K) = 0.5$ imply

$$\text{br}(f_2(1710) \rightarrow gg) < 30\%; \quad (69)$$

which would be consistent with this state being a qq .

By contrast, if $f_J(1710)$ is a single state with $J = 0$, we use eq. (38):

$$10^3 \text{br}(J=0 \rightarrow 0^{++}) = \left(\frac{m}{1.5 \text{ GeV}}\right) \left(\frac{\Gamma_{R \rightarrow gg}}{96 \text{ MeV}}\right) \frac{x \mathcal{H}_s(x) f_J^2}{35};$$

which implies

$$\text{br}(f_0(1710) \rightarrow gg) = \frac{(78 - 10) \text{ MeV}}{\text{br}(f_0(1710) \rightarrow K K)} \quad (70)$$

and hence

$$\text{br}(f_0(1710) \rightarrow gg) = 0.52 - 0.07; \quad (71)$$

in accord with eq. 14 of ref. [18]. In this case the $f_0(1710)$ would be a strong candidate for a scalar glueball. Knowing the spin and $K K$ branching fraction of $f_J(1710)$ is of great importance for a more detailed quantitative understanding of the composition of this state.

These questions have become central in view of new data from BEPC [42] which, for the first time, separates a $J = 2$ and $J = 0$ signal from the (1710) region. They find an $f_2(1696)$ with $\Gamma = 103 \pm 18 \text{ MeV}$ and $\text{br}(J=0 \rightarrow f_2)$ $\text{br}(f_2 \rightarrow K^+ K^-) = 2.5 - 0.4 (10^{-4})$. They also find an $f_0(1780)$ with $\Gamma = 85 \pm 25 \text{ MeV}$ and $\text{br}(J=0 \rightarrow f_0) = \text{br}(f_0 \rightarrow K^+ K^-) = 0.8 - 0.1 (10^{-4})$. These signals are weak, $O(10^{-4})$, in contrast to the $O(10^{-3})$ reported in the earlier literature cited above. The BEPC $J = 2$ state is consistent with a (radial excited) qq . Their $J = 0$ state strength appears too feeble for a glueball, unless $K^+ K^-$ is a minor decay mode.

If the BEPC data are definitive, then the possibility that $\text{br}(f_0(1780) \rightarrow K^+ K^-)$ is small merits investigation. In this context we note that ref. [3] analyzes $J=0 \rightarrow 4$ and finds a signal at about 1750 MeV consistent with 0^{++} , although 2^{++} is not absolutely excluded. If interpreted as f_0 , the width is $\Gamma =$

160 MeV and branching fraction in $J = 1$ $f_0(1500) \rightarrow 4$ is $(0.9 - 0.13) \times 10^{-3}$ [3]. Thus the analysis of ref [3] indicates that there is a scalar signal in $J = 1$ 4 at strength characteristic of gluonic states. We urge that BEPC investigate the 4 channel to see if their scalar state is visible at a level consistent with the above analysis of ref.[3].

A possible explanation of the observations, if both $f_0(1500)$ and $f_0(1710^{00})$ are produced at the 10^{-3} level in $J =$ radiative decay, is that both of them contain both qq and gg components [2, 43, 44, 45]. Better data, especially on the decay branching fractions and production in radiative $J =$ decay is crucial for resolving this question. We will consider complementary tests for this, through $\pi\pi$ production, in section 5. We examine mixing phenomenology in section 6.

4.3 (2230) Tensor Glueball candidate

The appearance of a narrow state (2230) in $J = 1$ $^{++}$; K^+K^- ; $K_s^0K_s^0$; pp has created considerable interest [6]. In each of these channels the branching ratios are typically $br(J = 1 \rightarrow XX) \sim 3 \times 10^{-5}$ for each of the channels where $X = K^+$ or K_s^0 , and 1.5×10^{-5} for pp . In all channels the signal is consistent with $\Gamma_{tot} \sim 20$ MeV. After allowing for associated neutral modes such as $\pi^0\pi^0$, and nn by isospin, this gives

$$br(J = 1 \rightarrow gg) \sim 0.1 \times 10^3 : \quad (72)$$

When combined with our formulae eqs. 38 and 39 this implies that $br(1 \rightarrow gg) \sim 0.4$ for $J = 0$ and 0.15 for $J = 2$. However eq. (72) is likely to be a gross underestimate because $\pi\pi$ and multibody channels were not included. Indeed, the absence of a signal in PS185 at CERN [46], suggests that two body final states constitute no more than 10% of the total. In view of the uncertainties in the measured quantities this gives $br(1 \rightarrow gg)$ consistent with unity for $J = 2$; for $J = 0$, it unacceptably exceeds the unitarity bound.

Thus we suggest that if evidence for the (2230) survives increases in statistics, the case $J = 2$ would be consistent with (2230) being a tensor glueball.

Such a result would have significant implications for the emergence of a glueball spectroscopy in accord with lattice QCD. It would also raise tantalising questions about the 0^+ sector, where lattice finds a glueball mass $\sim 2 \text{ GeV}$. This is interesting in view of the appearance of a clear 0^+ signal in the 1450 MeV region, which would then be difficult to reconcile with being the 0^+ glueball. We now turn to this question.

4.4 0^+ signals in $J=1$ R

The branching ratio for $(QQ)_V \rightarrow R$ in terms of the total gluonic decay width of the pseudoscalar state is given by eqs. 1 and 37

$$b_{\text{rad}}((QQ)_V \rightarrow R) = \frac{\alpha_s^2 \chi_{PS}^2(x) f^2 m}{8 (2-9) M^2} \quad (73)$$

As noted in eq. (40), the above formula may be scaled as follows

$$10^3 \text{br}(J=1 \quad 0^+) = \left(\frac{m}{1.5 \text{ GeV}}\right) \left(\frac{\alpha_s}{50 \text{ MeV}}\right) \frac{\alpha_s^2 \chi_{PS}^2(x) f^2}{45} \quad (74)$$

This subsumes eqs 2,3,5,6, 9 and 10 of ref.[18]. (Note that the dashed curve in fig 8 of ref. [18] corresponds to the above; the figure caption has typographical errors).

As a consistency test of this methodology for 0^+ states, we consider the production of the radial qq anticipated in this mass region⁶. The state (1295) ($m = 536 \text{ MeV}$, with dominant decay into $\pi\pi$) is a candidate on the grounds of mass and width [31]. The DM2 collaboration [47] may have evidence for the (1295) in their $J=1$ data, which contains a peak in $\pi\pi$ with the parameters $J^{PC} = 0^+$, $m = 1265$, $\Gamma = 44 \pm 20 \text{ MeV}$; $\text{br} = (0.26 \pm 0.06) \times 10^{-3}$. If this is the (1295), the scaled formula (40) then implies that $\text{br}((1295) \rightarrow gg) \approx 0.25$ if $\pi\pi$ is the dominant decay mode. This is consistent with a 0^+ (qq) because $\pi\pi$ dominance is expected [31].

⁶Note that the model cannot safely be applied to the 0^+ , whose total width is "accidentally" strongly suppressed because (apart from $\pi\pi$, which is suppressed by three body phase space) only electromagnetic decays are non-negligible

At slightly higher mass, the $(1440)^+$ [5] is more prominently produced. Historically this was seen in $J^P = 1^- K\bar{K}$ with $br = (4.3 \pm 1.7) \times 10^{-3}$ [48]. The prominence of this state caused it to be identified as a potential glueball [7]. Subsequently it was realised that there are two states contained within this structure [5] whose individual production rates were smaller than the earlier, apparently large value. This development, together with improved lattice QCD estimates of the 0^+ glueball mass which place it above 2 GeV rather than in the 1.4 GeV region, caused the glueball interpretation to fall from favour. As noted in ref. [18], such a large production rate as originally reported [48] would seriously oversaturate $br((1440)^+ \rightarrow gg)$, but subsequent separation of the signal into two resonances results in physically acceptable values for the individual gg widths.

The first analyses [49, 47] indicating the existence of additional structure in the $(1440)^+$ region were, however, not in agreement. Recent data in $pp \rightarrow (1440)^+ \rightarrow gg$ help us to identify the problematic measurement and to propose a consistent picture that experiments should now pursue. We shall suggest that there are two states, L and H (for "Low" and "High" mass respectively), where L has significant coupling to glue while H is dominantly the ss member of the nonet, mixed with glue. Before giving the theoretical analysis, we survey the evidence from various experiments for L ($1410)^+$ and $K\bar{K}$, $\Gamma_{tot} = 50$ MeV and for H ($1480)^+ \rightarrow K\bar{K}$, $\Gamma_{tot} = 100$ MeV.

Oberlies [13] sees two states in $pp \rightarrow L, H \rightarrow (K\bar{K})$ with the properties

$$L(1416 \pm 2)^+ \rightarrow K\bar{K}; \Gamma_{tot} = 50 \pm 4 \text{ MeV} \quad (75)$$

$$H(1460 \pm 10)^+ \rightarrow K\bar{K}; \Gamma_{tot} = 105 \pm 15 \text{ MeV} \quad (76)$$

These values agree with the central values for the sighting by Mark III [49] in $J^P = 1^-$ radiative decay. Combining errors in quadrature Mark III finds

$$L(1416 \pm 10)^+ \rightarrow a_0 \rightarrow K\bar{K}; \Gamma_{tot} = 54^{+40}_{-30} \text{ MeV} \quad (77)$$

$$H(1490 \pm 18)^+ \rightarrow K\bar{K}; \Gamma_{tot} = 91 \pm 68 \text{ MeV} \quad (78)$$

Further evidence for the low mass state, in the decay channel $\rightarrow \pi\pi$, comes from Mark III [50] who find $(1400 \pm 6)^+$, $\Gamma_{tot} = 47 \pm 13$ MeV; from DM2 [47] who find

(1398 6), $\Gamma_{\text{tot}} = 53 \pm 11 \text{ M eV}$; and the Crystal Barrel Collaboration [14]. The latter see pp π^0 (1410 ± 3) π^0 (π^0) with significant contribution in the glue favoured partial wave $[2]$. Their value $\Gamma_{\text{tot}} = 86 \pm 10 \text{ M eV}$ is however substantially larger than the $\Gamma_{\text{tot}} = 50 \text{ M eV}$ found by the other experiments [13, 47, 50]. Possibly the differing production mechanisms of these experiments affects the apparent width of the resonance due to differing interference effects.

We now compute the production rate for these states in $J^P = 0^-$ radiative decay. For the π_L, π_M arkIII [49] see the decay mode $a_0 \rightarrow K^+ K^-$ with

$$\text{br}(J^P = 0^- \pi_L (1410) \rightarrow K^+ K^-) = (0.66^{+0.29}_{-0.22}) \cdot 10^{-3}; \quad (79)$$

while a clear signal is found also in the $a_0 \rightarrow \pi^+ \pi^-$ channel by ref. [50]:

$$\text{br}(J^P = 0^- \pi_L (1410) \rightarrow \pi^+ \pi^-) = (0.34 \pm 0.08) \cdot 10^{-3}; \quad (80)$$

These two channels are expected to dominate the decays. Adding them together and inserting into eq. (74) implies

$$\Gamma(\pi_L (1410) \rightarrow gg) = (54 \pm 13) \text{ M eV}; \quad (81)$$

Combining the various width measurements, adding errors in quadrature, gives $\Gamma_{\text{tot}} = 54.2 \pm 3.4 \text{ M eV}$, and hence

$$\text{br}(\pi_L (1410) \rightarrow gg) = 0.9 \pm 0.2; \quad (82)$$

For comparison, using just the larger width from Crystal Barrel [14] would give $\text{br}(\pi_L (1410) \rightarrow gg) = 0.65 \pm 0.2$. The data clearly indicate a strong coupling to gluons which argues against $\pi_L (1410)$ being pure $q\bar{q}$. Rather, it couples like a glueball, perhaps mixed with the nearby $q\bar{q}$ nonet.

Now we consider the $\pi_H (1480)$. This state decays into $K^+ K^-$ and is not seen in $\pi^0 \pi^0$. Combining errors in quadrature, as above, MarkIII finds [49]

$$\text{br}(J^P = 0^- \pi_H (1490) \rightarrow K^+ K^-) = 1.03^{+0.33}_{-0.26} (10^{-3}) \quad (83)$$

⁷For this reason the DM 2 analysis [47] indicating $(1460) \rightarrow a_0 \rightarrow K^+ K^-$ is suspect, since a genuine resonance decaying to a_0 should also show up in $a_0 \rightarrow \pi^+ \pi^-$. Instead, in the $\pi^0 \pi^0$ channel only the (1410) is seen. Eliminating this DM 2 state produces a harmonious picture given the remaining observations. We thank A. Kirk for discussions of this point.

whereby with $\Gamma_{\text{tot}} = 100 \text{--} 20 \text{ MeV}$ [13], eq. (74) implies

$$\text{br}(\chi(1490) \rightarrow gg) = (0.5 \text{--} 0.2) = \text{br}(K \bar{K}) \quad (84)$$

Ref[31] anticipates that the radially excited ss should have a total width of up to 100 MeV, dominated by the channel $K \bar{K}$. The above result, eq. (84), is compatible with such a state. Ref.[31] also finds that nn has suppressed width, decaying into $\pi\pi$ through a_0 ; this is compatible with the results on $\chi(1295)$ above. Thus a tentative interpretation of the pseudoscalar states is as follows:

$$\begin{aligned} (1295) \quad & ^{nn}; \text{br}_{gg} \approx 0 \left(\frac{2}{3} \right) \approx 0.25 \\ & \chi_L(1410) \quad G(+qq); \text{br}_{gg} \approx 1 \\ & \chi_H(1480) \quad ^{ss}(+G); \text{br}_{gg} \approx 0.5 \end{aligned} \quad (85)$$

These conclusions can be sharpened if the widths and decays from Crystal Barrel and Obelix converge and if $J = 0^+$ is pursued further. We note also a new measurement from BEPC [35] which sees only a single state with a mass of $1467 \pm 3 \text{ MeV}$, $\Gamma = 89 \pm 6 \text{ MeV}$ and $\text{br}(\chi(1467) \rightarrow K \bar{K}) = 1.86 \pm 0.10 \pm 0.4(10^3)$. Since $\text{br}(\chi(1467) \rightarrow K \bar{K}) \approx 1$, this gives $\text{br}(\chi(1467) \rightarrow gg) \approx 1.1 \pm 0.2$. We urge that BEPC continue to investigate this state with a view to separating two signals: $\chi_H \rightarrow K \bar{K} \rightarrow K \bar{K}$ and $\chi_L \rightarrow a_0 \rightarrow K \bar{K}$.

The experimental data on 0^+ production in radiative $J = 0$ decays in this mass region need clarification before strong conclusions can be drawn, but if the existence of two states in the 1400–1500 MeV range, and their relative production (one or both much more strongly produced than $\chi(1295)$) is confirmed, we have a serious challenge to theoretical expectations. The experiments would appear to be telling us that the lightest pseudoscalar glueball is much lighter than predicted in quenched lattice QCD, (2.16–0.27 GeV [43]). In view of the apparent possible success (within uncertainties noted above) of the lattice QCD predictions for the 0^{++} and 2^{++} glueball masses, such a discrepancy between lattice QCD and nature would be of great interest. We note

that the mass and properties of the (1410) are consistent with predictions for a gluino-gluino bound state [17, 18], possibly mixed with nearby pseudoscalar qq states. If nature were supersymmetric and SUSY breaking did not violate R-invariance, the gluino mass would be $O(100)$ MeV [51, 52]. In that case the 0^{++} glueball would be in an approximate supermultiplet with the pseudoscalar gluino-gluino ($\tilde{g}\tilde{g}$) and spin-1/2 gluon-gluino bound states. This would lead to an "extra" isosinglet pseudoscalar in the spectrum, with mass around $1\frac{1}{2}$ GeV. Decay of such a $\tilde{g}\tilde{g}$ system would necessarily go through gluons, since its direct couplings to quarks would be suppressed by heavy squark masses and hadrons containing a single gluino would be too massive to be pair produced by the $\tilde{g}\tilde{g}$. Thus $br(\tilde{g}\tilde{g} \rightarrow gg) \approx 1$.

Improving the data on these states would provide important constraints. It is now a clear challenge for experiment to separate and quantify these signals.

5 Constraints on Glueballs from Widths

5.1 $R \rightarrow 1$ and $J=0 \rightarrow R$

If a state R_J is a glueball (or light gluinoball), it will occur in $1 \rightarrow R_J$ as a singleton and be strongly suppressed in $R_J \rightarrow 1$. By contrast, if R_J is an $I=0$ member of a qq nonet there will be two orthogonal states in the singlet-octet flavour basis available for production both in $J=0 \rightarrow R_J$ and $R_J \rightarrow 1$. Flavour mixing angles may suppress one or the other of the pair in either $R_J \rightarrow 1$ or in $J=0 \rightarrow R_J$ but there are strong correlations between the two processes so that a comparison of the two processes can help to distinguish glueball from qq . In particular, if a qq state is flavour "favoured" in $J=0 \rightarrow qq$, so that it is prominent and super-cially somewhat "glueball-like", it will also be flavour favoured in $1 \rightarrow R(qq)$ (see below) in dramatic contrast to a glueball.

For heavy QQ resonances such as the χ_c states, the amplitudes $\langle gg | R \rangle$, which enter the computation of $b_{rad}(QQ \rightarrow 1 + R)$, and $\langle R | j \rangle$, which

in principle can be measured as a function of photon virtualities, are identical except for the value of the overall coefficient A_{J^PC} . The relative rates (see eq. (36) and sec. 2.4) would be (in leading order)

$$(R_{\text{I}}) = \frac{9e_q^4}{2} \left(\frac{1}{s}\right)^2 (R_{\text{I}})_{gg}$$

where e_q is the relevant quark charge. Bearing in mind the limitations to use of this relation for light qq mesons discussed in sec. 2.4, (see also ref.[53]) we shall tentatively adopt this relation and test it against the known $f_2(1270;1525)$. Finding it to be qualitatively reasonable, we apply it in section 6 to $f_0(1370;1500;1710)$, allowing for mixing between nn , ss , and gg .

5.2 Orthogonal qq mesons coupling to $\gamma\gamma$ and gg

Define

$$R_{\text{I}} = \cos \beta |i\rangle + \sin \beta |i\rangle \quad (86)$$

$$R_{\text{II}} = \cos \beta |i\rangle - \sin \beta |i\rangle \quad (87)$$

where $|i\rangle, |8_i\rangle$ denote the $SU(3)$ flavor qq states. (This is a more natural basis for what follows than the ideal flavour basis used in section 3.2). Then in terms of the intrinsic rates for a single qq flavour ($uu; dd$ or ss assumed to be of equal strength)⁸

$$(R_{\text{I}})_{gg} = (qq)_{gg} \cdot 3 \cos^2 \beta \quad (88)$$

and

$$(R_{\text{II}})_{gg} = (qq)_{gg} \cdot 3 \sin^2 \beta; \quad (89)$$

while the widths are in a different proportion. Defining $(qq)_{\gamma\gamma}$ to be the width for quarks of unit electric charge:

$$(R_{\text{I}})_{\gamma\gamma} = (qq)_{\gamma\gamma} \left(\cos^2 \frac{2}{3} + \sin^2 \frac{1}{3} \right)^2 \quad (90)$$

⁸The factor 3 in this equation reflects the $1 = \frac{1}{3}$ projection of each of the three qq flavours in the flavour singlet state.

and

$$(R_{II}!) = (qq!) \left(\sin^2 \frac{2}{3} + \cos^2 \frac{1}{3} \right)^2: \quad (91)$$

In a form that shows the relation to the gg widths,

$$(R_I!) = (qq!) \frac{1}{6} \cos^2(\quad) \quad (92)$$

and

$$(R_{II}!) = (qq!) \frac{1}{6} \sin^2(\quad); \quad (93)$$

where $\tan^{-1} \frac{1}{2} = 19.5^\circ$.

It is clearly possible for an individual qq to decouple "accidentally" in gg or if 0 or 19.5. However for the orthogonal system we have the sum rule

$$(R_I!) + (R_{II}!) = \frac{1}{6} (qq!) \quad (94)$$

and

$$(R_I! gg) + (R_{II}! gg) = 3 (qq! gg): \quad (95)$$

Thus using eq. 36 and including the next order QCD corrections,

$$\frac{(R_I!) + (R_{II}!)}{(R_I! gg) + (R_{II}! gg)} = \frac{2}{4} (1 + c_s)^{-1} \quad (96)$$

where $c_s = 0.4$ for tensors and $8/6$ for scalars [29, 54].

In the case of tensors the input data are

$$(f_2(1270) + f_2(1525))! = 3.0 \quad 0.4 \text{ keV} \quad (97)$$

and, from our analysis in section 3.2,

$$(f_2(1270) + f_2(1525))! gg = 58 \quad 8 \text{ M eV} : \quad (98)$$

With these widths, eq. 96 gives $\frac{eff}{s} = 0.48 \pm 0.05$, not unreasonable for this mass region [55].

Considering the ratio of the and gg widths of the entire orthogonal system allowed us to extract $\frac{eff}{s}$, with little sensitivity to . We can instead

employ ratios of the R_I and R_{II} and gg widths to extract with little sensitivity to s_s^{eff} . For an orthogonal qq pair, eqs. 90-93 imply

$$\frac{(R_I \rightarrow gg)}{(R_{II} \rightarrow gg)} = \frac{1}{\tan^2} \quad (99)$$

while

$$\frac{(R_I \rightarrow \text{glueball})}{(R_{II} \rightarrow \text{glueball})} = \frac{1}{\tan^2(\theta)} \quad (100)$$

Note that the two sets of equations will not give the same value of θ if our procedure is not valid. As a consistency check, we determine θ both ways for the f_2 states. The $J=1$ R data gave us

$$\frac{(f_2(1270) \rightarrow gg)}{(f_2(1525) \rightarrow gg)} = \frac{41 \text{ } 7\text{M eV}}{17 \text{ } 2\text{M eV}} \Rightarrow \theta = (33 \pm 2)^\circ; \quad (101)$$

while θ data give

$$\frac{(f_2(1270) \rightarrow \text{glueball})}{(f_2(1525) \rightarrow \text{glueball})} = (26.2 \pm 2.8) \pm 26\% \Rightarrow \theta = (30.5 \pm 2)^\circ; \quad (102)$$

The consistency of these results encourages us to apply the ideas to scalar mesons. However, the presence of possibly three scalar states in close proximity, $f_0(1370;1500)$ and $f_{J=0?}(1710)$, and in the vicinity of the lattice scalar glueball, suggests that mixing involving both qq and gg will be essential. We shall now consider this situation.

5.3 qq nonet and glueball coupling to glueball and gg

If the qq nonet, $R_{I,II}$ is in the vicinity of a glueball, G , the above analysis requires generalisation. Three isoscalars arise. With $R_{I,II}$ as above, the mixed states may be written

$$\begin{aligned} 3 &= \cos \theta_{II} i \sin \theta_{I} j \\ 2 &= \cos \theta_{I} i \sin \theta_{II} j + \cos \theta_{II} i + \sin \theta_{I} j \\ 1 &= \sin \theta_{I} i + \cos \theta_{II} j + \cos \theta_{II} i + \sin \theta_{I} j \end{aligned} \quad (103)$$

If we ignore mass and phase space effects, and any differences between the nn and ss wavefunctions, then proceeding as in the previous section we obtain

$$\begin{aligned} ((\gamma_1 + \gamma_2)!) &= (\alpha\alpha!) \left(\frac{1}{6} \cos^2(\gamma) + \frac{1}{6} \sin^2(\gamma) \right) \\ (\gamma_3!) &= (\alpha\alpha!) \frac{1}{6} \sin^2(\gamma) \cos^2(\gamma) \end{aligned} \quad (104)$$

Defining $\sum_{i=1}^3 (\gamma_i!)$ and later γ_{gg} analogously, the generalisation of eq.(94) becomes

$$= \frac{1}{6} (\alpha\alpha!): \quad (105)$$

The generalisation of the relation for the gluon couplings, eq(95), becomes

$$\gamma_{gg} = 3 (\alpha\alpha! \gamma_{gg}) + (G! \gamma_{gg}): \quad (106)$$

Consequently

$$= \frac{2}{4} (1 + \cos^2(\gamma))^{-1} (\gamma_{gg} (G! \gamma_{gg})): \quad (107)$$

Thus, specialising to 0^{++} mesons, the experimentally measurable quantities and γ_{gg} obey the relation

$$[\text{keV}] \frac{(0.5 \gamma_s)^2}{20 (1 + 8 \cos^2(\gamma))} \gamma_{gg} [\text{MeV}]: \quad (108)$$

A major uncertainty comes from the large higher order QCD correction for the 0^{++} sector which reduces the right hand side by a factor of approximately 2.2 To be conservative we therefore work to leading order. If f_0 (1370) is one of the trinity of glue associated states, then we infer from $(f_0!) = 5.4 \pm 2.3$ keV [5] that

$$\gamma_{gg} = 108 \pm 46 \text{ MeV} \quad (109)$$

or from eq(38), neglecting mass dependence:

$$\text{br}(J = ! + \sum_{i=1}^3 f_0^i) = (1.1 \pm 0.5) 10^{-3}: \quad (110)$$

Since $\text{br}(J = ! f_1(1710)) = 0.97 \pm 0.12 \cdot 10^{-3}$ [5], this bound is satisfied by present data if the 1710 has $J = 0$, even if $\text{br}(f_0(1710)) = 1$

and there is negligible production of $f_0(1500)$ in radiative decay. However the limit it is only barely respected so unless $\text{br}(J=0 \rightarrow f_0(1500)) + \text{br}(J=0 \rightarrow f_0(1710))$ is very small, either $\text{br}(J=0 \rightarrow f_0(1500))$ and or $\text{br}(J=0 \rightarrow f_0(1370))$ must be non-negligible, or $\text{br}(f_0(1710) \rightarrow KK) < 1$. If the 1710 state proves to have $J=2$, the bound (110) will be very stringent indeed. We now consider specific examples of mixing in the $f_0(1370;1500;1710)$ system.

6 Three-State Mixings

An interesting possibility is that three f_0 's in the 1.4 – 1.7 GeV region are admixtures of the three isosinglet states gg , ss , and nn [2]. Recently there have been two specific schemes proposed which are based on lattice QCD and the emergent phenomenology of scalar mesons. In this section we present a simplified formalism for treating a three component system of this type.

At leading order in the glueball- qq mixing, ref[2] obtained

$$\begin{aligned} N_g \mathcal{G}_i &= \mathcal{G}_{0i} + \left(\frac{p}{2} \right) (j_{nni} + ! j_{ssi}) \\ N_s j_{ssi} &= j_{ssi} + ! \mathcal{G}_{0i} \\ N_n j_{nni} &= j_{nni} + \frac{p}{2} \mathcal{G}_{0i} \end{aligned} \quad (111)$$

where the N_i are appropriate normalisation factors, $! = \frac{E(G_0) E(dd)}{E(G_0) E(ss)}$ and the mixing parameter $\frac{p}{2} = \frac{E(dd) \mathcal{G}_{0i}}{E(G_0) E(dd)}$. Our analysis suggests that the $gg \rightarrow qq$ mixing amplitude manifested in $! = R(qq)$ is $O(\epsilon)$, so that qualitatively $O(\epsilon) \approx 0.5$. Such a magnitude implies significant mixing in eq.(111) and is better generalised to a 3×3 mixing matrix. Mixing based on lattice glueball masses lead to two classes of solution of immediate interest:

- (i) $! = 0$, corresponding to G_0 in the midst of the nonet [2]
- (ii) $! > 1$, corresponding to G_0 above the qq members of the nonet [45].

We shall denote the three mass eigenstates by R_i with $R_1 = f_0(1370)$, $R_2 = f_0(1500)$ and $R_3 = f_0(1710)$, and the three isosinglet states \mathcal{G}_i with

$f_1 = nn$, $f_2 = ss$ and $f_3 = gg$ so that $R_i = f_{ij} \psi_j$. Recent data on the decay $f_0(1500) \rightarrow K\bar{K}$ [38] may be interpreted within the scheme of ref[2] as being consistent with the parameter $f_2 = 0.5$. This enables simple analysis; if for illustration we adopt $f_1 = 0.5$, the resulting mixing amplitudes are (scheme "A"):

	f_{i1}	f_{i2}	f_{i3}
$f_0(1370)$	0.86	0.13	0.50
$f_0(1500)$	0.43	0.61	0.61
$f_0(1710)$	0.22	0.76	0.60

By contrast, Weingarten [45] has considered the case where the bare glueball lies above the ss member of the nonet. His mixing matrix is (scheme "B"):

	f_{i1}	f_{i2}	f_{i3}
$f_0(1370)$	0.87	0.25	0.43
$f_0(1500)$	0.36	0.91	0.22
$f_0(1710)$	0.34	0.33	0.88

The solutions for the lowest state are similar, as are the relative phases and qualitative importance of the G component in the high mass state. Both solutions exhibit destructive interference between the nn and ss favours for the middle state.

If we make the simplifying assumption that the photons couple to the nn and ss in direct proportion to the respective e_q^2 (i.e. we ignore mass effects and any differences between the nn and ss wavefunctions), then the corresponding two photon widths can be written in terms of these mixing coefficients:

$$(R_i) = f_{i1} \frac{5}{9} + f_{i2} \frac{1}{9} \quad ; \quad (112)$$

where Γ is the width for a qq system with $e_q = 1$. One can use eq. (112) to evaluate the relative strength of the two photon widths for the three f_0 states with the input of the mixing coefficients. These are (ignoring mass dependent effects)

$$f_0(1370) : f_0(1500) : f_0(1710) \quad 12 : 1 : 3 \quad (113)$$

in scheme A, to be compared with

$$f_0(1370) : f_0(1500) : f_0(1710) \quad 13 : 0.2 : 3 \quad (114)$$

in scheme B. At present the only measured width in this list is that of the $f_0(1370) = 5.4 \pm 2.3$ keV [5]. Using this to normalise the above, we anticipate $\Gamma(f_0(1500)) \approx 0.5$ keV (scheme A) or 0.1 keV (scheme B). Both schemes imply $\Gamma(f_0(1710)) \approx 1.2$ keV.

This relative ordering of widths is a common feature of mixings for all initial configurations for which the bare glueball does not lie nearly degenerate to the nn state. As such, it is a robust test of the general idea of nn and ss mixing with a lattice motivated glueball. If, say, the width of the $f_0(1710)$ were to be smaller than the $f_0(1500)$, or comparable to or greater than the $f_0(1370)$, then the general hypothesis of significant three state mixing with a lattice glueball would be disproven. The corollary is that qualitative agreement may be used to begin isolating in detail the mixing pattern.

Now we turn to $J=$ radiative decay rates. Since in either scheme

$$\Gamma = 7.5 \pm 2.8 \text{ keV}; \quad (115)$$

the discussion of the previous section implies,

$$\text{br}(J= \rightarrow f_0) \approx (1.5 \pm 0.6) \times 10^{-3}; \quad (116)$$

However each scheme makes a more specific prediction. By our hypothesis that qq coupling to gg is suppressed at $O(\alpha_s)$ relative to the corresponding glueball amplitude, we may scale the $J= \rightarrow f_0$ production amplitudes for the mixed states as follows. For simplicity we shall assume that $A(J= \rightarrow nn) = \sqrt{2} A(J= \rightarrow ss) = c_s A(J= \rightarrow G)$, where c is some constant whose magnitude and phase are in general model dependent. In this approximation, we have for scheme A

$$\begin{aligned} A(f_0(1370) \rightarrow gg) &= (0.5 + c_s 1.3) A_0 \\ A(f_0(1500) \rightarrow gg) &= 0.6 A_0 \\ A(f_0(1710) \rightarrow gg) &= (0.6 + c_s 1.1) A_0 \end{aligned}$$

In general we see that for mixing scheme A:

(i) The absence of a dominant signal in $J= \rightarrow f_0(1370)$ suggests that c is not negative and that the G - qq interference there is destructive.

(ii) The qq admixture in the $f_0(1500)$ is nearly pure flavour octet and hence decouples from gg. This leaves the strength of $\text{br}(J=0 \rightarrow f_0(1500))$ at about 40% of the pure glueball strength, which is consistent with the mean of the two analyses in section 4.1.

(iii) The destructive interference in the $f_0(1370)$ case implies a constructive effect for the $f_0(1710)$ and hence this picture predicts that $\text{br}(J=0 \rightarrow f_0(1710)) > \text{br}(J=0 \rightarrow f_0(1500)) > \text{br}(J=0 \rightarrow f_0(1370))$. If as a particular example for comparison between the two schemes we take $c_s = 0.5=1.3$ to decouple $f_0(1370)$ entirely in radiative $J=0$ decay, we find $\text{br}(J=0 \rightarrow f_0(1710)) : \text{br}(J=0 \rightarrow f_0(1500)) : \text{br}(J=0 \rightarrow f_0(1370)) = 1:1 : 0.4 : 0$.

Mixing scheme B, corresponding to an ideal glueball lying above the nonet, leads to the following amplitudes:

$$\begin{aligned} A(f_0(1370) \rightarrow gg) &= (0.4 + c_s 1.5)A_0 \\ A(f_0(1500) \rightarrow gg) &= (0.2 + c_s 0.4)A_0 \\ A(f_0(1710) \rightarrow gg) &= (0.9 + c_s 0.8)A_0 \end{aligned}$$

Here both $f_0(1370)$ and $f_0(1500)$ production are suppressed due to the destructive interference of the glueball and qq components; the $f_0(1710)$ being enhanced as in the previous example. For the example $c_s = 0.4=1.5$, (chosen to decouple the $f_0(1370)$ and enable comparison with scheme A as above) we find $\text{br}(J=0 \rightarrow f_0(1710)) : \text{br}(J=0 \rightarrow f_0(1500)) : \text{br}(J=0 \rightarrow f_0(1370)) = 1.2 : 0.01 : 0$.

Thus, in conclusion, both these mixing schemes imply a similar hierarchy of strengths in $J=0$ production which may be used as a test of the general idea of three state mixing between glueball and a nearby nonet. Prominent production of $J=0 \rightarrow f_0(1710)$ is also a common feature. When the experimental situation clarifies on the $J=0 \rightarrow f_0$ branching fractions, we can use the relative strengths to distinguish between the case where the glueball lies within a nonet, ref[2], or above the ss member, ref[45].

7 Summary

We have clarified the relationship between $b_{\text{rad}}(Q\bar{Q}_V \rightarrow \gamma + R)$ and $\text{br}(R \rightarrow gg)$ proposed by Cakir and Farrar [18]. In particular, we have examined its dependence on the $\langle gg | R \rangle$ form factors and discussed theoretical and experimental constraints on these form factors. We conclude that the relation can be used, possibly with generalized $H_J(x)$ functions, for light- $q\bar{q}$ mesons and glueballs as well as heavy $q\bar{q}$ mesons. Using this relation, we find

The $f_0(1500)$ is at least half-glueball if the Bugg et al analysis [3] of the 4π channel is confirmed, but is less so according to the BES results. Analysis of Mark III data on $J = 0$ is urgently needed. At this moment the experimental determinations of $\text{br}(J = 0 \rightarrow f_0(1500))$ are inconsistent.

The $f_0(1710)$ is also at least half-glueball, if $J = 0$; if $J = 2$ it is a $q\bar{q}$ meson. Experimental determinations of the $f_{0,2}$ spectra in the 1.6–1.8 GeV region are presently inconsistent.

The $f_0(2330)$ is unlikely to have $J = 0$, if present experimental data are correct. If it has $J = 2$ it strongly resembles a glueball.

The $f_0(1440)$ is separated into two states. The lower mass state, $f_{0L}(1410)$, has strong affinity for glue; the higher mass $f_{0H}(1480)$ is consistent with being the ss member of a nonet, perhaps mixed with glue.

It is of urgent importance to (a) arrive at an experimental consensus on the f_0 and f_2 masses and widths in the 1600–1800 region and (b) resolve the discrepancies in the present determinations of $\text{br}(J = 0 \rightarrow f_0(1500))$. Measurement of production branching fractions of the f_0 and f_2 mesons in radiative decay should be quite easy and yield useful additional information. We also outlined a procedure to use data on $\gamma \rightarrow R$ and $\gamma \rightarrow R$ together, to help unravel the $q\bar{q}$ and gg composition of mesons. To accomplish this, measurement of $\text{br}(f_0(1370;1500;1710) \rightarrow \gamma \gamma)$ is an essential ingredient.

An emerging mystery is the (1440) region. Its properties seem to differ in $J^=$ radiative decay and pp annihilation, and it has not been seen in central production. Possibly these differences are due to the different interplay of gluon and qq annihilation in the various production processes. This merits further investigation, both experimental and theoretical. The strong production of the (1410) in radiative $J^=$ decay indicate that it could be a glueball. However its low mass is difficult to reconcile with lattice gauge predictions. Its properties and mass are consistent with those expected for a bound state of light gluinos. Given that the (1410) may be evidence of a new degree of freedom in QCD, or evidence of dynamics beyond quenched lattice gauge theory in the 0^+ sector, more detailed experimental investigation of the pseudoscalar sector is a high priority.

8 Acknowledgements

We are indebted to D.V. Bugg, W. Dunwoodie and A. Kirk for discussions. This work supported by U.S. National Science Foundation grant nos. NSF-PHY-94-23002 (GRF) and NSF-PHY-90-23586 (ZPL). FEC is partially supported by the European Community Human Mobility Program Eurodafone, Contract CHRX-CT 92-0026

Appendix: Analytical Expressions for $H_J(x)$

The analytical expressions for the loop integral $H_J(x)$ are given in Ref. [19]. In the normalisation of the present paper they are:

$$H_{0^+}(x) = \frac{4}{x} L(1-2x) - L(1) - \frac{1-x}{2x} (2L(1-x) - \frac{2}{3} + \frac{1}{2} \ln^2(1-x)) - \frac{x}{1-2x} \ln(2x) + i4 \frac{1-x}{(2-x)x} \ln(1-x) \quad (117)$$

for $J = 0^+$, and

$$H_{0^{++}} = \frac{2}{3} \frac{2-3x}{x^2} + 10 \frac{1-x}{x^3} + 4 \frac{1-2x}{x^2} \ln(2) - \ln(1-x)$$

$$\begin{aligned}
& + \frac{8}{x^2} + 2 \frac{1-x}{x(1-2x)} \ln(2x) : 3 \frac{1-x}{x(2-x)} \ln^2(1-x) \\
& + \frac{8-6x+x^2-6x^3}{x^3(2-x)} \stackrel{2=6}{=} \frac{4-5x+2x^2}{x^3} L(1-2x) \\
& \quad 4 \frac{2-2x-x^2}{x^2(2-x)} L(1-x) + i \frac{1-x}{x(2-x)} \ln(1-x) \quad \# \quad (118)
\end{aligned}$$

for 0^{++} , where $L(x)$ is a Spence function, defined as

$$L(x) = \int_0^x \frac{dx}{x} \ln(1-x) : \quad (119)$$

There are three helicity amplitudes for the tensor state, and they are related to the total $H_{2^{++}}$ by

$$H_{2^{++}}(x) f^2 = H_{2^{++}}^0(x) f^2 + H_{2^{++}}^1(x) f^2 + H_{2^{++}}^2(x) f^2 : \quad (120)$$

The helicity amplitudes $H_{2^{++}}^1$ in Eq. 120 are

$$\begin{aligned}
H_{2^{++}}^0 &= \frac{2^p}{x^3} \frac{3}{3} x(6-5x) + \frac{2}{3} \frac{6-19x+18x^2}{x} (1-x) \ln(1-x) \\
& \quad \frac{10-12x+5x^2}{3(2-x)} g_1 + \frac{2}{3} \frac{6-38x+71x^2-37x^3}{1-2x} \ln(2x) \\
& \quad 8 \frac{(1-x)^2}{x^2(2-x)} g_2 + \frac{4}{3} \frac{6-6x-x^2}{x} \ln(2) - \frac{1}{2} i \\
& \quad \frac{4}{3} (12-26x+13x^2) g_3 ; \quad (121)
\end{aligned}$$

$$\begin{aligned}
H_{2^{++}}^1 &= \frac{2^p}{x^3} \frac{1}{3} (38-9x)x - \frac{2}{x} (4-13x+16x^2-4x^3) \ln(1-x) \\
& \quad 2 \frac{x(1-x)}{2-x} g_1 - \frac{4}{1-2x} (2-11x+16x^2-4x^3) \ln(2x) \\
& \quad + 8 \frac{(1-x)(2-2x+x^2)}{x^2(2-x)} g_2 - \frac{16}{3} \frac{3-x^2}{x} \ln(2) - \frac{1}{2} i \\
& \quad + 4(8-12x+3x^2) g_3 \quad (122)
\end{aligned}$$

and

$$\begin{aligned}
 H_{2^{++}}^2 = & \frac{p}{2} \frac{(1-x)^2}{x^3} + \frac{16}{3}x + \frac{4}{x}(1-6x+6x^2) \ln(1-x) + 2 \frac{5}{2} \frac{6x+2x^2}{x} g_1 \\
 & + 4(1-6x) \ln(2x) + 4 \frac{2}{x^2} \frac{4x+6x^2-4x^3+x^4}{(2-x)} g_2 \\
 & + \frac{4}{3} \frac{6-6x+11x^2}{x} \ln(2) + \frac{1}{2} i \sqrt{16(1-x)} g_3 \quad (123)
 \end{aligned}$$

where

$$g_1 = L(1-x) - L(1-2x); \quad (124)$$

$$g_2 = L(1-2x) - 2L(1-x) + L(1-x) \frac{1}{2} \ln^2(1-x) + i \ln(1-x) \quad (125)$$

and

$$g_3 = L(1-x) - L(1-2x) - \ln(2) \ln(1-x) \quad (126)$$

References

- [1] V.V. Anisovich et al, Phys. Lett. 323B, 233 (1994); C. Amisler et al, *ibid.* 342B, 433 (1994); 291B, 347 (1992); 340B, 259 (1994).
- [2] C. Amisler and F.E. Close, Phys Lett B 353-385 (1995); Phys Rev D 53 295 (1996)
- [3] D.V. Bugg, et al, Phys. Lett. 353B, 378 (1995).
- [4] D. Weingarten, Nucl. Phys. B (Proc. Suppl.) 34-29 (1994)
- [5] Particle Data Group, Phys. Rev. D 50, 1173 (1994).
- [6] J. Bai et al, BES Collaboration, Phys. Rev. Lett. 76, 3502 (1996); Kuang-Ta Chao, Commun. Theor. Phys 24 (1995) 373
- [7] K. Ishikawa, Phys Rev Letters 46 978 (1981); M. Chanowitz, *ibid* 981 (1981).

- [8] F.E. Close, *Rep. Prog. Phys.* 51 833 (1988)
- [9] S. Abatzis et al., *Phys. Lett. B* 324 509 (1994)
- [10] D. Alder et al., *Phys. Lett. B* 201 160 (1988);
- [11] V.V. Anisovich et al., *Phys. Lett. B* 323 233 (1994) ; C. Amisler et al.,
ibid B 342 433 (1995); *B* 291 347 (1992); *B* 340 259 (1994)
- [12] T.A. Armstrong et al., *Phys. Lett. B* 307 394 (1993); *ibid* 399
- [13] A. Tennikov (Obelix Collaboration) p.325 in "Hadron95" (World Scientific; M. Birse et al. eds, 1995); A. Bertin et al. *Physics Letters B* 361 187 (1995)
- [14] C. Amisler et al. (Crystal Barrel Collaboration) *Physics Letters B* 358, 389 (1995)
- [15] G. Bali et al. (UKQCD), *Phys. Lett. B* 309 378 (1993)
- [16] J. Sexton et al., *Phys Rev. Letters*, 75, 4563 (1995)
- [17] G.R. Farrar, *Phys Rev. Letters*, 76, 4111 (1996) and references therein.
- [18] M.B. Cakir and G. Farrar, *Phys. Rev. D* 50, 3268 (1994).
- [19] J.G. Komer, J.H. Kuhn, M. Kramer, and H. Schneider, *Nucl. Phys. B* 229, 115 (1983), J.G. Komer, J.H. Kuhn, and H. Schneider, *Phys. Lett.* 120B, 444 (1983).
- [20] A. Billore et al., *Phys. Lett.* 80B, 381 (1979)
- [21] K. Koller and T. Walsh, *Nucl. Phys. B* 140, 449 (1978).
- [22] J.H. Kuhn, J. Kaplan, and E. Sani, *Nucl. Phys.* 157, 125 (1979).
- [23] S.J. Brodsky and G. Farrar, *Phys Rev Letters* 31 1153 (1973)
- [24] R. Barbieri, R. Gatto, and R. Kogerler, *Phys. Lett.* 60B, 183 (1976).

- [25] A. I. Aleksëev, Zh. Eksp. Teor. Fiz. 34, 1195 (1958) [Sov. Phys. JETP, 7, 826 (1958)].
- [26] V. A. Novikov, M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Nucl. Phys. B 191, 301 (1981).
- [27] T. Barnes, F. Close and S. Monaghan, Nucl. Phys. B 198, 380 (1982).
- [28] Zhenping Li, F. E. Close and T. Barnes, Phys. Rev. D 43, 2621 (1991).
- [29] R. Barbieri, M. Cao, R. Gatto and E. Remiddi, Nucl. Phys. B 192, 61 (1981)
- [30] E. Braaten et al., Phys. Rev. D 46, 1914 (1992); Phys. Rev. Letters 71, 1673 (1993); F. E. Close and Zhenping Li, Phys. Rev. Lett. 66, 3109 (1991); Z. Phys. C 54, 147 (1992).
- [31] T. Barnes, F. E. Close, P. Page and E. Swanson; "Higher Quarkonia", RAL-96-039, hep-ph/9609339
- [32] B. Guberina, and J. H. Kuhn, Nuovo Cim. Lett. 32, 295 (1981).
- [33] S. J. Brodsky, T. A. de Grand, R. R. Horgan and D. G. Coyne, Phys. Lett. 73B, 203 (1978); K. Koller and T. Walsh, Nucl. Phys. B 140, 449 (1978)
- [34] A. Seiden et al., Phys. Rev. D 38, 824 (1988)
- [35] J. Z. Bai et al., (BES Collaboration) "Amplitude analysis of the (1440) in $J^P = 1^-$ Radiative Decay to the $K\bar{K}$ Final State", BHEP-EP1 95-04 July 1996. (unpublished)
- [36] D. V. Bugg, A. V. Sarantsev and B. S. Zou, "New Bounds on $\pi\pi$ Phase Shifts between 600 and 1900 MeV", Nucl. Phys. B (in press)
- [37] D. Bugg, private communication
- [38] R. Landua; Proc of XXVIII International Conference on High Energy Physics, Warsaw, July 1996.

- [39] S.J.Lindenbaum and R.S.Longacre, Phys. Lett. B 274 492 (1992)
- [40] R.Baltrusaitis et al (M k III), Phys Rev D 35 2077 (1987)
- [41] T.A m strong et al Phys. Lett. B 227 186 (1989)
- [42] J.Z Bai et al (BES Collaboration) \Structure analysis of the $f_J(1710)$ in the radiative decay $J = \frac{1}{2} \rightarrow K^+ K^-$ BEPC report, July 18 1996
- [43] F.E.Close and M.J.Teper, \On the lightest Scalar Glueball" Rutherford Appleton Laboratory report no. RAL-96-040; Oxford University report no. O U T P -96-35P
- [44] M.Genovese, Phys. Rev. D 46 5204 (1992)
- [45] D.W eingarten, hep-lat/9608070.
- [46] P.D Bames et al, Phys. Lett. B 309 469 (1993)
- [47] J.Augustin et al, Phys. Rev. D 42, 10 (1990)
- [48] D.Scharre et al (M ark 2) Physics Letters, 97B , 329 (1980)
- [49] Z Bai et al (M ark III Collaboration), Phys. Rev. Letters 65, 2507 (1990)
- [50] M.Burchell, Nucl. Phys. B 21, 132 (suppl) (1991) T Bolton et al (M ark III Collaboration), Phys Rev Letters 69, 1328 (1992)
- [51] G.R.Farrar, RU -95-17, hep-ph/9504295.
- [52] G.R.Farrar and G.G abadadze, RU -96-37, hep-ph/9608330.
- [53] M.Chanowitz, LBL-18701; P roc of V I International W orkshop on Photon-photon collisions, Lake Tahoe Sept 1984.
- [54] W.Kwong, P M ackenzie, R.Rosenfeld and J.Rosner, Phys Rev. D 37 3210 (1988); A.Petrelli, hep-ph/9603439.
- [55] B.W ebber, P roc of 27 Internat. Conf. on HEP, Glasgow 1994, (IOP Publishers, P.Bussey and I.Knowles eds.) p.213

[56] D Morgan and M Pennington, Phys Rev D 48 1185 (1993)

[57] H Behrend et al (CELLO Collaboration), Z Phys C 43, 91 (1989)

[58] Particle Data Group, Phys. Rev. D 54,1 (1996).




Figure 1: Magnitude of the loop integral, $x|H(x)|^2$ versus x for 0^{++} (dotted), 0^{+-} (dashed) and 2^{++} (solid); $x = 1 - \left(\frac{m_R}{m_V}\right)^2$.