

Time reversal odd fragmentation functions ¹

R. Jakob¹ and P.J. Mulders^{1,2}

(1) NIKHEF, P.O.Box 41882, 1009 DB Amsterdam, The Netherlands

(2) Free University, De Boelelaan 1081, 1081 HV Amsterdam, The Netherlands

ABSTRACT

The combination of transverse momentum and polarization effects in hard scattering processes, e.g polarized deep-inelastic leptonproduction, results in a rich variety of information on the hadronic structure. This information is encoded in a correspondingly large number of functions depending on both, longitudinal fractional momenta x and transverse momenta p_T . Integration over transverse momenta establishes the connection to the usual distribution and fragmentation functions. Constraints from hermiticity and invariance under parity operation induce some relations between different functions.

One specific aspect of the information on the hadronic structure is the appearance of *time reversal odd* fragmentation functions due to the non-applicability of time-reversal invariance for the hadronization of a quark. *T-odd* fragmentation functions are experimentally accessible via the measurement of, for instance, asymmetries in semi-inclusive leptonproduction.

The basic ingredient of the diagrammatic approach to hard scattering processes [1] is the assumption that the hadronic tensor *factorizes* in *hard* and *soft* parts separated by different powers of a hard scale Q . The information on the hadronic structure is encoded in hadronic matrix elements of quark (and, in general, gluon) fields, conveniently parametrized in distribution (DF) and fragmentation functions (FF). The leading contributions to 1-particle inclusive leptonproduction from the 'handbag' diagram (Fig.1) involve the non-local hadronic matrix elements

$$\Phi_{ij}(p, P, S) = \frac{1}{(2\pi)^4} \int d^4x e^{ip \cdot x} \langle P, S | \bar{\psi}_j(0) \psi_i(x) | P, S \rangle \quad (1)$$

for the distribution of quarks in the target hadron (with momentum P and spin vector S) and

$$\Delta_{ij}(k, P_h, S_h) = \sum_X \frac{1}{(2\pi)^4} \int d^4x e^{ik \cdot x} \langle 0 | \psi_i(x) | P_h, S_h; X \rangle \langle P_h, S_h; X | \bar{\psi}_j(0) | 0 \rangle \quad (2)$$




Figure 1: Diagram of the leading contributions to 1-particle inclusive leptonproduction.

¹Contributed paper at the 12th International Symposium on High Energy Spin Physics, Amsterdam, Sept. 10-14, 1996

describing the decay of a quark, thereby, amongst others producing the one observed hadron in the final state (characterized by P_h and S_h). Both objects, Φ and Δ , are hermitian and reveal symmetries under discrete transformations: both are parity invariant, whereas only Φ is invariant under *time reversal*, as well. The *time reversal* operation relates *in*-states to *out*-states and thus Δ has no simple symmetry properties under this operation. The most general ansatz for the quark-quark correlation function $\Delta(k; P_h, S_h)$ consistent with the hermiticity properties of the field and invariance under parity operation is (k is the momentum of the quark) [2]

$$\begin{aligned} \Delta(k; P_h, S_h) = & B_1 M_h + B_2 \not{P}_h + B_3 \not{k} + (B_4/M_h) \sigma_{\mu\nu} P_h^\mu k^\nu \\ & + i B_5 (k \cdot S_h) \gamma_5 + B_6 M_h \not{S}_h \gamma_5 + (B_7/M_h) (k \cdot S_h) \not{P}_h \gamma_5 + (B_8/M_h) (k \cdot S_h) \not{k} \gamma_5 \\ & + i B_9 \sigma_{\mu\nu} \gamma_5 S_h^\mu P_h^\nu + i B_{10} \sigma_{\mu\nu} \gamma_5 S_h^\mu k^\nu + i (B_{11}/M_h^2) (k \cdot S_h) \sigma_{\mu\nu} \gamma_5 k^\mu P_h^\nu \\ & + (B_{12}/M_h) \epsilon_{\mu\nu\rho\sigma} \gamma^\mu P_h^\nu k^\rho S_h^\sigma. \end{aligned} \quad (3)$$

Hermiticity requires all the coefficient functions B_i to be real. The additional constraint of time reversal invariance, if applicable, would imply $B_i^* = -B_i$ for B_4 , B_5 , and B_{12} . In this sense, those coefficient functions are referred to as *time reversal odd* (or, in short, *T-odd*).

The object appearing in the diagrammatic expansion is the k^+ -integrated hadronic matrix element, conventionally, parametrized in terms of generalized FF's (up to $\mathcal{O}(1/Q)$)

$$\begin{aligned} \frac{1}{4z} \int dk^+ \Delta(k, P_h, S_h) \Big|_{k^- = \frac{P_h^-}{z}, \mathbf{k}_T} &= \frac{M_h}{4P_h^-} \left\{ E + D_1 \frac{\not{P}_h}{M_h} + D_{1T}^\perp \frac{\epsilon_{\mu\nu\rho\sigma} \gamma^\mu P_h^\nu k_T^\rho S_{hT}^\sigma}{M_h^2} \right. \\ &+ D^\perp \frac{\not{k}_T}{M_h} + D_T \epsilon_{\mu\nu\rho\sigma} n_+^\mu n_-^\nu \gamma^\rho S_{hT}^\sigma + \lambda_h D_L^\perp \frac{\epsilon_{\mu\nu\rho\sigma} n_+^\mu n_-^\nu \gamma^\rho k_T^\sigma}{M_h} - E_s i \gamma_5 - G_{1s} \frac{\not{P}_h \gamma_5}{M_h} \\ &- G'_T \not{S}_{hT} \gamma_5 - G_s^\perp \frac{\not{k}_T \gamma_5}{M_h} - H_{1T} \frac{i \sigma_{\mu\nu} \gamma_5 S_{hT}^\mu P_h^\nu}{M_h} - H_{1s}^\perp \frac{i \sigma_{\mu\nu} \gamma_5 k_T^\mu P_h^\nu}{M_h^2} + H_1^\perp \frac{\sigma_{\mu\nu} k_T^\mu P_h^\nu}{M_h^2} \\ &\left. - H_T^\perp \frac{i \sigma_{\mu\nu} \gamma_5 S_{hT}^\mu k_T^\nu}{M_h} + H \sigma_{\mu\nu} n_-^\mu n_+^\nu - H_s i \sigma_{\mu\nu} \gamma_5 n_-^\mu n_+^\nu \right\}. \end{aligned} \quad (4)$$

The generalized FF's depend on the longitudinal momentum fraction z and on the transverse momentum \mathbf{k}_T and consist of (integrated) linear combinations of the coefficient functions B_i . Since, there are more FF's — 21 up to $\mathcal{O}(1/Q)$ — than the 12 independent coefficient functions B_i , a number of relations is expected. Indeed, exploiting the connection between the FF's and the B_i leads to nine relations. Focussing here on the seven *T-odd* functions D_{1T}^\perp , H_1^\perp , D_L^\perp , D_T , E_L , E_T and H there are four relations between them

$$D_L^\perp(z, -z\mathbf{k}_T) = -D_{1T}^\perp(z, -z\mathbf{k}_T) \quad (5)$$

$$D_T(z) = z^3 \frac{d}{dz} \left[\frac{D_{1T}^{\perp(1)}(z)}{z} \right] \implies \int_0^1 dz \left[\frac{D_T(z)}{z} + 2D_{1T}^{\perp(1)}(z) \right] = 0 \quad (6)$$

$$H(z) = z^3 \frac{d}{dz} \left[\frac{H_1^{\perp(1)}(z)}{z} \right] \implies \int_0^1 dz \left[\frac{H(z)}{z} + 2H_1^{\perp(1)}(z) \right] = 0 \quad (7)$$

$$E_L(z) = z^3 \frac{d}{dz} \left[\frac{E_T^{(1)}(z)}{z} \right] \implies \int_0^1 dz \left[\frac{E_L(z)}{z} + 2E_T^{(1)}(z) \right] = 0 \quad (8)$$

Here, the upper index (1) denotes the first moment in \mathbf{k}_T . The relations follow directly from the hermiticity properties of the fields and parity reversal invariance; the RHS's of (6), (7), and (8) assume not too singular end-point behaviour.

One particularly important example for the occurrence of *T-odd* FF's is the differential cross section for unpolarized electrons scattering off a transversely polarized target provided the transverse momentum of the produced hadron is measured [3,4].

$$\begin{aligned} \frac{d\sigma_{OT}}{dx dy dz d^2P_{h\perp}} &= \frac{4\pi\alpha s}{Q^4} |\mathbf{S}_\perp| \sum_{a,\bar{a}} e_a^2 (1-y) \left\{ \sin(\phi_h + \phi_s) I \left[\frac{\hat{\mathbf{h}} \cdot \mathbf{k}_T}{M_h} x h_1^a H_1^{\perp a} \right] \right. \\ &+ \left. \sin(3\phi_h - \phi_s) I \left[\frac{(4(\hat{\mathbf{h}} \cdot \mathbf{p}_T)^2 - \mathbf{p}_T^2) \hat{\mathbf{h}} \cdot \mathbf{k}_T - 2\hat{\mathbf{h}} \cdot \mathbf{p}_T \mathbf{p}_T \cdot \mathbf{k}_T}{M^2 M_h^2} x h_{1T}^{\perp a} H_1^{\perp a} \right] \right\} + \mathcal{O}\left(\frac{1}{Q}\right) \end{aligned} \quad (9)$$

with $I \left[(\hat{\mathbf{h}} \cdot \mathbf{p}_T) f D \right] \equiv \int d^2p_T d^2k_T \delta^2(\mathbf{p}_T + \mathbf{q}_T - \mathbf{k}_T) (\hat{\mathbf{h}} \cdot \mathbf{p}_T) f(x, \mathbf{p}_T) D(z, \mathbf{k}_T)$.

Eq.(9) contains two sine functions depending on different combinations of the azimuthal angles ϕ_s and ϕ_h of the target spin vector and the momentum of the produced hadron, respectively. The asymmetry caused by the first term is known as the ‘Collins effect’ [3]. Both azimuthal asymmetries are of leading order in an expansion in powers of $1/Q$. Leading order asymmetries involving *T-odd* FF's are possible only in experimental quantities sensitive to transverse momentum (for a more detailed discussion on the role of transverse momentum in deep-inelastic processes see [5]).

Another example of observables involving *time reversal odd* fragmentation functions is the number asymmetry of produced hadrons in the scattering of unpolarized leptons off a transversely polarized target.

$$\frac{\sigma(\ell \vec{H}) - \sigma(\ell \overleftarrow{H})}{\sigma(\ell \vec{H}) + \sigma(\ell \overleftarrow{H})} = \frac{2(2-y)\sqrt{1-y}}{1-y+y^2/2} \frac{M_h}{Q} \frac{\sum_{a,\bar{a}} e_a^2 h_1^a(x) (H^a(z) + 2zH_1^{\perp(1)a}(z))}{\sum_{a,\bar{a}} e_a^2 f_1^a(x) D_1^a(z)} \quad (10)$$

Using Eq.(7) the occurring linear combination of *T-odd* FF's, $H(z) + 2zH_1^{\perp(1)}(z)$ can be simplified to the form $z \frac{d}{dz} \left[z H_1^{\perp(1)}(z) \right]$. This is a typical example for the usefulness of the relations (5), (6),(7) and (8).

This work was supported by the foundation for Fundamental Research on Matter (FOM) and the Dutch Organization for Scientific Research (NWO).

- 1 H.D. Politzer, Nucl. Phys. B172, 349 (1980);
R.K. Ellis, W. Furmanski, R. Petronzio, Nucl. Phys. B212, 29 (1983).
- 2 J.P. Ralston, D.E. Soper, Nucl. Phys. B152, 109 (1979);
R.D. Tangerman, P.J. Mulders, Phys. Rev. D51, 3357 (1995).
- 3 J.C. Collins, Nucl. Phys. B396, 161 (1993).
- 4 P.J. Mulders, R.D. Tangerman, Phys. Lett. B352, 129 (1995); and NPB 461, 197 (1996).
- 5 R. Jakob, A. Kotzinian, P.J. Mulders, J. Rodrigues, these proceedings.