

Further Analysis on η -particle Properties

Shin Ishida, Taku Ishida, Muneaki Ishida,
Kunio Takamatsu, and Tsuneaki Tsuru

Atomic Energy Research Institute, College of Science and Technology,
Nihon University, Tokyo 101 National Laboratory for High Energy Physics
(KEK), Tsukuba 305

Department of Physics, University of Tokyo, Tokyo 113
Department of Engineering, Miyazaki University, Miyazaki 988-12

(Received May 25, 1997)

In a previous work, we have reanalyzed the $I=0$ S-wave $\pi\pi$ -scattering phase-shift through a new method of Interfering Breit-Wigner amplitudes, and have shown the existence of a light scalar, η particle, accompanied by a negative "repulsive core"-type background phase-shift, whose origin may have some correspondence to a "compensating" contact interaction. In this work we make further analysis on the phase shift under $K\bar{K}$ threshold from the same standpoint, to determine precise values of mass and width, so far as present experimental data are concerned.

x1. Introduction

Whether a light iso-scalar resonance ($J^{PC} = 0^{++}$), called " η "-particle, exists or not is one of the most interesting and important problems in hadron spectroscopy. Phenomenologically, since of its light mass and of its "vacuum" quantum number, it may affect various processes. Theoretically, for example, in the Nambu-Jona-Lasinio-type models^{2);3);4);5)}, realizing the situation of dynamical breaking of chiral symmetry and believed to be a low-energy effective theory of QCD, existence of η -meson is predicted as a chiral partner of the Nambu-Goldstone π -boson. Correspondingly the extensive experimental investigations in the $\pi\pi$ -channel have been made for many years. The $I=0$ S-wave $\pi\pi$ phase shift δ_0^0 is now well known to rise smoothly to 90° at around 900 MeV, then shows a rapid step-up by 180° near the $K\bar{K}$ threshold, and reaches only to 270° at $m \approx 1200$ MeV. This behavior of δ_0^0 was thought, in the 1976-through-1994 editions of PDG, to be due mainly to the narrow $f_0(980)$ and the broad $f_0(1370)$,¹⁾ since there remains no phase shift for light η -particle.

In contrast with this interpretation, we have shown in the previous work¹⁾ (to be referred as I), re-analyzing the phase shift δ_0^0 systematically up to 1300 MeV, a possibility for the existence of $\eta(555)$ -resonance with a rather narrow width of a few hundred MeV in addition to these two resonances.

The reasons which led us to a different result from the conventional one, even

¹⁾ However in the latest edition,⁶⁾ the behavior of δ_0^0 is understood as due to, in addition to the $f_0(980)$ and $f_0(1370)$, a very broad $f_0(400-1200)$ or η . See also the analyses suggesting the existence,^{7);8)} which introduce a repulsive $\pi\pi$ BG, similarly as in our case.

with the use of the same data of phase shifts, are twofold: On one hand technically, we have applied a new method of Interfering Breit-Wigner Amplitude (IA method) for the analyses, where the T-matrix (instead of K-matrix in the conventional treatment) for multiple resonance case is directly represented by the respective Breit-Wigner amplitudes in conformity with unitarity, thus parametrizing the phase shifts directly in terms of physical quantities, such as masses and coupling constants of the relevant resonant particles.

On the other hand physically, we have introduced a "negative background phase" B_{BG} : of hard core type, (with a core radius of about pion size,) making enough room for π -resonance. Here it is suggestive to remember that a similar type of B_{BG} is well-known to exist in the nucleus-nucleus scattering and in the nucleon-nucleon scattering⁹⁾. A possible origin of B_{BG} in the $\pi\pi$ system seems¹⁰⁾ to have some correspondence to the "compensating" repulsive 4S_1 interaction in the NJL-model or the linear σ -model, which is required from the viewpoint of the current algebra and the PCAC.

In this work, we shall extend the re-analysis of the phase shift in $I=0$ to determine the mass and coupling constant of $\pi\pi$ and core radius as precise as possible, from the present experimental data of $I=0$, inspecting especially the data under $K\bar{K}$ threshold.

2. Applied formulas

For our purpose we shall analyze the $I=0$ between the $\pi\pi$ -threshold and an energy slightly below the $K\bar{K}$ -threshold (980 MeV), taking into account the effects of two resonances, $\rho(770)$ and $f_0(980)$, and of the B_{BG} . Effects from other higher-mass resonances ($f_0(1370)$, $f_0(1500)$...) and from other channels than $\pi\pi$ are ignored.

The applied formulas of IA method (in the case of one channel with two resonances) are as follows.

The relevant partial S-wave S matrix element S in the $\pi\pi$ system is represented by the phase shift $\delta(s)$ and the amplitude $a(s)$. The $\delta(s)$ is given by the sum of $\delta^{Res}(s)$ and $B_{BG}(s)$, respectively, due to the resonance and background. The $\rho(770)$ and $f_0(980)$ resonances contribute additively to $\delta^{Res}(s)$.

$$S = e^{2i\delta(s)} = 1 + 2ia(s) \quad (1)$$

$$\delta(s) = \delta^{Res}(s) + B_{BG}(s); \quad \delta^{Res}(s) = \delta^{(\rho)}(s) + \delta^{(f_0)}(s); \quad (2)$$

Correspondingly the total S matrix is given by the product of individual S matrices.

$$S = S^{Res} S^{B_{BG}} = S^{(\rho)} S^{(f_0)} S^{B_{BG}}; \quad (3)$$

The unitarity of the total S matrix is now easily seen to be satisfied by the "unitarity of individual S matrices". Each of S^{Res} is given by corresponding amplitudes $a^{(R)}$'s, which is taken as the following relativistic Breit-Wigner (BW) form

$$S^{(R)}(s) = e^{2i\delta^{(R)}(s)} = 1 + 2ia^{(R)}(s); \quad R = \rho, f_0 \quad (4)$$

$$a^{(R)}_{\pi}(s) = \frac{p_{\pi} \bar{s}_R(s)}{(s - M_R^2) + i \frac{p_{\pi}}{s_R(s)}}; \quad (5)$$

$$p_{\pi} \bar{s}_R(s) = \frac{1}{8} g_R^2; \quad 1 - \frac{p_1}{8} \frac{p_1}{s}; \quad p_1 = \frac{q}{s=4 - m^2}; \quad (6)$$

where $\bar{s}_R(s = M_R^2)$ represents the peak width $\bar{s}_R^{(p)}$ of the resonance R , g_R is the π -coupling constant, $\frac{1}{8}$ is the π -state density and p_1 is the CM momentum of the pion. Here it is to be noted that the total resonance amplitude $a^{Res}(\sqrt{s}^{Res} - 1) = 2i$ is represented by the respective amplitudes as

$$a^{Res} = \frac{1}{a} + \frac{(f_0)}{a} + 2i \frac{1}{a} \frac{(f_0)}{a}; \quad (7)$$

where the last term, looking like an "interference" of the amplitudes, guarantees our amplitude to satisfy the unitarity constraint.

The $B_{\pi\pi}$ is supposed to be of the repulsive hard core type;

$$B_{\pi\pi}(s) = p_1 r_c; \quad (8)$$

In the actual analysis we have applied, in order to have a global fit in all relevant energy region, the relativistic BW formula (5) with a revised width

$$\begin{aligned} \tilde{\gamma}_R(s) &= \gamma_R(s) F(s) & (a); \\ F(s) &= 1 - \frac{s_0}{s + s_0} e^{-s/M_0^2}; \quad (M_0 = 400 \text{ MeV}) & (b); \end{aligned} \quad (9)$$

instead of $\gamma_R(s)$ defined in Eq.(6). As a matter of fact there has been no established form of relativistic BW formula, and the concrete form of $F(s)$ is considered to be determined following the dynamics of strong interactions. The conventional BW form (5) ($\tilde{\gamma}_R(s)$ with $F(s) = 1$ in Eq.(9)) gives, in the case of broad width, an unphysical mass spectrum in the low energy region. The phenomenological form of $F(s)$ with a parameter s_0 given in (9) is so designed as to give a good fit in the low-energy region affecting it only in the energy region below 400 MeV. Thus s_0 plays a similar role as a scattering length.

3. Mass and width of π and core radius

The θ_0 over $m_\pi > 600$ MeV is well determined by the analysis of the CERN-Munich experiment. Among 5 independent analyses performed¹¹⁾ in Gayer74, (i) the one originally presented in Hyams73 (b-analysis in Gayer74) is widely accepted, partly because of its agreement to those reported by Protopopescu73¹²⁾. (ii) Srinivasan75 gave the phase between 350-600 MeV, (iii) Rosselet77 and (iv) Belikov79 determined the phase between π -threshold and 400 MeV. (i)-(iv) will be used as the "standard phase shift θ_0 " in the present analysis. However, there are several works to report different behaviors of θ_0 within $\pm 20^\circ$ ambiguities in the $m_\pi = 400$ -800 MeV region. We will utilize them to estimate the allowed region of m_π and g in our analysis: (v) The c-analysis of Gayer74 and (vi) Cason82 are taken as the upper and lower bounds of θ_0 , respectively.




Fig. 1. $I=0$ scattering phase shift. (a) Best fit to the standard δ_0^0 . The respective contributions to δ_0^0 from $f_0(980)$ and $\delta_{B.G.}$ are also given. The dotted line with label of " $r_c=0$ " represents conventional fit without the repulsive background thus far made. (b) χ^2 , m_σ and $g_{\sigma nn}$ versus r_c . (c) Best fits to the upper and lower δ_0^0 (dashed lines), and fittings with 3-sigma deviation to the standard δ_0^0 corresponding to $r_c=2.6$ and 3.35 GeV^{-1} (dotted lines).

The properties of $f_0(980)$ have already been obtained in I by analyzing systematically the standard δ_0^0 , elasticity, and $\pi^+ K \bar{K}$ phase shift together in the range of $m_{\pi\pi} = 600 - 1300 \text{ MeV}$, with the 2-channel ($\pi, K \bar{K}$) three resonance ($\pi, f_0(980), f_0(1370)$) formula in the IA method. The obtained values are $M_{f_0} = 993.2^{+6.5}_{-6.9} \text{ MeV}$ and $g_{f_0} = 1680^{+91}_{-91} \text{ MeV}$, which are used as fixed parameters in the following analysis.¹⁾

Fig. 2 (a) shows the result of the best fit (solid line) to standard δ_0^0 by the formula (2) given above, which includes the sum of three contributions: $\pi, f_0(980)$, and $\delta_{B.G.}$. The best-fitted values of parameters are $M_\pi = 585 \text{ MeV}$, $g = 3.6 \text{ GeV}$ (corresponding

¹⁾ The $f_0(980)$ does not play essential role in the relevant mass region since of its small coupling (width).

Table I. Obtained values of parameters and their errors. $g^r = g \cdot F^{1/2} (s = M_\rho^2)$. As for the definitions of (p) and (d) , see the text.

	standard ρ_0 (i-iv)	upper bound (v)	lower bound (vi)
M	585 20 M eV	535 M eV	650 M eV
g^r	3600 350 M eV	3670 M eV	3465 M eV
(p)	385 70 M eV	430 M eV	335 M eV
(d)	340 45 M eV	370 M eV	315 M eV
r_c	3.03 0.35 GeV ⁻¹	3.03 (xed)	3.03 (xed)
	0.60 0.07 fm	(0.6 fm)	(0.6 fm)

to decay width of 340 M eV), and $r_c = 3.03 \text{ GeV}^{-1} (0.60 \text{ fm})^{-1}$ with $\chi^2 = 23.6$ for 30 degrees of freedom (34 data points with 4 parameters¹⁾). Note that the best-fitted core radius is nearly the same as the "structural size" (charge radius) of pion 0.7 fm.

In order to justify the preciseness of these values and to estimate their errors it is apparently necessary to take into account of the correlation between M , g and r_c . Several tests are performed for various fixed values of r_c between 0 and 4.0 GeV^{-1} . In Fig 2(b)²⁾, M and g are plotted as functions of r_c , where M and g decrease as r_c becomes larger. The χ^2 shows deep parabolic shape, and gives its minimum at 3.03 GeV^{-1} , where we obtain the above mentioned "best-fit" values. Note that the fit in the case of $r_c = 0$, which corresponds to the results of the conventional analyses without BG thus far made, gives $\chi^2 = 163.4$, about 140 worse than the best fit. In the first row of Table 1 the errors of relevant parameters are quoted which correspond to three sigma deviation in the χ^2 behavior in Fig 2(b). In Fig 2(c) the tests with $r_c = 2.6$ and 3.35 GeV^{-1} corresponding to the 3-sigma deviations are shown. The small bump around 750 M eV of the standard ρ_0 is not reproduced with small r_c (correspondingly with large g -width). To estimate the effect due to the ambiguities of experimental ρ_0 data mentioned above, data (v) and (vi) are analyzed with fixed $r_c = 3.03 \text{ GeV}^{-1}$. The results are also shown in Fig 2(c) and the obtained values of parameters are given in the second and the third rows of Table 1.

From all of these studies, we may conclude that the M is in the range of 535–650 M eV.

4. Supplementary discussions

We would add some comments on the results of our analysis:

i) First a possible influence of the tail of $f_0(1370)$ even below $m_\rho = 980 \text{ M eV}$ should

¹⁾ The significant difference of the value of M from that (553.3 0.5 M eV) in I is due to including a consideration on the correlation between M and r_c in the present work, and also, regrettably, due to mis-reading of the errors of some data points of (ii) in I.

²⁾ In (i), one point at $M = 910 \text{ M eV}$ seems to have a too small error and to disturb the continuity to adjacent data points. In fact, this point turned out to occupy large fraction of total χ^2 . The values in the text are obtained without this point, while $M = 600 \text{ M eV}$, $g^r = 3750 \text{ M eV}$, $r_c = 2.75 \text{ GeV}^{-1} (0.54 \text{ fm})$, and $P_{S_0} = 475 \text{ M eV}$ with $\chi^2 = 35.9$ for 31 degrees of freedom, in the case of including this point.

Table II. Pole positions on sheet II in present one-channel analysis. The errors corresponds to the 3-sigma deviation from the best fit to standard ρ^0 .

$s_{\text{pole}} = \text{GeV}^2$	$\sqrt{s_{\text{pole}}} = \text{GeV}$	$p_{\text{pole}} = \text{GeV}$
(0.324 0.020)	(0.602 0.026)	(0.271 0.015)
-i(0.236 0.044)	-i(0.196 0.027)	-i(0.109 0.014)

be mentioned. ¹⁾ Would $f_0(1370)$ have a large width, the value $r_c = 3.03 \text{ GeV}^{-1}$ in Table 1 should be regarded as including the $f_0(1370)$ contribution. ¹⁾ However, we may expect that it does affect very little the values of M and g , since these are determined mainly by the fine structure of the standard ρ^0 around 750 MeV.

ii) In our treatment the S-matrix is parametrized directly in terms of physically meaningful quantities, the masses and $\pi\pi$ -coupling constants of resonances. So it is not necessary to argue about pole positions on analytically continued complex Riemann sheets. However, for convenience to compare with other works, we give these values for the resonance in Table 2. Our S-matrix has the form of product of respective resonance-S-matrices. Each BW resonance (in our form of Eq.(5)) produces three poles, a virtual state pole, a BW pole and its complex conjugate pole. We are able to derive the approximate relations. $\text{Re } \sqrt{s_{\text{pole}}} \approx M_R$ and $\text{Im } \sqrt{s_{\text{pole}}} \approx \Gamma/2$ ^(p). This situation corresponds to the case of one-channel $(\pi\pi)$. In the two-channel case $(\pi\pi$ and $K\bar{K})$, one BW resonance induces four poles (except for virtual state pole), $s_{\text{p}}^{\text{III}}$, its complex conjugate $s_{\text{p}}^{\text{III}*}$, and the two in sheet II or IV. In I, $g_{K\bar{K}} = 0$ is obtained. This case leads to the four poles with the same s -values s_{p} and s_{p}^* on sheets II and III.

Here the decay width ^(d) should be discriminated from the peak width ^(p). The former is defined by the formula

$$\begin{aligned} \Gamma^{(d)} &= N^{-1} \int_{-\infty}^{\infty} ds \quad (s) = [(s - M^2)^2 + s \Gamma^2]; \\ N &= \int_{-\infty}^{\infty} ds \quad 1 = [(s - M^2)^2 + s \Gamma^2]: \end{aligned}$$

iii) In our analysis we have introduced a parameter s_0 in Eq.(9). The obtained value of it is $\sqrt{s_0} = 365 \text{ MeV}$, which corresponds to the scattering length $a_0^0 = 0.23$ in m^{-1} unit. We have introduced no ad hoc prescriptions for Adler-zero condition. However, we have found that our resultant amplitude has a zero at $s = 1.0 \text{ m}^2$.

iv) In the $I = 2$ system, there is no known/expected resonance, and accordingly it is expected that the repulsive core will show up itself directly. In Fig 2 the experimen-

¹⁾ $f_0(1370)$ is known to couple strongly with $\pi\pi$ channel and thus to give 180 contribution to ρ^0 .

¹⁾ In I we obtained the relevant values $M_{f_0} = 1310 \text{ MeV}$, $\Gamma_{\text{tot}}^{(p)} = 420 \text{ MeV}$ and $r_c = 3.7 \text{ GeV}^{-1}$ (0.73 fm). However, as Svec96¹³⁾ and Kaminski96¹⁴⁾ suggest, the analyses to obtain ρ^0 thus far made do not usually take a_1 exchange into account, which might affect ρ^0 in the mass region above $m_{\pi\pi} = 1 \text{ GeV}$. Also the width of $f_0(1370)$ is not actually known (see PDG 96⁶⁾). Exact estimation seems to be difficult at this stage.




Fig. 2. $I=2$ scattering phase shift. Fitting by hard core formula is also shown.

tal data¹⁵⁾ of the $I=2$ scattering S-wave phase shift δ_0^2 is shown from threshold to $m_{\pi\pi} = 1400$ MeV, which is apparently negative, and is fitted well also by the hard core formula $\delta_0^2 = -r_c^{(2)} p_1 j_w$ with the core radius of $r_c^{(2)} = 0.87 \text{ GeV}^{-1}$ (0.17 fm). The core radius is smaller by a factor of 3 than that in the $I=0$ system, which suggests the importance of quark-pair-annihilation force in the η system (see Ref.17) for possible explanation in the framework of the linear model).

5. Concluding remarks

In this work and in I we have shown the strong possibility of existence of η -particle with light mass ($\sim 580 \text{ MeV}$) and comparatively narrow width ($\sim 350 \text{ MeV}$). The value of M is close to $M \approx 2m_q$ (m_q being constituent quark mass) expected in the NJL type of models. Recently one of the present authors has argued¹⁰⁾ that our obtained values¹⁾ of m and Γ are consistent with the relation predicted in the linear model (LM). This fact seems to us to show that our "observed" η -particle is really a chiral partner of π -meson as a Nambu-Goldstone boson.

Here we also note an interesting argument^{10);17)} that the origin of repulsive core corresponds (at least in the low energy region where the structure of composite hadrons is negligible,) to the "compensating"⁴⁾ contact interaction, which is required from the current algebra and the PCAC.^{18);19)}

¹⁾ Recently Harada et al.¹⁶⁾ have made a similar analysis of η -scattering data as I leading to the η -existence with similar values of mass and width. However, they start from the viewpoint of Non-linear model and do not recognize it as a chiral partner of π -meson.

⁴⁾ The "effective" coupling, which includes both effects of an intermediate η -production and of the repulsive⁴⁾ interaction, becomes of a derivative type, while in the conventional Breit-Wigner formula, a non-derivative coupling of resonance is supposed. This seems to be a reason why η -meson existence has been overlooked for many years.

Historically the existence of η -particle, although it was anticipated from various viewpoints,^{2);4);5);20)} had been rejected for many years. One of the main reasons was that (i) the η has been missing in the $\pi\pi$ phase shift analyses,²¹⁾ and the other was due to (ii) the negative results of applications²²⁾ of L M to the low energy scattering and to the K_{14} decay form factors.

Concerning to (i), the present work may give a possible and clear solution. Recently there are several other phase shift analyses^{7);8);16)} resulting in η -particle existence. Also the problem (ii) should be reexamined¹⁰⁾ under the light of these recent progress of phenomenological search for η -particle. Finally we should like to note that recently a rather strong evidence²⁴⁾ for direct η production has been obtained in the central pp-collision. As a matter of fact, it was the motivation of our investigation.

References

- [1] S. Ishida, M.Y. Ishida, H. Takahashi, T. Ishida, K. Takamatsu and T. Tsuru, Prog.Theor.Phys. 95 (1996), 745.
- [2] R. Delbourgo and M.D. Scadron, Phys.Rev.Lett. 48 (1982), 379.
- [3] T. Hakioglu and M.D. Scadron, Phys.Rev.D 42 (1990), 941.
- [4] T. Hatsuda and T. Kunihiro, Prog.Theor.Phys. 74 (1985), 765; Phys.Rep. 247 (1994), 221.
- [5] Y. Kohyama, M. Takizawa et al., Phys.Lett.B 208 (1988), 165; Nucl.Phys.A 507 (1990), 617.
- [6] Review of Particle Properties, Phys.Rev.D 54 Part I (1996) 329 and also 355.
- [7] R. Kaminski, L. Lesniak and J.-P. Mallet, Phys.Rev.D 50 (1994), 3145, private communication.
- [8] N.A. Tomqvist and M. Roos, Phys.Rev.Lett. 76 (1996), 1575.
- [9] M. Taketani et al., Prog.Theor.Phys. Suppl.No.39 (1967); No.42 (1968). In particular see Chapter 7 (S.Otsuki, No.42, p.39) and also Chapter 6 (N.Hoshizaki, No.42, p.1).
- [10] M.Y. Ishida, Prog.Theor.Phys. 96 (1996), 853.
- [11] (i) G.G. Gayer et al., Nucl.Phys.B 75 (1974), 189, B. Hyams et al., Nucl.Phys.B 64 (1973), 134, (ii) V. Srinivasan et al., Phys.Rev.D 12 (1975), 681, (iii) L. Rosset et al., Phys.Rev.D 15 (1977), 574, (iv) A.A. Bel'kov et al., JETP Lett. 29 (1979), 579, (v) N.M. Casson et al., Phys.Rev.D 28 (1983), 1586.
- [12] S.D. Protopoulos et al., Phys.Rev.D 7 (1973), 1279.
- [13] M. Svec, Phys.Rev.D 53 (1996), 2343.
- [14] R. Kaminski, L. Lesniak and K. Rybicki, Henryk Niewodniczanski Institute of Nuclear Physics Report No.1730/PH (1996).
- [15] W. Hoogland et al., Nucl.Phys.B 126 (1977), 109; Nucl.Phys.B 69 (1974), 266, J.P. Pukop et al., Phys.Rev.D 10 (1974), 2055, M.J. Losty et al. Nucl.Phys.B 69 (1974), 185, D.Cohen et al., Phys.Rev.D 7 (1973), 661, E. Colton et al., Phys.Rev.D 3 (1971), 2028, W.D. Walker et al., Phys.Rev.Lett. 18 (1967), 630.
- [16] M. Harada, F. Sannino and J. Schechter, Phys.Rev.D 54 (1996), 1991.
- [17] M.Y. Ishida, Proc. of YITP conference, 1996, Kyoto, Yukawa Hall, to be published in Soryuusiron-kenkyuu; Proc. of the Workshop on Particle Physics at K arena with 50-GeV PS, KEK, Tsukuba, Japan Dec 19-20, 1996, edited by N. Sasao, S. Sugimoto, Y. Kuno and T. Komatsubara, Inst. for Nuclear Study, Univ. of Tokyo.
- [18] S. Weinberg, Phys.Rev. 166 (1968), 1568, T. Shiozaki, Prog.Theor.Phys. Suppl. Extra Number (44), 1968.
- [19] M. Bando, T. Kugo, and K. Yamawaki, Phys.Rep. 164 (1988), 217.
- [20] T. Morozumi, C.S. Lim and A.I. Sanda, Phys.Rev.Lett. 65 (1990), 404, M. Takizawa, T. Inoue and M. Oka, Prog.Theor.Phys. Suppl. 120 (1995), 335.
- [21] K.L. Au, D.Morgan and M.R. Pennington, Phys.Rev.D 41 (1987), 1633.

¹⁾ Ref.23) argues that the L M describes successfully both of the mentioned processes.

- [22] J. Gasser and H. Leutwyler, *Ann Phys.* 158 (1984), 142.
- [23] E. P. Shabalin, *Yad Fiz.* 49 (1989), 588 (*Sov. J. Nucl. Phys.* 49 (1989), 365).
- [24] D. A. Ide et al., *Phys. Lett. B* 397 (350), 1997; T. Ishida, Doctor Thesis of Tokyo University (1996).