

Spin asymmetries from Non Standard Physics with polarized proton beams at RHIC

P. Taxil and J.-M. Virey¹

Centre de Physique Théorique*, C.N.R.S. - Luminy, Case 907
F-13288 Marseille Cedex 9, France

and

Université de Provence, Marseille, France

Abstract

A new handed interaction between quarks could be at the origin of some small parity violating effects in the inclusive production of one-jet in polarized pp collisions. This interaction could originate from compositeness or from the presence of a new massive vector boson. The measurement of such asymmetries could be performed within a few years by using the RHIC collider as a polarized proton-proton collider.

*To appear in the proceedings of the
12th International Symposium on High Energy Spin Physics.*

PACS Numbers : 12.60.Cn; 13.87.-a; 13.88.+e; 14.70.Pw

Key-Words : New Gauge bosons, Jets, Polarization.

Number of figures : 1

October 1996

CPT-96/P.3391

anonymous ftp or gopher : cpt.univ-mrs.fr

*Unité Propre de Recherche 7061

¹ Moniteur CIES and allocataire MESR

email : Taxil@cpt.univ-mrs.fr

In a few years from now, the RHIC Spin Collaboration (RSC) will use the RHIC machine as a polarized pp collider. The center of mass energy will be as high as 500 GeV, the luminosity $\mathcal{L} = 2.10^{32} \text{ cm}^{-2} \cdot \text{s}^{-1}$ and the degree of beam polarization around 70% [1]. With these figures, spin asymmetries as small as 1% should be measurable in a few months running. For example, the parity conserving (PC) double-spin longitudinal asymmetry A_{LL} in inclusive one-jet production will be obtained with a very small error, hence allowing to test the spin structure of QCD. Adding the measurement of some parity violating (PV) asymmetries in W and Z boson production, it will be possible to determine very accurately the various polarized partonic distributions inside the proton (see [1,2] and also e.g. [3]). On the other hand, as was noticed some time ago [4], non conventional PV effects in hard-hadron reactions have never been searched for. Focusing on the inclusive production of one jet, one can define a double helicity PV asymmetry A_{LL}^{PV} as:

$$A_{LL}^{PV} = \frac{d\sigma_{(-)(-)} - d\sigma_{(+) (+)}}{d\sigma_{(-)(-)} + d\sigma_{(+) (+)}} \quad (1)$$

where $d\sigma_{\lambda,\lambda'}$ means the cross section in a given helicity configuration for the production of a single jet with transverse energy E_T , integrated over some rapidity interval around $y = 0$.

According to the Standard Model, this process is essentially governed by QCD plus a small contribution due to electroweak boson exchanges. The latter induces a small A_{LL}^{PV} which should be visible at RHIC. On the other hand, a new interaction between quarks could be at the origin of some deviations from the expected A_{LL}^{PV} , provided that it presents a particular chiral structure.

For instance, in the framework of compositeness, an effective handed interaction due to a contact term between quarks can be present [5] :

$$\mathcal{L}_{qqqq} = \epsilon \frac{\pi}{2\Lambda_{qqqq}^2} \bar{\Psi} \gamma_\mu (1 - \eta \gamma_5) \Psi \cdot \bar{\Psi} \gamma^\mu (1 - \eta \gamma_5) \Psi \quad (2)$$

where Ψ is a quark doublet, Λ is the compositeness scale, ϵ and η taking the values ± 1 . Furthermore, the presence of a new massive vector boson Z' , with right-handed and left-handed couplings to each quark of flavour i :

$$\mathcal{L}_{Z'} = \frac{g}{2 \cos \theta_W} Z'^\mu \bar{q}_i \gamma_\mu [C_{iL}(1 - \gamma_5) + C_{iR}(1 + \gamma_5)] q_i \quad (3)$$

could generate new amplitudes and lead to similar effects on the PV spin asymmetry provided its couplings to quarks are enhanced. Such "hadrophilic" or "leptophobic" neutral vector bosons Z' have been recently considered from the phenomenological point of view [6], in order to explain some possible deviations from the Standard Model predictions observed both at LEP and at CDF.

In two recent papers [7,8], we have studied the possibilities offered by the polarized RHIC collider for disentangling these new effects from the Standard spin asymmetry $A_{LL}^{PV}(SM)$ which is mainly due to the interference between the one-gluon exchange amplitude and the Z^0 exchange amplitude. An illustration is given in Fig. 1. One can see

that the Standard asymmetry (plain curve) rises with the transverse jet energy. This is simply due to the increasing importance of quark quark scattering relatively to other terms involving real gluons. The asymmetry is small but clearly visible at RHIC (the error bars refer to the statistical error with an integrated luminosity $L = 3.2fb^{-1}$). The dot-dashed curves show the effect of a contact term with a compositeness scale $\Lambda = 1.6$ TeV which corresponds to the present CDF bound. We have also assumed that parity were maximally violated by this interaction. The effect on A_{LL}^{PV} is clearly visible : if the contact term is indeed present this measurement should allow to get a unique information on its chirality structure since the sign of A_{LL}^{PV} is sensitive to the sign of the product $\epsilon.\eta$ ($\epsilon = -1$ means constructive interference, see [5,7]). From the absence of deviations from the expected A_{LL}^{PV} value, it will be possible to place a bound on Λ around 4 TeV, which competes with the Tevatron for an integrated luminosity $L = 100fb^{-1}$ (with unpolarized beams).

Concerning the effect of a new hadrophilic Z' , we restrict for illustration to the first model in [6] in which the new current exhibits a particular chiral structure with a dominance of the right-handed couplings on the left-handed couplings (for u quarks $C_{uR} \approx 2.C_{uL}$). As a consequence, A_{LL}^{PV} is negative : the dotted curve corresponds to a best fit value for the parameters of the model and the dashed curve to the maximal effect (the mass of the Z' is taken to be $M_{Z'} = 1$ TeV). Finally, the "SU(5)" curve corresponds to a "flipped SU(5)" gauge model inspired by superstring theories [9]. In this case, leptophobia occurs naturally as well as maximal parity violation in the up-quark sector. Leptophobia implies that the traditional Z' searches at future collider through the leptonic decay channel will be hopeless. Therefore, the PV spin asymmetry in the jet channel should be the best observable to pin down the presence of such an exotic object.

In conclusion, we have seen that some deviations from the expected Standard PV asymmetry which could be induced by New Physics beyond the Standard Model should be visible at RHIC, despite the relatively low value of the c.m energy. This is mainly allowed by the high luminosity and by the high degree of polarization of the beams. Therefore, the very precise measurements performed by the RSC could contribute significantly to the analyses of some possible manifestations of New Physics beyond the Standard Model.

References

- [1] G. Bunce et al., *Polarized protons at RHIC*, Particle World, **3**, 1 (1992)
- [2] see H. En'yo, N. Hayashi, S.F. Heppelman, Y. Makdisi, contributions to this conference.
- [3] C. Bourrely, J. Ph. Guillet and J. Soffer, Nucl. Phys. **B361**, 72 (1991)
- [4] M. Tannenbaum, in *Polarized Collider Workshop*, J. Collins, S.F. Heppelmann and R.W. Robinett eds., AIP Conf. Proceedings **223**, AIP, New York, 1990, p. 201.
- [5] E. Eichten, K.Lane and M. Peskin, Phys. Rev. Lett. **50**, 811 (1983) ;
E. Eichten et al., Rev. Mod. Phys. **56** (1984) 579
- [6] G. Altarelli et al., Phys. Lett. **B375**, 292 (1996);
P. Chiappetta et al. Phys. Rev. **D54**, 789 (1996)
- [7] P. Taxil and J.M. Virey, Phys. Lett. **B364**, 181 (1995).
- [8] P. Taxil and J.M. Virey, Phys. Lett. **B383**, 355 (1996).
- [9] J.L. Lopez and D.V. Nanopoulos [hep-ph 9605359]
- [10] M. Gluck, E. Reya and W. Vogelsang, Phys. Lett. **B359**, 201 (1995)

Figure captions

Fig. 1 The asymmetry A_{LL}^{PV} for inclusive one-jet production at RHIC as a function of the transverse jet energy E_T . The calculations are performed with GRV [10] distributions.




Fig 1