

QCD Description of Particle Spectra up to LEP-1.5 Energies and the Running of α_s

WOLFGANG OCHS

*Max-Planck-Institut für Physik
(Werner-Heisenberg-Institut)
Föhringer Ring 6, D-80805 Munich, Germany*

Abstract

Recent results on the energy spectra in QCD jets are reported. Within the Modified Leading Logarithmic Approximation (MLLA) and the Local Parton Hadron Duality (LPHD) model one finds a very good description of the e^+e^- data from the lowest up to the LEP 1.5 energies. A model with fixed α_s can be excluded, already in certain finite energy intervals. The fits also extrapolate smoothly into the region of small particle energies, in particular, the data follow a scaling prediction for the low energy limit derived from the colour coherence of the soft gluon emission.

to appear in the Proc. of the 28th International Conference on High Energy Physics, 25-31 July 1996, Warsaw, Poland.

QCD DESCRIPTION OF PARTICLE SPECTRA UP TO LEP-1.5 ENERGIES AND THE RUNNING OF α_s

WOLFGANG OCHS

*Max-Planck-Institute für Physik (Werner-Heisenberg-Institut)
Föhringer Ring 6, D-80805 Munich, Germany*

Recent results on the energy spectra in QCD jets are reported. Within the Modified Leading Logarithmic Approximation (MLLA) and the Local Parton Hadron Duality (LPHD) model one finds a very good description of the e^+e^- data from the lowest up to the LEP 1.5 energies. A model with fixed α_s can be excluded, already in certain finite energy intervals. The fits also extrapolate smoothly into the region of small particle energies, in particular, the data follow a scaling prediction for the low energy limit derived from the colour coherence of the soft gluon emission.

1 Introduction

One of the important predictions of perturbative QCD on the intrinsic structure of jets concerns the energy spectra of particles. The particles of low energy E are not multiplied with increasing jet energy E_{jet} because of colour coherence in the cascading process and this yields the bell shaped spectrum in the variable $\xi = \log \frac{E_{jet}}{E}$, the so-called “hump-backed plateau”.^{1,2}

Predictions on the spectrum have been carried out in the MLLA which takes into account the leading double logarithmic (DLA) results and all next-to-leading corrections of order $\sqrt{\alpha_s}$. Terms of higher order are included as well, although not completely, which allow to fulfill the initial condition for the parton cascade at threshold. These predictions, at the parton level, involve only two parameters: the QCD scale Λ which determines the running of the coupling α_s and the transverse momentum cut-off Q_0 of the gluon emission. Remarkably, the observed hadron spectra are rather well described by the spectrum of partons if a low value $Q_0 \sim 250$ MeV of the order of hadronic masses is used and this observation has led to the LPHD hypothesis³. As a justification of this approach is not yet available at a fundamental level it seems important to determine its range of applicability and its limitations. Also one would like to know to what extent the predictions are sensitive to QCD as the underlying theory. In this contribution the sensitivity of the energy spectra to the running α_s and the colour coherence - two properties specific to QCD as a field theory - are addressed. More details can be found in refs.^{4,5,6}.

2 Moment Analysis of Particle Spectra

To analyse the effect of the running α_s it is convenient to work with the moments of the ξ -spectrum of particles which evolve independently of each other with energy. They are defined by $\langle \xi^q(Y, \lambda) \rangle = \int d\xi \xi^q D(\xi, Y, \lambda) / \mathcal{N}$ where we use the logarithmic variables $Y = \log(E_{jet}/Q_0)$ and $\lambda = \log(Q_0/\Lambda)$; the normalization is by the multiplicity \mathcal{N} . We consider finally the cumulant moments K_q which are obtained from the $\langle \xi^q \rangle$ moments by $K_1 \equiv \langle \xi \rangle = \bar{\xi}$, $K_2 \equiv \sigma^2 = \langle (\xi - \bar{\xi})^2 \rangle$, $K_3 = \langle (\xi - \bar{\xi})^3 \rangle$, $K_4 = \langle (\xi - \bar{\xi})^4 \rangle - 3\sigma^4$, ...; also one introduces the reduced cumulants $\kappa_q \equiv K_q / \sigma^q$, in particular the skewness $s \equiv \kappa_3$ and the kurtosis $k \equiv \kappa_4$. The cumulant moments $K_q(Y, \lambda)$ behave at high energies like

$$K_q(Y, \lambda) = K_q(Y_0, \lambda) + \int_{Y_0}^Y dy \left(-\frac{\partial}{\partial \omega} \right)^q \gamma_\omega[\alpha_s(y, \lambda)] \Big|_{\omega=0} \quad (1)$$

where Y_0 is the initial energy. Here $\gamma_\omega[\alpha_s(y, \lambda)]$ denotes the anomalous dimension of the Laplace transform of the ξ -spectrum $dn/d\xi \equiv D(\xi, Y, \lambda)$. Eq.(1) shows directly the sensitivity of the cumulant moments to the running of α_s . In particular, for fixed coupling $K_q \sim Y$ at high energies. In our applications, we take into account the initial condition at threshold for the jet to consist of only one parton which implies $K_q = 0$ at $Y_0 = 0$.

The MLLA prediction for the ξ^q moments can be written for arbitrary Q_0 and Λ as⁷:

$$\langle \xi^q \rangle = \frac{1}{\mathcal{N}} \sum_{k=0}^q \binom{q}{k} (N_1 L_k^{(q)} + N_2 R_k^{(q)}) \quad (2)$$

where $N_1, N_2, L_k^{(q)}$ and $R_k^{(q)}$ are known functions of $a = 11N_c/3 + 2n_f/3N_c^2$, $b \equiv (11N_c - 2n_f)/3$, and λ where n_f, N_c denote the numbers of flavours and colours. Comparing these formulae to the experimental data on moments,⁴ best agreement is found for the “limiting spectrum”, where the two parameters coincide, i.e., $Q_0 = \Lambda$, or $\lambda = 0$. In this case the formulae simplify and the moments can be expressed⁷ in terms of the parameter $B \equiv a/b$ and the variable $z \equiv \sqrt{16N_c Y/b}$ as:

$$\begin{aligned} \frac{\langle \xi^q \rangle}{Y^q} &= P_0^{(q)}(B+1, B+2, z) + \\ &+ \frac{2}{z} \frac{I_{B+2}(z)}{I_{B+1}(z)} P_1^{(q)}(B+1, B+2, z) \end{aligned} \quad (3)$$

where $P_0^{(q)}$ and $P_1^{(q)}$ are polynomials of order $2(q-1)$ in z .

These expressions extrapolate smoothly to threshold, where they are determined by the initial condition for a single parton. Similarly one can derive the complete MLLA results for fixed α_s ⁴, which simplify for high energies to

$$\bar{\xi}_{fix} = \left[1 + \frac{\eta}{\bar{\gamma}_0} \right] \frac{Y}{2}, \quad \sigma_{fix}^2 = \frac{\gamma_0^2}{4\bar{\gamma}_0^3} Y \quad (4)$$

$$s_{fix} = -\frac{3\eta}{\gamma_0} \frac{1}{\sqrt{\gamma_0 Y}}, \quad k_{fix} = \frac{3(4\eta^2 - \gamma_0^2)}{\gamma_0^2 \bar{\gamma}_0} \frac{1}{Y} \quad (5)$$

where $\eta = a\gamma_0^2/8N_c$, $\bar{\gamma}_0 \equiv \sqrt{\gamma_0^2 + \eta^2}$ and $\gamma_0 = \sqrt{2N_c\alpha_s/\pi}$.

3 Relating parton and hadron spectra

Strictly speaking the QCD results are only reliable for $E \gg Q_0$. In order to determine the moments one has to integrate the spectra over the full range and to extrapolate to the soft limit. The experimental particle densities are usually given as function of momenta. The same kinematic limits of parton and hadron spectra is obtained if the same effective mass Q_0 is assigned to the charged particles which also limits the energy of the partons ($E_h = E_p \geq Q_0$).

Furthermore, a common behaviour of the parton and hadron spectra near the boundary can be obtained if the spectra are related by^{1,4}

$$E_h \frac{dn(\xi_E)}{dp_h} = K_h E_p \frac{dn(\xi_E)}{dp_p} \equiv K_h D(\xi_E, Y, \lambda) \quad (6)$$

at the same energy E or $\xi_E \equiv \xi = \log(E_{jet}/E)$, where $E_h = \sqrt{p_h^2 + Q_0^2}$ and $E_p = p_p$; K_h denotes a normalization parameter. With this choice both spectra vanish linearly in ξ ; this corresponds to the invariant distribution of hadrons $E \frac{dn}{d^3p}$ approaching a constant value in the soft limit $p \rightarrow 0$, as suggested by the experimental data. The prescription (6) represents a minimal extension of the QCD prediction for $E \gg Q_0$ towards the boundary at $E = Q_0$ but is not unique; other prescriptions using a smaller value of $Q_0 \sim m_\pi$ have also been applied (see discussion in⁶).

4 Comparison with experimental data

The moments $\langle \xi^q \rangle$ are determined⁴ from the spectra $E dn/dp$ vs. ξ_E after appropriate transformation of the measured $x_p = 2p/\sqrt{s}$ spectra of charged particles using $E = \sqrt{p^2 + Q_0^2}$ and therefore depend on Q_0 . First a fit of the two parameters Q_0 and Λ is obtained by comparing the moments for a selected Q_0 with the theoretical predictions from Eq.(2) for different Λ . The best agreement with the data was obtained for $Q_0 \approx \Lambda \approx (270 \pm 20)$ MeV. It was not possible to obtain a satisfying description for the fixed α_s case. Choosing $\gamma_0 = 0.64$ and the same $Q_0 = 270$ MeV a good description of the multiplicity \bar{N} and the slope of $\bar{\xi}$ vs. Y could be obtained but not for the other quantities. In Fig. 1 we show the evolution of the first cumulant moments K_q with energy for running α_s and fixed α_s . Note that these predictions depend only on the two parameters Q_0 and Λ which actually coincide in the fit. The absolute normalization of the moments is given at threshold $Y_0=0$ by $K_q=0$.

As this is an application of perturbative QCD to an extreme limit one could make the weaker (more conventional) assumption and choose a higher starting energy, say $Y_0 = 2$ ($\sqrt{s} \approx 4$ GeV). In this case for each moment an extra free parameter $K_q(Y_0)$ had to be introduced which, however, would not improve the fit for running α_s essentially; a backward evolution from Y_0 would again reproduce approximately the initial condition at threshold. Therefore, the initial condition yields the highly constrained fit with only two parameters (the normalization parameter K_h only enters the multiplicity \bar{N} , not shown here).

In case of fixed α_s one may ask the same ques-




Figure 1: The first four cumulant moments K^q of the charged particles' energy spectra Edn/dp vs. ξ_E , shown as a function of Y for $Q_0 = 270$ MeV. Predictions of the Limiting Spectrum of MLLA with running α_s (solid lines), of MLLA with fixed α_s ($= 0.214$) (dashed lines) and of MLLA with fixed α_s normalized by hand to the experimental point at $\sqrt{s} = 44$ GeV (dotted lines) are also shown; in all cases $n_f = 3$ (from Ref. 4).




Figure 2: Charged particle inclusive momentum distribution at LEP-1.5 from ALEPH (diamonds), DELPHI (squares) and OPAL Collaborations (triangles) in comparison with theoretical predictions of the Limiting Spectrum with $Q_0 = 270$ MeV (solid line). Dashed lines show the predictions of the Limiting Spectrum after correction for kinematical effects (from Ref. 5).

tion, whether the fit could be improved by choosing a higher starting value Y_0 , i.e. to assume a transition from a running coupling to a fixed coupling regime at Y_0 (This would correspond to a parallel shift of the dashed curve to the data at Y_0). As can be seen in Fig. 1 the energy dependence of the fixed and running α_s curves is quite different so that an improvement cannot be obtained in a finite energy interval. Rather, such a procedure can be excluded for a Y interval of about 1-2 units. An example is shown by the dotted curve which corresponds to $Y_0 = 4.3$ ($\sqrt{s} = 44$ GeV). The trend at yet larger energies is again different, in particular for the higher moments.

The dependence on the number of flavours n_f has been studied as well ^{4,5}. As the jet evolution is dominated at all energies by the gluon emission at the lower transverse momenta, the choice $n_f = 3$ is a good approximation up to LEP energies.

The recent LEP-1.5 data are also well accounted for ⁵ by the same theoretical scheme as is shown in Fig. 2. Here the experimental ξ -spectrum is shown in comparison with the limiting spectrum (setting $\xi_p = \xi_E$) which terminates at $Y \sim 5.3$ and after rescaling $\xi_E \rightarrow \xi_p$ with relation (6) which takes into account the boundary effects. Also the moments of the distribution and the position of the maximum are well reproduced by the theoretical predictions at this energy.




Figure 3: Invariant density $E dn/d^3p$ of charged particles in e^+e^- annihilation as a function of the particle energy $E = \sqrt{p^2 + Q_0^2}$ at $Q_0 = 270$ MeV at various cms energies in comparison with the MLLA predictions (from Ref. 6).

5 Soft limit of energy spectrum

Having observed the good fit of the QCD-LPHD prediction not only towards high but also towards low cms energies \sqrt{s} we now turn to the behaviour of the fits at low particle energy E . As the emission of very soft gluons from all other partons in the jet is coherent the production rate of such particles is nearly energy independent³. Indeed, the analytic results in the DLA and MLLA show the scaling behaviour of the spectrum in the soft limit and the production rate is essentially determined by the colour charge of the initial partons⁶. Remarkably, the observed hadrons follow the trend of this prediction as can be seen in Fig. 3 which shows the invariant particle density as a function of particle energy E (assuming the effective charged particle mass $Q_0 = 270$ MeV). The data from all energies in e^+e^- annihilation approach a common limit for $E \rightarrow Q_0$ ($p \rightarrow 0$). The curves which represent the MLLA calculation show again the scaling limit and reproduce the dependence on energy E and cms -energy \sqrt{s} rather well. This suggests that LPHD persists towards very low particle energies. Various further tests of this picture have been proposed⁶ by studying the dependence of the soft limit I_0 on the colour charge of the primary partons in $e^+e^- \rightarrow 3$ jets, deep inelastic scattering processes and hadronic collisions.

6 Conclusions

The particle energy spectra are well described by the QCD parton shower in the MLLA assuming a close similarity between partons and hadrons according to the LPHD approach. This similarity appears to work not only for the highest but also down to rather low cms energies as well as to low particle energies. This agreement is by no means trivial. For example, replacing the running α_s by the fixed α_s yields predictions quite incompatible with experiment, also if the threshold for the onset of a fixed α_s regime is increased. The scaling prediction derived from colour coherence for the soft particles works well for hadrons.

This may suggest⁴ to view the final stage of hadron production through resonances, as explicitly incorporated into Monte Carlo hadronization models, as being dual to the partonic evolution with running coupling down to the low scale of a few hundred MeV.

Acknowledgements

I would like to thank Valery Khoze and Sergio Lupia for many discussions and the collaboration on the subjects of this talk.

References

1. Yu. L. Dokshitzer, V. S. Fadin, V. A. Khoze, *Phys. Lett. B* **115**, 242 (1982); *Z. Phys. C* **15**, 325 (1982).
2. A. Bassetto, M. Ciafaloni, G. Marchesini and A. H. Mueller, *Nucl. Phys. B* **207**, 189 (1982).
3. Ya. I. Azimov, Yu. L. Dokshitzer, V.A. Khoze, S.I. Troyan, *Z. Phys. C* **27**, 65 (1985); *Z. Phys. C* **31**, 213 (1986).
4. S. Lupia and W. Ochs, *Phys. Lett. B* **365**, 339 (1996) and paper in preparation.
5. V. A. Khoze, S. Lupia and W. Ochs, *Phys. Lett. B* **386**, 451 (1996).
6. V. A. Khoze, S. Lupia and W. Ochs, "QCD coherence and the soft limit of the energy spectrum", hep-ph/9610204 (Sept. 1996).
7. Yu. L. Dokshitzer, V. A. Khoze and S. I. Troyan, *Int. J. Mod. Phys. A* **7**, 1875 (1992).