

QUARKONIUM PRODUCTION VIA FRAGMENTATION: A REVIEW ¹

K. Sridhar, Theory Group, Tata Institute of Fundamental Research,
Homi Bhabha Road, Bombay 400 005, India

Abstract

We review the physics of quarkonium production at high energy colliders, with a specific emphasis on production *via* fragmentation. We discuss the fragmentation picture for the production of quarkonia at large- p_T at the Tevatron and emphasise the importance of the colour-octet components of the quarkonium wave function which are included in the fragmentation functions. Applications of these theoretical ideas to other processes, both at the Tevatron and at other experiments like LEP, HERA and the LHC are important as they will provide stringent tests of these ideas.

The production of quarkonia has conventionally been described in terms of the colour-singlet model [1, 2]. In this model, a non-relativistic approximation is used to describe the binding of the heavy-quark pair produced via parton-fusion processes, into a quarkonium state. The heavy-quark pair is projected onto a physical quarkonium state using a colour-singlet projection and an appropriate spin-projection. It is also necessary to account for J/ψ 's produced *via* electromagnetic decays of the P -states, *viz.*, the χ 's, since inclusive J/ψ production is measured in experiments. In the model, both the direct S -state production and the indirect production *via* P -state decays are accounted for. This model has been successfully applied [2] to describe large- p_T J/ψ production at the relatively low energies in the ISR experiment. In going from the ISR energy ($\sqrt{s} = 63$ GeV) to the UA1 energy ($\sqrt{s} = 630$ GeV), it was found [3] that the production of b -quarks and their subsequent decay becomes an important source of J/ψ production. While at the UA1 experiment it was not possible to separate

¹Presented at the Fourth Workshop on High Energy Physics Phenomenology (WHEPP IV), Calcutta, India, January, 1996.

the b -quark contribution, the use of the silicon-vertex detector at the CDF experiment in studying J/ψ production in $\bar{p}p$ collisions at the Tevatron ($\sqrt{s} = 1.8$ TeV) allows the subtraction of the b -quark contribution from the total yield, thereby providing a measurement of direct inclusive J/ψ production at large- p_T . The inclusive J/ψ production cross-section measured by the CDF experiment [4] turned out to be an order of magnitude larger than the prediction of the colour-singlet model.

In fact, even before the experimental results from the CDF collaboration were available, it was shown by Braaten and Yuan [5] that in addition to the parton fusion contributions taken into account in the colour-singlet model, fragmentation of gluons and charm quarks could be an important source of large- p_T J/ψ production. In the fragmentation process, one considers the production of charmonia from final-state gluons or charm quarks which have large p_T , but almost zero virtuality. The fragmentation contribution is computed by factorising the cross-section for the process $AB \rightarrow (J/\psi, \chi_J)X$ (where A, B denote hadrons) into a part containing the hard-scattering cross-section for producing a gluon or a charm quark and a part which specifies the fragmentation of the gluon or the charm quark into the required charmonium state, i.e.

$$d\sigma(AB \rightarrow (J/\psi, \chi_J)X) = \sum_i \int_0^1 dz \, d\sigma(AB \rightarrow iX) D_{i \rightarrow (J/\psi, \chi_J)}(z, \mu), \quad (1)$$

where i is the fragmenting parton (either a gluon or a charm quark). $D(z, \mu)$ is the fragmentation function and z is the fraction of the momentum of the parent parton carried by the charmonium state. Because the gluon or the c -quark fragments into a heavy quarkonium state, the fragmentation function can be computed perturbatively, in the same spirit as in the colour-singlet model. This yields the fragmentation function at an initial scale μ_0 which is of the order of m_c . If the scale μ is chosen to be of the order of p_T , then large logarithms in μ/m_c appear which have to be resummed using the usual Altarelli-Parisi equation:

$$\mu \frac{\partial}{\partial \mu} D_{i \rightarrow (J/\psi, \chi_i)}(z) = \sum_j \int_z^1 \frac{dy}{y} P_{ij}\left(\frac{z}{y}, \mu\right) D_{j \rightarrow (J/\psi, \chi_i)}(y), \quad (2)$$

where the P_{ij} are the splitting functions of a parton j into a parton i . Only the fragmentation of gluons and charm quarks need be considered since the light quark contributions are expected to be very small.

The fragmentation function at the initial scale for the production of a quarkonium state \mathcal{O} from a gluon, $D_{g \rightarrow \mathcal{O}}(z, 2m_c)$, can be obtained by factorising the process $gg \rightarrow \mathcal{O} + n$ gluons into two parts. The first part specifies the production of a large- p_T but low-virtuality gluon. In the limit that the virtuality can be completely neglected with respect to the p_T of the gluon, this factor reduces to the on-shell scattering amplitude for the process $gg \rightarrow gg$. The second part is the amplitude for the process $g^* \rightarrow \mathcal{O} + n - 1$ gluons. The real gluon production process has the form $\mathcal{M}_\alpha \epsilon^\alpha(q)$, whereas the fragmentation process has the form $\mathcal{M}_\alpha(-ig^{\alpha\beta}/q^2)\mathcal{A}_\beta$, where the q^2 in the denominator is of the order of m_c^2 . The fragmentation probability is the ratio of the rates for these two processes. The fragmentation function is obtained after integrating over the invariant mass of the fragmenting gluon

from $4m_c^2$ to infinity. The charm-quark fragmentation functions are also calculated in a similar manner. While this calculation goes through in a straightforward manner for S -state quarkonia, for the P -states it is more complicated. This has to do with a non-factorising logarithmic infrared divergence that is encountered in the colour-singlet treatment of P -state fragmentation functions [6, 7]. The divergence arises from soft-gluon radiation that produces the 3P_J bound state. The problem arises because of assuming that the $c\bar{c}$ pair can exist only in a colour-singlet state. This assumption is tantamount to neglecting the relative velocity v between the c and the \bar{c} . However v is, in general, not negligible and $O(v^2)$ corrections need to be taken into account. A formulation based on non-relativistic QCD (NRQCD), using the factorisation method has been recently carried out [8], and in this formulation the quarkonium wave-function admits of a systematic expansion in powers of v in terms of Fock-space components : for example, the χ states have the colour-singlet P -state component at leading order, but there exist additional contributions at non-leading order in v , which involve octet S -state components; i.e.

$$|\chi_J\rangle = O(1)|Q\bar{Q}[^3P_J^{(1)}]\rangle + O(v)|Q\bar{Q}[^3S_1^{(8)}]g\rangle + \dots \quad (3)$$

The importance of the colour-octet components had already been seen in the analysis of the decays of the χ states [9] where the colour-singlet analysis [10] revealed a similar problem with the logarithmic infrared singularity. But the octet component allows the infrared singularity to be absorbed *via* a wave-function renormalisation, without having to introduce an arbitrary infrared cut-off. Thus, a consistent perturbation theory for the P -states necessitates the inclusion of the octet component of the wave-function. It should be pointed out that the colour-singlet term for the production of a P -state already starts at order v , and the colour-octet production *via* an octet 3S_1 state also appears at the same order in v .

The gluon- and charm-fragmentation functions have been calculated [5, 6, 7, 11, 12, 13] and using these as inputs several authors [14] have computed the contributions coming from fragmentation to the total J/ψ yield and have found that the gluon fragmentation contribution significantly increases the cross-section, and the order-of-magnitude discrepancy between the theory and the data from the CDF experiment can be resolved. The analyses of the CDF data, thus, demonstrates the importance of the fragmentation mechanism as an important source of J/ψ production at large p_T .

For S -state resonances like the J/ψ and the ψ' , the octet contribution is suppressed by powers of v . Further, the S -wave amplitude is not infrared divergent and can, therefore, be described in terms of a single colour-singlet matrix-element. But the CDF measurement [15] of the ratio of J/ψ 's coming from χ decays to those produced directly shows that the direct S -state production is much larger than the theoretical estimate. It has been suggested [16] that a colour octet component in the S -wave production coming from gluon fragmentation as originally proposed in Ref. [17], can explain this J/ψ anomaly. This corresponds to a virtual gluon fragmenting into an octet 3S_1 state which then makes a double E1 transition into a singlet 3S_1 state. While this process is suppressed by a factor of v^4 as compared to the colour-singlet process, it is enhanced by a factor of α_s^2 . One can fix the value of the colour-octet matrix-element by normalising to the data on direct J/ψ production cross-section from the

CDF experiment. The colour-octet contribution to S -state production has also been invoked [17] to explain the large ψ' cross-section measured by CDF [4], but there can be a sizeable contribution to this cross-section coming from the decays of radially excited P -states [18].

The long-distance octet matrix elements are not calculable, and have to be fitted by comparing with various experiments. After the suggestion of Ref. [17] that the S -state production could also be dominated by octet components, the production of charmonium and bottomonium resonances as direct octet production amplitudes via parton fusion has been computed in Ref. [19], rather than using the fragmentation approximation which holds only for large- p_T . It is found that for $p_T \geq 10$ GeV the fusion production of quarkonia *via* octet channels matches quite well with the predictions of the fragmentation hypothesis, whereas in the range $5 < p_T < 10$ GeV non-fragmentation contributions also turn out to be quite important. In effect, as shown in Ref. [19], only a linear combination of octet matrix-elements can be determined by fitting to the Tevatron data. However, since the colour-octet matrix elements are universal, they will also appear in other quarkonium production processes. So it is important to have independent tests of the colour-octet production mechanisms in other experiments ².

Several suggestions have been made in the literature which could provide more stringent quantitative tests of the colour-octet contributions and the fragmentation picture. In particular these processes can be used to fix the colour-octet non-perturbative matrix elements and determine the shape and size of the fragmentation functions. The following predictions of the theory for various experimental observables have been made:

1. Charmonium and bottomonium production at LEP provide a laboratory for testing several features of the theory of quarkonium production. The colour-singlet production process $Z \rightarrow J/\psi c\bar{c}$ [22] turns out to be two orders of magnitude larger than the process $Z \rightarrow J/\psi gg$ [23] and this is due to the fact that the former is enhanced by a fragmentation contribution which is not suppressed by m_c^2/m_Z^2 [11]. The resulting prediction for prompt J/ψ production is of the order of 3×10^{-5} , which is almost an order of magnitude below the experimental number for the branching fraction obtained from LEP [24]. Another fragmentation process $Z \rightarrow q\bar{q}g^*$ with the virtual gluon then fragmenting into a J/ψ through the colour-singlet processes $g^* \rightarrow J/\psi gg$ and $g^* \rightarrow \chi g$ has been studied [25]. Recently, the colour-octet contributions to J/ψ production in this channel have been studied [26, 27] and it is found that the inclusion of the colour-octet contributions in the fragmentation functions results in a predictions for the branching ratio which is 1.4×10^{-4} which is compatible with the measured values of the branching fraction from LEP [24]. Also, the energy distribution of the J/ψ coming from the colour-octet process is very soft and is peaked at around $z = 0.1$, which can be used to distinguish it from the colour-singlet production which gives a hard energy spectrum. The predictions from the colour-octet fragmentation process for the

²Recent papers, which have appeared after this talk was presented, show that a different linear combination of the same colour octet-matrix elements that appear in the Tevatron analysis also appears in the analyses of photoproduction [20] and hadroproduction experiments [21]. These analyses provide an important cross-check on the colour-octet contributions.

production of bottomonia [27] are also in reasonable agreement with the measurements from LEP [28].

2. The production of J/ψ in low energy e^+e^- machines can also provide a stringent test of the colour-octet mechanism [29]. In this case, the colour-octet contributions dominate near the upper endpoint of the J/ψ energy spectrum, and the signature for the colour-octet process is a dramatic change in the angular distribution of the J/ψ near the endpoint. The angular distribution has a general form $1 + A(E)\cos^2\theta$. While the colour-singlet model predicts that $A \approx -0.84$ for $E \approx E_{\max}$, the colour-octet prediction is that $A(E)$ should change sign suddenly near the endpoint of the spectrum and approach a value of A greater than $\sim +0.6$.
3. One striking prediction of the colour-octet fragmentation process both for $p\bar{p}$ colliders and for J/ψ production at the Z -peak, is that the J/ψ coming from the process $g \rightarrow J/\psi X$ is produced in a transversely polarised state [30]. For the colour-octet $c\bar{c}$ production, this is predicted to be a 100% transverse polarisation, and heavy-quark spin symmetry will then ensure that non-perturbative effects which convert the $c\bar{c}$ to a J/ψ will change this polarisation only very mildly. This spin-alignment can, therefore, be used as a test of colour-octet fragmentation.
4. The fragmentation process has also definite predictions for the large- p_T inelastic J/ψ photoproduction at HERA [31]. In the photoproduction process, the colour-singlet fusion process dominates over the charm-quark and gluon fragmentation process upto a p_T of 8–10 GeV. The fragmentation contributions become important only for $p_T > 10$ GeV. Even here, it is the charm-quark fragmentation that dominates over the gluon fragmentation process. The gluon fragmentation process is negligibly small over the entire range of p_T considered. The J/ψ cross-section at large- p_T will, therefore, provide a handle on the charm fragmentation functions, and provide information which is complementary to that obtained from Tevatron which is gluon-dominated ³.
5. The associated production of a $J/\psi + \gamma$ also provides a test of the fragmentation picture [33]. This process is dominated by the colour-singlet fusion process. If the colour-octet fragmentation picture is the true description of the Tevatron J/ψ cross-section enhancement, then we would expect that there would not be a similar enhancement of the $J/\psi + \gamma$ cross-section, and this would be very close to the colour-singlet fusion prediction ⁴.
6. Double J/ψ production via double gluon fragmentation has been studied in the context of the Tevatron [35]. The colour-octet fragmentation mechanism tends to increase the production rate for this process and the ratio of the cross-section for this process to

³A recent paper [32] deals with photoproduction of quarkonia at HERA via colour-octet production processes in detail.

⁴For a more recent discussion of the colour-octet contributions to this process, see [34].

that of single J/ψ production is of the order of 10^{-5} . Given the large number of single J/ψ events seen at the Tevatron, double J/ψ production may also be observable.

7. J/ψ and ψ' production in pp collisions at centre-of-mass energies of 14 TeV at the LHC also provides a crucial test of colour-octet fragmentation [36]. As in the Tevatron, gluon fragmentation is again the single most important source of quarkonium production – except that the magnitude of the gluon fragmentation contribution at the LHC completely overwhelms the contributions coming from either direct production or from charm fragmentation. Also given the large p_T coverage possible at the LHC, the fragmentation picture becomes almost exact and any non-fragmentation contribution is negligibly small. An experimental determination of the J/ψ and ψ' cross-sections at the LHC will provide a verification of the fragmentation picture and will also help to measure the colour-octet non-perturbative parameters that dominate the gluon fragmentation functions.

In summary, the theory of quarkonium production has changed quite dramatically in an attempt to understand the anomalously large production rates for charmonia at the Tevatron. Fragmentation and the inclusion of colour-octet components of the quarkonium wave-function have provided a way to resolve the discrepancy between theory and experiment, but at the expense of introducing undetermined non-perturbative parameters. However, these parameters have a rigorous definition as matrix elements of colour-octet operators in NRQCD, and are shown to be universal. This universality helps us to predict the rates for quarkonium production processes once these matrix-elements have been determined by fitting to one experiment. The success of the program is, therefore, crucially dependent on indentifying what physical observable determines a specific combination of matrix-elements. We have reviewed some proposals in the literature, which show how the colour-octet fragmentation picture used to describe the Tevatron data can be tested in other experiments. A preliminary study [37] of higher-order corrections to the fragmentation contribution in the case of 1S_0 production has been made, and shows the possible importance of the higher-order corrections to fragmentation. More work needs to be done particularly in the understanding of higher-order corrections in α_s and v .

References

- [1] E.L. Berger and D. Jones, *Phys. Rev.* **D 23** (1981) 1521.
- [2] R. Baier and R. Rückl, *Z. Phys.* **C 19** (1983) 251.
- [3] E.W.N. Glover, A.D. Martin and W.J. Stirling, *Z. Phys.* **C 38** (1988) 473.
- [4] F. Abe et al., *Phys. Rev. Lett.* **69** (1992) 3704; *Phys. Rev. Lett.* **71** (1993) 2537; K. Byrum, CDF Collaboration, Proceedings of the 27th International Conference on High Energy Physics, Glasgow, (1994), eds. P.J. Bussey and I.G. Knowles (Inst. of Physics Publ.) p.989.
- [5] E. Braaten and T.C. Yuan, *Phys. Rev. Lett.* **71** (1993) 1673.
- [6] E. Braaten and T.C. Yuan, *Phys. Rev.* **D 50** (1994) 3176.
- [7] J.P.Ma, *Nucl. Phys.* **B 447** (1995) 405.
- [8] G.T. Bodwin, E. Braaten and G.P. Lepage, *Phys. Rev.* **D 51** (1995) 1125.
- [9] G.T. Bodwin, E. Braaten and G.P. Lepage, *Phys. Rev.* **D 46** (1992) R1914.
- [10] R. Barbieri, R. Gatto and E. Remiddi, *Phys. Lett.* **B 61** (1976) 465.
- [11] E. Braaten, K.Cheung and T.C. Yuan, *Phys. Rev.* **D 48** (1993) 4230.
- [12] Y.Q. Chen, *Phys. Rev.* **D 48** (1993) 5181.
- [13] T.C. Yuan, *Phys. Rev.* **D 50** (1994) 5664.
- [14] E. Braaten, M.A. Doncheski, S. Fleming and M. Mangano, *Phys. Lett.* **B 333** (1994) 548; D.P. Roy and K. Sridhar, *Phys. Lett.* **B 339** (1994) 141; M. Cacciari and M. Greco, *Phys. Rev. Lett.* **73** (1994) 1586.
- [15] G. Bauer, CDF Collaboration, presented at the "Xth Topical Workshop on $p\bar{p}$ collisions", Fermilab (1995).
- [16] M. Cacciari, M. Greco, M. Mangano and A. Petrelli, *Phys. Lett.* **B 356** (1995) 553.
- [17] E. Braaten and S. Fleming, *Phys. Rev. Lett.* **74** (1995) 3327.
- [18] F.E. Close, *Phys. Lett.* **B 342** (1995) 369; D.P. Roy and K. Sridhar, *Phys. Lett.* **B 345** (1995) 537.
- [19] P. Cho and A.K. Leibovich, *Phys. Rev.*, **D 53** (1996) 150; *Phys. Rev.*, **D 53** (1996) 6203.

- [20] N. Cacciari and M. Krämer, *Phys. Rev. Lett.* **76** (1996) 4128; J. Amundson, S. Fleming and I. Maksymyk, UTTG-10-95, hep-ph/9601298.
- [21] S. Gupta and K. Sridhar, *Phys. Rev.* **D 54** (1996) 5545; M. Beneke and I. Rothstein, *Phys. Rev.* **D 54** (1996) 2005; W.-K. Tang and M. Vanttinen, NORDITA 96/18 P, hep-ph/9603266; S. Gupta and K. Sridhar, TIFR-TH/96-48, hep-ph/9608433.
- [22] V. Barger, K. Cheung and W.-Y. Keung, *Phys. Rev.* **D 41** (1990) 1541.
- [23] W.-Y. Keung, *Phys. Rev.* **D 23** (1981) 2072; J.H. Kühn and H. Schneider, *Phys. Rev.* **D 24** (1981) 2996.
- [24] O. Adriani et al. (L3 collaboration), *Phys. Lett.* **B 288** (1992) 412; P. Abreu et al. (DELPHI collaboration), *Phys. Lett.* **B 341** (1994) 109; G. Alexander et al. (OPAL collaboration), CERN Preprint CERN-PPE-95-153.
- [25] K. Hagiwara, A.D. Martin and W.J. Stirling, *Phys. Lett.* **B 267** (1991) 527.
- [26] K. Cheung, W.-Y. Keung and T.C. Yuan, *Phys. Rev. Lett.* **76** (1996) 877.
- [27] P. Cho, *Phys. Lett.* **B 368** (1996) 171.
- [28] G. Alexander et al. (OPAL collaboration), CERN Preprint CERN-PPE-95-181.
- [29] E. Braaten and Y. Chen, *Phys. Rev. Lett.* **76** (1996) 730.
- [30] P. Cho and M. Wise, *Phys. Lett.* **B 346** (1995) 129.
- [31] R.M. Godbole, D.P. Roy and K. Sridhar, *Phys. Lett.* **B 373** (1996) 328.
- [32] N. Cacciari and M. Krämer, Presented at the Workshop on Future Physics at HERA, DESY, May 1996, hep-ph/9609500.
- [33] D.P. Roy and K. Sridhar, *Phys. Lett.* **B 341** (1995) 413.
- [34] C.S. Kim, J. Lee and H.S. Song, KEK Preprint KEK-TH-474, hep-ph/9610294.
- [35] V. Barger, S. Fleming and R.J.N. Phillips, *Phys. Lett.* **B 371** (1996) 111.
- [36] K. Sridhar, *Mod. Phys. Lett.* **A 11** (1996) 1555.
- [37] P. Ernström, P. Hoyer and M. Vanttinen, NORDITA-96/3 P, hep-ph/9601333.