

Suppression of $\bar{N}N$ (1535) Coupling in the QCD Sum Rule

D. Jido¹ and M. Oka

Department of Physics, Tokyo Institute of Technology
Meguro, Tokyo 152 Japan

A. Hosaka

Numazu College of Technology
3600 Ooka, Numazu 410 Japan

Abstract

The $\bar{N}N$ and $N\bar{N}$ coupling constants are studied based on the two point functions between the vacuum and a one meson state in the soft meson limit. In terms of a suitably constructed interpolating field for the nucleon and resonances, we have found that the $\bar{N}N$ couplings vanish, while $N\bar{N}$ couplings remain finite. This result explains a relatively suppressed coupling of $\bar{N}N$ (1535) as compared with others. We compare the present result with predictions of low energy effective models.

PACS: 11.55.Hx 12.38.+t 13.75.Gx 14.20.Gk 14.40.Aq

¹email address: jido@th.phys.titech.ac.jp

Baryon resonances provide a good testing ground of effective models for QCD at low energy. Not only the resonance masses but also transition matrix elements are subject to recent research interest. New data for these quantities will soon become available from facilities such as TJNL (former CEBAF), ELSA, MAMI and possibly Spring8 [1]. In particular, transition matrix elements are very sensitive to details of resonance wave functions, and therefore, they will provide strong constraints on the low energy effective models.

Among various baryon resonances, negative parity resonances N^- have particularly interesting properties. For example, $N^-(1535)$ has relatively large branching ratio for the decay $N^-(1535) \rightarrow N \pi$ which is comparative to that of $N^-(1535) \rightarrow N \eta$. Considering the difference in the available phase spaces, this leads to a relatively large coupling constant for $N^- N \pi$ (1535). One may also look at the problem in the following way. Using the experimental decay widths of the resonance, we obtain $g_{N^- N \pi} \approx 0.7$ and $g_{N^- N \eta} \approx 2$. These values are in fact much smaller than those in the $N N \pi$ sector: $g_{N N \pi} \approx 13$ and $g_{N N \eta} \approx 5$. Furthermore, the pion couples weaker than the eta in the $N N \pi$ sector, as opposed to the $N N \eta$ sector. Thus, one may ask why the coupling $g_{N^- N \pi}$ is suppressed so much as compared with other couplings. We will look at the problem of the couplings from this point of view in this paper.

Previously, Jido, Kodama and Oka [3] have studied masses of negative parity octet baryons in the QCD sum rule [4, 5]. A method to extract the information of N^- from a correlation function has been formulated. The resulting masses are generally in good agreement with data. One of the motivations in the previous work was to study chiral properties of the negative parity baryons. The behavior of the masses as functions of the quark condensate shows that the negative parity baryons may be regarded as chiral partners of the ground state baryons. It is then very interesting to study the coupling constants in the same framework.

Our starting point is the observation that an interpolating field can couple to both positive and negative parity baryons [6]. The general interpolating field without derivatives for the nucleon J_N is given by a superposition of two independent terms [7]:

$$J_N(x; t) = \epsilon^{abc} [(u_a(x) C d_b(x)) \gamma_5 u_c(x) + t(u_a(x) C \gamma_5 d_b(x)) u_c(x)]; \quad (1)$$

where a, b and c are color indices, $C = i \gamma_2 \gamma_0$ (in the standard notation) is the charge conjugation matrix, and t is the mixing parameter of the two independent interpolating fields. $J_N(x; t=1)$, which is called the "Ioffe's interpolating field" [8], is commonly used for the study of the ground state nucleon in the QCD sum rule since it is almost optimized for the lowest-lying nucleon [7]. For the study of N^- we found in ref. [3] that the field $J_N(x; t=0.8)$ couples strongly to the negative-parity states.

The correlation function constructed from J_N , therefore, contains information of both parity baryons. In ref. [3], a method to separate the term of either positive or negative parity state from the other exclusively has been developed. Since the correlation function consists of two terms with opposite chirality, their sum and difference have a definite parity.

In order to calculate the meson-baryon coupling constants, we follow the method used by Shiomi and Hatsuda [9]. They studied the $N N \pi$ coupling constant $g_{N N \pi}$ by using the two point function between the vacuum and a one meson state in the soft meson limit ($q \rightarrow 0$). The relevant correlation function is

$$\begin{aligned} \chi(q) &= i \int d^4x e^{i q \cdot x} \langle 0 | T J_N(x) J_N(0) | 0 \rangle \\ &= i \left(\chi_0(q^2) + \chi_1(q^2) \not{q} \right); \end{aligned} \quad (2)$$

where J_N is defined in (1), and m denotes either π or η . The parameter t will be chosen suitably depending on whether J_N should be coupled strongly to positive or negative parity baryons.

Let us first look at the phenomenological side of the correlation function to see how information of the negative parity nucleon can be extracted. We shall see this for the NN coupling ($m = \pi$). The phenomenological NN interaction lagrangian is defined by

$$L_{NN} = g_{NN} \bar{N} \gamma_5 \gamma_i N \pi^i; \quad (3)$$

where N and \bar{N} are the field operators for the positive and negative parity nucleons, π^i is the pion field, and γ_i ($i = 1; 2; 3$) are the Pauli matrices for isospin. From the Lagrangian (3), the NN contribution in the $\langle T \rangle$ is given in the soft pion limit by

$$g_{NN} = \frac{p^2 + m_N^2}{(p^2 - m_N^2)(p^2 - m_\pi^2)} + \frac{\mathcal{P}(m_N + m_\pi)}{(p^2 - m_N^2)(p^2 - m_\pi^2)} \gamma_5; \quad (4)$$

where N and \bar{N} are defined by $\langle 0 | J_N | N \rangle = u_N$ and $\langle 0 | J_N | \bar{N} \rangle = \bar{u}_N \gamma_5$, respectively, with u_N and \bar{u}_N being the Dirac spinor for N and \bar{N} . We note that there appear two terms in (4); one proportional to γ_5 and the other proportional to $\mathcal{P} \gamma_5$. In contrast, the NN contribution has only one term,

$$g_{NN} = \frac{\gamma_5}{p^2 - m_N^2}; \quad (5)$$

as is derived from the NN interaction lagrangian

$$L_{NN} = g_{NN} \bar{N} \gamma_5 N \pi^i \gamma_i; \quad (6)$$

We note that (5) is also obtained by replacing M_N by M_N in (4).

In the soft pion limit, Shiomi and Hatsuda studied the sum rule using the non-vanishing term of (5) and found that the resulting NN coupling constant satisfies the Goldberger-Treiman relation with $g_A = 1$ [9]. Recently, Birse and Krippa also studied the coupling constant g_{NN} at a nonzero pion momentum [10]. In our case for the NN coupling constant, we study the term proportional to $\mathcal{P} \gamma_5$, since that term exclusively contains transitions of $N \rightarrow N$. There is a problem, however, that it contains not only the transitions from negative parity resonances but also those from positive parity resonances, the lowest of which is $N(1440)$. Such a term is, however, proportional to the mass difference, e.g. $M_{N(1440)} - M_N$, unlike the sum as in the second term of (4). Thus the contribution from positive parity resonances is expected to be relatively suppressed as compared with that of negative parity resonances, at least in the low mass region. Moreover, by choosing the mixing parameter $t = 0.8$, the interpolating field (1) is made to couple strongly to negative parity states, and the contribution from positive parity resonances is expected to be further suppressed.

The sum rule for the NN coupling is similarly constructed by replacing the isospin matrices in the NN coupling by the unit matrix. For example, the interaction lagrangian for the NN coupling is written as

$$L_{NN} = g_{NN} \bar{N} \gamma_5 N \pi^0; \quad (7)$$

where $\pi^0 = 1$.

The correlation function is now computed by the operator product expansion (OPE) perturbatively in the deep Euclidean region. The result for the terms of $\mathcal{P} \gamma_5$ takes the

following form

$$\begin{aligned} \chi_{\text{OPE}}(p) = & i \int_0^Z d^4x e^{ip \cdot x} \langle \bar{\psi}(0) J_N(x; s) J_N(0; t) \psi(0) \rangle \\ & \equiv C_4 \ln(p^2) + C_6 \frac{1}{p^2} + C_8 \frac{1}{p^4} + \dots + i \int_0^Z C_3 p^2 \ln(p^2) + \dots; \end{aligned} \quad (8)$$

where we allow to use the different mixing parameters for the interpolating fields such that $J_N(0; t=0)$ couples dominantly to the N state, while $J_N(x; s=1)$ to the \bar{N} state. Note that the terms of χ_{OPE} are of even dimension. The correlation function (8) has been calculated up to dimension 8, ignoring higher order terms in m_q and s . The results are

$$C_4 = m_q \langle \bar{\psi} \psi \rangle_5 \langle \bar{\psi} \psi \rangle_1 \frac{m_q!}{0!} \quad (9)$$

$$C_6 = \frac{s}{4} \langle \bar{\psi} \psi \rangle_5 \langle \bar{\psi} \psi \rangle_1 + \langle \bar{\psi} \psi \rangle_5 \langle \bar{\psi} \psi \rangle_1 \quad (10)$$

$$C_8 = \frac{s}{144} \langle \bar{\psi} \psi \rangle_5 \langle \bar{\psi} \psi \rangle_1 + \langle \bar{\psi} \psi \rangle_5 \langle \bar{\psi} \psi \rangle_1 \quad (11)$$

where $\langle \bar{\psi} \psi \rangle_5 = \langle \bar{\psi} \psi \rangle_5$ and we assume the vacuum saturation for four-quark matrix elements. For the calculation of the matrix element of the operators between the vacuum and one meson state, we use the soft meson theorem :

$$\langle \bar{\psi}(0) \psi(0) \rangle_5 \langle \bar{\psi}(0) \psi(0) \rangle_1 \xrightarrow{Z} \frac{1}{2f_m} \int d^3x \langle \bar{\psi}(0) J_{50}^i(x); O(0) \rangle \psi(0); \quad (12)$$

where $J_5^i(x) = \bar{q}(x) \gamma_5 \gamma^i q(x)$ and f_m is the decay constant of the meson m . We apply this formula to both the pion and the eta, obtaining the following relations:

$$\langle \bar{\psi} \psi \rangle_5 \langle \bar{\psi} \psi \rangle_1 = \frac{m}{f_m} \langle \bar{\psi} \psi \rangle; \quad (13)$$

$$\langle \bar{\psi} \psi \rangle_5 \langle \bar{\psi} \psi \rangle_1 = \frac{m}{f_m} \langle \bar{\psi} \psi \rangle; \quad (14)$$

$$\langle \bar{\psi} \psi \rangle_5 \langle \bar{\psi} \psi \rangle_1 = \frac{m}{f_m} \langle \bar{\psi} \psi \rangle; \quad (15)$$

$$\langle \bar{\psi} \psi \rangle_5 \langle \bar{\psi} \psi \rangle_1 = \frac{m}{f_m} \langle \bar{\psi} \psi \rangle; \quad (16)$$

where $\frac{1}{2} = 1 - \frac{1}{6}$ and $\frac{1}{6} = 1 - \frac{1}{2}$. Note that the sign change in (14) and (16) is the only source for the difference between the pion and eta matrix elements. This originates from the different isospin structure: $\frac{1}{2} = \frac{1}{2}(uu - dd)$, while $\frac{1}{6} = \frac{1}{6}(uu + dd - 2ss)$ by neglecting small mixing angle effects. We note that the ss component in $\frac{1}{6}$ is irrelevant up to dimension 8 if s corrections are ignored, since the interpolating field (1) does not contain strange quarks. From (8) { (16), we find that the correlation function for the $N\bar{N}$ coupling vanishes in the chiral limit $m_q \rightarrow 0$, and therefore $g_{N\bar{N}} = 0$. It is also found that this result remains unchanged when s corrections are included. In contrast, the correlation function for the $\bar{N}N$ coupling does not vanish, and so the coupling constant $g_{N\bar{N}}$ remains finite.

Vanishing of the $\frac{1}{2}$ term in the correlation function for $N\bar{N}$ is, in fact, a general consequence of chiral symmetry. We might have applied the soft meson theorem to the correlation function (2) from the beginning. Using the transformation property

$$[Q_5^a; J_N] = i \gamma_5 \gamma^a J_N; \quad (17)$$

we find

$$\begin{aligned}
\langle \bar{q}^a(p) | &= \lim_{q \rightarrow 0} \int d^4x e^{ipx} \langle 0 | T J_N(x) J_N(0) | q^a \rangle \\
&= \frac{i}{2f} \int d^4x e^{ipx} \langle 0 | T J_N^a(x) J_N(0) | 0 \rangle \\
&= \frac{1}{2f} \int d^4x e^{ipx} f_5^a \langle 0 | T J_N(x) J_N(0) | 0 \rangle
\end{aligned} \tag{18}$$

Due to the Lorentz structure $\langle 0 | T J_N(x) J_N(0) | 0 \rangle = A \delta + B \gamma_5$, the f_5 term disappears in (18).

The result obtained from the commutation relation (18) is more general than that in the OPE, since the momentum p in (18) can be arbitrary, while that in OPE (8) is in the deep Euclidean region. Therefore, Eq. (18) implies that all transitions of $N \rightarrow N$ vanish, if N are the states that couple to J_N satisfying (17). In the real world, there are finite quark mass corrections, but as far as the pion sector is concerned, one may expect such transitions are strongly suppressed. Indeed, such a suppression seems to be realized phenomenologically except for a few cases, i.e. $N(1440)$ and $N(1650)$ [2].

To summarize shortly, our main conclusion here is that the resonance states which couple to the interpolating field J_N of (1) have strongly suppressed couplings for $N \rightarrow N$ in the chiral limit, if J_N satisfies the commutation relation (17). Since J_N couples strongly to $N(1535)$ when $t \rightarrow 0$ [3], the present observation is applied to this state. For the eta case, we do not find a relation similar to (18), because the interpolating field J_N is not a good eigen state of the $U(1)$ axial charge. Thus we find a non-vanishing contribution to the correlation function, which in turn is used in the sum rule analysis to extract a finite coupling constant g_{NN} .

Now, we compare the present results with those of various low energy models. We briefly discuss the nonrelativistic quark model, the large- N_c method and effective chiral lagrangian approach.

In the nonrelativistic quark model, the negative parity nucleon is formed by exciting one of the valence quarks into the $p(l=1)$ orbit [11]. Then there are two independent states for $J^P = 1/2^-$: $|1, 1; S=1/2\rangle^{J=1/2}$ and $|1, 1; S=3/2\rangle^{J=1/2}$, where S is the total intrinsic spin of the three quarks. The physical state for $N(1535)$ is a linear combination of these two states. The coupling constants are the matrix elements of the operators, $O = \sum_{i=1}^3 \vec{\sigma}_i \cdot \vec{r}_i$ for the NN and $O = \sum_{i=1}^3 \vec{\sigma}_i \cdot \vec{r}_i$ for the NN . The relative phase of the two states $|1, 1$ and $|1, 1$ are then determined by the sign of the tensor force. In the Isgur-Karl model, it is brought by the one gluon exchange potential, while in a more sophisticated model, there is a significant contribution from the one pion exchange potential also [12]. In both cases the phase is given such that the interference acts destructively for NN while constructively for NN . This explains the relatively suppressed g_{NN} .

The suppression of g_{NN} is also observed in the large- N_c limit. Assuming that the lowest baryon state develops the hedgehog intrinsic state with $K = J + I = 0$ (here the hedgehog has negative parity as $J^P = 1/2^-$), it is possible to show that the matrix element for the g_{NN} coupling $\langle N | \bar{q} q | N \rangle$ is of higher order in $1/N_c$ as compared with that of $\langle N | \bar{q} \gamma_5 q | N \rangle$. This $1/N_c$ counting is in fact an example of the $I_t = J_t$ rule for large- N_c baryons [13].

One may wonder if such a suppression of g_{NN} could be in some way related to symmetry properties. There have been several attempts to treat positive and negative parity baryons as chiral partners of spontaneous chiral symmetry breaking. DeTar and Kunikida considered the parity doublet nucleons in the linear sigma model of $SU(2)$

SU(2) [14]. In addition to the standard chiral invariant interaction terms, they introduced a chiral invariant mass term between the positive and negative parity baryons. The strength m_0 for the non-standard mass term reduces to the mass of the would-be chiral doublet nucleons when the chiral symmetry restores. In the spontaneously broken phase, the mass splitting is proportional to the non-zero expectation value of the sigma field. In this model, it has been shown that g_{NN} is proportional to m_0 to the leading order in m_0 . In the previous QCD sum rule study [3], the masses of N and N^* seem to get degenerate and decrease as the quark condensate $\langle \bar{q}q \rangle$ is decreased. This implies a small $m_0 \rightarrow 0$, which could be a possible explanation why the coupling constant g_{NN} vanishes in the QCD sum rule.

Earlier, Christos investigated an effective model for mesons and baryons by identifying the interpolating field $B^1 = (q\bar{c} - s\bar{q})q$ with the positive parity nucleon and $B^2 = (q\bar{c} + q)\bar{q}$ with the negative parity nucleon [15]. Using the transformation properties of the B^1 and B^2 field, he wrote down a chiral invariant effective Lagrangian, which leads to the relation $g_{NN} = 0$. His Lagrangian, however, does not contain the m_0 term of the model of DeTar and Kunihiro, which is the reason for the identically vanishing g_{NN} .

In summary, we have studied the NN and NN^* coupling constants using two point correlation functions in the soft meson limit. In the OPE, the correlation function for NN was calculated up to dimension eight and was shown to be the quantity of order $O(m_q)$. Thus the NN coupling constant is strongly suppressed. This suppression turns out to be a general consequence of chiral symmetry when the interpolating field has the suitable transformation property. The present results are consistent with the predictions of various low energy effective approaches. It is interesting that such a suppression of g_{NN} is supported by the chiral effective theory with parity doublet assumption.

Acknowledgments

The authors would like to thank M. Birse for kind correspondences to our original manuscript and comments. They also thank S.H. Lee, T. Hatsuda and T. Kunihiro for comments and discussions.

References

- [1] CEBAF/INT workshop on N physics, Seattle, Sep. (1996).
- [2] Particle Data Group, Phys. Rev. D 54 (1996) 1.
- [3] D. Jido, N. Kodama and M. Oka, Phys. Rev. D 54 (1996) 4532; D. Jido and M. Oka, TIT-preprint "QCD sum rule for $\frac{1}{2}^-$ Baryons", hep-ph/9611322.
- [4] M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Nucl. Phys. B 147 (1979) 385, 448.
- [5] L. J. Reinders, H. Rubinstein and S. Yazaki, Phys. Rep. 127 (1985) 1.
- [6] Y. Chung, H. G. Dosch, M. Krammer and D. Schall, Nucl. Phys. B 197 (1982) 55.
- [7] D. Espriu, P. Pascual and R. Tarrach, Nucl. Phys. B 214 (1983) 285.
- [8] B. L. Ioane, Nucl. Phys. B 188 (1981) 317, (E) B 191 (1981) 591.
- [9] H. Shiomi and T. Hatsuda, Nucl. Phys. A 594 (1995) 294.
- [10] M. Birse and B. Krippa, Phys. Lett. B 373 (1996) 9.
- [11] N. Isgur and G. Karl, Phys. Rev. D 18 (1978) 4187; R. Konik and N. Isgur, Phys. Rev. D 21 (1980) 1868.

- [12] M .Arim a, K .Shim izu and K .Yazaki, Nucl. Phys. A 543 (1992) 613.
- [13] M .P.M attis and M .M ukerjee, Phys. Rev. Lett. 61 (1988) 1344. M .P.M attis, Phys. Rev. D 39 (1989) 994.
- [14] C .DeTar and T .Kunihiro, Phys. Rev. D 39 (1989) 2805.
- [15] G .A .Christos, Z .Phys. C 21 (1983) 83; Phys. Rev. D 35 (1987) 330.