

Small P_{\perp}^2 Elastic Proton-Proton Polarization near 300 GeV

A. D. Krisch¹ and S. M. Troshin²

(1) *Randall Laboratory of Physics, University of Michigan, Ann Arbor, MI 48108-1120*

(2) *Institute for High Energy Physics, Protvino, Moscow Region, 142284 Russia*

Using two hopefully reasonable assumptions plus existing data on the analyzing power in high energy proton-proton elastic scattering, we estimate that the hadronic helicity-flip amplitude at P_{\perp}^2 of 0.003 (GeV/c)^2 is negligible near 300 GeV. This would allow the absolute calibration of a CNI polarimeter.

To successfully accelerate a polarized proton beam to a very high energy, one must somehow measure the beam polarization. Unfortunately, above about 100 GeV, the hadronic spin asymmetries used in most present polarimeters are small and not well known. Fortunately, in 1948 Schwinger [1] proposed that the neutrons from neutron-nucleus elastic scattering would have a large and energy-independent polarization due to the interference between the hadronic (Nuclear) and electromagnetic (Coulomb) scattering amplitudes. This Coulomb-Nuclear Interference (CNI) polarization was subsequently studied for proton-proton elastic scattering by Kopeliovich and Lapidus [2] and later by Buttimore, Gotsman and Leader [3]. The simplest version of the CNI formula has zero parameters.

However, this simplest version does assume that the hadronic single-helicity-flip amplitude is zero. In using the CNI mechanism for a precise high energy polarimeter, one must question the validity of this assumption. This paper attempts to make a reliable estimate of the CNI analysing power A_n near 300 GeV, by trying to set a limit on the hadronic single-flip amplitude. There have been previous attempts to estimate this single-flip amplitude using other techniques [4, 5] and one attempt to measure the CNI polarization at 200 GeV [6].

The total proton-proton elastic scattering amplitude in the helicity state i is

$$f_i = f_i^C + f_i^N; \quad (1)$$

the electromagnetic amplitudes f_i^C can be easily calculated using QED. At small P_{\perp}^2 or $-t$, the dominant hadronic amplitudes f_i^N seem to fit an exponential; its parameters are well determined by measurements of the total cross-section, the slope of the diffraction peak, and the ratio of the real to imaginary amplitudes.

The hadronic part of the proton-proton interaction seems to be helicity conserving; the CNI analysing power would be then due to the interference between a real electromagnetic helicity-flip amplitude and an imaginary hadronic helicity-nonflip amplitude. However, the hadronic interaction may not conserve helicity in small angle scattering. Helicity conservation does not follow from QCD in a region where

chiral symmetry is spontaneously broken. In Regge theory the Pomeron is usually assumed to be helicity conserving; however this assumption has not been tested. Moreover it was shown that unitarity, which is the absorptive correction, generates different phases in the helicity-flip and nonflip Pomeron contributions [7]. The CNI analysing power would certainly change if there were any significant hadronic single-helicity-flip amplitude. Thus, we will try to use the existing experimental data on the analysing power in proton-proton elastic scattering to estimate the hadronic single-helicity-flip amplitude near $E_{lab} = 300$ GeV, where one might use a CNI polarimeter with a polarized proton beam [8, 9].

In standard calculations, the CNI analyzing power arises through the interference between the Coulomb helicity-single-flip amplitude f_5^C and the hadronic (Nuclear) helicity-nonflip amplitudes f_1^N and f_3^N . However, there could also be a non-zero hadronic helicity-single-flip amplitude f_5^N , which can be parameterized by

$$\tau(s, t) \equiv \frac{m}{\sqrt{-t}} \frac{|f_5^N|}{|f_1^N|}. \quad (2)$$

We assume that, at small $-t$, the ratio τ does not depend strongly on $-t$. One can then estimate τ from the high energy A_n data near $-t = 0.2$ (GeV/c)². At small t , one can neglect the three hadronic helicity-flip amplitudes f_2^N , f_4^N and f_5^N in comparison to the helicity-nonflip amplitudes f_1^N and f_3^N . Indeed, at small t , due to conservation of angular momentum, f_4^N is proportional to $-t$ while f_5^N is proportional to $\sqrt{-t}$. Moreover, at small t , the imaginary part of f_2^N is proportional to $\Delta\sigma_t$, which is the difference between the transverse-spin total cross-sections, $\sigma^{tot}(\uparrow\uparrow) - \sigma^{tot}(\uparrow\downarrow)$; this difference is probably small at high energies [10].

We shall also assume that $f_1^N = f_3^N$ at small t , since the existing 12 GeV data on A_U is small, and the existing $\Delta\sigma_t$ data is small and decreases with energy. Thus, for purely hadronic processes at small t ,

$$A_n \frac{d\sigma}{dt} \simeq \frac{-8\pi}{s(s-4m^2)} \text{Im}[f_1^N f_5^{N*}], \quad (3)$$

where

$$\frac{d\sigma}{dt} \simeq \frac{4\pi}{s(s-4m^2)} |f_1^N|^2. \quad (4)$$

Note that, at high energy and small $-t$, the spin-nonflip amplitude f_1^N is primarily imaginary because the elastic hadronic nonflip scattering is primarily the diffractive shadow scattering due to the dominant inelastic processes; thus, $f_1^N \simeq i|f_1^N|$. However, the spin-flip amplitude f_5^N can have both a real and an imaginary part; writing it as $f_5^N = |f_5^N|e^{i\phi_5}$, we have

$$A_n \frac{d\sigma}{dt} \simeq \frac{-8\pi}{s(s-4m^2)} |f_1^N| |f_5^N| \cos \phi_5, \quad (5)$$

where ϕ_5 is the phase. Note that $\phi_5 = \pi/2$ corresponds to a pure imaginary helicity-single-flip amplitude while the phases 0 and π correspond to a pure real f_5^N .

Now by combining Eqs. (4) and (5), we obtain

$$A_n = -2 \frac{|f_5^N|}{|f_1^N|} \cos \phi_5 \quad (6)$$

Combining Eqs. (2) and (6), we then obtain a simple approximation for τ , which is valid at small $-t$.

$$\tau \cos \phi_5 \simeq -\frac{A_n m}{2\sqrt{-t}}. \quad (7)$$

For any hadronic scattering process, angular momentum conservation implies that A_n must go to zero at $t = 0$; for the scattering of two spin-1/2 protons, A_n should be proportional to $\sqrt{-t}$ near $t = 0$. Moreover, the experimental data near $-t = 0.2$ (GeV/c)² imply an s^{-1} energy dependence for A_n ; therefore, we have

$$A_n \simeq \frac{A\sqrt{-t}}{s}. \quad (8)$$

The constant $A \simeq 3.6$ GeV can be obtained from the $A_n = 1.6/s$ fit to the experimental data. Then the estimate for the parameter τ is

$$\tau \cos \phi_5 \simeq -\frac{Am}{2s} \simeq -\frac{1.7}{s}. \quad (9)$$

At 300 GeV we then obtain $\tau \cos \phi_5 \simeq 0.003$, which we assumed can be extrapolated from $-t = 0.2$ to 0.003 (GeV/c)².

Now we can estimate the hadronic contribution to the full single helicity-flip amplitude $f_5 = f_5^C + f_5^N$, where

$$f_5^C \simeq -\frac{\alpha s g F_1(t) F_2(t)}{2m\sqrt{-t}}. \quad (10)$$

where $\alpha = 1/137$ and $g = 1.79$ is the proton's anomalous magnetic moment, while $F_1(t)$ and $F_2(t)$ are the proton's form factors. We use the standard exponential parameterization for the hadronic amplitude at high energies and small t [2, 3, 5]

$$f_5^N = \frac{\tau(\cos \phi_5 + i \sin \phi_5) s\sqrt{-t} (\rho^2 + 1)^{1/2} \sigma_{tot} e^{bt/2}}{8\pi m} \quad (11)$$

At 300 GeV, the ratio of the real to the imaginary forward scattering amplitude squared ρ^2 is experimentally well below 0.003 [11], while σ_{tot} is about 40 mb. Moreover, the $e^{bt/2}$ term and both form factors are all very close to 1 at $-t = 0.003$ (GeV/c)². Thus, at 300 GeV and $-t = 0.003$ (GeV/c)² we obtain

$$\frac{f_5^N}{f_5^C} \simeq 1.9\tau(\cos \phi_5 + i \sin \phi_5) \quad (12)$$

Now we assume that $\tau \sin \phi_5$ is not much larger than $\tau \cos \phi_5$. This is because the helicity-flip amplitudes certainly can not be diffractive shadow scattering, since the final and initial spin states are different. Then we estimate that

$$\tau(\cos \phi_5 + i \sin \phi_5) = 0.003(1 + i), \quad (13)$$

thus we have

$$\frac{|f_5^N|}{|f_5^C|} \simeq 5.7 \cdot 10^{-3} |1 + i| < 10^{-2} \quad (14)$$

Thus, at $-t = 0.003 \text{ (GeV/c)}^2$ and 300 GeV, the hadronic amplitude is at least 100 times smaller than the Coulomb amplitude. Therefore, the two assumptions:

1. τ depends weakly on $-t$
2. ϕ_5^N is not pure imaginary

allow one to neglect f_5^N and use the standard calculation of the CNI analysing power [2, 3]. This result does not contradict the earlier estimate [5], but it does set a more stringent limit on the hadronic helicity-flip amplitude.

This work was supported by research grants from the U.S. Department of Energy and the Russian Ministry of Science. One of us (S.T.) is grateful to the Organizing Committee of SPIN96 Symposium for financial support. We would like to thank N. E. Tyurin, B. B. Blinov, and the other members of the SPIN Collaboration for their help and advice. We are especially grateful to N. H. Buttimore and T. L. Trueman for their comments.

- [1] J. Schwinger, Phys. Rev. **73**, 407 (1948).
- [2] B. Z. Kopeliovich and I. I. Lapidus, Sov. J. Nucl. Phys. **19**, 114 (1974).
- [3] N. H. Buttimore, E. Gotsman and E. Leader, Phys. Rev. **D18**, 694 (1978).
- [4] K. Hinotani *et al.*, Nuov. Cim. **A52**, 363 (1979).
- [5] N. Akchurin, N. H. Buttimore and A. Penzo, Phys. Rev. **D51**, 3944 (1995).
- [6] N. Akchurin *et al.*, Phys. Rev. **D48**, 3026 (1993).
- [7] S. M. Troshin and N. E. Tyurin, Spin Phenomena in Particle Interactions, World Sci. Press (Singapore), 1994.
- [8] SPIN Collaboration A. D. Krisch *et al.*, *Acceleration of Polarized Protons to 120 GeV and 1 TeV*, unpub. Univ. of Michigan Report UM HE 95-09 (July 24, 1995).
- [9] M. Beddo *et al.*, RHIC Spin Proposal, Brookhaven National Laboratory (1992), (unpublished).
- [10] W. de Boer *et al.*, Phys. Rev. Letts. **34B**, 558 (1975).
- [11] J. P. Burq *et al.*, Phys. Lett. **109B**, 124 (1982).