

Preprint K O B E - F H D - 96-01

O c t o b e r, 1996

# A n o m a l o u s Q u a r k C h r o m o m a g n e t i c M o m e n t Induced by Instantons <sup>1</sup>

N . I. K o c h e l e v

B o g o l i u b o v L a b o r a t o r y o f T h e o r e t i c a l P h y s i c s, J o i n t I n s t i t u t e f o r N u c l e a r R e s e a r c h,  
141980 D u b n a, M o s c o w R e g i o n, R u s s i a

## A b s t r a c t

It is shown that the quark-gluon interaction induced by instantons leads to the quark anomalous chromomagnetic moment. In the instanton-liquid model for the QCD vacuum, the chromomagnetic moment is  $\mu_a = \frac{f}{2s}$ , where  $f$  is the packing fraction of instantons in the vacuum.

---

<sup>1</sup> Talk presented at the X III International Seminar on High Energy Physics Problems, 2-7 September, Dubna, Russia

A possible existence of the large quark anomalous chromomagnetic moment (QACHM) is widely under discussion in these days [1]–[5]. In [1], an assumption on the existence of the QACHM was made to explain the observed anomalies in the cross section for reactions with polarized particles. In [2], the QACHM was introduced to improve the description of the spectroscopy of hadrons including heavy quarks. It was shown in [3] that the QACHM allows us to explain the large value of the cross section for  $t\bar{t}$  productions near threshold and  $R_c, R_b$  ratio in the  $Z^0$  peak. A very interesting explanation [5], based on the QACHM, for the excess of the jet cross sections at high  $E_T$  observed by CDF [4] was proposed.

It should be mentioned that, in perturbative QCD, the appearance of the QACHM is related with higher order  $\alpha_s$  corrections, and, therefore, its value needs to be small.

In this paper, the quark anomalous chromomagnetic moment is calculated in non-perturbative QCD by using the liquid instanton model for the QCD vacuum [6].

The effective Lagrangian induced by instantons has the following form [7]<sup>2</sup>

$$L_{\text{eff}} = \sum_q \int d^4x \left( \bar{\psi}_q \left( m_q - \frac{1}{2} \sigma_{\mu\nu} G_{\mu\nu}^a \right) \psi_q \right) \exp \left( \frac{2}{g} \int d^4x \bar{\psi} \psi \right) \int d\Omega \, d_0(\Omega) \, d\Omega^3 + R + L; \quad (1)$$

where  $\rho$  is the instanton size,  $U_a$  are the matrices of the  $SU(2)_c$  subgroup of the  $SU(3)_c$  colour group,  $d_0(\Omega)$  is the density of the instantons,  $d\Omega$  stands for integration over the instanton orientation in colour space,  $d\Omega^3 = 1$ ,  $U$  is the orientation matrix of the instanton,  $\sigma_{\mu\nu}$  is the numerical 't Hooft symbol and  $\sigma_{\mu\nu} = [\gamma_\mu, \gamma_\nu] = 2\epsilon_{\mu\nu\alpha\beta} \sigma_{\alpha\beta}$ .

The effective quark-gluon vertex can be obtained by expanding (1) in the set of the powers of the gluon strength and by integrating over  $d\Omega^3$ . In the first order of  $G^b$ , we found out that

$$L_A = \sum_q \int d^4x \frac{n_{\text{eff}}(\rho)}{m_q g} i \bar{\psi}_q \gamma_5 \psi_q G^a; \quad (2)$$

where  $n_{\text{eff}}(\rho) = d_0(\rho) \sum_q (\bar{\psi}_q \psi_q) = 5$  is the effective instanton density in the QCD vacuum,  $m_q = m_q - \frac{1}{2} \sigma_{\mu\nu} G_{\mu\nu}^a < 0$  for  $q = u, d, s$  is the effective quark mass [6] and  $t^a = \sigma_{\mu\nu} G_{\mu\nu}^a$  are  $SU(3)_c$  matrices.

<sup>2</sup>The Lagrangian (1) has been obtained for the case when the scale of the gluon-field fluctuations was significantly larger than the size of the instanton, i.e., for  $q < 1/\rho$ , where  $q$  is the gluon virtuality.

<sup>3</sup>To integrate over  $d\Omega$ , one can use the method of [8].

It should be mentioned that the Lagrangian (2) was derived as a result of factorization of the Lagrangian (1). The factorization procedure is consisted with the contraction of some quark legs from the instanton vertex through the quark condensate (see [9] and the first reference in [6]).

In the instanton liquid model for the QCD vacuum, the effective density is

$$n_{\text{eff}}(\rho) = n_c(\rho_c); \quad (3)$$

where  $\rho_c$  is the average size of the instanton and  $n_c$  is determined by the value of the gluon condensate  $n_c = \langle 0 | \frac{1}{2} G^a G^a | 0 \rangle = 16$ .

Using (3) we obtain

$$L_A = i \frac{f}{2 \rho_s(\rho_c)} \sum_q \frac{g}{2m_q} \bar{q} \gamma_5 G^a q; \quad (4)$$

where  $f = n_c \rho_c^2$  is the so-called packing fraction of instantons in the vacuum.

The definition for the anomalous chromomagnetic moment is (see [3])

$$L_A = i \gamma_5 \sum_q \frac{g}{2m_q} \bar{q} \gamma_5 G^a q; \quad (5)$$

Therefore, the QACHM is given by the formula

$$a = \frac{f}{2 \rho_s(\rho_c)}; \quad (6)$$

It should be stressed that the final result for the QACHM includes the value of the quark-gluon coupling constant in the denominator. Therefore, it is impossible to get the similar result from perturbative QCD.

For the numerical calculation, we use the NLO approximation for the strong coupling constant

$$\rho_s(\rho) = \frac{2}{1t} \left( 1 + \frac{2 \log t}{1t} \right); \quad (7)$$

where

$$1 = \frac{33 - 2N_f}{6}; \quad 2 = \frac{153 - 19N_f}{12} \quad (8)$$

and

$$t = \log \left( \frac{1}{2\rho^2} + \rho \right); \quad (9)$$

In Equation (9), the parameter  $1 = \frac{2}{c} \tau^2$  provides a smooth interpolation of the value of  $\chi_s(\tau)$  from the perturbative ( $\tau \rightarrow 0$ ) to the nonperturbative region ( $\tau \rightarrow 1$ ) [10].

For  $N_f = 3$ ,  $\mu = 230 \text{ MeV}$ ,  $\bar{c} = 1.6 \text{ GeV}^{-1}$ , by using the standard value for the gluon condensate

$$\langle 0 | \text{tr} G^a G^a | 0 \rangle = 0.012 \text{ GeV}^4; \quad (10)$$

we have the estimate

$$\alpha_a = 0.2 \text{ for } \bar{c} = 1.6 \text{ GeV}^{-1}; \quad (11)$$

Unfortunately, the average size of instantons is not well fixed in the model [6]. For example, if one use the same model to calculate the masses of the pseudoscalar nonet mesons based on QCD sum rules method (see [6]), the more suitable value for average size of the instantons will be  $\bar{c} = 2 \text{ GeV}^{-1}$ . In this case, the value of the QACHM will be larger

$$\alpha_a = 0.4 \text{ for } \bar{c} = 2 \text{ GeV}^{-1}; \quad (12)$$

In any case, the instanton contribution to the QACHM is quite large.

From the above observation, we can say that the quark-gluon interaction through the QCD vacuum leads to the rise of the quark anomalous chromomagnetic moment.

The applications of this result to the hadron spectroscopy and to the high-energy lepton-hadron and hadron-hadron scattering are in progress.

The author is sincerely thankful to M. Anselmino, A.M. Baldin, A.E. Dorokhov, J. Ellis, R.N. Faustov, S.B. Gerasimov, E. Leader, P. Mulders, M. Oka, P. Ratcliffe, and E.V. Shuryak for useful discussions, to T. Morii for warm hospitality at Kobe University, where the part of the work was done, and to the JSPS and the Heisenberg-Landau program for financial support.

## References

- [1] B.A. Arbuzov, E.E. Boss, O.V. Kravchenko, and S.S. Kurennoi, Proc. SPIN '89 Conference, (September, 1986, Protvino, Russia).
- [2] D. Ebert, R.N. Faustov, V.O. Galkin, A.P. Martynenko, and V.A. Saleev, HUP-EP-96/25, hep-ph/9607314.
- [3] D. Atwood, A. Aeppli, and A. Soni, Phys. Rev. Lett. D 69 2754 (1992);  
D. Atwood, A. Kagan, and T.G. Rizzo, Phys. Rev. Lett. D 52 6254 (1995);

- T G Rizzo, Phys. Rev. Lett. D 53 2326 (1996);  
P Haberl, O Nachtmann, and A Wilch, HD-THEP-95925 (1995), hep-ph/9505409.
- [4] CDF Collaboration, F Abe et al., FERMILAB-PUB-96-020-E, hep-ex/9601008.
- [5] D Silverman, SLAC-PUB-7166, hep-ph/9605318.
- [6] E V Shuryak, Phys. Rep. 115 151 (1984);  
D Dyakonov and V Yu Petrov, Nucl. Phys. B 272 457 (1986).
- [7] C G Callan, R Dashen, and D J Gross, Phys. Rev D 17 2717 (1978);  
C G Callan, R Dashen, and D J Gross, Phys. Rev D 19 1826 (1979);  
V A Novikov, M A Shifman, V I Vainshtein, and V I Zakharov, Sov. Phys. Usp. 25 195 (1982).
- [8] M A Shifman, V I Vainshtein, and V I Zakharov, Nucl. Phys. B 163 46 (1980).
- [9] N I Kichelev, Sov. J. Nucl. Phys. 41 291 (1985).
- [10] M A Shifman, V I Vainshtein, and V I Zakharov, Phys. Lett. B 76 471 (1978).