

RESUMMATION OF INFRARED RENORMALONS IN THE PION ELECTROMAGNETIC FORM FACTOR CALCULATIONS

Shahin S. Agaev

High Energy Physics Lab., Baku State University,
Z. Khalilov st.23, 370148 Baku, Azerbaijan

Abstract

The pion electromagnetic form factor $F(Q^2)$ is calculated at the leading order of pQCD using the running coupling constant $(\alpha_s(Q^2(1-x)(1-y)))$ approach. The resummed expression for $F(Q^2)$ is found. It is shown that the effect of the infrared renormalons can be taken into account by scale-setting procedure $\alpha_s(Q^2) \rightarrow \alpha_s(e^{f(Q^2)}Q^2)$ in the leading order expression.

1. It is known that one of the important problems in the perturbative QCD (pQCD) calculations is a choice of the renormalization scheme and scale in a truncated perturbative series [1,2]. Of course, when working to all orders in $s(Q^2)$ the choice of scale and scheme is irrelevant, because a measurable physical quantity is the same for all choices. But this freedom can be a source of difficulties, when one truncates a perturbative series for f . Namely, a bad choice of the coupling constant scale may result in large higher-order corrections in such truncated series. The method proposed by Brodsky, Lepage and Mackenzie (BLM) in Refs.[1,2] for determination of the coupling constant scale seems to be a good tool for the handling of these problems. The BLM scale-setting procedure amounts to absorbing certain vacuum polarization corrections appearing at higher-order into the one-loop QCD running coupling constant. In other words, the essence of this method is to find an average virtuality of a gluon in a loop diagram and to use it as the scale in the running coupling constant. As was emphasized in [1,2], this method may be applied to both inclusive and exclusive processes.

Recently, the BLM method has been generalized by performing the calculations with the running coupling constant $s(k^2)$ at the vertices of Feynman diagrams, where k is the momentum flowing through the virtual gluon line [3,4]. This approach is equivalent to a resummation of an infinite number of terms in the perturbative series for the quantity f .

In our previous works [5,6] we have used the running coupling constant approach for calculation of the pion and kaon electromagnetic form factors at the leading order of pQCD. As a result we have found the perturbative series in $s(Q^2)$ for $Q^2 F_M(Q^2)$. In this letter we shall resum the series obtained in Ref.[5] for the pion form factor using the technique of the Borel transform and show that an effect of such resummation can be taken into account by redefining the scale of $s(Q^2)$ in the leading order expression.

2. Let us start from the factorized QCD expression for the pion electromagnetic form factor

$$F(Q^2) = \int_0^1 \int_0^1 dx dy (y; Q^2) T_H(x; y; Q^2; s(Q^2)) (x; Q^2); \quad (1)$$

where $Q^2 = -q^2$ is the four momentum square of the virtual photon, Q^2 is the renormalization and factorization scale. In (1) terms of order m^2/Q^2 , as well as higher Fock state components of the pion wave function have been neglected. Here the pion wave function $(x; Q^2)$ is non-perturbative quantity and gives the amplitude for finding the quark (antiquark) carrying the longitudinal fractional momentum $x(1-x)$ within the pion. The function $T_H(x; y; Q^2; s(Q^2))$ is the hard scattering amplitude for the subprocess $q\bar{q}^0 + \gamma^* \rightarrow q\bar{q}^0$ and can be found by using pQCD,

$$T_H(x; y; Q^2; s(Q^2)) = s(Q^2) T_H^1(x; y; Q^2) + s(Q^2) T_H^2(x; y; Q^2) + \dots \quad (2)$$

The leading order term $s(\phi^2)T_H^1(x;y;Q^2)$ here is given by [7]

$$s(\phi^2)T_H^1(x;y;Q^2) = \frac{16 C_F s(\phi^2)}{Q^2(1-x)(1-y)}; \quad C_F = 4/3; \quad (3)$$

The next-to-leading order correction $T_H^2(x;y;Q^2)$ was found in [8]. The one-loop QCD running coupling constant $s(\phi^2)$ in (2), (3) is defined as

$$s(\phi^2) = \frac{4}{\ln(\phi^2/\mu_0^2)}; \quad (4)$$

where

$$\mu_0 = 11 - \frac{2}{3}n_f;$$

and n_f being the number of quark flavors.

In Ref.[1] the authors argued that in a meson form factor calculations the scale in (3) has to be chosen equal to the square of a momentum transfer of the exchanged gluon $\phi^2 = Q^2(1-x)(1-y)$. But it is evident that $s(Q^2(1-x)(1-y))$ suffers from singularities associated with the behavior of s in the soft region $x \rightarrow 1; y \rightarrow 1$. In other words, integrations over $x; y$ in (1) extended to $x \rightarrow 1; y \rightarrow 1$ indicate the appearance of non-perturbative effects. Therefore, until now in all pQCD calculations the scale ϕ^2 was taken equal to some fixed quantity (as a rule, $\phi^2 = Q^2 = 4$) and results for $F(Q^2)$ were obtained in the so-called "frozen coupling constant" approximation. In our previous paper [5] for calculation of $F(Q^2)$ in the running coupling constant approach we have expressed $s(Q^2(1-x)(1-y))$ by means of the renormalization group equation in terms of $s(Q^2)$ and calculated obtained integrals over $x; y$ as principal value ones (see, [9,10]). For $Q^2 F(Q^2)$ we have found the following perturbative series in $s(Q^2)$,

$$Q^2 F(Q^2) = (16\pi f)^2 \sum_{n=1}^{\infty} \frac{s(Q^2)^n}{4} S_n; \quad (5)$$

where r and S_n are the model dependent coefficients. In calculations we have used the asymptotic $\phi_{asy}(x)$ and the two-humped Chernyak-Zhitnitsky $\phi_{CZ}(x;\phi^2)$ wave functions [11],

$$\begin{aligned} \phi_{asy}(x) &= \sqrt{\frac{3}{2}} f \sqrt{x(1-x)}; \\ \phi_{CZ}(x;\phi^2) &= \phi_{asy}(x) \left[1 + \frac{2}{3} C_2^{3=2} (2x-1) \frac{s(\phi^2)^{\frac{9}{2}-\gamma_0}}{s(\frac{2}{3})} \right]; \end{aligned} \quad (6)$$

Here $f = 0.093 \text{ GeV}$ is the pion decay constant, $C_2^{3=2}(2x-1)$ is the Gegenbauer polynomial, $\gamma_0 = 50/9$ is the anomalous dimension, $\mu_0 = 0.5 \text{ GeV}$ is the normalization point. The phenomenological pion wave function $\phi_{CZ}(x;\phi^2)$ depends on

the scale Φ^2 . But in this article we ignore this dependence and take it equal to $c_Z(\mathbf{x}; \frac{2}{0})$. The coefficients S_n of the perturbative series (5) and r for $\gamma_{asy}(\mathbf{x})$ and $c_Z(\mathbf{x}; \frac{2}{0})$ are given by the following expressions, respectively

$$S_n = c_n \frac{n-1}{0} = (n-1)! n^{-2} + \frac{n+4}{2^{n+1}} \frac{n-1}{0}; \quad r=1; \quad (7)$$

and

$$S_n = c_n \frac{n-1}{0} = (n-1)! \left[\frac{14}{3} + n^{-1} + \frac{25}{2^{n+1}} + \frac{64}{3^{n+1}} + \frac{1}{4^{n-1}} \right. \\ \left. \frac{25}{2^{n-1}} + \frac{8}{3^n} + \frac{35}{3 \cdot 4^{n-1}} \right] \frac{n-1}{0}; \quad r=5 \quad (8)$$

The S_n in both cases are proportional to $(n-1)!$, as a result the series (5), (7), (8) are ill-defined and must be resummed using some procedures for removing divergences. The procedure for such resummation is well known; perform the Borel transform in (5) with respect to the inverse coupling constant (see, for example, [3,12]) and after that calculate the inverted integral applying some regularization methods.

3. The Borel transform of the series (5) is

$$B[\Phi^2 F](u) = \sum_{n=1}^{\infty} \frac{u^{n-1}}{(n-1)!} c_n; \quad (9)$$

Eq.(9) can be formally inverted to give

$$\int_0^{\infty} \Phi^2 F(\Phi^2) \stackrel{i}{res} = \frac{(16 \pi^2)^{Z-1}}{0} \int_0^{\infty} du \exp \left[-\frac{4}{u} \right] u = \int_0^{\infty} \Phi^2 B[\Phi^2 F](u) du; \quad (10)$$

In these cases, when one finds integral (10), it defines the Borel sum $[\Phi^2 F(\Phi^2)]^{res}$ of the series (5). For the solution of our problem we may follow this general recipe to find $[\Phi^2 F(\Phi^2)]^{res}$: But by means of the identity

$$\int_0^{\infty} \exp[-u(t+w+z)] du = \frac{1}{t+w+z}; \quad (11)$$

Eq.(1) can be written in a more convenient way [5,6]. Indeed, after substituting the leading order term in the renormalization group equation's solution [10]

$$s(\Phi^2(1-x)(1-y))' = \frac{s(\Phi^2)}{1 + \frac{1}{t} \ln[(1-x)(1-y)]}; \quad \frac{1}{t} = \frac{s(\Phi^2)_0}{4}; \quad (12)$$

into (1), and changing of the variables $x; y$ to $z = \ln(1-x); w = \ln(1-y)$, introducing the formula (11) and integrating over $z; w$ for the asymptotic wave

function we find

$$\begin{aligned} Q^2 F(Q^2) = & \frac{(16 f_0)^2}{0} \int_0^1 du \exp^{h_4 u} \exp^{i_4 u} (Q^2)^i \frac{1}{(1-u)^2} + \\ & \frac{1}{(2-u)^2} - \frac{2}{1-u} + \frac{2}{2-u} : \end{aligned} \quad (13)$$

It is evident that

$$B[Q^2 F]^{asy}(u) = \frac{1}{(1-u)^2} + \frac{1}{(2-u)^2} - \frac{2}{1-u} + \frac{2}{2-u} : \quad (14)$$

The Borel transform of the series (5), (8) is

$$\begin{aligned} B[Q^2 F]^{cz}(u) = & \frac{1}{(1-u)^2} + \frac{25}{(2-u)^2} + \frac{64}{(3-u)^2} + \frac{16}{(4-u)^2} \\ & \frac{14}{3(1-u)} - \frac{50}{2-u} + \frac{8}{3-u} + \frac{140}{3(4-u)} : \end{aligned} \quad (15)$$

It is easy by using

$$C_n = \frac{d}{du} \left[\frac{1}{n-1} B[Q^2 F](u) \right]_{u=0} ;$$

to recover the series (5), (7), (8).

The Borel transforms (14), (15) have singularities on the real u axis (at $u = 1; 2; 3; 4$), which are known as infrared renormalons. The infrared renormalons are responsible for a factorial growth of the expansion coefficients in Eqs.(7),(8). This fact confirms our conclusion in [5] concerning an infrared renormalon nature of divergences in the perturbative series (5), (7), (8).

The difficulties with the double and single poles at $u = 1; 2$ in (13) can be avoided by taking the integral in its principal value. Then we obtain

$$Q^2 F(Q^2)_{asy}^{ires} = \frac{(16 f_0)^2}{0} \left[\frac{3}{2} + (\ln 2) \frac{Li(-)}{2} + (\ln + 2) \frac{Li(-^2)}{2} \right] : \quad (16)$$

The similar calculations lead to

$$\begin{aligned} Q^2 F(Q^2)_{cz}^{ires} = & \frac{(80 f_0)^2}{0} \left[\frac{233}{6} + (\ln \frac{14}{3}) \frac{Li(-)}{3} + \right. \\ & 25 (\ln 2) \frac{Li(-^2)}{2} + 8 (8 \ln + 1) \frac{Li(-^3)}{3} + \\ & \left. 4 (4 \ln + \frac{35}{3}) \frac{Li(-^4)}{4} \right] : \end{aligned} \quad (17)$$

In (16),(17) $\text{Li}(\cdot)$ is the logarithmic integral [13]

$$\text{Li}(x) = \int_0^x \frac{dx}{\ln x}; \quad x = Q^2 = s: \quad (18)$$

The principal value prescription used above to regulate the infrared renormalon singularities at $u = 1; 2; 3; 4$ in Eq.(10) (or at $x = 1$ in (18)) produces an ambiguity in the perturbative series for the pion form factor $F_\pi(Q^2)$. Any other approach for defining the integral (10) would differ in the treatment of these singularities. The ambiguity introduced by our treatment, as has been shown in [5], is a higher twist and for $Q^2 F_\pi(Q^2)$ behaves as $\sim Q^2$. It is well known that appearance of infrared renormalon singularities in the perturbative calculations indicate an importance of non-perturbative higher twist effects [14]. A nature of the non-perturbative higher twist corrections is process dependent [9,10,14]. In the case of the pion form factor they, probably, may be associated with contributions from the pion's higher Fock state wave functions $j\bar{q}\bar{q}g$, etc.

4. Some comments are in order concerning these results. First of all, it is instructive to compare sources of the infrared renormalons in our example and in other QCD processes considered in [3,4]. In these articles the running coupling constant approach was used in one-loop order calculations for resummation any number of fermion bubble insertions in the gluon propagator. This technique corresponds to partial resummation of the perturbative series for a quantity under consideration. In fact, in pQCD an infrared-safe and renormalization scheme invariant quantity has the perturbative series in s^{-1} similar to that shown in (5). But now the coefficients S_n of such series have the following expansion in powers of α_s

$$S_n = c_n \alpha_s^{n-1} + r_n \alpha_s^{n-2} + \dots \quad (19)$$

The Borel transform of such perturbative series defined as in (9) (that is, using only c_n from (19)) corresponds to partial resummation of the series (5) and the only source of terms of order α_s^{n-1} in (5) is the running coupling constant [3,4].

We have emphasized in Sect.2 that in our calculations we use only the leading order term $\alpha_s(Q^2) T_H^1$. Therefore, it is not accidental that S_n in (7),(8) have exactly α_s^{n-1} dependence and, hence, the expressions (16),(17) are exact sums of the corresponding series. The source of term α_s^{n-1} in (5) is the running coupling $\alpha_s(Q^2(1-x)(1-y))$, but here it runs due to the integration in (1) over the pion quark's (antiquark's) longitudinal fractional momentum. May in our case terms α_s^{n-1} be also described by fermion vacuum polarization insertion into the exchanged between the pion constituents gluon propagator or not, require further investigations.

5. It is useful to find how much the resummed expressions (16),(17) differ from the "bare" ones. In Figs.1;2; the ratio $R_{WF} = \frac{Q^2 F_\pi(Q^2)_{WF}^{\text{res}}}{Q^2 F_\pi(Q^2)_{WF}^0}$,

where $\mathbb{Q}^2 F(\mathbb{Q}^2) \big|_{w.f.}^0$ is the pion form factor in the frozen coupling approximation, is depicted. As is seen, the contributions of the infrared renormalons to $\mathbb{Q}^2 F(\mathbb{Q}^2) \big|_{w.f.}^{res}$ are considerable for both wave functions. Nevertheless, in both cases these contributions can be hidden into the scale of $s(\mathbb{Q}^2)$ in $s(\mathbb{Q}^2) T_H^1(x; y; \mathbb{Q}^2)$. Indeed, let us look for a function $f(\mathbb{Q}^2) = c_1 + c_2 s(\mathbb{Q}^2)$, such that the leading order term $s(\mathbb{Q}^2) T_H^1(x; y; \mathbb{Q}^2)$ with $\mathbb{Q}^2 = e^{f(\mathbb{Q}^2)} \mathbb{Q}^2$ will give for $\mathbb{Q}^2 F(\mathbb{Q}^2)$ approximately the same results as Eqs.(16),(17). Numerical fitting allows us to get ($n_f = 3$),

$$\begin{aligned} f(\mathbb{Q}^2) &= 5.6 + 10.91 s(\mathbb{Q}^2); \quad \text{for }_{asy}; \\ f(\mathbb{Q}^2) &= 11.76 + 38.09 s(\mathbb{Q}^2); \quad \text{for }_{CZ}; \end{aligned} \quad (20)$$

It is worth noting that a dependence of $f(\mathbb{Q}^2)$ on $s(\mathbb{Q}^2)$ was introduced and argued in the next-to-leading order scale-setting calculations in [2].

The results of numerical calculations of $R_{w.f.}$ with rede ned scale in the leading order expression are shown in Figs.1;2. As is seen from Figs.1;2; $R_{w.f.} \approx 1$ in the whole range of \mathbb{Q}^2 .

The rede nition of the scale $\mathbb{Q}^2 \rightarrow e^{f(\mathbb{Q}^2)} \mathbb{Q}^2$ means, that an average virtuality of the exchanged gluon in the pion form factor diagrams equals to $e^{f(\mathbb{Q}^2)} \mathbb{Q}^2$. The coupling constant $s(e^{f(\mathbb{Q}^2)} \mathbb{Q}^2)$ for the functions under consideration (6) varies from $s = 0.39$ at $\mathbb{Q}^2 = 2 \text{ GeV}^2$ to $s = 0.49$ at $\mathbb{Q}^2 = 20 \text{ GeV}^2$ for $_{CZ}$, and from $s = 0.54$ at $\mathbb{Q}^2 = 2 \text{ GeV}^2$ to $s = 0.35$ at $\mathbb{Q}^2 = 20 \text{ GeV}^2$ for $_{asy}$. These values are typical for such kinds of studies [3,4].

It is important to notice that the functions (20) cannot be obtained directly from perturbative series (5), (7), (8) by means of the BLM next-to-leading order scale-setting prescription [2].

6. In this letter we have found the resummed expression for the pion electromagnetic form factor. For regularization of the infrared renormalons induced divergences in the perturbative series for $F(\mathbb{Q}^2)$ we have used the method of the Borel transform and the principal value prescription. We have demonstrated that the resummed result depends on the pion model wave functions used in calculations. Our investigations of the infrared renormalon effects in the pion electromagnetic form factor calculations allow us to conclude that exclusive processes have two sources of infrared renormalon contributions;

- 1) integration in an exclusive process amplitude over a hadron constituents' longitudinal momenta,
- 2) the running coupling constant in higher-order Feynman diagrams.

The self-consistent treatment of these two sources of the infrared renormalon effects, as well as their study by taking into account a dependence of $(x; \mathbb{Q}^2)$ on \mathbb{Q}^2 , are subject of forthcoming publications.

REFERENCES

1. S.J.Brodsky, P.G.Lepage and P.B.Mackenzie, PhysRevD 28 (1983) 228.
2. S.J.Brodsky and H.Lu, PhysRevD 51 (1995) 3652.
3. M.Neubert, PhysRevD 51 (1995) 5924.
4. P.Ball, M.Beneke and V.M.Braun, PhysRevD 52 (1995) 3929;
P.Ball, M.Beneke and V.M.Braun, NuclPhysB 452 (1995) 563;
C.N.Lovett-Turner and C.J.Maxwell, NuclPhysB 452 (1995) 118.
5. S.S.Agaev, PhysLettB 360 (1995) 117; E:PhysLettB 369 (1996) 379.
6. S.S.Agaev, ModPhysLettA 10 (1995) 2009.
7. G.P.Lepage and S.J.Brodsky, PhysLettB 90 (1979) 355; PhysRevD 22 (1980) 2157;
A.Duncan and A.H.Mueller, PhysLettB 90 (1980) 159; PhysRevD 21 (1980) 1636;
A.V.Efremov and A.V.Radyushkin, PhysLettB 94 (1980) 245.
8. R.D.Field, R.Gupta, S.Otto and L.Chang, NuclPhysB 186 (1981) 429.
9. A.H.Mueller, NuclPhysB 250 (1985) 327.
10. H.Contopanagos and G.Sternan, NuclPhysB 419 (1994) 77.
11. V.L.Chernyak and A.R.Zhitnitsky, PhysRep 112 (1984) 173.
12. V.I.Zakharov, NuclPhysB 385 (1992) 452.
13. A.Erdelyi, Higher transcendental functions, v.2, McGraw-Hill Book Company, New York, 1953.
14. I.I.Bigi, M.A.Shifman, N.G.Uraltsev and A.I.Vainshtein, PhysRevD 50 (1994) 2234;
A.V.Manochar and M.B.Wise, PhysLettB 344 (1994) 407;
B.R.Webber, PhysLettB 339 (1994) 148;
G.P.Korchemsky and G.Sternan, NuclPhysB 437 (1995) 415;
R.Akhoury and V.I.Zakharov, Saclay preprint SACLAY-SPHT-95/043, April 1995.

F U G U R E C A P T I O N S

Fig.1. The dependence of the ratio R_{asy} on Q^2 . Here the curves 1 and 2 correspond to R_{asy} with $[Q^2 F(Q^2)]_{asy}^0$ in the frozen coupling constant approximation and after scale-setting procedure $Q^2 \rightarrow e^{f(Q^2)} Q^2$, respectively. In calculations the QCD scale parameter Λ has been taken equal to 0.1 GeV:

Fig.2. The same as in Fig.1, but for R_{cz} :

Fig.1




Fig.2

