

Small Angle Bhabha Scattering at LEP 1. Analytical Results for Wide Angular Acceptance.

A.B. Arbuzov^a, G.I. Gach^b, V.Yu. Gontchar^b, E.A. Kuraev^a,
N.P. Merenkov^b and L. Trentadue^c

^aBogoliubov Laboratory of Theoretical Physics, JINR, Dubna, 141980, Russia

^bInstitute of Physics and Technology, Kharkov, 310108, Ukraine

^cDipartimento di Fisica, Università di Parma and INFN, Gruppo Collegato di
Parma, 43100 Parma, Italy

Abstract

Analytical formulae for small angle Bhabha scattering cross section at LEP 1 are given for the case of wide angular acceptance. Inclusive and calorimeter event selections are considered. Numerical results are presented.

PACS 12.20.+m Quantum electrodynamics, 12.20.Ds Specific calculations

Key words: Bhabha scattering, high energy, small angles, event selection

Introduction

The small angle Bhabha scattering (SABS) process is used to measure luminosity of electron-positron colliders. At LEP 1 an experimental accuracy on the luminosity better than one per mille has been reached [1]. To estimate the total accuracy a systematic theoretical error must be added. That is why in recent years a considerable attention has been devoted to theoretical investigation of SABS cross section [2-11].

The theoretical calculation of SABS cross section at LEP 1 has to cope with two problems. The first one is the description of experimental restrictions in terms of final particles phase space used for event selection. The second one concerns the computation of matrix element squared with the required accuracy. There are two approaches for the theoretical study of SABS at LEP 1: the one basing on Monte Carlo (MC) programs [2-6] and the other using semi-analytical calculations [7-11].

The advantage of MC method is the possibility to model different types of detectors and event selections [2]. But at this approach some problems with exact matrix element squared exist. Contrary, the advantage of the analytical approach is the possibility to use exact matrix element squared and its defect is a low mobility relative the change of an experimental conditions for event selection. Nevertheless, the analytical calculations are of great importance because they allow to check numerous MC calculations for different types of ideal detectors.

In this letter we list some analytical results for SABS cross{section at LEP1 suitable for inclusive (labeled in [2] as BARE1) and calorimeter (CALO1 and CALO2) event selections in the case of asymmetrical wide{narrow circular detectors. We give analytical formulae for the full first order correction to the cross{section as well as for leading second and third order ones. Our numerical estimations include also next{to{leading second order contribution in the case BARE1.

1 First order corrections

Let us consider at first BARE1 event selection. We introduce the dimensionless quantity $\sigma_1 = Q_1^2 \sigma_{obs} = (4\pi^2) \sigma_1$; where $Q_1^2 = s_1^2$ (s_1 is the beam energy and θ_1 is the minimal scattering angle for wide circular detector) and σ_{obs} is an experimentally observable (by means of ideal detectors) cross{section. Then the first order QED correction can be written as follows:

$$\sigma_1 = \frac{1}{2} \int_{x_c}^1 \frac{dz}{z^2} \left(\frac{z^2}{x_c} \right) \left(\frac{1}{2} (1-x) + (L-1)P^{(1)}(x) \left(\frac{1}{42} + \frac{(x)}{42} \right) + \frac{1+x^2}{1-x} K \right) dx + \frac{1}{2} \int_{x_c}^1 \frac{dz}{z^2} \left(\frac{z^2}{x_c} \right) \left(\frac{1}{2} (1-x) + (L-1)P^{(1)}(x) \left(1 + \frac{(x)}{3} \right) + \frac{1+x^2}{1-x} K \right) dx; \quad (1)$$

where

$$\begin{aligned} \bar{K}(x; z; \frac{1}{4}; \frac{1}{2}) &= \frac{(1-x)^2}{1+x^2} \left(\frac{1}{42} + \frac{(x)}{42} \right) + \frac{1}{42} \bar{E}_1 + \frac{(x)}{42} \bar{E}_2 + \left(\frac{(x)}{4} - \frac{(x)}{2} \right) \bar{E}_3 \\ &+ \left(\frac{(x)}{4} - \frac{(x)}{2} \right) \bar{E}_4; \\ K(x; z; \frac{1}{3}; 1) &= \frac{(1-x)^2}{1+x^2} \left(1 + \frac{(x)}{3} \right) + L_1 + \frac{(x)}{3} L_2 + \frac{(x)}{3} L_3; \end{aligned} \quad (2)$$

and $L = \ln(zQ_1^2/m_e^2)$ is the large logarithm; parameter x_c puts the restriction on the energies of the final electron and positron: $s_1^2 \geq x_c^2$. In Eqs.(1) and (2) we used the following notation for \bar{E}_i functions and E_i :

$$\bar{E}_i^{(x)} = (x^2 \frac{1}{x_c} - z); \quad E_i = (x^2 \frac{1}{x_c} - z); \quad \bar{E}_i^{(x)} = 1 - \frac{(x)}{x_c};$$

$$\begin{aligned}
\bar{z}_i &= 1 - z_i; \quad z_{42} = z_4 - z_2; \quad \bar{z}_{42} = \bar{z}_4 - \bar{z}_2; \\
\mathbb{E}_1 &= \ln \frac{(z - \frac{z^2}{2})(\frac{z^2}{4} - z)x^2}{(x - \frac{x^2}{4} - z)(x - \frac{x^2}{2} - z)}; \quad \mathbb{E}_2 = \ln \frac{(z - x^2 - \frac{z^2}{2})(x^2 - \frac{z^2}{4} - z)}{x^2(x - \frac{x^2}{4} - z)(x - \frac{x^2}{2} - z)}; \\
\mathbb{E}_3 &= \ln \frac{(z - x^2 - \frac{z^2}{2})(x - \frac{x^2}{4} - z)}{(x^2 - \frac{z^2}{4} - z)(x - \frac{x^2}{2} - z)}; \quad \mathbb{E}_4 = \ln \frac{(z - \frac{z^2}{2})(x - \frac{x^2}{4} - z)}{(\frac{z^2}{4} - z)(x - \frac{x^2}{2} - z)};
\end{aligned}$$

The quantities L_i can be obtained from \mathbb{E}_i by the substitution $z_4 \rightarrow z_3; z_2 \rightarrow 1$. Here we use the same notations as in [10]. Note only that for wide (narrow) angular acceptance

$$z_3 > z_4 > z_2 > z_1 = 1; \quad z_i = \frac{1}{1}; \quad (3)$$

where z_i are the limiting angles of the circular detectors (see Section 3). Function $P^{(1)}(x)$ defines the iterative form of the non-singlet electron structure function (see for example [6]). The first (second) line in the r.h.s. of Eq.(1) is the contribution due to photon emission by positron (electron). The terms accompanied with x -dependent \mathcal{F} -functions under the integral sign correspond to the initial-state correction while the rest belongs to the final-state one.

The CALO1 cluster is the cone with angular radius $\theta = 0.01$ around the direction of the final electron (or positron) momentum. If photon belongs to the cluster the whole cluster energy is measured, and electron may have any possible energy. Therefore, the limits of x -integration for σ_{obs} extend from 0 to 1 here. If photon escapes the cluster the event looks the same as for BARE1. The above restrictions on x -integration limits can be written symbolically as follows:

$$\int_{x_c}^{z_1} dx + \int_0^{z_c} (|j|) dx = \int_0^{z_1} dx + \int_0^{z_c} (|j|) dx; \quad r = \frac{k}{1} \frac{q_1^2}{n_1}; \quad (4)$$

where $k(q_1^2)$ and $1/n_1$ are the transverse momentum and the energy of the hard photon (electron).

As we saw on the example of BARE1 event selection it is necessary to distinguish the contributions into σ_1 due to electron and positron radiation:

$$\sigma_1 = \sigma_e + \sigma_p; \quad (5)$$

According to (4) we have

$$= \sigma_i + \sigma_f + \sigma_i^c + \sigma_f^c; \quad = \sigma_e + \sigma_f + \sigma_e^c + \sigma_f^c; \quad (6)$$

where index $i(f)$ labels the initial (final) state and index c points on a cluster-form dependence.

Quantities γ_i and ϵ_i coincide with the corresponding initial state correction for BARE1 (see comments on Eq.(1)). For γ_f and ϵ_f we can use the form of differential cross section suitable for inclusive event selection with extended x-integration limits:

$$\begin{aligned}\gamma_f &= \frac{1}{2} \int_{\frac{1}{2}}^{\frac{1}{2}} \frac{dz}{z^2} \int_0^1 \left(1 - x + \frac{1+x^2}{1-x} L_1\right) dx; \\ \epsilon_f &= \frac{1}{2} \int_{\frac{1}{2}}^{\frac{1}{2}} \frac{dz}{z^2} \int_0^1 \left(1 - x + \frac{1+x^2}{1-x} E_1\right) dx + \frac{1+x^2}{1-x} \left(\frac{1}{4} - \frac{1}{2}\right) E_4 dx : \quad (7)\end{aligned}$$

In order to find the additional contributions into γ_i which depend on the cluster form it is enough to use the simplified differential cross section of single photon radiation, neglecting electron mass, and taking into account the restrictions $|j| < \dots$ (for the initial state) and $|j| > \dots$ (for the final state). The contribution due to photon emission by the initial state electron can be written as follows:

$$\gamma_i^c = \frac{1}{2} \int_0^{z_c} \frac{1+x^2}{1-x} dx \int_{\frac{1}{2}}^{\frac{1}{2}} \frac{dz}{z^2} F(z_1; z; x;) = \dots; \quad (8)$$

where $\int_{\frac{1}{2}}^{\frac{1}{2}}$ defines the integration limits over z (in the square brackets) and over z_1 (in the parenthesis):

$$\begin{aligned} &= [a^2; a_0^2](x^2 z_+; x^2) + [b^2; a^2](x^2 z_+; x^2 z_-) + [b_0^2; b^2](x^2 \frac{2}{3}; x^2 z_-); \quad a_0 = \frac{1}{2}; \quad b_0 = \frac{1}{4}; \\ a &= \max(\frac{1}{2}; 1 - (1-x)); \quad b = \min(\frac{1}{4}; \frac{1}{3} - (1-x)); \quad z = \left(\frac{1}{2} - (1-x)^2\right); \end{aligned}$$

and function F is defined below:

$$\begin{aligned} F(z_1; z; x;) &= \frac{2}{z_1 - xz} + \frac{1}{z - z_1} \arctan \frac{(z - z_1)Q}{\left(\frac{1}{z} - \frac{1}{z_1}\right)^2}; \\ Q &= \frac{\sqrt{2x^2(1-x)^2} \left(\frac{1}{z_1} - x\frac{1}{z}\right)^2}{\left(\frac{1}{z_1} + x\frac{1}{z}\right)^2 - 2x^2(1-x)^2} : \quad (9)\end{aligned}$$

The cluster dependent contribution due to photon emission by the final state electron reads

$$\gamma_f^c = \frac{1}{2} \int_0^{z_c} \frac{1+x^2}{1-x} dx \int_{\frac{1}{2}}^{\frac{1}{2}} \frac{dz}{z^2} F(z_1; z; x;) + \frac{1}{a_0^2} \frac{dz}{z^2} \ln \frac{x^2 \frac{2}{3} - z}{z} + 1$$

$$+ \frac{z_0^2}{z^2} \ln \frac{z_1^2 x}{z} + 1 \quad ; \quad 1 = \ln \frac{(2 \frac{p}{z} - \frac{p}{z_1} (1-x))}{z - 2x \frac{p}{z} - \frac{p}{z_1} (1-x)} ;$$

$$F = -\frac{2}{a^2} \frac{1}{z_1 x z} - \frac{1}{z_1 x^2 z} \arctan \frac{(\frac{p}{z_1} - x \frac{p}{z})^2}{(z_1 - x^2 z) Q} :$$

To obtain e_i^c it is enough to substitute in the expression for the parameters $a; b; a_0$ and b_0 by $a; b; a_0$ and b_0 respectively,

$$a = z_2 + (1-x); \quad a_0 = \max(z_2, (1-x));$$

$$b = z_4 - (1-x); \quad b_0 = \min(z_4 + (1-x), z_3):$$

Finally, the cluster-dependent contribution due to photon emission by the final state positron can be written as follows:

$$e_f^c = \frac{z^c}{2} \int_0^1 \frac{1+x^2}{1-x} dx \int_1^z \frac{dz}{z^2} [(a_0^2 - z) - (z - b_0^2)] E_4$$

$$+ \int_f^c (a; b; a_0; b_0) ! (a; b; a_0; b_0; z_3; z_1) ! (z_4; z_2) : \quad (10)$$

The CALO 2 event selection differs from the CALO 1 one by the form of the cluster (see [2]). Only cluster-dependent contributions into γ_1 will be changed here. Analytical formulae are very cumbersome, and we give the result for symmetrical wide case only ($\theta = 0$) but our numerical calculations include narrow angular acceptance too.

$$e_i^c = \frac{z^c}{2} \int_0^1 \frac{1+x^2}{1-x} dx \int_1^z \frac{dz}{z^2} dz_1 \left[\frac{2}{z_1 x z} + \frac{1}{z - z_1} \right] [\gamma_1 \gamma_1 + \gamma_2 \gamma_2 + \gamma_3 \gamma_3]; \quad (11)$$

$$\gamma_1 = \arctan(Q_i^{(-)}) - \arctan(\gamma); \quad \gamma_2 = \arctan(\gamma^{(-)}); \quad \gamma_3 = \arctan \frac{1}{Q_i^{(+)}} ;$$

$$= r_i \cot \frac{\gamma}{2} ; \quad r_i = \frac{(\frac{p}{z} - \frac{p}{z_1})^2}{z - z_1} ;$$

$$Q_i^{(-)} = r_i \frac{\sqrt{x^2 (\frac{p}{z} + \frac{p}{z_1})^2 - (1-x)^2 (\frac{p}{z_1} - x)^2}}{(1-x)^2 (\frac{p}{z_1} - x)^2 - x^2 (\frac{p}{z} - \frac{p}{z_1})^2} ;$$

$$\gamma_1 = [z_3^{(-)}; 1](x^2 J_+^2; x^2 z_+) + [(z_3 - (1-x))^2; z_3^{(-)}](x^2 \frac{z}{z_3}; x^2 z_+);$$

$$\gamma_2 = [z_1^{(+)}; 1](x^2 z_+; x^2) + [(z_3 - (1-x))^2; z_1^{(+)}](x^2 z_+; x^2 J^2)$$

$$+ [z_3^2; (z_3 - (1-x))^2](x^2 \frac{z}{z_3}; x^2 J^2);$$

$$\gamma_3 = [z_1^{(+)}; (1 + (1-x))^2](x^2 J_+^2; x^2) + [z_3^2; (1 + (1-x))^2](x^2 J^2; x^2 z_-):$$

We present also the corresponding formula for photon emission by the final state electron:

$$\begin{aligned}
\sigma_f^{(c)} = & \frac{1}{2} \int_0^{z_c} \frac{1+x^2}{1-x} dx \int_{z_1}^z \frac{dz}{z^2} dz_1 \frac{2}{z_1} \frac{1}{xz} \frac{1}{z_1 x^2 z} [{}_1F_1 + {}_2F_2 + {}_3F_3] \\
& + \int_0^{z_1^{(+)}} \frac{dz}{z^2} \ln \frac{(x^2 z_3 - z)(J_+^2 - z)}{(x^2 z_3 - z)(x J_+^2 - z)} + \int_{z_1^{(+)}}^{z_3^{(+)}} \frac{dz}{z^2} \ln \frac{x}{1-z} + 1; \quad (12) \\
F_1 = & \arctan \left(\frac{1}{Q_f^{(+)}} \right); \quad F_2 = \arctan \left(\frac{1}{Q_f^{(+)}} \right); \quad F_3 = \arctan \left(Q_f^{(+)} \right); \quad \theta = \arctan \left(\frac{r_f}{z} \right); \\
r_f = & \frac{z_1}{x^2 z}; \quad 1 = 1 - \left(\frac{z_1}{x^2 z} \right)^2; \quad Q_f^{(+)} = \frac{r_f}{r_i} Q_i^{(+)}; \quad \sin(\theta) = \frac{r_f}{z} \sin(\theta_i); \\
\sigma_2 = & [z_1^{(+)}; 1](x^2 J_+^2; x^2) + [z_3^{(+)}; z_1^{(+)}](x^2 J_+^2; x^2 J_+^2) + [z_3^{(+)}; z_3^{(+)}](x^2 z_3^2; x^2 J_+^2);
\end{aligned}$$

Angle θ and quantity r_f , which enter into Eq.(12), define the form and the size of CALO 2 cluster. Namely [2]

$$\theta = \frac{3}{32}; \quad r_f = \frac{0}{1}; \quad \theta_0 = \frac{0.051}{16};$$

Finally, functions $J_i^{(+)}$ and $z_i^{(+)}$ are defined as follows:

$$\begin{aligned}
J_i^{(+)} = & \frac{1}{z} \sqrt{\frac{1-x^2}{1-x^2(1-x)^2 - 2 \sin^2 \frac{\theta}{2}}} (1-x) (1-2x \sin^2 \frac{\theta}{2}); \\
= & 1 - 4x(1-x) \sin^2 \frac{\theta}{2}; \quad z_i^{(+)} = (z_i - (1-x))^2 - 4x(1-x) \sin^2 \frac{\theta}{2};
\end{aligned}$$

2 Second and third order corrections

In this Section we give the analytical form of the leading second and third order corrections to SABS cross section suitable for both, inclusive and calorimeter, event selections. The contribution connected with pair production in the singlet channel is negligible for LEP1 conditions, while the one in the non-singlet channel can be taken into account by means of effective QED coupling [6]. Therefore, we will consider here the photonic corrections only.

The second order correction can be presented in the form

$$\sigma_2 = \sigma_2^{(1)} + \sigma_2^{(2)} + \sigma_2^{(3)}; \quad (13)$$

The first term in r.h.s. of Eq.(13) is responsible for two-photon (real and virtual) emission by the electron. The second one describes two-photon emission by the positron. And the third one considers the situation when both the electron and the positron radiate.

The leading contributions in the case of inclusive event selection read

$$\begin{aligned}
 L &= \frac{1}{4} \int_0^1 \frac{dz}{z^2} L^2 \bigg|_{x_c}^{x_1} dx \frac{1}{2} (1 + \frac{x}{x_1}) P^{(2)}(x) + \int_0^1 \frac{dt}{t} P^{(1)}(t) P^{(1)}(\frac{x}{t}) \frac{x}{t} \frac{1}{x_1} ; \\
 L &= \frac{1}{4} \int_0^1 \frac{dz}{z^2} L^2 \bigg|_{x_c}^{x_1} dx \frac{1}{2} (1 + \frac{x}{x_2}) P^{(2)}(x) + \int_0^1 \frac{dt}{t} P^{(1)}(t) P^{(1)}(\frac{x}{t}) \frac{x}{t} \frac{1}{x_2} ; \\
 L &= \frac{1}{4} \int_0^1 \frac{dz}{z^2} L^2 \bigg|_{x_c}^{x_1} dx_1 \int_{\frac{x_c}{x_1}}^{x_2} dx_2 P^{(1)}(x_1) P^{(1)}(x_2) (1 + \frac{x_1}{x_1}) (1 + \frac{x_2}{x_2}) : \quad (14)
 \end{aligned}$$

where

$$P^{(2)}(x) = \int_0^1 \frac{dt}{t} P^{(1)}(t) P^{(1)}(\frac{x}{t}) \frac{x}{t} ; \quad \int_0^1 P^{(2)}(x) dx = 0 :$$

The contribution of initial (final) state radiation for L and L are accompanied with x -dependent (x -independent) functions and $x_1; x_2$ -dependent ($x_1; x_2$ -independent) ones for L . The terms with additional integration over t -variable describe the simultaneous radiation of one photon from the initial state and the other from final state (initial/final state radiation).

In the case of calorimeter event selection we have to take in the r.h.s. of Eqs.(14) the terms corresponding to initial state radiation only. The elimination of final state one exhibits itself by means the last equality. As concerns the contribution due to initial/final state radiation it may be understood as follows.

In fact t -variable in Eqs.(14) is the energy fraction carried out by both the final state radiated photon and the final state electron (or positron). Just this value defines the cluster energy for calorimeter event selection. The x -variable which is the energy fraction of the final electron by definition can change here from 0 up to 1. That is why initial/final state radiation of the electron for calorimeter event selection will be proportional to

$$\int_{x_c}^1 \int_0^1 \frac{dx}{t} P^{(1)}(t) P^{(1)}(\frac{x}{t}) \frac{x}{t} \frac{1}{x_1} = \int_{x_c}^1 \int_0^1 P^{(1)}(t) \frac{1}{t} P^{(1)}(\frac{x}{t}) \frac{x}{t} \frac{1}{x_1} dy dy = 0 : \quad (15)$$

The same is valid of course for the corresponding part of positron radiation.

Thus, we see that it is enough to have only single nal{state radiated photon to elim inate the leading contribution due to initial{ nal{state radiation. This conclusion reflects the essence of a reduced Lee{Nauenberg theorem [12] and is valid for all higher order corrections.

The leading third order correction reads

$$\begin{aligned}
 L_3 &= \frac{1}{2} \int_0^Z \frac{dz}{z^2} L_3^{Z^1} Z_1 + \int_{x_c=x}^{Z^1} Z_2 dx_1 \quad ; \quad (16) \\
 Z_1 &= \frac{1}{6} (2 \langle x \rangle_{42} + \langle x \rangle_{42} \langle x \rangle_{31} + \langle x \rangle_{31} \langle x \rangle_{42}) P^{(3)}(x) \\
 &+ \frac{1}{2} \int_x^{Z^1} \frac{dt}{t} (\langle t \rangle_{42} \langle x \rangle_{31} + \langle t \rangle_{31} \langle x \rangle_{42}) P^{(1)}(t) P^{(2)}\left(\frac{x}{t}\right) + P^{(2)}(t) P^{(1)}\left(\frac{x}{t}\right) \quad ; \\
 Z_2 &= \frac{1}{2} [(\langle x \rangle_{31} + \langle x \rangle_{31} \langle x \rangle_{42}) (\langle x \rangle_{42} + \langle x \rangle_{42} \langle x \rangle_{31}) + (\langle x \rangle_{42} + \langle x \rangle_{42} \langle x \rangle_{31}) (\langle x \rangle_{31} + \langle x \rangle_{31} \langle x \rangle_{42})] P^{(1)}(x) P^{(2)}(x_1) \\
 &+ P^{(1)}(x) \int_{x_1}^{Z^1} \frac{dt}{t} [\langle t \rangle_{31} (\langle x \rangle_{42} + \langle x \rangle_{42} \langle x \rangle_{31}) + \langle t \rangle_{42} (\langle x \rangle_{31} + \langle x \rangle_{31} \langle x \rangle_{42})] P^{(1)}(t) P^{(1)}\left(\frac{x_1}{t}\right) \quad ; \\
 P^{(3)} &= \int_x^{Z^1} \frac{dt}{t} P^{(1)}(t) P^{(2)}\left(\frac{x}{t}\right) \quad :
 \end{aligned}$$

For calorimeter event selection it is needed to take

$$Z_1 = \frac{1}{6} (\langle x \rangle_{42} \langle x \rangle_{31} + \langle x \rangle_{31} \langle x \rangle_{42}) P^{(3)}(x); \quad Z_2 = \frac{1}{2} (\langle x \rangle_{31} \langle x \rangle_{42} + \langle x \rangle_{42} \langle x \rangle_{31}) P^{(1)}(x) P^{(2)}(x_1) :$$

Note that in this case the leading second and third order contributions have a universal character and do not depend on cluster form. Thus, they are suitable for both, CALO 1 and CALO 2.

Quantity Z_1 describes the situation when only one fermion (electron or positron) radiate (one{side emission), while Z_2 is responsible for simultaneous radiation of the electron and the positron (opposite{side emission).

The formulae for leading second and third order contributions are written in the form with different {functions under integral sign. One can eliminate these {functions using such kind relations as, for example

$$\int_{x_c}^Z \int_{x_c}^{\langle x \rangle_4} \int_{x_c}^{\langle x \rangle_3} dz dx dx_1 = \int_{x_c}^{\frac{Z}{2}} \int_{x_c}^{\frac{P_{\bar{Z}}}{Z}} \int_{x_c}^{\frac{P_{\bar{Z}}}{Z}} dx dx_1 : \quad (17)$$

It is needed to keep in mind only that every integral has to be equal to zero if the lower limit

of z -integration becomes more than the upper one. The last statement is valid for the first order correction too.

3 Numerical results

In our calculations we restrict ourselves with pure QED corrections supposing Z (exchange, vacuum polarization and up/down interference) are switched off. As shown in papers by W. Beenakker and B. Pietrzyk [7], a sufficiently accurate luminosity determination requires the full Born plus complete order corrected cross section. Nevertheless our numerical results can be used for comparisons and cross checks with numerous Monte Carlo and semi analytical computations [2].

We performed the numerical calculations for the beam energy $\sqrt{s} = 46.15$ GeV and the following sets of limiting angles of circular detectors:

- i) BARE1, CALO1: $\theta_1 = 0.024$; $\theta_3 = 0.058$; $\theta_2 = \theta_1 + h$; $\theta_4 = \theta_3 - h$; $h = \frac{0.017}{8}$;
- ii) CALO2: $\theta_1 = 0.024 + h$; $\theta_3 = 0.058 - h$; $\theta_2 = \theta_1 + h$; $\theta_4 = \theta_3 - 3h$;

The Born cross section

$$\sigma_B = \frac{4\pi\alpha^2 Z}{Q_1^2} \frac{dz}{z^2} \left(1 - \frac{z^2}{2} \right) \quad (18)$$

(limits of integration are $(\frac{2}{4}; \frac{2}{2})$ for NN and WW angular acceptances and $(\frac{2}{3}; 1)$ for WW one) equals to

| | |
|------------|-------------------------------|
| 175.922 nb | { BARE1, CALO1 WW ; |
| 139.971 nb | { BARE1, CALO1 NN; CALO2 WW ; |
| 103.299 nb | { CALO2 NN : |

Formula (18) takes into account the contributions of scattering diagram and interference of scattering and annihilation ones. The contribution of pure annihilation diagram is proportional to $\frac{4}{1}$. It is negligible even at the Born level. When calculating the QED corrections to cross section (18) we ignore systematically the terms proportional $\frac{2}{1}$. Terms of this kind have double logarithmic asymptotic behavior [13] and are equal parametrically to $(\frac{2}{1}) \ln^2(\frac{2}{1}) = (\frac{2}{1})$, what is about 0.1 per mille as compared with unit for LEP1 conditions.

The results of our calculations of QED correction with the switched off vacuum polarization are shown in the Table 1. The centre of mass energy is $\sqrt{s} = 92.3$ GeV. The second order

Table 1

The results of our analytical calculations for the SABS cross{section

| x_c | BARE1 | | | CALO1 | | | CALO2 | | |
|---|---------|---------|---------|---------|---------|---------|---------|--------|--------|
| | W W | NN | W N | W W | NN | W N | W W | NN | W N |
| (nb) with $O(1)$ corrections | | | | | | | | | |
| 0.1 | 166.008 | 130.813 | 134.504 | 166.285 | 131.032 | 134.270 | 130.997 | 94.666 | 98.354 |
| 0.3 | 164.702 | 129.797 | 133.416 | 166.006 | 130.833 | 134.036 | 130.705 | 94.491 | 98.127 |
| 0.5 | 162.203 | 128.001 | 131.428 | 165.244 | 130.416 | 133.466 | 130.141 | 94.177 | 97.720 |
| 0.7 | 155.390 | 122.922 | 125.809 | 161.749 | 128.044 | 130.542 | 127.491 | 92.981 | 95.874 |
| 0.9 | 134.334 | 106.478 | 107.945 | 149.866 | 118.822 | 120.038 | 117.491 | 86.303 | 87.696 |
| (nb) with $O(1)$ and $O(2)$ photonic corrections | | | | | | | | | |
| 0.1 | 166.958 | 131.674 | 134.808 | 167.073 | 131.740 | 134.572 | 131.705 | 95.334 | 98.609 |
| 0.3 | 165.447 | 130.534 | 133.583 | 166.686 | 131.467 | 134.231 | 131.339 | 95.118 | 98.314 |
| 0.5 | 162.574 | 128.474 | 131.127 | 165.718 | 130.903 | 133.471 | 130.628 | 94.731 | 97.793 |
| 0.7 | 155.597 | 123.206 | 125.255 | 162.042 | 128.361 | 130.378 | 127.808 | 93.377 | 95.782 |
| 0.9 | 137.153 | 108.820 | 109.677 | 150.732 | 119.560 | 120.411 | 118.229 | 86.931 | 87.961 |
| absolute values of the $O(2)$ pair production correction (nb) | | | | | | | | | |
| 0.1 | 0.007 | 0.004 | 0.015 | 0.046 | 0.045 | 0.024 | 0.045 | 0.047 | 0.024 |
| 0.3 | 0.033 | 0.033 | 0.020 | 0.046 | 0.045 | 0.024 | 0.045 | 0.047 | 0.024 |
| 0.5 | 0.058 | 0.050 | 0.041 | 0.048 | 0.046 | 0.025 | 0.046 | 0.047 | 0.024 |
| 0.7 | 0.090 | 0.074 | 0.069 | 0.069 | 0.059 | 0.042 | 0.059 | 0.051 | 0.036 |
| 0.9 | 0.142 | 0.115 | 0.115 | 0.137 | 0.111 | 0.102 | 0.111 | 0.085 | 0.075 |
| absolute value of the $O(3)$ leading correction (nb) | | | | | | | | | |
| 0.1 | 0.055 | 0.047 | 0.006 | 0.041 | 0.036 | 0.002 | 0.036 | 0.034 | 0.001 |
| 0.3 | 0.065 | 0.053 | 0.018 | 0.046 | 0.040 | 0.007 | 0.040 | 0.037 | 0.003 |
| 0.5 | 0.038 | 0.040 | 0.004 | 0.044 | 0.039 | 0.006 | 0.039 | 0.037 | 0.005 |
| 0.7 | 0.089 | 0.058 | 0.124 | 0.023 | 0.022 | 0.012 | 0.022 | 0.027 | 0.008 |
| 0.9 | 0.291 | 0.220 | 0.331 | 0.021 | 0.013 | 0.049 | 0.013 | 0.002 | 0.038 |

correction in the case BARE1 contains both leading and next-to-leading contributions. In the rest cases the higher order corrections are taken in the leading approximation. For a comparison we give in Table 2 also the corresponding numbers derived by the help of Monte Carlo generator BHLUMI [2]. Parameters are the same as for Table 1. The results of the non-exponentiated version of BHLUMI for BARE1 differs from the exponentiated ones by three digits after decimal point, which are given in parenthesis. The numbers in square brackets are

Table 2

BHLUM I results for the small angle Bhabha cross section (in nb)

| x_c | $O(\alpha^1)$ | | | beyond the first order | | | | | | |
|-------|---------------|---------|---------|------------------------|---------|---------|---------|--------|--------|--|
| | BARE1 | CALO1 | CALO2 | BARE1 | | CALO1 | | CALO2 | | |
| 0.1 | 166.046 | 166.329 | 131.032 | 166.892 (988) | [0.008] | 167.203 | 131.835 | 95.458 | 98.834 | |
| 0.3 | 164.740 | 166.049 | 130.739 | 165.374 (471) | [0.014] | 166.795 | 131.450 | 95.233 | 98.539 | |
| 0.5 | 162.241 | 165.287 | 130.176 | 162.530 (594) | [0.018] | 165.830 | 130.727 | 94.841 | 98.020 | |
| 0.7 | 155.431 | 161.794 | 127.528 | 155.688 (620) | [0.018] | 162.237 | 127.969 | 93.520 | 96.054 | |
| 0.9 | 134.390 | 149.925 | 117.541 | 137.342 (191) | [0.018] | 151.270 | 118.792 | 87.359 | 88.554 | |

absolute difference (in nb) between our second order photonic correction and the one of the non(exponentiated BHLUM I version. BHLUM I numbers beyond the first order for CALO2 case correspond to WW , NN and WN angular acceptances respectively; and the rest is for WW case. Beyond the first order all BHLUM I numbers, except the ones in parenthesis for BARE1, correspond to the version based on the Yennie{Frautchi{Suura exponentiation.

On the level of the first order correction BHLUM I numbers exceed our ones approximately on 0.3 per mille for all variants of event selection. We think that this is due to the difference in our approaches: BHLUM I computes the first order correction [14] according to complete $O(\alpha)$ formulae, while we take into account only t-channel graphs as discussed above.

To be consequent we have to compare our results due to second order photonic contribution with BHLUM I ones which belongs to the non(exponentiated version only. These are the numbers into the parenthesis for BARE1 (three figure after point in the cross section). To compare it needs to remove the difference due to the first order contribution. After this we find that our second order photonic correction which includes leading and next-to-leading contributions exceeds a little bit the BHLUM I one and conclude about very expressive agreement in the case of BARE1 WW .

As concerns calorimeter event selection we have not explicit calculation of the second order next-to-leading contribution. That is why now we can speak about the first order correction only. As one can see from the Tables the agreement of our numbers for WW variant of CALO1 and CALO2 with BHLUM I ones is on the same level as for BARE1 WW one.

Acknowledgements

Authors thank S. Jadach, G. Montagna, B. Pietrzyk and B. Ward for fruitful discussions. This work was supported in part by the INTAS Grant 93{1867.

References

- [1] LEP Electroweak Working Group, CERN Report LEPEW WG/95{02;
LEP Collaboration notes: ALEPH 95{093 PHYSICS 95{086; DELPHI 95{137 PHYS 562; L3
Note 1814; OPAL Technical Note TN 312 (1995);
I.C. Brock et al., preprint CERN {PPE/96{89' CMU {hep/96{04 (1996).
- [2] H. Anlauf et al., Events Generator for Bhabha Scattering, In: conv. S. Jadach and O. Nicrosini,
CERN Yellow Report CERN 96{01, V 2, (1996) 229.
- [3] S. Jadach, E. Richter-Was, B.F.L. Ward and Z. Was, Comput. Phys. Commun. 70 (1992) 305.
- [4] G. Montagna et al., Comput. Phys. Commun. 76 (1993) 328; Nucl. Phys. B 401 (1993) 3;
M. Cacciari, G. Montagna, O. Nicrosini and F. Piccinini, Comput. Phys. Commun. 90 (1995)
301; preprint CERN {TH/95{169.
- [5] S. Jadach, E. Richter-Was, B.F.L. Ward and Z. Was, Phys. Lett. B 353 (1995) 349, 362;
S. Jadach, M. Melles, B.F.L. Ward and S.A. Yost, Phys. Lett. B 377 (1996) 168.
- [6] S. Jadach, M. Skrzypek and B.F.L. Ward, Phys. Rev. D 47 (1993) 3733;
S. Jadach, E. Richter-Was, B.F.L. Ward and Z. Was, Phys. Lett. B 260 (1991) 438.
- [7] W. Beenakker, F.A. Berends and S.C. van der Marck, Nucl. Phys. B 355 (1991) 281;
W. Beenakker and B. Pietrzyk, Phys. Lett. B 296 (1992) 241; *ibidem* B 304 (1993) 366.
- [8] M. Gao, H. Czyz and E. Remiddi, Nuovo Cim. 105 A (1992) 271; Int. J. Mod. Phys. 4 (1993)
591; Phys. Lett. B 327 (1994) 369.
- [9] G. Montagna, O. Nicrosini and F. Piccinini, preprint FNT/T {96/8.
- [10] A. Arbuzov et al., CERN Yellow Report, CERN 95{03, 369;
A.B. Arbuzov et al., preprint CERN {TH/95{313, hep{ph/9512344, Nucl. Phys. B 485 (1997)
457.
- [11] A.B. Arbuzov, E.A. Kuraev, N.P. Merenkov and L. Trentadue, JETP 81 (1995) 638.
- [12] T.D. Lee and M. Nauenberg, Phys. Rev. 133 (1964) 1549.
- [13] V.G. Gorshkov, Usp. Fis. Nauk 110 (1973) 45;
F.A. Berends et al., Nucl. Phys. B 57 (1973) 371;
E.A. Kuraev and G.V. Melehin, Nucl. Phys. B 122 (1977) 485.
- [14] S. Jadach and B.F.L. Ward, Phys. Rev. D 40 (1989) 3582.