

CERN-TH/96-325

hep-ph/9612209

OUTLOOK ON NEUTRINO PHYSICS

John Ellis

TH Division, CERN, 1211 Geneva 23, Switzerland

ABSTRACT

Some of the hot topics in neutrino physics are discussed, with particular emphasis on neutrino oscillations. After proposing credibility criteria for assessing various claimed effects, particular stress is laid on the solar neutrino deficit, which seems unlikely to have an astrophysical explanation. Comments are also made on the possibility of atmospheric neutrino oscillations and on the LSND experiment, as well as cosmological aspects of neutrinos and neutralinos. Several of the central issues in neutrino physics may be resolved by the new generation of experiments now underway, such as CHORUS, NOMAD and Superkamiokande, and in preparation, such as SNO and a new round of accelerator- and reactor-based neutrino-oscillation experiments. At the end, there is a brief review of ways in which present and future CERN experiments may be able to contribute to answering outstanding questions in neutrino physics.

Invited Talk presented at the Conclusion of the Neutrino 96 Conference, Helsinki, June 1996.

CERN-TH/96-325

November 1996

OUTLOOK ON NEUTRINO PHYSICS

John Ellis

TH Division, CERN, CH-1211 Geneva 23, Switzerland

Abstract

Some of the hot topics in neutrino physics are discussed, with particular emphasis on neutrino oscillations. After proposing credibility criteria for assessing various claimed effects, particular stress is laid on the solar neutrino deficit, which seems unlikely to have an astrophysical explanation. Comments are also made on the possibility of atmospheric neutrino oscillations and the LSND experiment, as well as cosmological aspects of neutrinos and neutralinos. Several of the central issues in neutrino physics may be resolved by the new generation of experiments now underway, such as CHORUS, NOMAD and Superkamiokande, and in preparation, such as SNO and a new round of accelerator- and reactor-based neutrino-oscillation experiments. At the end, there is a brief review of ways in which present and future CERN experiments may be able to contribute to answering outstanding questions in neutrino physics.

1 Preamble

The story so far is that we know there are three (and only three) light neutrino species¹: apart from that, all we have are upper limits on neutrino properties². According to the Standard Model, each neutrino flavour carries its own lepton number, and these are all separately conserved: $L_e = L_\mu = L_\tau = 0$, and all the light neutrinos are in fact massless: $m_e = m_\mu = m_\tau = 0$. One may either accept the Standard Model's opinion, and use neutrinos as tools for probing nuclear/particle physics, astrophysics and cosmology, or one may question the Standard Model's wisdom, and study neutrinos in their own rights as possible arenas for new physics beyond the Standard Model. Most of this meeting has been, and most of this talk is, concerned with searches for such new physics, which is also the likely focus of most future neutrino experiments, but let us first review briefly some Standard Model physics with neutrinos.

2 Testing the Standard Model

Neutrinos may be used in nuclear physics to measure interesting matrix elements, which may cast light on nuclear wave functions. In this connection, it was reported here³ that the LSND experiment finds a discrepancy with calculations of $(\nu + {}^{12}\text{C} \rightarrow \nu + \text{N})$. Also, it should not be forgotten that neutrino-proton elastic scattering may also cast light on the presence of hidden strangeness in the proton⁴. It would be a pity if these mundane issues got forgotten completely in the enthusiasm for possible new physics with LSND.

Neutrinos can also be used to make measurements of fundamental Standard Model parameters, such as $\sin^2 \theta_W$ and s_ν . We heard here⁵ that the present error on the deep-inelastic $\nu\text{-N}$ scattering measurement of $\sin^2 \theta_W$ is 0.0048, to be compared with the LEP/SLC error of about 0.0003. However, this comparison is not really fair, since different definitions of $\sin^2 \theta_W$ are used. The $\nu\text{-N}$ measurement can be considered as effectively a determination of $M_W = M_Z$. The promised⁵ error of 0.0025 is equivalent to $M_W = 130 \text{ GeV}$, comparable to the present error on the direct measurements of M_W ^{6,7}, and should be compared with the errors of 50 MeV or so expected in the future from LEP 2 and the Tevatron collider. These measurements are in turn important for the indirect determination of the Higgs mass. A global fit to all the available electroweak measurements, including present $\nu\text{-N}$ scattering data, gives⁸

$$\log_{10}(M_H/\text{GeV}) = 2.16 \pm 0.33 \quad (1)$$

corresponding to a factor of 2 error in M_H , which we may hope future $\nu\text{-N}$ scattering experiments could help reduce.

Deep-inelastic $\nu\text{-N}$ scattering also provides an interesting determination of s_ν , in particular from the Gross-Llewellyn-Smith sum rule⁹:

$$s_\nu(M_Z) = 0.111 \pm 0.003 \pm 0.004 \quad (2)$$

where the dominant experimental systematic error comes from the beam energy calibration⁵. Taking this into account, the measurement (2) is not incompatible with the world average of $s_\nu(M_Z) = 0.118 \pm 0.003$ ¹⁰. There are other low-energy determinations of $s_\nu(M_Z)$ which yield relatively high values, e.g., from τ decay¹¹ and the Bjorken sum rule¹², and the available measurements seem to me to scatter normally around this central value, with no significant trend for lower-energy determinations to yield lower values, as has sometimes been suggested. To make a graceful transition to physics beyond the Standard Model, it is worth recalling that measurements of s_ν and $\sin^2 \theta_W$ test theories of Grand Unification, so these neutrino experiments bear on fundamental theoretical issues.

3 Massive Neutrinos?

As already mentioned, the Standard Model abhors masses for the neutrinos, and all we have so far from experiments are upper limits²:

$$m_e < 4.5\text{eV}; m_\mu < 160\text{KeV}; m_\tau < 23\text{MeV} \quad (3)$$

Theoretically, there is no good reason for the neutrino masses to vanish, and non-zero masses are expected in generic grand unification theories. There are probably more models for the neutrino mass matrix than there are theorists who have worked on the problem: here I shall be guided by the simplest see-saw form¹³:

$$(\bar{L}; R) \begin{pmatrix} 0 & m_q \\ m_q & M_{\text{GUT}} \end{pmatrix} \begin{pmatrix} L \\ R \end{pmatrix} \quad (4)$$

whose diagonalization leads naturally to light neutrinos

$$m_i \sim \frac{m_{q_i}^2}{M_{\text{GUT}}} \ll m_q, \quad (5)$$

These non-zero masses would be accompanied by mixing via angles θ_{ij} : $i, j = e, \mu, \tau$; analogous (and in some models related) to the Cabibbo-Kobayashi-Maskawa angles in the quark sector.

Our best direct information about m_e comes from measurements of the end point of the ^3He β -decay spectrum. The latest results of two current experiments were presented here. As seen in Fig. 1, neither data set is fit well by the standard theory. Relative to the standard Kurie plot, the Mainz spectrum¹⁴ shown in Fig. 1(a) has a banana shape that raises question about the presence of final-state ionic and/or molecular effects that are not well understood. In the case of the Troitsk experiment¹⁵ (see Fig. 1(b)), there is a mysterious bump near the end of the spectrum that appears at different energies in the data from different years. Both experiments report^{14,15} a feature around 250 to 300 eV from the end point, with an apparent branching probability of 1 or 2%. However, before accepting a very stringent upper limit on m_e from the end-point measurements, or crying λ on the basis of this lower-energy feature, the community will require more evidence that the theoretical understanding of the spectrum is adequate.

The Mainz group does plan¹⁴ to measure final-state effects by monitoring additional π and molecular break up in a more calorimetric approach. There is also an interesting suggestion¹⁶ to use ^{187}Re (which has $Q = 2\text{KeV}$) in a calorimetric experiment based on superconducting bolometry to remove final-state effects.

A powerful indirect way to search for m_e is to look for $0 \nu \beta\beta$ decay. However, the interpretation of these experiments is subject to uncertainties: a positive effect could be due to some other mechanism such as R

violation in supersymmetric models, and, even if the results continue to be negative, there could be a Dirac mass or some other mechanism for a cancellation in $\langle m_e \rangle$. Nevertheless, the upper limit¹⁷

$$\langle m_e \rangle < 0.5 \text{ eV} \quad (6)$$

is very impressive. As we heard here¹⁸, there are prospects for incremental improvements, leading perhaps to $m_e < 0.2 (0.1) \text{ eV}$ by the year 2000 (2005), and many other horses will be competing in this race.




Figure 1: Features at the ends of the tritium β -decay spectra seen (a) in the Mainz experiment¹⁴, and (b) in the Troitsk experiment¹⁵, which are not fitted well by standard models.

4 Neutrino Oscillations?

This is a very difficult experimental field, and the stakes are very high. Therefore, we must ask for a very high standard of proof: my personal credibility criteria are that any claimed effect should be confirmed by more than one experiment using more than one technique. From this standpoint, the possibility of Solar Neutrino Oscillations holds up very well: the solar neutrino deficit has now been seen by 5 experiments (Homestake¹⁹, Kamikande²⁰, SAGE²¹, GALLEX²² and now Superkamikande²⁰) using 3 1=2 techniques (Cl, H₂O and two somewhat different Ge techniques). The claim of Atmospheric Neutrino Oscillations, on the other hand, is supported by just 2 1=2 experiments (Kamikande²⁰ and IMB²³, with Soudan²⁴ as yet inconclusive), using just 1 1=2 techniques (water Cherenkov and 1=2 for calorimetry/tracking, since other tracking detectors (NUSEX, Frejus) do not see the effect). Finally, the claim of possible Accelerator Neutrino Oscillations is made by only one experiment (LSND²⁶) using a scintillator technique.

In my view, the fact of a solar neutrino deficit is well established²⁷, and the essential question now is whether it is due to astrophysics or neutrino properties. On the other hand, the existence of an atmospheric neutrino deficit cannot be regarded as well established, but requires further checks. There are still considerable uncertainties in the absolute cosmic-ray neutrino fluxes²⁸ which cloud the interpretation (too few or too many?), and confirmation by different techniques is essential. The LSND²⁶ claim of an accelerator neutrino effect is in even greater need of confirmation, and it is encouraging that the LSND Collaboration itself, as well as an upgrade of the KARMEN experiment³⁰, should be able to address this issue.

We saw at this meeting new, and in some cases definitive, results from the four experiments that see a solar neutrino deficit. Homestake now reports¹⁹ a flux of $2.54 \pm 0.16 \pm 0.14$ SNU, and the previous suggestion of some time dependence (an anti-correlation with the sunspot number) has declined substantially in statistical significance. The definitive Kamikande result of $0.424 \pm 0.029 \pm 0.05$ of the Bahcall-Pinsonneault 1995 flux also exhibits no significant time dependence²⁰:

$$\frac{\text{Data}}{\text{Standard Solar Model}} = 0.398^{+0.088}_{-0.078} + (9.4^{+7.2}_{-7.0})10^{-4} N_{ss} \quad (7)$$

SAGE²¹ reported here a new result: 72^{+15+5}_{-10-7} SNU and the preliminary results of a calibration test: $\phi = 0.54 \pm 0.11$. We also heard about new data from GALLEX²², which are somewhat below their previous average, though consistent within the errors. Their latest result is 70 ± 8 SNU, which includes statistical and systematic errors, the latter incorporating the results from their two calibration runs, which together yield

$= 0.92 \pm 0.07$ ²². These two new results tend even to sharpen the dilemma of the solar neutrino deficit.

As was discussed here by Smirnov³², and seen in Fig. 2, the indications are that the deficit is not monotonic with neutrino energy, in the way that would be expected, for instance, if it was due simply to a reduction in the central temperature of the Sun. Nor does the deficit look energy-independent³³, as was suggested here by Conforto³⁴. The latest Standard Solar Model calculations do include diffusion effects and the latest available opacities²⁷. Personally, I am at least reassured about the accuracy of the assumed values of the nuclear interaction rates, but it still seems very difficult to accommodate the apparent (near?) absence of intermediate-energy solar neutrinos coming mainly from ⁷Be.

Since there was no talk here on helioseismology, I would like to underline that this does not help solve the solar neutrino problem³⁵, but rather the reverse³⁶. Quoting from³⁷: "\substantial mixing... solution of... deficit of ³⁷Cl... seems unlikely", "\inclusion of settling... tends to increase neutrino fluxes", "\helioseismological results accentuate the neutrino problem", "\models that... reduce the neutrino flux... are generally inconsistent with the observed frequencies", and "\it appears unlikely that the solar neutrino problem will find an astrophysical solution".




Figure 2: A planar presentation³³ of the solar neutrino deficits seen in different experiments, compared with a selection of different models. The data do not support a suppression that is independent of energy³⁴.

As is well known, and was emphasized here by Smirnov³² and by Petoov³⁸, matter-enhanced Mikheyev-Smirnov-Wolfenstein (MSW)³⁹ oscillations fit the data very well: a fit⁴⁰ including the latest data is shown in Fig. 3 (a). An issue raised here was the possibility of a diminution of the MSW effect by fluctuations in the solar density^{41;42}: it will be important to take helioseismological constraints into account when evaluating this possibility. We should also not forget the possibility of vacuum oscillations: the results of a fit⁴⁰ including all the latest data are shown in Fig. 3 (b). Perhaps our biggest peril is theoretical seduction by the MSW³⁹ idea: caveat emptor, and do not forget that "the Sun is not a piece of cake"⁴³!




Figure 3: Fits⁴⁰ to the solar neutrino data available before and after this meeting: (a) within an MSW interpretation, and (b) using vacuum oscillations.

The outlook for progress on the solar neutrino problem is good. The first experiment in the next generation, Superkamiokande, is already operating, and its preliminary data confirm the previous measurements of the Kamiokande experiment²⁰. It should soon supplement this with additional information on the ⁸B neutrino spectrum, and possible seasonal and day/night effects, thereby providing us with many tests of the MSW interpretation of the solar neutrino deficit. The Homestake Iodine experiment is about to start taking data¹⁹. During 1997, SNO should enter into operation⁴⁴, and provide us with the first information on the neutral/charged current ratio, as well as more detailed information on the possible spectrum distortion and the possible seasonal effect, providing many tests of the MSW and vacuum oscillation interpretations. Also approved is the BOREXINO experiment⁴⁵, which should provide us with definitive information on the intermediate-energy (⁷Be) neutrino

nos, and hence on the MSW interpretation, starting in 1999. Other projects for the Gran Sasso laboratory are under active discussion. One is the Gallium Neutrino Observatory (GNO)²², which would continue the work of GALLEX, eventually increasing the detector mass to 100 t. The ICARUS experiment⁴⁶ has already been approved for a prototype module, and should eventually be able to contribute a measurement of the ^8B spectrum. The HELIAZ project⁴⁷ would be able to contribute a measurement of the pp spectrum and test the vacuum oscillation interpretation. It may take a few years, but we are on the way to an experimental resolution of the solar neutrino problem.

As already mentioned, a deficit in the atmospheric $\bar{\nu}_e$ ratio has been reported by the H₂O Cerenkov detector Kamikande:

$$\frac{(\bar{\nu}_e)_{\text{data}}}{(\bar{\nu}_e)_{\text{Monte Carlo}}} = 0.60^{+0.06}_{-0.05} \quad (8)$$

in their sub-GeV data, and an angle-dependent effect has been seen in their multi-GeV data²⁰. A deficit in this ratio has also been seen by the other H₂O detector IMB²³, but is not confirmed by the electronic detectors NUSEX and Frejus²⁵, whilst the Soudan result²⁴

$$\frac{(\bar{\nu}_e)_{\text{data}}}{(\bar{\nu}_e)_{\text{Monte Carlo}}} = 0.72 \pm 0.19^{+0.05}_{-0.07} \quad (9)$$

is ambiguous. Because of uncertainties of as much as 30% in the absolute neutrino flux normalizations²⁸, as seen in Fig. 4, the correct interpretation of the ratios (8, 9) is not immediately obvious: are there too few $\bar{\nu}_e$, or too many ν_e , or both? The ambiguity could be removed by more accurate data on $\bar{\nu}_e$ production in laboratory p-Nucleus collisions, which should soon be available from the SPY experiment at CERN⁴⁸, and on the cosmic-ray muon flux, which could be obtained with the L3 experiment at CERN⁴⁹.

There are several prospects for experimental progress on the atmospheric neutrino problem. Superkamikande²⁰ will soon have an order of magnitude more data than Kamikande, with a correspondingly better control of systematics, less leakage from the side, more stopping muons, and better multi-GeV information. However, it does not use a new technique. There are also several prospects of checks using accelerator neutrino beams over long base lines. First among these will be the LBL experiment²⁰ in which a neutrino beam is sent from KEK to Superkamikande, a distance of 250 km, produced initially by a 12 GeV p beam in 1999, which can look directly for $\bar{\nu}_e$ oscillations, and subsequently with a 50 GeV p beam in 2003, which will permit the observation of ν_e . In the United States, the MINOS experiment⁵⁰ involves sending the beam produced by a 120 GeV p beam over 730 km to

Soudan, which is currently scheduled to start in 2002. This experiment should be sensitive to $m^2 = 0.001 \text{ eV}^2$ and $\sin^2 2\theta = 0.02$. Also under discussion is a possible CERN-Gran Sasso experiment⁵¹, which I will discuss later.




Figure 4: Comparison²⁹ of the atmospheric μ and e fluxes observed in different experiments with theoretical calculations²⁸.

Some aspects of the atmospheric neutrino problem can be checked with reactor experiments that are sensitive to $\bar{\nu}_e$ disappearance. The Chooz experiment⁵² with a base line of 1 km starts taking data in 1996, and the Palo Verde experiment⁵³ with a 750-m base line in 1997: they should each reach $m^2 = 0.001$ and $\sin^2 2\theta = 0.05$.

One of the most dramatic recent claims in neutrino physics has been that by the LSND collaboration²⁶ to have observed an excess of $\bar{\nu}_e p$!

e^+n events (tagged by a second signature from $np \rightarrow d\gamma$) which they interpret as evidence for $\bar{\nu}_e$ oscillations. The largest irreducible background is that due to a contamination of 8×10^{-4} e^- in the beam. There is in addition a reducible background due to cosmic rays, which is fought by shielding, veto counters, etc. The LSND collaboration reports a raw e^+ excess for $36 \text{ MeV} < E < 60 \text{ MeV}$ (recall that the end point is at 53 MeV) of $300 \pm 160.5 \pm 76.2 = 63.3 \pm 20$ events (where the first subtraction is for the beam \rightarrow background, the second for the beam \rightarrow background), a 3σ effect. When they further select events with a coincidence, as determined by a cut on a function $R(t; N_{PMT}; r) > 30$, as seen in Fig. 5, they are left with $22 \pm (2.5 \pm 0.4) \pm (2.1 \pm 0.4) = 17.6 \pm 4.7$ events, somewhat more than 3σ .




Figure 5: The tail of the R distribution for events with $20 \text{ MeV} < E_e < 60 \text{ MeV}$ in the LSND experiment has an apparent excess²⁶.

There were initially some worries about the possibility of neutrons leaking into the outer regions of the detector⁵⁴. If one restricts the fiducial volume to the "safest" 45%, the signal is reduced to 6 events with a background of 1.7 ± 0.3 ²⁶. It should also be noted²⁶ that the excess in R shows up only in the last two bins shown in Fig. 5. Finally, the LSND result is only marginally consistent with previous experiments, most importantly E 776, KARMEN³⁰ and the Bugey reactor experiment, and the most recent CCFR limit³¹, as seen in Fig. 6.

The LSND speaker here²⁶ talked of a "significant oscillation-like excess that needs confirmation". Many people here would echo heartily the latter phrase. Fortunately, efforts to confirm the claimed effect are underway by LSND itself²⁶, which is searching in its data for $\bar{\nu}_e$ oscillations, and should have results by 1997, and by the KARMEN collabor-

oration³⁰, which is improving its n detection efficiency and installing veto counters in its surrounding steel blockhouse to veto cosmic-ray muons that might produce neutrons, and should have results by 1998^a.




Figure 6: Comparison of the LSND result²⁶ with those of E 776, KARMEN³⁰, Bugey and CCFR³¹.

Finally, let us recall the motivation for a new round of accelerator neutrino-oscillation experiments. Theorists of cosmological structure formation would like some Hot Dark Matter⁵⁶, which is most plausibly one or more massive neutrinos with mass 1 to 10 eV, which is compatible with the simplest possible seesaw model:

$$m = \left(\frac{m_t}{m_c}\right)^2 m \quad (10)$$

It is, furthermore, plausible that the m might have observable mixing with the ν_e . These considerations have motivated the CHORUS⁵⁷ and NOMAD⁵⁸ experiments, which will be running at CERN until the end of 1997, and should attain sensitivities 0.0003 to \sin^2 for $m^2 \leq 50$ eV². In my view, there is certainly interest in extending this search down to lower m^2 , as well as to smaller mixing angles, by using either

^aThe KARMEN collaboration presented³⁰ their own "anomaly", namely an apparent excess of 112 ± 32 events occurring after 3.6 ± 0.25 s, that might be due to the radiative decay of some state X with a mass of 33.9 MeV. However, this effect has not been confirmed by an experiment at PSI⁵⁵.

a longer baseline or a lower-energy beam. Indeed, in the longer term, the COSMOS experiment⁵⁹ plans to take data for 4 years starting in 2001, and expects to reach a sensitivity to $\sin^2 \theta_{12} \sim 1:4 \cdot 10^{-5}$, and get down to $m^2 \sim 0.08 \text{ eV}^2$.

5 Cosmological Constraints on Neutrinos

We heard here⁶⁰ that the success of Big-Bang Nucleosynthesis constrains the number of neutrino species:

$$N = 3.0 \pm 0.23 \pm 0.38^{+0.11}_{-0.57} \quad (11)$$

where the last uncertainty is that due to the baryon-to-entropy ratio. We also heard⁶⁰ that there is no crisis for Big-Bang Nucleosynthesis, but possibly for over-simplified theories of the chemical evolution of the Galaxy⁶¹. The range (11) corresponds to an upper bound

$$N < 3.9 \text{ (90\% c.l.)} \quad (12)$$

to be compared with the LEP determination $N = 2.989 \pm 0.012^1$.

We also heard⁶² that Big-Bang Nucleosynthesis constrains the mass of any metastable neutrino, assumed to be the ν_{τ} neutrino:

$$m_{\nu_{\tau}} > 32 \text{ MeV or } < 0.95 \text{ MeV} \quad (13)$$

if it has a Dirac mass, and

$$m_{\nu_{\tau}} > 25 \text{ MeV or } < 0.37 \text{ MeV} \quad (14)$$

if it has a Majorana mass. Both of the lower limits in (13,14) conflict with the upper bound $m_{\nu_{\tau}} < 23 \text{ MeV}$ from ALEPH⁶³, indicating that $m_{\nu_{\tau}}$ must actually lie below 1 MeV. We also know that any stable neutrino must weigh less than

$$m_{\nu} = 92h^2 \text{ eV} \quad (15)$$

where h is the present expansion rate of the Universe in units of 100 km/s/Mpc, and it is assumed that the present mass density of the Universe is no larger than the critical density, $\Omega = 1$, as suggested by theories of cosmological inflation. There are also cosmological constraints on unstable neutrinos⁶², which I will not go into here.

There has recently been discussion⁵⁶ about the observational value of h and its compatibility with the age of the Universe as determined by astrophysicists. We see from Fig. 7 that there is no significant discrepancy⁵⁶, provided h is not much larger than about 0.5^b , corresponding

^bThis is not incompatible with recent astrophysical determinations of h , for example from time delays in quasar lensing⁶⁴.

to $m < 23 \text{ eV}$. On the other hand, it may well be that most of the critical density is not in the form of massive neutrinos.




Figure 7: The $(\Omega; H_0)$ plane exhibits no serious discrepancy between the average measured value of H_0 (indicated by the vertical error bar), $\Omega = 1$ and an age for the Universe of 10^{10} years. This plot also shows the estimates of the present baryon density obtained from visible features in the Universe, from Big Bang Nucleosynthesis and from rich clusters.

The favoured theory of the formation of structures in the early Universe is that they are due to the gravitational instability of overdense regions as they come within the horizon. These overdense regions would have originated from density perturbations created by quantum fluctuations during the inflation. However, it is generally thought that the growth of such inflationary perturbations needs to be accelerated by matter that is non-relativistic (cold) during the epoch of structure formation⁵⁶, as illustrated in Fig. 8.

It is generally thought that most of the present mass density must be in the form of cold dark matter. However, the detailed comparison of data on microwave background fluctuations as first observed using the COBE satellite⁶⁵, other observational data on large-scale astrophysical structures, and data on smaller scales, indicate that the pure cold dark matter model requires modification. One possibility is that there may be energy density in the vacuum (a cosmological constant), another is that the spectrum of inflationary perturbations may be scale-dependent

(tilted), and a third is that there is an admixture of hot dark matter⁶⁶, i.e., matter that was relativistic during the epoch of structure formation:

$$\Omega_{\text{Cold}} = 0.7; \quad \Omega_{\text{Hot}} = 0.2 \quad (16)$$

The only plausible candidate for the hot component of the dark matter is one or more species of massive neutrino.




Figure 8: A sketch indicating the rôle of Cold Dark Matter in magnifying primordial density perturbations via gravitational instability, while the baryons are still coupled to radiation.

Simulations of structure formation indicate that any cosmological constant is unlikely to be large enough to alleviate significantly the (non-existent) age problem. Moreover, peculiar motions on the scale of galactic clusters seem to be larger than expected in cold dark matter models with tilt. As summarized in Fig. 9, mixed dark matter models (16) seem to fit the observational data best⁵⁶. Although it has been suggested that there may be more than one species of neutrino with similar masses⁶⁷, the simplest hypothesis, which is consistent with the seesaw matrix (4), is that one neutrino, most likely the ν_τ , dominates the hot dark

matter density. In such a mixed dark matter model, one would have

$$m \approx 5 \text{ eV} \quad (17)$$

Moreover, such a mass (17) is quite consistent with the value of $m \approx 2 \times 10^{-3} \text{ eV}$ expected in the MSW interpretation of the solar neutrino deficit, and the ratio $(m_t = m_c)^2$ expected in the simple seesaw model (4). If true, the estimate (17) suggests an optimization of accelerator searches for neutrino oscillations down to $m^2 \approx 10 \text{ eV}^2$.




Figure 9: A sketch indicating the relative successes of different models of structure formation, as compared with different types of astrophysical and cosmological data⁵⁶.

6 Neutrinos

We have already addressed the possibility that a massive neutrino might constitute hot dark matter in the Universe. Whether this is necessary is still an open question: what seems much better established is the need for a large amount of cold dark matter, at least on a cosmological scale if inflation is to be accepted, and also on a galactic scale if models of structure formation are to be believed. The question of immediate experimental interest is how much dark matter of what type may be present in the galactic halo that surrounds us. This is presumably not constituted of neutrinos, because they would not cluster on such a small scale⁶⁸.

Recently a couple of observational programmes have seen microlensing events interpreted as due to sub-solar-mass objects in the galactic halo⁶⁹. The best estimate of the MACHO collaboration⁷⁰ is that about half our halo could be in this baryonic form, and perhaps even all of it, whereas the EROS collaboration⁷¹ seems to rule out this probability. Even if 60% or more of our galactic halo is in the form of microlensing objects, one may anticipate a local density of cold dark matter particles of 10^{-25} g cm^{-3} or more⁷².

My favourite candidate for this cold dark matter is the lightest supersymmetric particle, usually thought to be a neutralino, i.e., some mixture of the spin-1=2 supersymmetric partners of the Z^0 , and Higgs⁷³. Fig. 10 shows the lower limit on the neutralino mass established (modulo certain loopholes) by the ALEPH collaboration⁷⁴ on the basis of unsuccessful sparticle searches at LEP 1 and 1.5. There is a good likelihood⁷⁶ that the cosmological relic density of neutralinos may lie in the range of interest to stationary cosmologists: $0.1 < h^2 < 0.3$ ⁷⁷. As also seen in Fig. 10, the ALEPH limit⁷⁴ may be strengthened (and its loopholes removed) if one postulates that the neutralino density falls within this range^{75 c}.

Both of these analyses^{74;75} are based on the assumption that sparticle masses have certain universality properties which may not be valid. If these assumptions are relaxed, the lower bound on the mass of the lightest neutralino may not be altered qualitatively, but its phenomenology may be significantly modified⁷⁷, perhaps changing from a mainly gaugino composition to a mainly higgsino composition as seen in Fig. 11. Such a change can also alter the prospects for neutralino searches, as we discuss next.

Three strategies for such searches are favoured: neutralino annihilation in the galactic halo which yields stable particles ($p; e^+; \dots$) in the cosmic rays⁷⁹, which will be explored by the AMS satellite, annihilation after capture within the Sun or Earth, which may yield high-energy neutrinos detectable in underground detectors (either directly or via the γ s generated by interactions in rock)⁸⁰, and elastic scattering on a nuclear target in the laboratory⁸¹.

^cFor updates of this analysis to include the preliminary results of higher-energy LEP 2 data, see⁷⁸.




Figure 10: The ALEPH lower limit⁷⁴ on the lightest neutralino mass $m_{\tilde{\chi}_1^0}$ (short-dashed line) as a function of the ratio of Higgs vacuum expectation values $\tan \beta$, which has an important loophole indicated by the double arrow, is compared with the absolute lower limits obtained from a combination with other e^+e^- experiments (dotted line), with that inferred from the D0 experiment (long-dashed line), and those obtained from combining phenomenological and cosmological considerations, both with (solid line) and without (dot-dashed line) the supplementary theoretical assumption of dynamical electroweak symmetry breaking⁷⁵.




Figure 11: The preferred composition of the lightest neutralino may change from being mainly a gaugino to mainly a higgsino, if the normal assumptions of universality are relaxed⁷⁷.

In the second case, which is of particular interest to neutrino physicists, annihilation follows the passage of relic neutralinos through the Sun or Earth accompanied by scattering and the loss of recoil energy, which is what causes the neutralinos to become trapped⁸⁰. Among the neutralino annihilation products will be some energetic neutrinos from heavy-quark decays, which can escape from the core of the Sun or Earth and be detected in underground experiments, either directly or indirectly as mentioned above. Underground search experiments such as Baksan are now imposing significant constraints on models^{77;82}, as seen in Fig. 12. This type of search is promising for the future generation of experiments that includes Baikal, Nestor and Amund⁸³. One point that needs to be watched is the possibility that also these high-energy neutrinos oscillate⁸⁴. In particular, there could be a significant suppression of the solar μ flux and a corresponding enhancement of the e flux if the large-angle MSW solution to the low-energy solar-neutrino deficit is correct. This implies the need for some caution in interpreting upper limits on the flux of muons generated by high-energy neutrinos from the Sun.




Figure 12: Upper limits on energetic neutrino emission from the Sun due to neutralino annihilations are already beginning to exclude certain models⁸².

The direct detection of the elastic scattering of neutralinos on nuclei in the laboratory⁸¹ is also a promising technique. There are important contributions to the scattering matrix elements from both spin-dependent and spin-independent forces. The former are determined by axial-current matrix elements related⁸⁵ to that appearing in neutron decay^d, whereas the latter are related to the different quark contributions to the nucleon mass. These spin-independent forces are coherent for heavy nuclei, and likely to dominate for favoured detector materials such as Ge⁷⁶. Recent upper limits on the spin-dependent and spin-independent elastic dark-matter scattering rates were shown at this meeting⁸⁷. Eventually, such experiments should improve significantly in sensitivity, and may be able to compete with, and complement, accelerator searches for supersymmetric particles⁸⁵.

7 The CERN Experimental Programme

The large accelerator laboratories such as Fermilab and CERN have not been very much in evidence at this meeting, reflecting the long-term trend of neutrino physics towards non-accelerator experiments. Nevertheless, accelerator laboratories do have important roles to play, and I would like to mention briefly some of the neutrino and neutrino-related activities at my home laboratory, CERN.

As you know, the present CERN short-baseline neutrino oscillation experiments, CHORUS⁵⁷ and NOMAD⁵⁸, will continue taking data until the end of 1997. Two other neutrino-related experiments have also been taking data during 1996. One is SPY⁴⁸, which has measured charged-particle yields in p-Be collisions, with a view to the better calibration of accelerator beams and perhaps better predictions for atmospheric fluxes. The other is NA 55⁸⁸, which has been measuring neutron production in collisions, with a view to understanding better this possible background for atmospheric experiments⁸⁹.

Another activity at CERN is COSMOLLEP, which seeks to use the detectors of LEP experiments to look at extended air showers⁹⁰. The use of L3 to measure the atmospheric flux has also been approved recently⁴⁹. This could reduce significantly the current uncertainties in the flux, enabling one to determine whether Kamikande and IMB have been seeing too few and/or too many events²⁸.

There are also extensive discussions taking place at CERN about the possibility of one or more future experiments following CHORUS and NOMAD. One of these possibilities is an idea to use a beam at

^dIt should be emphasized that the naive quark model is not a good guide to the magnitudes of these matrix elements, in particular since there is a significant contribution from the strange quark current⁸⁶

the PS accelerator to probe the LSND claim⁹¹. Another possibility is a short-baseline experiment at the CERN SPS, as a follow-up to CHORUS and NOMAD⁹². Talk of a medium-baseline experiment using a detector in the Jura mountains, about 20km away from CERN, has also been revived recently⁹³. Another attractive option would be a long-baseline experiment⁵¹ sending a beam to the Gran Sasso laboratory, 730 km away, which might be accompanied by a nearby detector near or under the Geneva airport.

In my view, one of the most interesting options on this list is the short-baseline follow-up to CHORUS and NOMAD. The COSMOS experiment⁵⁹ planned for Fermilab is very promising, but it is a long time in the future, and, as I emphasized earlier, one should always require confirmation in the neutrino business. On the other hand, there are already two entries in the long-baseline race, LBL E²⁰ and MINOS⁵⁰. Moreover, the motivations for neutrino oscillations detectable in a suitable short-baseline experiment, namely hot dark matter, the seesaw mechanism and the solar neutrino deficit, remain just as strong as ever. The only change, perhaps, is a trend towards a slightly lower neutrino mass for the cosmological hot dark matter. As already mentioned, this might suggest a somewhat lower neutrino energy and/or longer baseline for any follow-up experiment, which is motivated whether or not CHORUS and NOMAD find anything in their present data sets.

8 Conclusion

To conclude this talk, I present a possible chronology of future neutrino experiments and others of potential interest to this community. An exciting era is opening up, with major new experiments such as Superkamiokande and LEP 2 starting to take data. Some of the major issues in particle physics have a chance of being resolved by the time of Neutrino 98, with the opening up of new domains of exploration for the Higgs boson, supersymmetric particles and neutrino masses. Let us all cross all our enable body parts, and hope for progress by the next meeting in this series!

Possible Chronology of Future Neutrino and Related Experiments

Year	Accelerator Neutrino Experiments	Other Accelerator Experiments	Non-accelerator Experiments
1996	CHORUS, NOMAD	LEP 2, SPY, COSMOLEP, NA 55	AMANDA, Baikal, SKK, Chooz, Homestake Iodine
1997	LSND	CosmoL3	SNO, Palo Verde
1998	KARMEN upgrade		AMS, GNO?
1999	LBL E (12 GeV)	B factories, HERA-B	BOREXINO, ICARUS 600t
2000	ALADINO/NOE/ TENOR/HELLAZ/ full ICARUS?		0.2 ! 0.2 eV?
2001	COSMOS, MINOS		MAP?
2002			
2003	LBL E (50 GeV)		COBRAS/SAMBA?
2004			BAND?
2005		LHC	0.2 ! 0.1 eV?

1. A. Blondel, Plenary talk at the International Conference on High Energy Physics, Warsaw, 1996, reporting the analysis of the LEP Electroweak Working Group and the SLD Heavy Flavor Group, CERN Report No. LEPEWWG/96-02, available at the URL: <http://www.cern.ch/LEPEWWG>.
2. Particle Data Group, R.M. Barnett et al, Phys.Rev. D 54 (1996) 1.
3. P. Vogel, V. Sandberg, talks at this meeting.
4. J. Ellis and M. Karliner, Phys.Lett. B 213 (1988) 73;
D.B. Kaplan and A. Manohar, NuclPhys. B 310 (1988) 527.
5. R. Bernstein, talk at this meeting;
A.L. Kataev, A.V. Kotikov, G. Parente and A.V. Sidorov, Phys.Lett. B 388 (1996) 179.
6. M. Demarteau, Fermilab preprint Conf96/354, hep-ex/9611019,

and references therein.

7. ALEPH, DELPHI, L3 and OPAL collaborations, presentations at open session of the CERN LEP Experiments Committee, Oct. 8th, 1996;
See, for example, OPAL collaboration, K. Ackerstaedt et al., CERN Preprint PPE/96-141.
8. J. Ellis, G.L. Fogli and E. Lisi, CERN preprint TH/96-216 (1996), hep-ph/9608329;
see also W. de Boer, A. Dabelstein, W. Hollik, W. Mosle and U. Schwicklerath, hep-ph/9607286;
S. Dittmaier and D. Schildknecht, hep-ph/9609488.
9. A.L. Kataev and A.V. Sidorov, Phys.Lett. B 331 (1994) 179;
CCFR-NUTEV collaboration, D. Harris et al., hep-ex/9506010;
see also L.S. Barabash et al., hep-ex/9611012.
10. M. Schmelling, rapporteur talk at International Conference on High-Energy Physics, Warsaw 1996.
11. M. Girone and M. Neubert, Phys.Rev.Lett. 76 (1996) 3061, and references therein.
12. J. Ellis, E. Gardi, M. Karliner and M. Samuel, Phys.Lett. B 366 (1996) 268 and Phys.Rev. D 54 (1996) 6986.
13. T. Yanagida, Proc. Workshop on the Unified Theory and the Baryon Number in the Universe (KEK, Japan, 1979);
R. Slansky, Talk at Sanibel Symposium, Caltech Preprint CALT-68-709 (1979).
14. J. Bonn, talk at this meeting. The possibility of a crumpled configuration of the (supposedly) monolayer source, leading to multiple scattering, is a concern in this experiment.
15. V.M. Lobashev, talk at this meeting. I understand from A. Kusenko that a re-evaluation of multiple scattering in the source of this experiment may lead to a revised upper limit on the possible high-mass branch of about 0.5%.
16. A. Swift, talk at this meeting.
17. H.V. Klapdor-Kleingrothaus, talk at this meeting.
18. F.T. Avignone, A. Barabash, H. Ejiri, J. Farine and E. Fiorini, talks at this meeting.
19. K. Lande, talk at this meeting;
20. Y. Suzuki, talk at this meeting:
Kamiohka collaboration, Y. Fukuda et al., preprint ICRR-372-96-23 (1996);
see also J.N. Bahcall, P.I. Krastev and E. Lisi, Princeton preprint IASSNS-AST-96/43.
21. V.N. Gavrin, talk at this meeting;
SAGE Collaboration, D.N. Abdurashitov et al., Phys.Lett. B 328 (1994) 234.

22. T. Kirsten, talk at this meeting;
GALLEx Collaboration, W. Hampel et al., Saclay preprint DAPNIA-SPP-96-10 (1996), and references therein.
23. IMBC Collaboration, R. Becker-Szendy et al., Phys.Rev. D 46 (1992) 3720.
24. E.A. Peterson, talk at this meeting;
Soudan IC Collaboration, W.W.M. Allison et al., hep-ex/9611007.
25. NUSEX Collaboration, M. Aglietta et al., Europhys. Lett. 8 (1989) 611;
Frejus Collaboration, C. Berger et al., Phys.Lett. B 227 (1989) 489 and B 245 (1990) 305.
26. D.O. Caldwell, talk at this meeting;
LSND Collaboration, C. Athanassopoulos et al., nucl-ex/9605001 and nucl-ex/9605003.
27. J. Bahcall, talk at this meeting;
J. Bahcall and M.H. Pinsonneault, hep-ph/9610542.
28. T. Gaisser, talk at this meeting, hep-ph/9611301;
T. Gaisser, M. Honda, K. Kasakara, H. Lee, S. Midorikawa, V. Naumov and T. Stanler, hep-ph/9608253.
29. G.L. Fogli and E. Lisi, Phys.Rev. D 52 (1995) 2775.
30. J. Kleinfeller, talk at this meeting;
KARMEN Collaboration, B. Bodmann et al., Phys.Lett. B 332 (1994) 251 and B 348 (1995) 19.
31. CCFR Collaboration, A. Romosan et al., Columbia preprint NEVIS-1529, hep-ex/9611013.
32. A. Smimov, talk at this meeting
33. See, e.g., N. Hata, S.A. Bludman and P. Langacker, Phys.Rev. D 49 (1994) 3622;
V. Castellani, S. Degl'Innocenti and G. Fiorentini, Astron. and Astrophys. 271 (1993) 601.
34. G. Conforto, talk at this meeting.
35. A. Dar, talk at this meeting, astro-ph/9611014.
36. J.N. Bahcall, M. Pinsonneault, S. Basu and J. Christensen-Dalsgaard, astro-ph/9610250.
37. J. Christensen-Dalsgaard, Nucl.Phys. B (ProcLSuppl.) 48 (1996) 325.
38. S. Petcov, talk at this meeting.
39. L. Wolfenstein, Phys.Rev. D 17 (1978) 2369;
S.P. Mikheyev and A.Yu. Smimov, Yadernaya Fizika 42 (1985) 1441.
40. P. Krashev, private communication.
41. C. Burgess, talk at this meeting;
C.P. Burgess and D. Michaud, hep-ph/9606295 and hep-ph/9611368.

42. A . Rossi, talk at this meeting;
H . Nunokawa, A . Rossi, V . Sem ikoz and J W F . Valle, NuclPhys.
B 472 (1996) 495 and hep-ph/9610526.
43. K . Row ley, public com m unication .
44. A . M cDonald, talk at this meeting;
SNO Collaboration, G . T . Evan et al., SNO -87-12 (1987);
see also J N . Bahcall and E . Lisi, Princeton Preprint IASSNS-AST
96/33 (1996).
45. E . Bellotti, talk at this meeting;
BOREXINO Collaboration, C . A rpesella et al., BOREXINO pro-
posal (University of Milano, Milano, 1992);
see also J N . Bahcall and P . I . K rastev, astro-ph/9607013.
46. ICARUS Collaboration, P . Cennini et al., ICARUS proposal
(1993).
47. HELLAZ Collaboration, G . Laurenti et al., Proc. Fifth Int. W ork-
shop on Neutrino Telescopes, Venice 1993, ed. M . Baldo-Ceolin
(University of Padova, Padova, 1994) 161.
48. SPY Collaboration, <http://www.cern.ch/NA56/> .
49. P . Le Coultre, on behalf of the L3 collaboration, private com m uni-
cation .
50. S . W ojcicki, talk at this meeting;
MINOS Collaboration, E . A bles et al., Ferm ilab proposal P-875
(1995).
51. NOE Collaboration, M . Ambrosio et al., Nucl. Inst. Meth. A 363
(1995) 604;
T . Ypsilantis, Nucl. Inst. Meth. A 371 (1996) 330;
see also ⁴⁶ .
52. C . B en porad, talk at this meeting;
see also R . I . Steinberg, hep-ph/9306282.
53. G . G ratta, talk at this meeting;
Palo Verde Collaboration, F . Bohm et al., Stanford preprint HEP-
96-04 (1996).
54. J . E . H ill, Phys.Rev.Lett. 75 (1995) 2654.
55. M . D aum , talk at this meeting.
56. J . P rim ack, talk at this meeting;
see also J . P rim ack, hep-ph/9610321;
J . P rim ack and A . K lypin, astro-ph/9607061.
57. K . N iwa, talk at this meeting;
CHORUS Collaboration, <http://choruswww.cern.ch/> .
58. J . D um archez, talk at this meeting;
NOMAD Collaboration, <http://nomadinfo.cern.ch/> .
59. R . A . Sidwell, talk at this meeting;
COSMOS Collaboration, P 803 proposal to Ferm ilab (1993).
60. K . O live, talk at this meeting;

- see also: K. Olive, astro-ph/9609071;
 K. Olive and D. Thomas, hep-ph/9610319;
 for the latest words on the primordial lithium abundance, see P.
 Bonifacio and P. Monaco, astro-ph/9611043;
 and on the controversial primordial deuterium abundance, see
 A. Songaila, E. J. Wampler and L. L. Cowie, astro-ph/9611143.
61. N. Hata et al., Phys.Rev.Lett. 77 (1995) 3977;
 G. Steigman, astro-ph/9608084.
 62. K. Kainulainen, talk at this meeting;
 63. A. Gregorio, talk at this meeting;
 ALEPH Collaboration, D. Buskulic et al., Phys.Lett. B 349 (1995)
 585.
 64. T. Kundic et al., astro-ph/9610162.
 65. C. L. Bennett et al., astro-ph/9601067;
 future satellite projects are MAP and COBRAS/SAMBA.
 66. R. Schaefer and Q. Sha, Nature 359 (1992) 119;
 M. Davis, F. J. Summers and D. Schlegel, Nature 359 (1992) 393;
 A. N. Taylor and M. Rowan-Robinson, Nature 359 (1992) 396.
 67. J. Primack, J. Holtzman, A. Klypin and D. O. Caldwell,
 Phys.Rev.Lett. 74 (1995) 2160;
 see also S. Ghigna et al., astro-ph/9611103.
 68. J. Ellis and P. Sikivie, Phys.Lett. B 321 (1994) 390.
 69. A. Milsztajn, talk at this meeting;
 see also: F. Adams and G. Laughlin, astro-ph/9602006;
 G. Chabrier, L. Segretain and D. Mera, astro-ph/9606083;
 A. Gould, astro-ph/9611185.
 70. MACHO collaboration, C. A. Lcock et al., astro-ph/9606165.
 71. EROS collaboration, R. Ansari et al., Astron. and Astrophys. 299
 (1995) L21.
 72. M. S. Turner, E. J. Gates and G. Gyuk, astro-ph/9601168.
 73. J. Ellis, J. S. Hagelin, D. V. Nanopoulos, K. Olive and M. Sred-
 nicki, Nucl.Phys. B 238 (1984) 453.
 74. ALEPH collaboration, D. Buskulic et al., CERN preprint PPE/96-
 83 (1996).
 75. J. Ellis, T. Falk, K. Olive and M. Schmitt, Phys.Lett. B 388
 (1996) 97.
 76. G. Jungman, M. Kamionkowski and K. Griest, Phys. Rep. 267
 (1996) 195.
 77. V. Berezinskii, talk at this meeting, astro-ph/9610263;
 see also A. Bottino, N. Fornengo, G. Mignola, M. Olechowski and
 S. Scopel, astro-ph/9611030.
 78. J. Ellis, T. Falk, K. Olive and M. Schmitt, CERN preprint
 TH/96-284, hep-ph/9610410.
 79. See, for example, J. Silk and M. Srednicki, Phys.Rev.Lett. 53

- (1984) 624;
 J. Ellis et al., Phys.Lett. B 214 (1988) 403.
80. J. Silk, K. A. Olive and M. Srednicki, Phys.Rev.Lett. 55 (1985) 259.
 81. M. Goodman and E. Witten, Phys.Rev. D 31 (1985) 3059.
 82. L. Bergstrom, talk at this meeting;
 L. Bergstrom, J. Edsjö and P. Gondolo, hep-ph/9607237.
 83. F. Halzen, for the AMANDA collaboration, hep-ex/9611014.
 84. J. Ellis, R. A. Flores and S. Masood, Phys.Lett. B 294 (1992) 229.
 85. J. Ellis and R. A. Flores, Phys.Lett. B 263 (1991) 259 and B 300 (1993) 175; Nucl.Phys. B 307 (1988) 375 and B 400 (1993) 25.
 86. J. Ellis and M. Karliner, Lectures at the Int. School of Nucleon Spin Structure, Erice 1995, CERN preprint TH/95-334, hep-ph/9601280.
 87. F. T. A. Vignone, talk at this meeting;
 J. J. Quenby et al., Imperial College, London, Astrophysics Group preprint 5 (1996).
 88. NA55 Collaboration, F. Bohm et al., CERN SPSLC/95-62, Proposal P 293 (1995).
 89. O. G. Ryazhskaya, Nuovo Cimento C 18 (1995) 77;
 J.E.T.P. Lett. 61 (1995) 237;
 see, however, Kamikande Collaboration, Y. Fukuda et al., Preprint ICRR-373-96-24 (1996).
 90. COSMOLEP Collaboration, <http://alephwww.cern.ch/COSMOLEP/>.
 91. P. Zucchelli, in preparation.
 92. ALADINO Collaboration, M. Bonesini et al., CERN SPSLC/95-37, Letter of Intent L205 (1995);
 J.-J. Gomez-Cardenas et al., Nucl. Inst. Meth. A 378 (1996) 196;
 TENOR Collaboration, A. Ereditato et al., CERN preprint PPE/96-106 (1996).
 93. See T. Ypsilantis, second paper in ⁵¹.