

Chirality and Reliability of Baryon QCD Sum Rules*

Xuemin Jin[†] and Jian Tang[‡]

Center for Theoretical Physics

Laboratory for Nuclear Science

and Department of Physics

Massachusetts Institute of Technology

Cambridge, Massachusetts 02139, USA

(MIT-CTP-2601 hep-th/9701230 January 1997)

Abstract

The QCD sum-rule method has been widely used in studying various baryon properties. For a given problem, there are usually more than one sum rules and they do not work equally well. In this paper, we point out that chirality plays an important role in determining the reliability of a baryon sum rule. The contributions of positive- and negative-parity excited baryon states partially cancel each other in the chiral-odd sum rules, but add up in the chiral-even sum rules. As such, the chiral-odd sum rules are generally more reliable than the chiral-even sum rules. This allows one to identify the more reliable sum rules and use them in extracting the ground-state baryon property of interest. This is illustrated in an explicit example.

Submitted to: *Physical Review D*

[†]Email address: jin@ctp02.mit.edu

[‡]Email address: jtang@ctp02.mit.edu

*This work is supported in part by funds provided by the U.S. Department of Energy (D.O.E.) under cooperative research agreement #DF-FC02-94ER40818.

The QCD sum-rule approach [1] has been used extensively in extracting baryon observables [2,3]. It is found in these studies that for a given problem there are more than one sum rules and they do not work equally well. In particular, some sum rules work well while the others may fail. This pattern has been seen in both the baryon mass sum rules and the sum rules for other baryon properties [3]. (Here we consider the baryons containing only light flavors: up, down, and strange.)

In this paper, we point out that chirality plays an important role in determining the reliability of baryon QCD sum rules. In the “chiral-odd” sum rules where chiral-odd operators dominate, the contributions of positive- and negative-parity excited baryon states partially cancel each other, which significantly reduces the excited-state contamination. On the other hand, the “chiral-even” sum rules where chiral-even operators dominate, suffer from large excited-state contamination because of the addition of the contributions from positive- and negative-parity excited states. As such, the chiral-odd sum rules are generally more reliable than the chiral-even sum rules. This allows one to identify the more reliable sum rules in a given problem and use them in extracting the ground-state baryon property of interest. We will illustrate this point in an explicit example.

To keep our discussion succinct, we will use the octet baryon mass sum rules to explain the idea, which applies to general baryon sum rules based on two-point and three-point correlation functions. The baryon mass sum rules usually study the correlation function

$$\Pi(q) \equiv i \int d^4x e^{iq \cdot x} \langle 0 | T[\eta_B(x) \bar{\eta}_B(0)] | 0 \rangle , \quad (1)$$

where η_B is a baryon interpolating field constructed from local QCD operators, carrying the quantum numbers of the baryon of interest. For the octet baryons, there are two independent interpolating fields which contain only quark fields with no derivatives and couple to spin-1/2 states only [4,5]. The general expression of the proton interpolating field is

$$\eta_p(x) = \epsilon_{abc} \left\{ \left[u^{aT}(x) C \gamma_5 d^b(x) \right] u^c(x) + \beta \left[u^{aT}(x) C d^b(x) \right] \gamma_5 u^c(x) \right\} , \quad (2)$$

where $u(x)$ and $d(x)$ stand for up and down quark fields, a , b , and c are the color indices, $C = -C^T$ is the charge conjugation matrix, and β is an arbitrary real parameter. The corresponding interpolating fields for the other members of the octet can be obtained by applying appropriate SU(3) rotations [2]. The interpolating field advocated by Ioffe [4] and often found in QCD sum-rule calculations may be recovered by setting $\beta = -1$ and multiplying an overall factor of -2 . Lorentz covariance and symmetries then dictate that $\Pi(q)$ has two invariant structures

$$\Pi(q) = \Pi_s(q^2) + \Pi_q(q^2) q_\mu \gamma^\mu , \quad (3)$$

where Π_s and Π_q are two invariant functions. So, two sum rules, one for Π_s and one for Π_q , can be obtained. In principle, results obtained from these sum rules should be the same. In practice, however, one has to truncate the operator product expansion (OPE) on the QCD side of the sum rule and use a crude model for the excited-state (continuum) contribution on the phenomenological side of the sum rule. Thus, one expects that some sum rules are more reliable than the others.

One of the key ingredients of the QCD sum-rule approach is the use of the Borel transform, which introduces an auxiliary parameter–Borel mass. If a sum rule were perfect, one would expect that the two sides of the sum rule overlap for all values of the Borel mass. In practical calculations, the two sides of the sum rules overlap only in a limited range of the Borel mass (at best) because of the truncation of the OPE and the crudity of the continuum model. A common consensus is that if there exists a wide region of Borel mass where the two sides of a sum rule match, the sum rule is said to “work” and the sum-rule prediction is possible. This, however, needs to be interpreted carefully.

To maintain the predicative power of the sum-rule approach, the phenomenological side of a sum rule is usually parameterized by a pole describing the ground-state baryon of interest plus a continuum model accounting for the contribution of all excited states. The continuum model approximates the excited-state contributions in terms of the perturbative evaluation of the invariant functions, starting from an effective threshold; it is thus a crude model. In order to extract the spectral parameters of the pole by matching the sum rules, one should work in a region of Borel mass where the pole contribution dominates the phenomenological side. This usually sets an upper bound in the Borel mass space, beyond which the excited-state contribution dominates. On the other hand, the truncated OPE must be sufficiently convergent as to accurately describe the true OPE. This, in practice, sets a lower limit in the Borel mass space, beyond which higher order terms not present in the truncated OPE may be significant and important.

Therefore, we expect a sum rule to work if the two sides of the sum rule match in a *valid* window in Borel mass space where the pole contribution to the phenomenological side dominates and the higher order OPE terms are under control. The relative reliability of a sum rule is thus determined by the quality of overlap of the two sides, the size of valid Borel window, and the relative contribution of the excited states in the valid Borel window. Such criteria have been advocated in Refs. [6,7] and used in various sum rule calculations [6–11]. Here, we adopt these criteria to determine the relative reliability of a sum rule.

Let us now focus on the phenomenological representation of $\Pi(q)$, which can be obtained by inserting a complete set of eigenstates with the quantum numbers of the interpolating field. Since η_B couples to not only the positive-parity baryon states but also the negative-parity baryon states [5], $\Pi(q)$ can be expressed as

$$\Pi^{\text{phen}}(q) = -\lambda_0^2 \frac{q_\mu \gamma^\mu + M_0}{q^2 - M_0^2} - \sum_{i \neq 0} (\lambda_i^+)^2 \frac{q_\mu \gamma^\mu + M_i^+}{q^2 - (M_i^+)^2} - \sum_i (\lambda_i^-)^2 \frac{q_\mu \gamma^\mu - M_i^-}{q^2 - (M_i^-)^2}, \quad (4)$$

in the discrete-state approximation (i.e., neglecting the widths of baryon states, which is usually assumed in the QCD sum-rule approach), where “0” denotes the ground state baryon of interest with mass M_0 , and “+”/“−” the positive/negative parity excited baryon state with mass M_i^+/M_i^- . Here we have omitted the infinitesimal as we are only concerned with large and space-like q^2 . The $\lambda_i^\pm$ ($\lambda_0 \equiv \lambda_0^+$) stands for the coupling strength of η_B to the physical baryon state :

$$\langle 0 | \eta_B | i, q, s, + \rangle = \lambda_i^+ U(q, s), \quad \langle 0 | \eta_B | i, q, s, - \rangle = \lambda_i^- \gamma_5 U(q, s), \quad (5)$$

with $U(q, s)$ the baryon Dirac spinor. After the Borel transform, one obtains from Eq. (4)

$$\Pi_s^{\text{phen}}(M^2) = \lambda_0^2 M_0 e^{-M_0^2/M^2} + \sum_{i \neq 0} (\lambda_i^+)^2 M_i^+ e^{-(M_i^+)^2/M^2} - \sum_i (\lambda_i^-)^2 M_i^- e^{-(M_i^-)^2/M^2}, \quad (6)$$

$$\Pi_q^{\text{phen}}(M^2) = \lambda_0^2 e^{-M_0^2/M^2} + \sum_{i \neq 0} (\lambda_i^+)^2 e^{-(M_i^+)^2/M^2} + \sum_i (\lambda_i^-)^2 e^{-(M_i^-)^2/M^2}, \quad (7)$$

where M is the Borel mass. We observe that the positive- and negative-parity excited-state contributions partially cancel each other in $\Pi_s^{\text{phen}}(M^2)$, but add up in $\Pi_q^{\text{phen}}(M^2)$. This indicates that the continuum contamination is significantly smaller in the sum rule for Π_s than in that for Π_q .

This feature attributes to chirality. If there were no spontaneous and explicit chiral symmetry breaking, the chirality of quarks could not be changed during propagation, and there would be chiral doubling in the baryon states and $\Pi_s^{\text{phen}}(M^2) = 0$. In nature, the chiral symmetry is violated by QCD vacuum and (small) current quark masses, which can cause quark chirality flipping and hence the shift of the positive-parity states relative to the negative-parity states. The two invariant functions Π_s and Π_q correspond to the chirality-changing and chirality-conserving parts of Π , respectively. The chirality-conserving part $\Pi_q^{\text{phen}}(M^2) \neq 0$ even if there were no chiral symmetry violation. This gives rise to extra background noises in the sum rule for Π_q . Thus, the QCD side of the sum rule for Π_s must be dominated by chiral-odd operators (e.g., $\bar{q}q$, $\bar{q}g_s\sigma \cdot Gq$) as the current quark masses are small compared to the baryon masses and the QCD side of the sum rule for Π_q must be dominated by chiral-even operators (e.g., 1 , G^2). We define “chiral-odd” sum rules as those where the chiral-odd operators dominate and “chiral-even” sum rules where the chiral-even operators dominate. Therefore, the chiral-odd sum rules are generally more reliable than the chiral-even sum rules.

As an explicit example, we consider the sum rules for the proton matrix element of $H = \langle p | \bar{u}u - \bar{d}d | p \rangle / 2M_N$, which can be obtained from the linear response of Π to a constant external isovector-scalar field [12,13]. In the formalism suggested in Ref. [14], the two sum rules given in Ref. [13] can be rewritten as:

$$\begin{aligned} & \frac{c_1}{32} a \left(M_N^2 M^2 - M^4 \right) L^{-4/9} - \frac{c_2}{48} m_0^2 a M_N^2 L^{-24/27} \\ & + \frac{c_3}{12} \chi a^2 M_N^2 L^{4/9} - \frac{c_3 \chi}{48} \left(1 + \frac{M_N^2}{M^2} \right) m_0^2 a^2 L^{-2/27} \\ & = 2H \tilde{\chi}_N^2 M_N e^{-M_N^2/M^2} + \frac{c_1}{32} a \left(M_N^2 M^2 - M^4 \tilde{E}_1 \right) L^{-4/9} e^{-s_1/M^2}, \end{aligned} \quad (8)$$

$$\begin{aligned}
& -\frac{c_4}{16} (M_N^2 M^6 - 3M^8) L^{-8/9} + \frac{c_4}{16} \chi a (M_N^2 M^4 - 2M^6) \\
& -\frac{3c_5}{16} \chi m_0^2 a (M_N^2 M^2 - M^4) L^{-14/27} - \frac{c_6}{24} a^2 M_N^2 \\
& = 2H \tilde{\lambda}_N^2 M_N^2 e^{-M_N^2/M^2} - \frac{c_4}{16} (M_N^2 M^6 \tilde{E}_2 - 3M^8 \tilde{E}_3) L^{-8/9} e^{-s_2/M^2} \\
& + \frac{c_4}{16} \chi a (M_N^2 M^4 \tilde{E}_1 - 2M^6 \tilde{E}_2) e^{-s_2/M^2} \\
& - \frac{3c_5}{16} \chi m_0^2 a (M_N^2 M^2 - M^4 \tilde{E}_1) L^{-14/27} e^{-s_2/M^2} , \tag{9}
\end{aligned}$$

where s_1 and s_2 denote the continuum thresholds, $c_1 = 1 + 10\beta + \beta^2$, $c_2 = 4 + 7\beta + \beta^2$, $c_3 = 1 - 2\beta + \beta^2$, $c_4 = 5 + 2\beta - 7\beta^2$, $c_5 = 1 - \beta^2$, $c_6 = 7 + 4\beta + \beta^2$, $\tilde{E}_1 = \frac{s_i}{M^2} + 1$, $\tilde{E}_2 = \frac{s_i^2}{2M^4} + \frac{s_i}{M^2} + 1$, $\tilde{E}_3 = \frac{s_i^3}{6M^6} + \frac{s_i^2}{2M^4} + \frac{s_i}{M^2} + 1$, $a = -(2\pi)^2 \langle \bar{q}q \rangle$, $m_0^2 = \langle \bar{q}g_s \sigma \cdot Gq \rangle / \langle \bar{q}q \rangle$, $L = \ln(M/\Lambda_{\text{QCD}})/\ln(\mu/\Lambda_{\text{QCD}})$, and $\lambda_N = (2\pi)^2 \lambda_N$. For definiteness, we use $\beta = -1.2$ [7] and $\chi = 2.0 \text{ GeV}^{-1}$ [12,13]. Here, we have put the continuum contributions on the right-hand sides of the sum rules.

One notices that sum rule (8) is the chiral-odd sum rule, which, according to the above discussion, is more reliable than the chiral-even sum rule (9). This has been emphasized in Refs. [12,13]. Here, we give a numerical demonstration. We take the experimental value for the nucleon mass M_N and extract H and the continuum thresholds from the sum rules. To this end, we sample the sum rules in a valid Borel region, where the continuum contribution is less than 50% of the total phenomenological (right-hand) side and the contribution of the highest order OPE term is less than 10% of the total QCD (left-hand) side. The fit of the two sides is measured by $\delta(M^2) = |\text{LHS} - \text{RHS}|^2$ averaged over 150 points evenly spaced within the valid Borel region.

We find that the two sides of the chiral-odd sum rule (8) overlap very well in a large valid Borel region. In Fig. 1, we have plotted the relative continuum and highest order OPE term contributions in the valid Borel window. In contrast, there is no valid Borel window for the chiral-even sum rule (9). This is mainly due to the large continuum contamination. To stress this point, we have displayed in Fig. 1 the relative continuum and highest order OPE term of the chiral-even sum rule within the valid Borel window of (8), with the use of H and the continuum threshold extracted from the chiral-odd sum rule. One can see that the continuum contribution is dominant in the chiral-even sum rule, making it impossible to isolate the ground state signal of interest in this sum rule. This may also be understood from the different behavior of the continuum model in the two sum rules. While the leading Borel mass dependence is M^4 in (8), it is M^8 in (9). So, the continuum contribution grows much more rapidly in (9) than in (8) as M^2 increases, implying much larger continuum contamination in (9) than in (8).

Thus, if the OPE is truncated at the order usually considered in the literature, the chiral-even sum rules are likely to fail to have a valid Borel window. To improve this situation, one may have to include many higher order OPE terms in order to compensate for the smaller upper bound in Borel mass space compared to the chiral-odd sum rule. However, one usually




FIG. 1. The relative continuum (upper) and highest order OPE term (lower) contributions in the valid Borel window for the chiral-odd sum rule of (8), with H and the continuum thresholds extracted from (8). The solid curves are for the chiral-odd sum rule of (8) and the dashed curves for the chiral-even sum rule of (9). Note that the relative continuum contribution in the chiral-even sum rule are greater than 100%. This results from partial cancelation between the first term and the rest of the right-hand side of (9).

does not have much control over the values of such terms, which makes it difficult to extract any useful information about the ground state from the chiral-even sum rules.

In the literature it is often found that both chiral-odd and chiral-even sum rules are used in extracting baryon properties. In some cases, certain chiral-even sum rules are identified as the most reliable sum rules. This largely relies on the so-called ratio method widely adopted in the literature. There, one chooses the continuum threshold (to be the same for two different sum rules) to make the ratio of two *different* sum rules as flat as possible as a function of the Borel mass.

However, it is worthwhile emphasizing that the ratio method has serious drawbacks. (1) The ratio method does not check the validity and reliability of each individual sum rule. It may happen that individual sum rules are not valid (i.e., there does not exist a valid Borel window) while their ratio is flat as function of the Borel mass. (2) The ratio method

may lead to misleading results as the continuum contributions are not monitored in the ratio. It is obvious that the ratio is always perfectly flat for large Borel mass, which is trivial because the continuum contribution is dominant in a sum rule and modeled by the perturbative evaluation of the OPE. There, however, one should not expect to get any reliable information about the ground state. (3) The continuum threshold is a phenomenological parameter introduced to parameterize the spectral function. Hence, one should treat the continuum threshold as an independent parameter (just as the other parameters, masses, coupling strength, ect.) to be extracted from the sum rule. Fixing the continuum threshold will introduce artificial bias to the extracted baryon properties of interest.

To summarize, we have pointed out that chirality plays an important role in determining the reliability of baryon QCD sum rules. In particular, the contributions of positive- and negative-parity excited baryon states partially cancel each other in the chiral-odd sum rules, but add up in the chiral-even sum rules. Consequently, the continuum contamination is much smaller in the former than in the latter. Moreover, there are other (relatively) large uncertainties in the chiral-even sum rules arising from large perturbative corrections to the Wilson coefficients in the OPE, the factorization assumption for chiral-even operators, and the onset of nonfactorizable operators in relatively low dimension [7]. Thus, the chiral-odd sum rules are more reliable than the chiral-even sum rules. The chiral-odd sum rules can be easily identified for a given problem and should be used in extracting various baryon properties. On the other hand, without further improvement, the chiral-even sum rules are likely to fail in most cases and hence should be disregarded. Any misuse of the chiral-even sum rules could lead to inconsistent and misleading results.

REFERENCES

- [1] M. A. Shifman, A. I. Vainshtein, and V. I. Zakharov, Nucl. Phys. **B147**, 385 (1979); **B147**, 448 (1979).
- [2] L. J. Reinders, H. Rubinstein, and S. Yazaki, Phys. Repts. **127**, 1 (1985), and references therein.
- [3] There have been many works studying baryon properties in the QCD sum-rule approach. These works can be retrieved easily from the SLAC SPIRES.
- [4] B. L. Ioffe, Nucl. Phys. **B188**, 317 (1981); **B191**, 591(E) (1981).
- [5] Y. Chung, H. G. Dosch, M. Kremer, and D. Schall, Nucl. Phys. **B197**, 55 (1982); D. Espriu, P. Pascual, and R. Tarrach, Nucl. Phys. **B214**, 285 (1983).
- [6] D. B. Leinweber, Ann. Phys. (N.Y.) **198**, 203 (1990).
- [7] D. B. Leinweber, University of Washington Report No. UW-DOE/ER/40427-17-N95, nucl-th/9510051 (1995), to appear in Ann. Phys. (N.Y.) (1997).
- [8] X. Jin and D. B. Leinweber, Phys. Rev. C **52**, 3344 (1995).
- [9] M. J. Iqbal, X. Jin, and D. B. Leinweber, Phys. Lett. **B386**, 55 (1996).
- [10] R. J. Furnstahl, X. Jin, and D. B. Leinweber, Phys. Lett. **B387**, 253 (1996).
- [11] F. X. Lee, D. B. Leinweber, and X. Jin, TRIUMF Report No. TRI-PP-96-44, nucl-th/9611011 (1996), to appear in Phys. Rev. D (1997).
- [12] X. Jin, M. Nielsen, and J. Pasupathy, Phys. Rev. D **51**, 3688 (1995).
- [13] X. Jin, Phys. Rev. D **52**, 2964 (1995).
- [14] X. Jin, hep-ph/9608303 (1996), Phys. Rev. D **55**, (1997) in press.