

INTRODUCTION TO THE NRQCD FACTORIZATION APPROACH TO HEAVY QUARKONIUM ^a

E. Braaten

Department of Physics, Ohio State University, Columbus OH 43210 USA

I present an introduction to the NRQCD factorization method for calculating annihilation rates and inclusive production rates for heavy quarkonium. Using this method, annihilation decay rates and sufficiently inclusive cross sections are factored into long-distance NRQCD matrix elements and perturbative short-distance coefficients. I derive the velocity-scaling rules that are used to estimate the magnitudes of the nonperturbative NRQCD matrix elements and I describe perturbative matching methods for calculating their short-distance coefficients. Some simple applications for which NRQCD factorization methods have dramatic implications are discussed.

1 Introduction

The NRQCD factorization approach is a systematic framework for analyzing annihilation decay rates and sufficiently inclusive production cross sections for heavy quarkonium ¹. This method separates the effects of short distances that are comparable to or smaller than the inverse of the heavy quark mass from the effects of longer distance scales. Short-distance effects are calculated using perturbative QCD and long-distance effects are described by matrix elements in an effective field theory called nonrelativistic QCD (NRQCD). This paper is an introduction to the NRQCD factorization approach. In Section 2, the effective lagrangian for NRQCD is constructed, the velocity-scaling rules that are used to estimate the magnitude of NRQCD matrix elements are derived, and perturbative matching methods for calculating short-distance coefficients are described. The application of NRQCD factorization methods to annihilation rates and to inclusive production rates are discussed in Sections 3 and 4.

2 Heavy Quarkonium and NRQCD

2.1 Scales in Quarkonium Physics

Heavy quarkonium is a meson containing a heavy quark and its antiquark. There are a number of different energy scales that play an important role in the quarkonium physics. While the many scales make the physics complex, they also make it interesting. The scales of quarkonium include the mass

^abased on a series of lectures presented at the Third International Workshop on Particle Physics Phenomenology in Taipei, November 1996.

Table 1: Quarkonium energy scales

	cc	bb	tt
M	1.5 GeV	4.7 GeV	180 GeV
M v	0.9 GeV	1.5 GeV	16 GeV
M v ²	0.5 GeV	0.5 GeV	1.5 GeV

M of the heavy quark, its typical momentum M v, and its typical kinetic energy M v². The quark mass M sets the total energy scale for annihilation decays and the scale of the kinematic threshold for onium production. The inverse of the typical momentum M v is the length scale for the size of the onium state. The typical kinetic energy M v² is the scale of the splittings between radialexcitations and between orbital-angular-momentum excitations in the onium spectrum. For both charm onium and bottom onium, the splittings between the two lowest ³S₁ states (J= and ⁰ for charm onium, and ⁰ for bottom onium) are approximately 600 M eV. The splittings between the lowest ³S₁ and ³P_J states (J= and _{cJ} for charm onium, and _{bJ} for bottom onium) are approximately 400 M eV. We take 500 M eV as an estimate for the scale M v². In Table 1, we give our estimates for the scales M, M v, and M v² for charm onium and bottom onium. The estimate for M is half the mass of the lowest energy level of quarkonium. The estimate for the scale M v is the geometric mean of the estimates for M and M v². From Table 1, we see that v² is approximately 1/3 for charm onium and 1/9 for bottom onium. These values are small enough to justify a theoretical approach based on an expansion in powers of v.

Another important energy scale in quarkonium physics is Λ_{QCD} , the scale of nonperturbative effects involving gluons and light quarks. The potential energy V(R) between a quark and antiquark separated by a distance R varies from Coulombic at sufficiently short distances,

$$V(R) \approx \frac{4}{3} \frac{\alpha_s(l=R)}{R} \quad \text{as } R \rightarrow 0; \quad (1)$$

to linear at sufficiently long distances:

$$V(R) \approx \sigma R \quad \text{as } R \rightarrow \infty; \quad (2)$$

where empirically $\sigma \approx 450 \text{ MeV}$. Since Λ_{QCD} is independent of M, it must be proportional to Λ_{QCD} .

To clarify the relation between the scale Λ_{QCD} and the other scales in quarkonium physics, we consider two limiting cases. The first case is a heavy quark whose mass M is large enough that the onium wavefunction is dominated by the Coulombic term in the potential. The size of a bound state is determined by a balance between the kinetic energy and the potential energy, so we must have

$$M v^2 \sim \frac{4}{3} \frac{s(1=R)}{R} : \quad (3)$$

The size of the bound state is comparable to the inverse of the typical momentum $M v$ of the heavy quark in the bound state. Setting $R \sim 1/(M v)$ in (3), we obtain

$$v \sim s(M v) : \quad (4)$$

This equation can be solved self-consistently for v as a function of M . If M is sufficiently large, the resulting value of $M v^2$ is much greater than Λ_{QCD} . For the top quark with mass $M = 180 \text{ GeV}$, we obtain $v \sim 0.18$. An alternative estimate for v , which is closer in spirit to the estimates used for charm onium and bottom onium in Table 1, is obtained by taking $M v^2$ to be the splitting between the ground state and the first excited state in the potential (1). This splitting is approximately 1.5 GeV , which corresponds to $v \sim 0.09$. This value for $M v^2$ and the corresponding estimate for the scale $M v$ are included in Table 1. These would be the appropriate scales for toponium if the top quark were stable enough to form bound states.

The second limiting case is a heavy quark whose mass M is small enough that the wavefunction is dominated by the linear term (2) in the potential, but still large enough that the onium is nonrelativistic. The balance between the kinetic and potential energies requires

$$M v^2 \sim \Lambda^2 R : \quad (5)$$

Setting $R \sim 1/(M v)$, we find that $M v^3 \sim \Lambda^2$. Identifying Λ with the scale Λ_{QCD} , we find that the typical velocity of the heavy quark is such that this scale is intermediate between the scales $M v$ and $M v^2$. If the Coulombic and linear regions of the potential were both equally important, then the relations (3) and (5) would be satisfied simultaneously. The fact that these relations are compatible suggests that the scaling relation (4) might be applicable even for rather low values of M .

It is a remarkable coincidence of quarkonium physics that the scales $M v^2$ are almost identical for charm onium and bottom onium. The fact that this scale is so insensitive to the value of M suggests that Λ_{QCD} should be identified with the scale $M v^2$ for quark masses in this range. This identification is supported by the numerical value of the scale $M v^2$ given in Table 1.

Table 2: Value of the QCD coupling constant at the characteristic momentum scales for heavy quarkonium

	cc	bb	tt
$\alpha_s(M)$	0.35	0.22	0.11
$\alpha_s(Mv)$	0.52	0.35	0.16
$\alpha_s(Mv^2)$	1	1	0.35

If any of the scales M , Mv , and Mv^2 is large enough compared to Λ_{QCD} , then the effects of that scale can be calculated using QCD perturbation theory. The values of the running coupling constant of QCD at the scales M , Mv , and Mv^2 are given in Table 2 for charmonium, bottomonium, and toponium. All three scales are perturbative in the case of toponium. For charmonium and bottomonium, $\alpha_s(M)$ is small enough to justify perturbation theory at the scale M . The coupling constant $\alpha_s(Mv)$ is also small enough that perturbation theory seems reasonable. However, potential model calculations imply that the wavefunctions of charmonium and bottomonium have significant support in the linear region of the potential, and this suggests that nonperturbative effects may be significant at the scale Mv . As for the scale Mv^2 , it is hopelessly nonperturbative for charmonium and bottomonium.

In addition to the scales M , Mv , Mv^2 , and Λ_{QCD} , there are also kinematic energy scales that can play an important role in quarkonium physics. For example, in the production of quarkonium, the total center-of-mass energy \sqrt{s} and the transverse momentum p_T of the onium can be important. Unravelling the effects of the various energy scales is essential in order to understand quarkonium physics. This is particularly important for charmonium and bottomonium, because the coupling constant α_s runs rather dramatically with the momentum at scales of order M and smaller.

The NRQCD factorization approach is based on separating short-distance effects involving momenta of order M or larger from those effects that involve the smaller momentum scales Mv , Mv^2 , and Λ_{QCD} . The scale M is assumed to be perturbative, so that short-distance effects can be calculated using perturbation expansions in $\alpha_s(M)$. No assumption is made about the validity of perturbation theory at the scale Mv . Instead, we exploit the fact that in a nonrelativistic bound state, the typical velocity v provides a small expansion parameter.

2.2 Integrating out Relativistic Effects

One way to separate the effects of the momentum scale M from the lower momentum scales in a field theory is to integrate out all modes with momenta greater than some cutoff that is much less than M . The result of this renormalization group transformation is a Wilsonian effective field theory that describes the modes with momenta smaller than M . All effects of the scale M are encoded in the parameters of the effective field theory. In our case, the original field theory is QCD with a heavy quark. It is described by the lagrangian

$$L_{\text{QCD}} = L_{\text{light}} + \overline{(i \not{D} - m_Q)} ; \quad (6)$$

where L_{light} is the lagrangian that describe gluons and light quarks. The mass parameter m_Q of the heavy quark can be identified with M . It is implicit in the Lorentz-invariant lagrangian (6) that the cutoff is much larger than M . Integrating out the momentum scale M is equivalent to lowering the cutoff to a value lower than M . We will argue that if Λ is in the range $M \ll \Lambda \ll M^2$, the resulting effective field theory can still be described by a local lagrangian. In other words, the effects of modes with momenta of order M can be reproduced by local interactions among the lower momentum modes.

Suppose the onium is in a virtual state that includes a quark with relativistic momentum of order M . Then that state is off its energy-shell by an amount of order M , which is much larger than the scale M^2 of the splittings between onium energy levels. By the uncertainty principle, the lifetime of that highly virtual state is less than or of order $1/M$. In that short time, fields can propagate only over distances of order $1/M$ that are pointlike on the scale $1/(M v)$ of onium structure. Thus the effects of virtual states that are excluded by a momentum cutoff in the range $M \ll \Lambda \ll M^2$ can be reproduced by local interactions involving low momentum modes.

The above argument applies equally well to virtual states that contain a light parton with momentum of order M in addition to the $Q\bar{Q}$ pair. However, it does not apply to virtual states obtained by the annihilation of the $Q\bar{Q}$ pair. Such states can contain light partons with momenta of order M without being far off the energy shell. A momentum cutoff satisfying $\Lambda \ll M$ excludes these states, but their effects cannot be reproduced in detail by local interactions. For example, the annihilation decay of the onium produces light hadrons, some of which must have momenta of order M and therefore parton constituents with momenta of order M . With such modes excluded by the cutoff, we cannot hope to describe the annihilation decays accurately. Nevertheless, as we shall see in Section 3, the total annihilation width of an onium state can be described accurately. Thus the momentum cutoff can be extended to light partons at

the expense of a restriction on the physical observables that can be described within the effective theory.

With a momentum cutoff that excludes relativistic Q and \bar{Q} states, it is convenient to describe the heavy quark and antiquark by separate 2-component Pauli fields ψ and $\bar{\psi}$, rather than by a single 4-component Dirac field ψ . If we simply substitute $\psi = \begin{pmatrix} \psi \\ 0 \end{pmatrix}$ into the Lagrangian (6), we obtain off-diagonal terms that couple ψ to $\bar{\psi}$ and $\bar{\psi}$ to ψ , allowing the creation and annihilation of $Q\bar{Q}$ pairs. We will argue that terms that change the numbers of heavy quarks and antiquarks can be removed from the Lagrangian and compensated by terms that conserve the numbers of Q 's and \bar{Q} 's. This is obvious for terms that create $Q\bar{Q}$ pairs, because a virtual state containing an additional $Q\bar{Q}$ must be on its energy shell by an amount of order M . As mentioned above, this is not completely true for terms that allow the $Q\bar{Q}$ pair in the onium to annihilate. The effects of states consisting of gluons and light quark pairs that are produced by $Q\bar{Q}$ annihilation cannot be reproduced in detail by local interactions. However, the effects of these states on sufficiently inclusive observables can be described accurately. Thus, with this restriction on physical observables, terms in the effective Lagrangian that change the numbers of Q 's and \bar{Q} 's can be eliminated from the effective Lagrangian.

For the Dirac term of the Lagrangian (6), the decoupling of the fields ψ and $\bar{\psi}$ can be accomplished by a unitary transformation called the Foldy-Wouthuysen-Tani transformation. For the case of a background gauge field, it is straightforward to construct the transformation that diagonalizes the Lagrangian to any desired order in the heavy quark velocity. The simplest form of this transformation in the Dirac representation is

$$U = \exp(i\alpha \gamma_5) : \quad (7)$$

After this transformation, the heavy-quark term in the Lagrangian (6) can be approximated by

$$\bar{\psi} \begin{pmatrix} m_Q + iD_0 + D^2 = 2m_Q & 0 \\ 0 & m_Q + iD_0 - D^2 = 2m_Q \end{pmatrix} \psi : \quad (8)$$

If we take D to scale like Mv , the corrections to the entries in the matrix scale like Mv^4 .

For the effective Lagrangian with momentum cutoff Λ , the elimination of terms that change the numbers of Q 's and \bar{Q} 's is more complicated than simply applying a unitary transformation. In addition to the terms that are quadratic in ψ and $\bar{\psi}$, the effective Lagrangian also includes terms that are quartic and

higher in the heavy quark fields. A further complication is that gluon interactions modify the coefficients of the terms produced by the Foldy-Wouthuysen-Tani transformation. Nevertheless, by the general arguments presented above, one can describe the low-energy $Q\bar{Q}$ sector of QCD by an effective field theory in which the numbers of heavy quarks and antiquarks are strictly conserved.

2.3 Effective Field Theory

One could in principle construct a nonrelativistic effective Lagrangian that describes the low-energy $Q\bar{Q}$ sector of QCD by starting with the Lagrangian (6) and carrying out the sequence of two transformations described in Section 2.2. The first is a renormalization group transformation that removes modes with momenta greater than Λ . The second is a transformation that removes interactions that change the numbers of heavy quarks and antiquarks. Both of these steps would be extremely complicated to carry out in practice. Fortunately, there is an alternative to the explicit construction of the effective Lagrangian and that is to use the strategy of "effective field theory".² In this approach, the construction of the effective Lagrangian proceeds through the following steps:

1. Identify the fields that are required to describe the low-energy excitations of the theory.
2. Identify the symmetries that one could maintain in the effective theory by using a suitable cutoff and making appropriate field redefinitions.
3. Specify the accuracy to which low energy observables in the original theory should be reproduced by the effective theory.
4. Write down the most general effective Lagrangian that is consistent with the symmetries, including all terms that are required to reproduce the physics to the specified level of accuracy.
5. Determine the coefficients of those terms by matching low-energy observables of the effective theory with those of the full theory.

The effective field theory that is obtained by applying the above strategy to the low-energy $Q\bar{Q}$ sector of QCD is called nonrelativistic QCD (NRQCD).³ The fields that are required to describe the low energy degrees of freedom are the heavy quark and antiquark fields Q and \bar{Q} , the SU(3) gauge fields A_μ^a , and the Dirac fields for the light quarks. The symmetries of NRQCD are the following:

SU(3) gauge symmetry. This local symmetry requires that the gluon fields enter into the effective lagrangian only through the gauge-covariant derivatives D_0 and D and the QCD field strengths E and B .

rotational symmetry. A nonrelativistic description of the heavy quark necessarily breaks the Lorentz symmetry of QCD down to its rotational subgroup.

charge conjugation and parity. These discrete symmetries of QCD are also symmetries of the effective theory. The charge conjugation transformations of the heavy quark and antiquark fields are

$$Q \rightarrow i(\gamma_2)^T \bar{Q}^c; \quad \bar{Q} \rightarrow i(\gamma_2)^T Q^c; \quad (9)$$

The parity transformations are

$$Q(t; \mathbf{r}) \rightarrow \gamma_0 Q(t; -\mathbf{r}); \quad \bar{Q}(t; \mathbf{r}) \rightarrow \gamma_0 \bar{Q}(t; -\mathbf{r}); \quad (10)$$

heavy-quark phase symmetry. This symmetry guarantees the separate conservation of the number of heavy quarks and antiquarks. Its action on the fields is

$$Q \rightarrow e^{i\theta} Q; \quad \bar{Q} \rightarrow e^{-i\theta} \bar{Q}; \quad (11)$$

Having identified the symmetries of NRQCD, we can write down the most general effective lagrangian that is consistent with these symmetries. It has the form

$$L_{\text{NRQCD}} = L_{\text{light}} + \psi \bar{\psi} iD_0 + \frac{D^2}{2M} \psi \bar{\psi} + \psi \bar{\psi} iD_0 \frac{D^2}{2M} + L; \quad (12)$$

where L_{light} is the usual lagrangian that describes gluons and light quarks. The desired level of accuracy is specified by the order in v with which the onium energy levels must be reproduced by the effective theory. The heavy quark terms that are shown explicitly in (12) are those that are required to calculate the energy levels up to errors of order M^{-4} . The term L in (12) includes the correction terms that must be added to decrease the errors to order M^{-6} or smaller.

The invariance of physical quantities under field redefinitions can be exploited to eliminate some terms in the NRQCD lagrangian. Gauge invariance requires that the gluon fields in L appear only through the covariant derivatives D_0 and D . However, a redefinition of the field can be used to eliminate terms in which D_0 acts on ψ , and similarly for $\bar{\psi}$. Thus field redefinitions

can be used to eliminate all occurrences of D_0 in \mathcal{L} except in the combination $[D_0; D] = igE$. Because of these field redefinitions, NRQCD will not reproduce the low-energy behavior of the Green's functions of QCD. It will only agree with full QCD for on-shell physical quantities.

The minimal form of NRQCD is obtained by setting $L = 0$ in (12). It contains two parameters, the heavy-quark mass parameter M and the gauge coupling constant g . These parameters can be tuned as functions of the QCD coupling constant g_s , the heavy-quark mass parameter m_Q , and the ultraviolet cutoff of NRQCD so that the splittings between the onium energy levels are reproduced up to errors of order $M v^4$. Since the energy splittings between radial excitations (such as $J=0$ and $0'$) and between orbital-angular-momentum excitations (such as $J=0$ and 0_J) scale like $M v^2$, these are reproduced up to errors of relative order v^2 . Spin splittings in heavy quarkonium, such as the splitting between the lowest 1S_0 and 3S_1 states (1S_0 and $J=0$ for charmonium), scale like $M v^4$. These splittings vanish in minimal NRQCD due to the following symmetry:

heavy-quark spin symmetry. Under this symmetry, the two spin components of the heavy quark and the two spin components of the antiquark are mixed by independent unitary transformations:

$$U \psi; \quad U \bar{\psi}; \quad (13)$$

where U and V are $SU(2)$ matrices. This is only an approximate symmetry of the complete NRQCD Lagrangian, holding up to corrections of relative order v^2 .

If we wish to reduce the errors in the quarkonium energy levels to smaller than order $M v^4$, it is necessary to add additional terms \mathcal{L} to the Lagrangian in (12). Using the velocity-scaling rules that are discussed in Section 2.4, it can be shown that the terms that are necessary and sufficient to reduce the errors to order $M v^6$ are

$$\begin{aligned} \mathcal{L} = & \frac{c_1}{8M^3} \psi (D^2)^2 \psi + \frac{c_2}{8M^2} \psi (D \cdot gE - gE \cdot D) \psi \\ & + \frac{c_3}{8M^2} \psi (iD \cdot gE - gE \cdot iD) \psi + \frac{c_4}{2M} \psi (gB \cdot \sigma) \psi \\ & + \text{charge conjugate terms}; \end{aligned} \quad (14)$$

where c_1, c_2, c_3 , and c_4 are dimensionless coefficients. We will refer to the terms in (14) as the v^2 -improvement terms in the NRQCD Lagrangian. The two terms in (14) that contain the Pauli matrix σ break the spin symmetry

of minimal NRQCD. They give spin splittings that scale like $M v^4$ and are accurate up to errors of relative order v^2 . Splittings between radial excitations and splittings between orbital-angular-momentum excitations are reproduced up to errors of relative order v^4 .

2.4 Velocity-scaling rules

The relative importance of the terms in the NRQCD Lagrangian can be deduced from the self-consistency of the quantum field equations for minimal NRQCD and from the basic qualitative features of quarkonium⁴. The results of this analysis are summarized by the velocity-scaling rules in Table 3. The magnitude of a matrix element of a local gauge-invariant operator between quarkonium states can be estimated by multiplying the appropriate factors from Table 3. The scaling with M follows simply from dimensional analysis, so the nontrivial content of Table 3 is the scaling with v .

The first few lines in Table 3 can be derived very easily. The expectation value of the number operator $\int d^3x \psi^\dagger \psi$ in a quarkonium state $|H\rangle$ is very close to 1:

$$\int d^3x \psi^\dagger \psi |H\rangle = 1: \quad (15)$$

We have normalized the quarkonium state so that $\langle H | H \rangle = 1$. From the fact that a quarkonium state can be localized to within a region $l = (M v)^{-3}$, we conclude that ψ must scale like $(M v)^{3/2}$. The expectation value of the kinetic energy term in the NRQCD Hamiltonian scales like $M v^2$:

$$\int d^3x \psi^\dagger (-D^2) \psi |H\rangle = M v^2: \quad (16)$$

This implies that D must scale like $M v$. The fact that D_0 scales like $M v^2$ when acting on ψ then follows immediately from the field equation for ψ :

$$iD_0 \psi - \frac{D^2 \psi}{2M} = 0: \quad (17)$$

The estimates for gA_0 and gA in Table 3 are specific to Coulomb gauge, which is defined by $\nabla \cdot A = 0$. As shown below, the field equations in this gauge indicate that the effects of the vector potential A are suppressed by a factor of v relative to the scalar potential A_0 . The dominant terms in the field equations for ψ and A_0 are therefore

$$i\partial_0 \psi - gA_0 + \frac{r^2 \psi}{2M} = 0; \quad (18)$$

$$r^2 gA_0 + g^2 \psi = 0: \quad (19)$$

Table 3: Estimates of the magnitudes of NRQCD operators for matrix elements between heavy-quarkonium states.

Operator	Estimate
	$(M v)^{3=2}$
	$(M v)^{3=2}$
D_0 (acting on ψ or $\bar{\psi}$)	$M v^2$
D	$M v$
gE	$M^2 v^3$
gB	$M^2 v^4$
gA_0 (in Coulomb gauge)	$M v^2$
gA (in Coulomb gauge)	$M v^3$

In (18), the balance between the kinetic energy and the potential energy represented by the A_0 term requires that gA_0 scale like $M v^2$. On the other hand, assuming that a gradient acting on A_0 scales like $M v$, (19) requires that gA_0 scale like $g^2 M v$. These two estimates are consistent if the effective coupling constant $g_s = g^2 = 4$ at the scale $M v$ scales like v . This is identical to the naive estimate (4) that followed from balancing the kinetic energy and the Coulomb term in the potential energy. Since this scaling relation follows simply from the consistency of the field equations, it applies to charmonium and bottomonium even though perturbation theory at the scale $M v$ is of questionable validity. The neglect of terms involving A in the field equations (18) and (19) is justified by the field equation for A , for which the dominant terms are

$$\partial_0^2 r^2 gA - gA_0 r gA_0 - \frac{g^2}{M} v r = 0: \quad (20)$$

The last two terms in (20) scale like $M^3 v^5$ and $g^2 M v^4$, respectively, and they are comparable if g^2 scales like v . Assuming that a gradient acting on A scales like $M v$, we obtain the estimate $M v^3$ for gA in Table 3.

Using the estimates for gA_0 and gA in Coulomb gauge, we can obtain estimates for the field strengths. In Coulomb gauge, the dominant term in the chromoelectric field strength E is $r A_0$, and the resulting estimate for gE is $M^2 v^3$. The dominant term in the chromomagnetic field strength B is $r A$, which leads to the estimate $M^2 v^4$ for gB . These estimates for gE and gB , although derived in Coulomb gauge, hold in general for matrix elements of gauge-invariant operators.

According to the velocity-scaling rules in Table 3, the terms in the Lagrangian density in (12) for minimal NRQCD scale like $M^4 v^5$. Multiplying

by a volume factor of $1=(Mv)^3$, we find that quarkonium energies scale like Mv^2 . Each of the terms in L given in (14) scales like M^4v^7 and therefore contributes to onium energies at order Mv^4 . All other terms that can be added to the NRQCD Lagrangian give contributions of order Mv^6 or smaller.

The validity of the velocity-scaling relations has been demonstrated convincingly by nonperturbative calculations of the bottomonium and charmonium spectrum using Monte Carlo simulations of lattice NRQCD⁵. The two parameters of minimal NRQCD can be tuned to give the spin-averaged spectrum to an accuracy of about 30% for charmonium and about 10% for bottomonium. When the v^2 -improvement terms are included, the errors decrease to about 10% for charmonium and to about 1% for bottomonium. These terms also give spin splittings that are accurate to about 30% for charmonium and to about 10% for bottomonium.

2.5 Fock state expansion

The simplest intuitive picture of quarkonium is that it is a bound state consisting of a Q and \bar{Q} with very little probability of containing additional gluons or $q\bar{q}$ pairs. This simple picture is in fact realized in Coulomb gauge. Using the velocity-scaling rules of section 2.4, one can quantify the probabilities of Fock states containing additional gluons by determining how they scale with v .

The Coulomb gauge ($\nabla \cdot \mathbf{A} = 0$) is a physical gauge with no negative norm states, a necessary condition for a sensible Fock space. In this gauge, the scalar potential A_0 does not propagate. Dynamical gluons are created and destroyed by the vector potential \mathbf{A} . In Coulomb gauge, the Lagrangian (12) can be reorganized as an expansion in powers of v . The powers of v can be made explicit by rescaling the space-time coordinates \mathbf{r} and t by $1=(Mv)$ and $1=(Mv^2)$, respectively, rescaling the elds \mathbf{A} and \mathbf{E} by $(Mv)^{3=2}$, and rescaling the elds A_0 and \mathbf{A} by $Mv^{3=2}$. The terms in the NRQCD Lagrangian that are of order v^0 after such a rescaling are

$$L_0 = L_{\text{light}} + \int d^3r \left[\frac{1}{2M} \psi^\dagger \nabla^2 \psi + \frac{1}{2M} \psi^\dagger \nabla^2 \bar{\psi} \right] + \int d^3r \left[\frac{1}{2} \mathbf{E}^2 + \frac{1}{2} \mathbf{B}^2 \right] : \quad (21)$$

This Lagrangian in Coulomb gauge can be used to calculate quarkonium energy levels to the same accuracy as the gauge-invariant Lagrangian of minimal NRQCD. The terms in the NRQCD Lagrangian that are of order v after the rescaling are

$$L_1 = \int d^3r \left[\frac{1}{M} \psi^\dagger (\mathbf{A} \cdot \nabla) \psi + \frac{C_4}{2M} \psi^\dagger (\nabla \cdot \mathbf{A}) \psi \right] + \text{charge conjugate terms} : \quad (22)$$

At first order in perturbation theory, these terms give transitions from $Q\bar{Q}$ Fock states to states that contain a dynamical gluon. The expectation values of the terms in (22) between $\mathcal{D}\bar{Q}i$ Fock states vanish, so L_1 first contributes to quarkonium energy levels at second order in perturbation theory, giving shifts of order Mv^4 . The terms in the Coulomb-gauge Lagrangian that are of order v^2 after rescaling are

$$\begin{aligned}
L_2 = & \frac{1}{2M} \int d^3x (gA_0)^2 + \frac{C_1}{8M^3} \int d^3x (r^2)^2 \\
& + \frac{C_2}{8M^2} \int d^3x (r^2 gA_0) + \frac{C_3}{4M^2} \int d^3x (r gA_0) \cdot r \\
& + \frac{C_4}{2M} \int d^3x (igA_0 \cdot gA) + \text{charge conjugate terms} \quad (23)
\end{aligned}$$

The first term comes from expanding out the covariant derivative D in (12), while the last four terms in (23) come from the v^2 -improvement terms in (14). Energy levels calculated in Coulomb gauge using the Lagrangian $L_0 + L_1 + L_2$ will differ only at order Mv^6 from the energy levels calculated using the gauge-invariant NRQCD Lagrangian (12) with L given by (14).

We now consider the Fock state expansion in Coulomb gauge for a quarkonium state $\mathcal{H}i$. It has the schematic form

$$\mathcal{H}i = \int d^3x \frac{H}{Q\bar{Q}} \mathcal{D}\bar{Q}i + \int d^3x \frac{H}{Q\bar{Q}g} \mathcal{D}\bar{Q}gi + \dots; \quad (24)$$

where spin and color indices and momentum arguments have all been suppressed. The dominant Fock state $\mathcal{D}\bar{Q}i$ consists of a Q and \bar{Q} in a color-singlet state with definite angular momentum quantum numbers $^{2S+1}L_J$. The higher Fock states, such as $\mathcal{D}\bar{Q}gi$, include dynamical gluons or light $q\bar{q}$ pairs. Since the Lagrangian L_0 in (21) does not include any terms that couple A to q , the probabilities for higher Fock states are suppressed by powers of v .

The $\mathcal{D}\bar{Q}gi$ states with the highest probabilities are those that couple to the dominant $\mathcal{D}\bar{Q}i$ state via the Lagrangian L_1 in (22). We first consider the term $\int d^3x (igA_0 \cdot r)$. We refer to a transition that proceeds via this term or its charge conjugate as an electric transition. An electric transition from the dominant $\mathcal{D}\bar{Q}i$ Fock state produces $\mathcal{D}\bar{Q}gi$ states for which the angular momentum quantum numbers of the $Q\bar{Q}$ pair satisfy the selection rules $L = 1$ and $S = 0$. The simplest way to determine the probabilities of these Fock states is to use the fact that a second-order perturbation in L_1 changes the mass of the onium state by an amount of order Mv^4 . This mass shift can be expressed as the product of the energy E of the virtual $\mathcal{D}\bar{Q}gi$ state multiplied by its probability P . If the energy of the dynamical gluon is of order Mv , then $E = Mv$ and we find that $P = v^3$. If the gluon has energy of order Mv^2 or less,

then $E \sim M v^2$ and we obtain $P \sim v^2$. We conclude that $\bar{Q}Q$ states which satisfy the selection rules $L = 1$ and $S = 0$ are dominated by very soft dynamical gluons with momenta of order $M v^2$ or less and have probabilities of order v^2 .

We next consider the term $\bar{\psi}(x) g A(x)$ in (22). We refer to a transition that proceeds via this term or its charge conjugate as a magnetic transition. A magnetic transition from the dominant $\bar{Q}Q$ Fock state produces $\bar{Q}Q$ states that satisfy the selection rules $L = 0$ and $S = 1$. We can use the same argument as before to determine the probabilities of these Fock states, except that we must take into account the fact that the transition amplitude from the term $\bar{\psi}(x) g A(x)$ is weighted by the momentum of the gluon. If the gluon has energy of order $M v$, the mass shift from a second-order perturbation in L_1 is given correctly by the velocity-scaling rules to be of order $M v^4$. Since the virtual $\bar{Q}Q$ state has energy $E \sim M v$, we obtain a probability $P \sim v^3$. The contribution to the mass shift from a gluon with energy of order $M v^2$ is suppressed by a factor of v^2 from the transition amplitudes and is therefore of order $M v^6$. Taking $E \sim M v^2$, we obtain $P \sim v^4$. We conclude that $\bar{Q}Q$ states which satisfy the selection rules $L = 0$ and $S = 1$ are dominated by dynamical gluons with momenta of order $M v$ and have probabilities of order v^3 .

Similar arguments can be used to determine the magnitudes of the probabilities for other Fock states. Any such state can be reached by a sequence of electric transitions and zero or one magnetic transition. Electric transitions obey the selection rules $L = 1$ and $S = 0$, while magnetic transitions satisfy $L = 0$ and $S = 1$. Both electric and magnetic transitions change the color state of a color-singlet $\bar{Q}Q$ pair to color-octet, and they change the color state of a color-octet $\bar{Q}Q$ pair to either color-singlet or color-octet. The probability of a particular Fock state is determined by the color and angular-momentum quantum numbers of the $\bar{Q}Q$ pair in that state. If that Fock state can be reached from the dominant $\bar{Q}Q$ Fock state by a sequence of electric transitions, then its probability scales like v^{2E} . If it can be reached by a sequence of electric transitions and a magnetic transition, then its probability scales like v^{2E+3} .

2.6 Matching of NRQCD and QCD

The NRQCD Lagrangian contains adjustable parameters that must be tuned in order that its predictions for low-energy observables in the $\bar{Q}Q$ sector agree with those of QCD. In the minimal NRQCD Lagrangian, there are two parameters: g and M . In addition to these parameters, the definition of NRQCD

requires an ultraviolet cutoff to remove ultraviolet divergences. In the v^2 -improved lagrangian obtained by adding the terms in (14), there are 6 parameters, $g, M, c_1, c_2, c_3,$ and $c_4,$ in addition to the ultraviolet cutoff. The determination of the parameters in the NRQCD lagrangian is called matching.

One could in principle determine the N parameters in the NRQCD lagrangian by tuning them so that the masses of N states in NRQCD match the corresponding masses in full QCD. For example, the parameters g and M of minimal NRQCD could be determined by matching the mass splittings between $J=0$ and $J=1$ and between $J=0$ and $J=2$. Since the masses are sensitive to long-distance effects, they must be calculated nonperturbatively. The only reliable nonperturbative method that is currently available is Monte Carlo simulations of lattice NRQCD. While the masses in full QCD could in principle be computed nonperturbatively using lattice simulations, it is easier to take them directly from experiment. Using masses to tune the NRQCD parameters is an example of nonperturbative matching.

Nonperturbative matching would become increasingly difficult as we strive for higher accuracy by adding more improvement terms. The determination of the parameters in the v^2 -improved lagrangian would require the nonperturbative calculation of 6 masses as functions of 6 independent parameters. Fortunately, the asymptotic freedom of QCD provides an alternative, and that is perturbative matching. This matching procedure is based on the fact that QCD and NRQCD are equivalent except on distance scales of order $1/M$ where perturbative QCD is by assumption accurate. The procedure for perturbative matching is the following:

1. Use perturbative QCD to calculate scattering amplitudes between asymptotic $Q, \bar{Q},$ and gluon states with momenta much less than M as functions of s and m_Q and expand them in powers of k/m_Q .
2. Use perturbative NRQCD to calculate the same scattering amplitudes in terms of the parameters in the NRQCD lagrangian and expand them in powers of k/M .
3. Adjust the NRQCD parameters so that the scattering amplitudes match to the desired order in k/m_Q , which we take to be of order v .

It is essential to match scattering amplitudes or other physical observables rather than Green functions, because the construction of NRQCD involves field redefinitions. Such redefinitions can change the off-shell Green functions of the theory, but they leave on-shell physical observables unchanged.

In present calculations in lattice NRQCD, the parameters are determined by a combination of nonperturbative and perturbative matching. The coefficients

coefficients $c_1, c_2, c_3,$ and c_4 of the v^2 -improvement terms are generally determined by perturbative matching, while the parameters g and M are determined by the nonperturbative matching of masses in the onium spectrum. Perturbative matching calculations can be used to relate these parameters to the fundamental parameters of QCD. By combining these perturbative matching relations with lattice NRQCD calculations of the bottomonium spectrum, the QCD coupling constant α_s and the bottom quark mass m_b have been determined with high precision.⁶

The method of perturbative matching is somewhat paradoxical. We have assumed that M is large enough that perturbation theory is accurate at the scale M . We allow for the scale Mv to be small enough that perturbation theory is not reliable at that scale. If that is the case, perturbative calculations in NRQCD would never give accurate results for physical observables, since NRQCD only reproduces full QCD accurately at scales of order Mv or less. Nevertheless, a comparison of perturbative calculations in NRQCD and full QCD can be used to accurately determine the parameters in the NRQCD lagrangian. The reason for this is that the tuning of the parameters of NRQCD that makes this theory equivalent to QCD at momenta of order Mv or smaller also makes the perturbative approximations to these theories equivalent. Perturbation theory breaks down in precisely the same way for both theories, predicting among other things, the existence of asymptotic states consisting of isolated quarks and gluons. Since the parameters in the NRQCD lagrangian are sensitive only to momenta on the order of M where perturbative QCD is accurate, they can be correctly determined by matching perturbative calculations in QCD and NRQCD.

As an illustration of perturbative matching, we consider the simplest perturbative observable. This is the energy-momentum relation for the heavy quark, which is given by the location of the pole in the heavy-quark propagator. At tree level in full QCD, the energy-momentum relation is

$$E = \sqrt{m_Q^2 + p^2} = m_Q + \frac{p^2}{2m_Q} - \frac{p^4}{8m_Q^3} + \dots \quad (25)$$

At tree level in NRQCD, we can read off the energy-momentum relation from the lagrangian (12):

$$E = \frac{p^2}{2M} - c_4 \frac{p^4}{8M^3} + \dots \quad (26)$$

By matching the expressions (25) and (26) we find

$$M = m_Q; \quad c_4 = 1: \quad (27)$$

If the energy-momentum relations are computed to higher order in perturbation theory, the matching will give perturbative corrections to the results in (27). Since the parameters M_c and c_1 are sensitive only to short distances of order $1/m_Q$ or smaller, the corrections can be expressed as power series in $1/m_Q$.

3 Annihilation Decays of Heavy Quarkonium

3.1 Decay of ψ_c in the Color-Singlet Model

A simple intuitive picture of the annihilation decay of a quarkonium state is that it proceeds through the annihilation of the $Q\bar{Q}$ pair in the dominant Fock state into gluons and light $q\bar{q}$ pairs. These light partons ultimately hadronize into the observed final states that consist of light hadrons. The inclusive annihilation rate of the $Q\bar{Q}$ pair can be plausibly calculated using perturbative QCD. By combining that perturbative calculation with a phenomenological wavefunction for the dominant $Q\bar{Q}$ Fock state, we can calculate the annihilation decay rate of the quarkonium. This model for annihilation decays is called the color-singlet model.

The simplest illustration of the color-singlet model is the calculation of the decay rate of the ψ_c . In the color-singlet model, the ψ_c is modeled by a $c\bar{c}$ pair in a color-singlet 1S_0 state. Its wavefunction is the product of a color factor $\delta_{ij} = \frac{1}{\sqrt{3}}$, a spin factor $(\uparrow\downarrow) = \frac{1}{\sqrt{2}}$, and a coordinate-space wavefunction $\psi(r) = R(r) = \frac{1}{\sqrt{4}}$. A $c\bar{c}$ pair in such a state can annihilate into two gluons. If we assume that the two gluons hadronize into light hadrons with probability 1, the decay rate can be written as

$$\Gamma(\psi_c) = \frac{1}{2M_c} \int \frac{d^3k}{(2\pi)^3} \frac{2(M_c - 2|k|)}{M_c} T[\psi_c \rightarrow g(k)g(-k)]^2 \quad (28)$$

The T-matrix element for this decay can be expressed in terms of the momentum-space wavefunction $\psi(q)$ of the ψ_c :

$$T[\psi_c \rightarrow g(k)g(-k)] = \frac{1}{2M_c} \int \frac{d^3q}{(2\pi)^3} \psi(q) T[c(q)\bar{c}(-q) \rightarrow g(k)g(-k)]; \quad (29)$$

where we have suppressed all color and spin indices. The $c\bar{c}$ annihilation amplitude T varies significantly with q only when $|q|$ is on the order of m_c or larger. The wavefunction has significant support only for $|q|$ of order $m_c v$. Since T is almost independent of q for such small values of $|q|$, we can set $q = 0$ in the annihilation amplitude. The resulting expression for the decay

rate has a factored form :

$$\Gamma(\chi_c) = \frac{1}{(2M_c)^2} \int \frac{d^3q}{(2\pi)^3} |\psi(q)|^2 \int \frac{d^3k}{(2\pi)^3} \frac{2(M_c - 2k)}{M_c} \text{Im} T[c(0)c(0)^\dagger g(k)g(-k)]^2 : \quad (30)$$

The integral over q gives the wavefunction evaluated at the origin, $R(0) = \frac{3}{4} \frac{P}{M_c}$. The integral over k in (30) can be calculated from the lowest order QCD Feynman diagrams for $cc \rightarrow gg$. The final expression for the decay rate is

$$\Gamma(\chi_c) = \frac{8}{3M_c^2} \frac{s}{c} R(0)^2 : \quad (31)$$

3.2 Decay of χ_c in NRQCD

The decay rate of the χ_c into individual final states consisting of light hadrons can not be described within the framework of NRQCD. One obstacle is that we have imposed a symmetry on the effective field theory that guarantees the separate conservation of the numbers of c 's and \bar{c} 's and therefore forbids the annihilation process $cc \rightarrow gg$. Furthermore, in the construction of NRQCD, we have integrated out gluons with momenta on the order of m_c . Even if we relax the definition of the effective theory to allow gluons with momenta of order m_c and interaction terms in the Lagrangian that allow cc annihilation, we cannot describe annihilation decays accurately, because the interaction of a c or \bar{c} with a gluon of momentum m_c cannot be described accurately in a local nonrelativistic theory.

While the decay rate of the χ_c into a specific final state consisting of light hadrons cannot be described within NRQCD, the total decay rate can. The conservation of the number of c and \bar{c} is not an obstacle, because the optical theorem can be used to express the inclusive annihilation rate in terms of an amplitude that conserves the numbers of c and \bar{c} . As an illustration, applying the optical theorem to the expression (28), we find that the decay rate can be written

$$\Gamma(\chi_c) = \frac{1}{(2M_c)^2} \int \frac{d^3q}{(2\pi)^3} \int \frac{d^3q^0}{(2\pi)^3} |\psi(q)|^2 \text{Im} T[c(q)c(-q)^\dagger c(q^0)c(-q^0)] : \quad (32)$$

The process $cc \rightarrow cc$ conserves the numbers of c 's and \bar{c} 's. The factored form (30) is recovered by using the fact that the imaginary part of the T-matrix

element for $c \bar{c}$ is insensitive to momenta q and q^0 on the order of $M v$ where the wavefunctions have their support:

$$(\bar{c}c) = \frac{1}{(2M v_c)^2} \int \frac{d^3q}{(2\pi)^3} (q) \int_0^2 \text{Im T}[c(0)c(0) \bar{c}(0)c(0)]: \quad (33)$$

The integral over q in (33) gives the square of the wavefunction evaluated at the origin. This reflects the fact that the initial c and \bar{c} must have spacetime separations of order $1=M v$ in order to annihilate, and this separation is small compared to the length scale $1=(M v)$ of the wavefunction. That the $c\bar{c}$ pair must have a spacetime separation of order $1=M v$ follows from the fact that the Feynman diagrams for the annihilation process involve a heavy quark propagator that is off its mass-shell by an amount of order $M v$. The same argument implies that the final c and \bar{c} in the T-matrix element in (33) must have a spacetime separation of order $1=M v$. While it is not quite as obvious, the spacetime separation of the annihilation points for the initial $c\bar{c}$ pair and the final $c\bar{c}$ pair must also be of order $1=M v$. This follows from the requirement that the wavefunctions of the annihilation gluons must overlap, which localizes the production point of a gluon to within its wavelength $1=M v$.

The fact that $c\bar{c}$ annihilation occurs within a region whose size is of order $1=M v$ provides a clue as to how the effects of annihilation can be taken into account in NRQCD. All modes with momenta of order $M v$ that can be sensitive to the length scale $1=M v$ have been removed from this effective theory. Thus the effects of the annihilation can be reproduced by including in the NRQCD Lagrangian (12) a local 4-fermion interaction term that destroys a $c\bar{c}$ pair and creates it again. The specific term that is relevant to $c\bar{c}$ decay is

$$\mathcal{L} = \frac{f}{M^2} \bar{\psi} \psi \bar{\psi} \psi; \quad (34)$$

where the coefficient f is dimensionless. The term (34) annihilates a $c\bar{c}$ pair in a color-singlet 1S_0 state and then creates a $c\bar{c}$ pair in the same state.

The dimensionless coefficient f in (34) can be determined by perturbative matching of the $c\bar{c} \rightarrow c\bar{c}$ scattering amplitudes in full QCD and NRQCD. In full QCD, the scattering amplitude includes box diagrams of order α_s^2 in which the scattering proceeds through intermediate states consisting of two gluons. In NRQCD, this contribution to the scattering amplitude can only be reproduced by 4-fermion interactions such as those in (34). By matching the NRQCD scattering amplitude from the term (34) with the annihilation part of the scattering amplitude in full QCD, we can determine the coefficient f . Since the c and \bar{c} can annihilate into two on-shell gluons, the QCD scattering

amplitude has an imaginary part. The coefficient f in (34) must therefore have an imaginary part and it is particularly simple to calculate. The result is

$$\text{Im } f = \frac{2}{9} \frac{s^2(m_c)}{s^2(m_c)}; \quad (35)$$

Since this coefficient is sensitive only to distances of order $1/m_c$ or smaller, the running coupling constant is evaluated at the scale m_c . The fact that coefficients in the NRQCD Lagrangian have imaginary parts implies that the Hamiltonian for NRQCD is not hermitian. This is perfectly natural, since we have removed states from the theory that are essential for exact unitarity. In particular, we have eliminated the light partons with momenta on the order of m_c that can be produced by the annihilation of the c and \bar{c} .

We now consider the effect of the correction terms (34) on the energy of the χ_{c0} . If that term is treated as a first-order perturbation, the resulting correction to the energy of the χ_{c0} is

$$E_{\chi_{c0}} = \frac{f}{M^2} \frac{\langle \chi_{c0} | h_c j^y j_{c0}^y | \chi_{c0} \rangle}{2M_c}; \quad (36)$$

where we have assumed that the state $|\chi_{c0}\rangle$ has the standard relativistic normalization. Since the coefficient f has an imaginary part, this energy shift has an imaginary part. A state whose energy has a small imaginary part $\Gamma/2$ should be interpreted as a resonance of width Γ . Thus the width of the χ_{c0} due to the term (34) is

$$\Gamma(\chi_{c0}) = \frac{1}{2M_c} \frac{4}{9m_c^2} \frac{s^2(m_c)}{s^2(m_c)} \langle \chi_{c0} | h_c j^y j_{c0}^y | \chi_{c0} \rangle; \quad (37)$$

The connection with the result (32) from the color-singlet model is made by inserting a complete set of states between h_c and j^y in the matrix element. Assuming that the sum over states is dominated by the vacuum, we have

$$\langle \chi_{c0} | h_c j^y j_{c0}^y | \chi_{c0} \rangle = \langle 0 | j^y j_{c0}^y | 0 \rangle; \quad (38)$$

In the color-singlet model, the χ_{c0} -to-vacuum matrix element on the right side of (38) can be expressed in terms of the wavefunction:

$$\langle 0 | j^y j_{c0}^y | 0 \rangle = \frac{3}{2} \frac{r}{2M_c} R(0); \quad (39)$$

Inserting (38) into (37) and identifying M_c with $2m_c$, we reproduce the result (32) from the color-singlet model.

3.3 NRQCD Factorization Formula

In the expression (37) for the decay rate of the ψ_c , short-distance and long-distance effects have been factored. Long-distance effects involving the quarkonium wavefunction appear only in the NRQCD matrix element, which scales like $M^{-4}v^3$ according to the velocity-scaling rules in Table 3. Short-distance effects involving the annihilation of the $Q\bar{Q}$ pair appear only in the coefficient $2\text{Im}f = M^{-2}$, which is expressed in terms of the fundamental parameters α_s and m_c of QCD. Thus the expression for the decay rate in (37) scales like $\alpha_s^2 v^3$.

The formula (37) can be generalized to all orders in α_s and to all orders in v . The general factorization formula for the annihilation decay rate of a quarkonium state H is

$$\Gamma(H) = \frac{1}{2M_H} \sum_{m,n} C_{m,n} \langle H | \mathcal{O}_{m,n} | H \rangle; \quad (40)$$

where the onium state $|H\rangle = |H(P=0)\rangle$ has the standard relativistic normalization. The sum in (40) extends over all operators that can appear in the NRQCD Lagrangian and that have the form

$$\mathcal{O}_{m,n} = \bar{\psi} K_m \psi K_n; \quad (41)$$

These operators must be gauge invariant, invariant under parity and charge-conjugation, and scalars under rotations. Each of the factors K_n and K_m is the product of a spin matrix (1 or γ^i), a color matrix (1 or T^a), and a polynomial in D and $[D_0, D] = gE$. The operator $\mathcal{O}_{m,n}$ in (41) annihilates a $Q\bar{Q}$ pair in a color and angular-momentum state determined by K_n and creates a $Q\bar{Q}$ pair at the same point in a state determined by K_m .

The NRQCD factorization formula (40) untangles the effects of short distances of order $1=M$ from those of long distances of order $1=(Mv)$ or larger. All long-distance effects involving the quarkonium wavefunction are factored into the NRQCD matrix elements. Short-distance effects involving the annihilation of the $Q\bar{Q}$ pair are contained in the coefficients. The coefficient $C_{m,n}$ in (40) is twice the imaginary part of the coefficient of the operator $\mathcal{O}_{m,n}$ in the NRQCD Lagrangian. If that operator has scaling dimension $d_{m,n}$, then $C_{m,n}$ is $1=m_Q^{d_{m,n}-4}$ multiplied by a power series in $\alpha_s(m_Q)$.

The coefficients $C_{m,n}$ can be calculated using perturbative matching methods. A general matching prescription, called the threshold expansion method, has been developed by Braaten and Chen.⁸ The matching calculations are carried out using perturbative asymptotic states $cc = c(q; \epsilon)$ that consist of a c and a \bar{c} with relative momentum q and in a spin/color state that is represented

by the Pauli spinors χ and $\bar{\chi}$. The matching prescription is

$$\sum_X (2)^4 \int \frac{d^4 p}{(2\pi)^4} (\bar{\chi} \gamma^\mu \chi) (T_{cc^0}^\dagger)_X T_{cc^0}^\dagger X_{pQCD} \\ \sum_{m,n} C_{m,n} h_{cc^0}^0 j^Y K_m^Y K_n^Y j^c i_{pNRQCD}; \quad (42)$$

where $P = (2 \sqrt{m_c^2 + q^2}; 0)$ is the four-momentum of the cc pair and $T_{cc^0}^\dagger X$ is the T-matrix element for its annihilation into a final state X consisting of light partons. The sum over X on the left side of (42) includes integration over the phase space of the light partons. The complete determination of the short-distance coefficients requires the use of different states cc and cc^0 in the T-matrix element and in its complex conjugate. In the matching procedure, the left side of (42) is calculated using perturbative QCD, and then expanded in powers of the relative momenta q and q^0 . The matrix elements on the right side are calculated using perturbative NRQCD, and then expanded in powers of q and q^0 . The coefficients $C_{m,n}$ are determined by matching these expansions order by order in v .

The relative importance of the various terms in the factorization formula (40) is determined by the order in v of the coefficient $C_{m,n}$ and by the order in v of the matrix element. The magnitude of the matrix element can be estimated using the velocity-scaling rules for operators given in Table 3 and the estimates for the probabilities of higher Fock states in Coulomb gauge given in Section 2.5. If the operator $O_{m,n}$ annihilates and creates a $Q\bar{Q}$ pair in the same color and angular-momentum state as in the dominant $\mathcal{D}\bar{Q}i$ Fock state of H , then the estimate for the matrix element is obtained by dividing the estimate for the operator $O_{m,n}$ from Table 3 by $M^2 v^3$. Otherwise, we must take into account suppression factors from the transitions required to go from the dominant $\mathcal{D}\bar{Q}i$ state to a Fock state in which the $Q\bar{Q}$ pair can be annihilated by the operator $O_{m,n}$ and then back to the dominant $\mathcal{D}\bar{Q}i$ state. There is a suppression factor of v for every electric transition and a suppression factor of $v^{3=2}$ for every magnetic transition. Let E and M be the total number of electric and magnetic transitions required. If K_m and K_n contain D factors of the covariant derivative D but no factors of gE or gB , then the matrix element $\langle H | \mathcal{D}_{m,n} | H \rangle$ scales like $v^{3+D+E+3M=2}$. For each factor of gE or gB , there is an additional suppression factor of v^3 or v^4 , respectively.

Spin symmetry relates NRQCD matrix elements for quarkonium states that differ only in their spin quantum numbers. An example involving the c and $J =$ (the lowest 1S_0 and 3S_1 states of charmonium) is

$$\langle H | j^Y j^Y | H \rangle = \langle H | j^Y j^Y | H \rangle = \langle H | j^Y j^Y | H \rangle = \langle H | j^Y j^Y | H \rangle; \quad (43)$$

Since spin symmetry is only an approximate symmetry of NRQCD that is broken at order v^2 , the equality (43) holds only up to corrections of relative order v^2 .

The vacuum-saturation approximation can be used to express some of the matrix elements in the factorization formula (40) in terms of vacuum-to-quarkonium matrix elements. This approximation can only be applied if the operator O_{mn} annihilates and creates $Q\bar{Q}$ pairs in the same color and angular momentum state as in the dominant $Q\bar{Q}$ Fock state of H . In this case, we can insert a complete set of states between \langle and \rangle :

$$\langle H_j^Y K_m^Y K_n^Y | H | i \rangle = \sum_X \langle H_j^Y K_m^Y | X \rangle \langle X | K_n^Y K_n^Y | i \rangle \quad (44)$$

The vacuum-saturation approximation consists of keeping only the vacuum term $|0\rangle$ in the sum over states, as illustrated in (38). The vacuum-saturation approximation is a controlled approximation with an error of relative order v^4 . This follows from the fact that in Coulomb gauge, the next most important term in the sum over states in (44) is a gg Fock state, which contains two dynamical gluons. The leading contributions to the matrix elements in (44) then come from $Q\bar{Q}gg$ Fock states whose probabilities are of order v^4 .

The NRQCD matrix elements are sensitive to long-distance effects, and therefore can only be calculated using nonperturbative methods. The only practical nonperturbative method that is presently available is Monte Carlo simulations of lattice NRQCD. The first such calculations have been carried out recently by Bodwin, Sinclair, and Kim.⁹ They demonstrated that the relation (38) implied by the vacuum-saturation approximation holds to within numerical accuracy. They calculated the matrix elements $\langle 0 | j^Y | j_c \rangle$ and $\langle 0 | j^{YD^2} | j_c \rangle$ and their analogues for bottomonium. These matrix elements contribute to decays of the lowest S-wave states at leading order in v and at relative order v^2 . They also calculated $\langle 0 | j^{YD} | j_0 \rangle$ and $\langle c_0 | j^Y | T^a$
 $\langle j_c \rangle$ and the analogues of these matrix elements for bottomonium. As discussed in Section 3.4, these are the matrix elements that contribute to decays of the lowest P-wave states at leading order in v . Thus far, the calculations of NRQCD matrix elements have been carried out only in minimal NRQCD and without dynamical quarks.

3.4 Annihilation Decays of P-wave States

The NRQCD factorization formula for annihilation decay rates has dramatic implications for the decays of P-wave states, such as h_c (the 1P_1 state of charmonium) and ψ_{cJ} , $J = 0; 1; 2$ (the 3P_J states). Calculations of their annihilation decay rates in the color-singlet model suffer from infrared divergences.

The NRQCD factorization approach not only resolves the problem of the infrared divergences, but it also leads to new qualitative insights about P-wave charmonium.

We first consider the Fock state expansion of h_c and ψ_{cJ} in Coulomb gauge. The dominant Fock state consists of a color-singlet cc pair in a 1P_1 state for h_c and a 3P_J state for ψ_{cJ} . An electric transition from the dominant Fock state produces a ψ_{cJ} state, with the cc pair in a color-octet state with angular momentum quantum numbers 1S_0 or 1D_2 for h_c and $^3S_1, ^3D_1, ^3D_2, \text{ or } ^3D_3$ for ψ_{cJ} . Therefore these Fock states have probabilities of order v^2 . A magnetic transition from the dominant Fock state produces a ψ_{cJ} state, with the cc pair in a color-octet state with angular-momentum quantum numbers 3P_J for h_c and 1P_1 for ψ_{cJ} . This Fock state has a probability of order v^3 . All other Fock states have probabilities of order v^4 or smaller.

We proceed to identify the most important matrix elements in the NRQCD factorization formula. The lowest dimension operator that can annihilate the cc pair in the dominant Fock state of the h_c is $\psi_{\mathcal{D}} \psi_{\mathcal{D}}$, where \mathcal{D} is defined by $\psi_{\mathcal{D}} = \psi(\mathbf{D}) \cdot (\mathbf{D} \cdot \mathbf{Y})$. For ψ_{cJ} , the lowest dimension operator has the form $\psi_{\mathcal{D}}^{m,n} \psi_{\mathcal{D}}^{i,j}$, where the indices are contracted in different ways for $J = 0, 1, \text{ and } 2$. The matrix elements of these operators for $h_c; \psi_{c0}; \psi_{c1}, \text{ and } \psi_{c2}$ are related by spin symmetry. Up to corrections of relative order v^2 , they satisfy

$$\begin{aligned} \langle h_c | j^{\mathcal{D}} \psi_{\mathcal{D}} \psi_{\mathcal{D}} | h_c \rangle &= \langle \psi_{c0} | j^{\mathcal{D}} \psi_{\mathcal{D}} \psi_{\mathcal{D}} | \psi_{c0} \rangle \\ &= \frac{1}{3} \langle \psi_{c0} | j^{\mathcal{D}} \psi_{\mathcal{D}} \psi_{\mathcal{D}} | \psi_{c0} \rangle \\ &= \frac{1}{2} \langle \psi_{c1} | j^{\mathcal{D}} \psi_{\mathcal{D}} \psi_{\mathcal{D}} | \psi_{c1} \rangle \\ &= \langle \psi_{c2} | j^{\mathcal{D}} \psi_{\mathcal{D}}^{(m,n)} \psi_{\mathcal{D}}^{(m,n)} | \psi_{c2} \rangle; \end{aligned} \quad (45)$$

where $T^{(m,n)}$ denotes the symmetric traceless part of a tensor $T^{m,n}$. The vacuum-saturation approximation can be used to express these matrix element in a simpler form. Up to corrections of relative order v^4 , the matrix element for the ψ_{c0} can be written

$$\langle \psi_{c0} | j^{\mathcal{D}} \psi_{\mathcal{D}} \psi_{\mathcal{D}} | \psi_{c0} \rangle = \langle h_c | j^{\mathcal{D}} \psi_{\mathcal{D}} \psi_{\mathcal{D}} | h_c \rangle : \quad (46)$$

In the color-singlet model, the vacuum-to- ψ_{c0} matrix element on the right side of (46) can be expressed in terms of the radial wavefunction $R(r)$ for the P-wave states:

$$\langle h_c | j^{\mathcal{D}} \psi_{\mathcal{D}} \psi_{\mathcal{D}} | h_c \rangle = \frac{9}{2M_{c0}} \int_0^\infty r^2 R^0(0) : \quad (47)$$

According to the velocity-scaling rules, the matrix elements in (45) scale like v^5 . For a consistent analysis, we must also include all other matrix elements that scale like v^5 . By enumerating the possibilities, one can see that the only other operators whose matrix elements scale like v^5 are $\bar{\psi}T^a\psi$ for h_c and $\bar{\psi}T^a\psi$ for c_J . These operators annihilate and create cc pairs in color-octet 1S_0 and 3S_1 states, respectively. Spin symmetry implies that the matrix elements for h_c ; c_0 , c_1 , and c_2 are equal, up to corrections of relative order v^2 :

$$\langle h_c j | \bar{\psi}T^a\psi | j c_0 i \rangle = \langle h_c j | \bar{\psi}T^a\psi | j c_1 i \rangle = \langle h_c j | \bar{\psi}T^a\psi | j c_2 i \rangle; \quad J = 0; 1; 2; \quad (48)$$

We have found that, up to corrections that are suppressed by v^2 , the annihilation decay rates of the P-wave states can all be expressed in terms of the following two independent matrix elements:

$$\langle h_0 | i \rangle = \frac{\langle h_0 j | \bar{\psi}T^a\psi | j_0 i \rangle}{2M_{c_0}}; \quad (49)$$

$$\langle h_8 | i \rangle = \frac{\langle h_8 j | \bar{\psi}T^a\psi | j_8 i \rangle}{2M_{c_0}}; \quad (50)$$

Their short-distance coefficients can be calculated as power series in $\frac{m_c}{s}$. The annihilation processes that contribute to the coefficients at order $\frac{m_c^2}{s}$ are $cc \rightarrow gg$ and $c\bar{c} \rightarrow q\bar{q}$. If we keep only those terms in the short-distance coefficients, the annihilation decay rates are

$$\langle h_c \rangle = \frac{5}{6m_c^2} \frac{m_c^2}{s} \langle h_8 | i \rangle; \quad (51)$$

$$\langle c_0 \rangle = \frac{4}{m_c^4} \frac{m_c^2}{s} \langle h_0 | i \rangle + \frac{n_f}{3m_c^2} \frac{m_c^2}{s} \langle h_8 | i \rangle; \quad (52)$$

$$\langle c_1 \rangle = \frac{n_f}{3m_c^2} \frac{m_c^2}{s} \langle h_8 | i \rangle; \quad (53)$$

$$\langle c_2 \rangle = \frac{16}{45m_c^4} \frac{m_c^2}{s} \langle h_0 | i \rangle + \frac{n_f}{3m_c^2} \frac{m_c^2}{s} \langle h_8 | i \rangle; \quad (54)$$

where $n_f = 3$ is the number of flavors of light quarks. At this order in $\frac{m_c}{s}$, the decay rates for h_c and c_1 receive contributions from $h_8 | i \rangle$ only, because Yang's theorem forbids the annihilation process $cc \rightarrow gg$ for a cc pair in a state with total angular momentum 1. At order $\frac{m_c^3}{s}$, all of the P-wave states have contributions from $h_0 | i \rangle$. At this order in $\frac{m_c}{s}$, the short-distance coefficients of $h_0 | i \rangle$ depend logarithmically on a factorization scale that can be interpreted

as an infrared cutoff on the energy of soft gluons. The matrix element $\langle 0 | \dots | 0 \rangle$ also depends logarithmically on Λ , which in this case can be identified with the ultraviolet cutoff of NRQCD. The Λ -dependence cancels between $\langle 0 | \dots | 0 \rangle$ and the coefficient of $\langle 0 | \dots | 0 \rangle$.¹⁰ In the color-singlet model, the $\langle 0 | \dots | 0 \rangle$ terms are absent and the decay rate depends logarithmically on the infrared cutoff Λ . The NRQCD factorization approach provides a simple and natural solution to this problem.

The NRQCD factorization formula for a P-wave state has a simple interpretation in terms of the Fock state expansion in Coulomb gauge. The color-singlet terms proportional to $\langle 0 | \dots | 0 \rangle$ are contributions from the dominant 3S_1 Fock state, while the color-octet terms proportional to $\langle 0 | \dots | 0 \rangle$ are contributions from a 3S_1 Fock state. The 3S_1 state has a small probability of order v^2 , and its effects on most observables are small compared to those of the 3S_1 state. However, in the case of the annihilation decay rate, the effects of the 3S_1 state are suppressed by v^2 due to the orbital angular momentum of the 3S_1 pair. The contribution of the 3S_1 Fock state has no angular-momentum suppression and therefore contributes at the same order in v . For the h_c and ψ_c , the effects of the 3S_1 Fock state actually dominate, because the short-distance coefficients of $\langle 0 | \dots | 0 \rangle$ are suppressed by a factor of $\alpha_s(m_c)$.

4 Inclusive Production of Heavy Quarkonium

4.1 Topological Factorization

The cross section for producing a quarkonium state H in a high energy process necessarily involves both "short distances" of order $1/M$ or smaller and "long distances" of order $1/(Mv)$ or larger. The creation of the $Q\bar{Q}$ pair involves short distances, because the parton processes that produce the $Q\bar{Q}$ pair always involve particles that are off their mass shells by amounts of order M and can therefore propagate only over short distances. The binding of the Q and \bar{Q} into the state H involves long distances, because gluons whose wavelengths are comparable to or larger than the size of the bound state, which is of order $1/(Mv)$, play a large role in the binding.

The production of quarkonium in a high energy physics process typically involves another hard momentum scale Q in addition to the scale M . If the production cross section is sufficiently inclusive, it can be described by an NRQCD factorization formula that separates short-distance effects involving the momentum scales Q and M from long-distance effects that involve lower momentum scales. It is convenient to separate the derivation of the factorization formula into two steps. In the first step, which we refer to as topological factorization, the standard factorization methods of perturbative QCD are used to separate the effects of the hard momentum scale Q from those of the

soft momentum scale Q_{CD} . These methods are applicable even if we identify the hard scale Q with the heavy quark mass M . An additional step is required to separate the effects of the scale M from those of the scale Mv .

The general expression for the inclusive cross section for the production of a quarkonium state H with four-momentum P is

$$\sum_X d(12!_{H(P)+X}) = \frac{1}{4E_1 E_2 v_{12}} \frac{d^3P}{(2\pi)^3 2E_P} \sum_X (2\pi)^4 \delta^4(k_1 + k_2 - P - k_X) T_{12!_{H(P)+X}}^J; \quad (55)$$

where $T_{12!_{H(P)+X}}$ is a T-matrix element for producing H and the additional particles X and the sum on the right side includes integration over the phase space of the additional particles. At the parton level, both $T_{12!_{H(P)+X}}$ and its complex conjugate can be expressed as sums of Feynman diagrams. The product of a single diagram in $T_{12!_{H(P)+X}}$ and a single diagram in $T_{12!_{H(P)+X}}$ is called a "cut Feynman diagram." Using the factorization methods of perturbative QCD, one can identify the cut diagrams that dominate in the limit $Q \gg 1$. After taking into account cancellations between real and virtual soft gluons, the dominant cut diagrams have the following structure:

- a hard-scattering subdiagram H to the left of the cut. The outgoing lines include a $Q\bar{Q}$ pair with small relative momentum of order Mv and additional hard partons. There can also be incoming hard-parton lines if the process involves hadrons in the initial state.

- a hard-scattering subdiagram H to the right of the cut that is just the mirror image of H .

- a jet-like subdiagram J for each of the hard partons attached to H . The subdiagram extends through the cut and is attached to H and to H by single hard parton lines.

- an onium subdiagram O that extends through the cut and is attached to H and to H by Q and \bar{Q} lines that have small relative momentum.

The cut diagrams that do not have the above structure are suppressed by powers of $1=Q$. With this topological factorization of the dominant cut diagrams, all effects involving the hard momentum scale Q are factored into the hard-scattering subdiagrams H and H , while all effects of the soft scale Q_{CD} are factored into O, J_1, J_2, \dots . The gluon interactions that bind the $Q\bar{Q}$ pair into the onium state H are also contained within the onium subdiagram O .

The proofs of the factorization theorems of perturbative QCD are very difficult, and explicit proofs are available only for a very few processes, such as inclusive hadron production in e^+e^- annihilation and the Drell-Yan process for lepton pair production in hadron collisions.¹¹ However there is no apparent obstacle to extending these proofs to inclusive onium production.

4.2 NRQCD Factorization

After topological factorization, the effects of the scale M_v are distributed in a complicated way between H , H_0 , and the Q and \bar{Q} propagators that connect them. The onium subdiagram O involves the scale M_v because gluons with momentum of order M_v play an important role in the binding of the $Q\bar{Q}$ pair into an onium state. The hard-scattering subdiagrams H and H_0 involve the scale M_v , because the outgoing Q and \bar{Q} lines have relative momenta on the order of M_v . It is in factoring the scale M_v out of H and H_0 that NRQCD enters into the picture.

Consider the part of the cut diagram that includes H , O , and the Q and \bar{Q} propagators that connect them. If the Q and \bar{Q} have four-momenta $\frac{1}{2}P + q$ and $\frac{1}{2}P - q$, then the diagram also involves an integral over the relative momentum q . A simple way to disentangle the momentum scale M_v from H is to expand it as a Taylor series in q and absorb the factors of q as well as the integration over q into O . Each term in the Taylor expansion corresponds to a local operator that creates a $Q\bar{Q}$ pair from the vacuum. By applying a similar procedure to disentangle the momentum scale M_v from H_0 , the onium subdiagram is reduced to vacuum-expectation values of local operators that create and annihilate $Q\bar{Q}$ pairs. After a renormalization group transformation and appropriate field redefinitions, the matrix elements can be expressed as expectation values in the NRQCD vacuum of the form

$$\langle 0 | \bar{\psi}_m^H \psi_n^H | 0 \rangle = \langle 0 | \bar{\psi}_m^H \psi_n^H P_H | 0 \rangle \quad (56)$$

where P_H projects onto states that in the asymptotic future contain the quarkonium state H plus soft partons S whose total energy is less than the ultraviolet cutoff of NRQCD:

$$P_H = \sum_s |H + S\rangle \langle H + S| \quad (57)$$

If we integrate the dominant cut diagrams over the phase space of all the hard partons in the process, we obtain the NRQCD factorization formula for

the inclusive cross section:

$$\sum_X d(12! H(P) + X) = \frac{1}{4E_1 E_2 v_{12}} \frac{d^3 P}{(2)^3 2E_P} \sum_{m,n} C_{m,n}(k_1; k_2; P) hO_{m,n}^H i; \quad (58)$$

The sum in (58) extends over all NRQCD matrix elements of the form (56). The product of the operators $\mathcal{Y}K_m$ and $\mathcal{Y}K_n$ must be gauge-invariant. It need not be rotationally invariant if the quarkonium state H is polarized. In the factorization formula (58), the hard-scattering subdiagrams H and H and the jet-like subdiagrams J_i have all been subsumed in the short-distance coefficients $C_{m,n}$.

Since the coefficients $C_{m,n}$ in (58) involve only short distances of order $1/M$ or larger, they can be expressed as perturbation series in $\alpha_s(M)$. The threshold expansion method provides a general prescription for calculating the short-distance coefficients.⁸ Denoting by $cc(P)$ a state consisting of a c and \bar{c} with relative momentum q that has been boosted to four-momentum P , the matching prescription is

$$\sum_X (2)^4 (k_1 + k_2 - P - k) (\Gamma_{12!}^{cc^0(P)+X} \Gamma_{12!}^{cc(P)+X})_{PQCD} \sum_{m,n} C_{m,n}(k_1; k_2; P) hO_j^{\mathcal{Y}K_m} P_{cc^0;cc} \mathcal{Y}K_n \rangle_{PNRQCD} i; \quad (59)$$

where the projection operator in the NRQCD matrix element is

$$P_{cc^0;cc} = \sum_s \mathcal{Y}c^0 + S i h c c + S j; \quad (60)$$

The left side of (59) is to be calculated using perturbative QCD, and then expanded in powers of the relative momenta q and q^0 of the cc pairs. The matrix elements on the right side are to be calculated using perturbative NRQCD, and then expanded in powers of q and q^0 . The coefficients $C_{m,n}$ are then determined by matching these expansions order by order in α_s .

The relative importance of the various terms in the factorization formula (40) is determined by the magnitudes of the coefficients $C_{m,n}$ and by the order in v of the matrix elements $hO_{m,n}^H i$. The size of the coefficient $C_{m,n}$ is determined not only by the order in α_s , but also by its dependence on dimensionless ratios of kinematic variables that are involved in the cc production process. The magnitudes of the matrix elements can be estimated by using the velocity-scaling rules for operators given in Table 3 and the scaling with v of the rates for electric and magnetic transitions. If the operator $O_{m,n}^H$ creates and annihilates

a $Q\bar{Q}$ pair in the same color and angular-momentum state as in the dominant $Q\bar{Q}$ Fock state of H , then the magnitude of the matrix element $\langle 0_{m,n}^H | i$ is estimated by multiplying the factors in Table 3 and dividing by $M^2 v^3$. For other matrix elements, we must take into account suppression factors from the transitions required to go from the $Q\bar{Q}$ state created by the operator ${}^Y K_n$ to a state in which the $Q\bar{Q}$ pair has the same quantum numbers as in the dominant $Q\bar{Q}$ Fock state and then to a state in which the $Q\bar{Q}$ pair can be annihilated by the operator ${}^Y K_m$. There is a suppression factor of v for every electric transition that is required and a suppression factor of $v^{3=2}$ for every magnetic transition. The scaling of the production matrix element $\langle 0_{m,n}^H | i$ with v is identical to that of the corresponding decay matrix element $\langle H | \mathcal{D}_{m,n} | H \rangle$.

The NRQCD matrix elements that appear in the NRQCD factorization formula (58) can be simplified by using symmetries of NRQCD. Rotational symmetry is an exact symmetry of NRQCD. It implies, for example, that

$$\langle H | {}^Y J_{T^a} | P_{J=0} \rangle = \frac{1}{3} \langle H | {}^Y K_{T^a} | P_{J=0} \rangle \langle H | {}^Y K_{T^a} | i \rangle; \quad (61)$$

Spin symmetry is an approximate symmetry of NRQCD that holds up to corrections of order v^2 . It implies, for example, that

$$\langle H | {}^Y J_{T^a} | P_{(\lambda)} \rangle = \langle H | {}^Y K_{T^a} | P_{(\lambda)} \rangle \langle H | U_i^Y | P_{(\lambda)} \rangle \langle H | {}^Y K_{T^a} | i \rangle; \quad (62)$$

where U_i specifies the polarization of the $J=0$ and U_i is the unitary matrix that transforms vectors from the spherical basis to the Cartesian basis.

The vacuum-saturation approximation can be used to simplify the matrix elements of operators that create and annihilate cc pairs in the dominant Fock state of the quarkonium. In the vacuum-saturation approximation, the projection operator P_H defined in (57) is replaced by the single term $|H\rangle\langle H|$. This is a controlled approximation in NRQCD, holding up to corrections that are of order v^4 . It implies, for example, that

$$\langle 0 | j^Y | P_{(\lambda)} \rangle \langle P_{(\lambda)} | j^Y | i \rangle = \langle 0 | j^Y | i \rangle \langle i | P_{(\lambda)} | j^Y | i \rangle; \quad (63)$$

The matrix elements in the NRQCD factorization formula involve long-distance effects and therefore can only be calculated using nonperturbative methods. Unfortunately, in contrast to the decay matrix elements $\langle H | \mathcal{D}_{m,n} | H \rangle$, there are no effective prescriptions for calculating the production matrix elements $\langle 0_{m,n}^H | i$ using lattice NRQCD. The problem lies in implementing on the lattice the projection defined by (57). Thus these NRQCD matrix elements must be treated as phenomenological parameters to be determined by experiment. The only exceptions are the matrix elements to which the vacuum-

saturation approximation can be applied. Vacuum-to-quarkonium matrix elements of the form $\langle H | j^{\nu} K_n | j \rangle$ can be calculated using Monte-Carlo simulations of NRQCD.

4.3 Prompt Charmonium at the Tevatron

The NRQCD factorization framework (58) has many applications, some of which are described in a recent review.¹² One application for which the implications are particularly dramatic is the production of prompt charmonium at large transverse momentum in pp collisions. At sufficiently large transverse momentum p_T , the cross section for $pp \rightarrow \chi + X$ is dominated by gluon fragmentation.¹³ It can be factored into the cross section for producing a gluon with transverse momentum $p_T = z$ and a fragmentation function $D_{g\chi}(z)$ that gives the probability that the jet initiated by the gluon includes a χ carrying a fraction z of the gluon momentum:

$$d(\sigma(pp \rightarrow \chi + X)) = \int_0^1 dz d^4(\sigma(pp \rightarrow g(p=z) + X)) D_{g\chi}(z); \quad (64)$$

Using the NRQCD factorization approach, the fragmentation function can be expressed in the form

$$D_{g\chi}(z) = \sum_{m,n} d_{m,n}(z) \langle O_{m,n} \rangle; \quad (65)$$

where all effects of the momentum scale μ_c have been factored into the short-distance coefficients $d_{m,n}(z)$. The relative importance of the various terms in the fragmentation function is determined by the order in v of the matrix element and the order in s of its coefficient.

The matrix element that is leading order in v is $\langle \chi | \bar{\psi} \psi | \chi \rangle$, which scales like v^3 . This term corresponds to the formation of a χ from a cc pair that is created in a color-singlet 3S_1 state, and it is the only contribution in the color-singlet model. The leading contribution to the short-distance coefficient of this term in the gluon fragmentation function (65) is of order $\frac{3}{s}$ and comes from the parton process $g \rightarrow ccg$. Keeping only this term in the fragmentation function (65), the cross section predicted by (64) is about a factor of 30 below recent data on prompt χ production at the Tevatron.

The color-singlet model term in the gluon fragmentation function scales like $\frac{3}{s}v^3$. All other terms have matrix elements that scale like v^5 or smaller. There are however terms whose short-distance coefficients are suppressed by fewer powers of s . There is one coefficient in particular that is of order s , because it receives a contribution from the parton process $g \rightarrow cc$. The matrix

element is $h^{\gamma k T^a} P^{\gamma k T^a} i$ and it scales like v^7 . This term corresponds to the formation of a ψ from a cc pair that is created in a color-octet 3S_1 state. At leading order in α_s , this term in the fragmentation function is

$$D_{g\psi}(z) = \frac{\alpha_s(m_c)}{96m_c^4} (1-z) h^{\gamma k T^a} P^{\gamma k T^a} i: \quad (66)$$

Braaten and Fleming proposed that the enhancement from the two fewer powers of α_s relative to the color-singlet model term can overcome the suppression by v^4 , and that this term might dominate the gluon fragmentation function.¹⁴ The p_T -dependence predicted by this mechanism is in agreement with the Tevatron data. The normalization depends on the unknown matrix element in (66), but the value of the matrix element required to fit the data is consistent with suppression by a factor of v^4 relative to the color-singlet-model matrix element $h^{\gamma k} P^{\gamma k} i$.

Cho and Wise pointed out that this production mechanism has dramatic implications for the polarization of the ψ .¹⁵ At leading order in α_s , the ψ 's produced by gluon fragmentation will be 100% transversely polarized. The radiative corrections to the fragmentation function were examined by Beneke and Rothstein, and they concluded that the spin alignment at large p_T will remain greater than 90%.¹⁶ The largest corrections to the spin alignment at values of p_T that can be measured at the Tevatron come from nonfragmentation contributions that fall like $1/p_T^2$, and these contributions have recently been calculated by Beneke and Kraemer.¹⁷ An experimental measurement of the spin alignment in agreement with these predictions would constitute a dramatic triumph of the NRQCD factorization approach.

5 Conclusions

The NRQCD factorization formulas (40) and (58) provide a firm theoretical foundation for analyzing annihilation decay rates and inclusive production rates of heavy quarkonium. The short-distance coefficients can be calculated as power series in the running coupling constant $\alpha_s(M)$, and the long-distance factors are defined in terms of NRQCD matrix elements that scale in a definite way with v . This approach not only provides a framework for carrying out systematic quantitative calculations of quarkonium processes, but it also leads to new qualitative insights into quarkonium physics.

Acknowledgments

This work was supported in part by the U.S. Department of Energy, Division of High Energy Physics, under Grant DE-FG 02-91-ER 40684. I thank G.T.

Bodwin and G.P. Lepage for valuable discussions and I.M. Aksenov for a careful and critical reading of the manuscript. I would also like to thank the organizers of the Third International Workshop on Particle Physics Phenomenology for their hospitality.

References

1. G.T. Bodwin, E. Braaten, and G.P. Lepage, *Phys. Rev. D* 51, 1125 (1995); erratum in hep-ph/9407339.
2. H. Georgi, *Ann. Rev. Nucl. Part. Sci.* 43, 209 (1993).
3. W.E. Caswell and G.P. Lepage, *Phys. Lett.* 167B, 437 (1986).
4. G.P. Lepage et al., *Phys. Rev. D* 46, 4052 (1992).
5. C.T.H. Davies et al., *Phys. Rev. Lett.* 73, 2654 (1994); *Phys. Lett. B* 345, 42 (1995).
6. C.T.H. Davies et al., *Phys. Rev. D* 50, 6963 (1994); *Phys. Rev. D* 52, 6519 (1995).
7. M. Luke and A.V. Manohar, hep-ph/9610534.
8. E. Braaten and Y.-Q. Chen, *Phys. Rev. D* 54, 3216 (1996); hep-ph/9610401 (to appear in *Phys. Rev. D*); hep-ph/9701242.
9. G.T. Bodwin, D.K. Sinclair, and S. Kinoshita, *Phys. Rev. Lett.* 77, 2376 (1996).
10. G.T. Bodwin, E. Braaten, and G.P. Lepage, *Phys. Rev. D* 46, R1914 (1992).
11. J.C. Collins and D.E. Soper, *Ann. Rev. Nucl. Part. Sci.* 37, 383 (1987)
12. E. Braaten, S. Fleming, and T.C. Yuan, *Ann. Rev. Nucl. Part. Sci.* 46, 197 (1996).
13. E. Braaten and T.C. Yuan, *Phys. Rev. Lett.* 71, 1673 (1993).
14. E. Braaten and S. Fleming, *Phys. Rev. Lett.* 74, 3327 (1995).
15. P. Cho and M. Wise, *Phys. Lett. B* 346, 129 (1995).
16. M. Beneke and I.Z. Rothstein, *Phys. Lett. B* 372, 157 (1996).
17. M. Beneke and M. Kramer, hep-ph/9611218.