

HIGGS PHYSICS AT A MUON COLLIDER *

JOHN F. GUNION

*Davis Institute for High Energy Physics, Department of Physics, University of California at Davis
Davis, CA 95616, USA***ABSTRACT**

A brief review of muon-collider s -channel Higgs physics in the Standard Model and its minimal supersymmetric extension is presented.

1. Introduction

One of the most important tasks in high energy physics in the next decades will be to determine if Higgs boson(s) exist, and to measure the properties of any that are found. Only then will a full understanding of the Higgs sector and its role in the complete underlying theory be possible. Here we outline the very substantial and, in some areas, unmatched capabilities of a muon collider for these tasks.¹ We restrict our discussions to the Standard Model (SM) with a single neutral Higgs boson, h_{SM} , and to the minimal supersymmetric extension of the SM (MSSM) with a light SM-like Higgs boson, h^0 , and two heavier neutral Higgs scalars, H^0 and A^0 .

Although muon colliders have only received substantial attention in the last few years, so far it seems² that there are “no barriers” to either of the following two designs:

- $\sqrt{s} \lesssim 500$ GeV, $\mathcal{L} \sim \text{few} \times 10^{33} \text{cm}^{-2} \text{s}^{-1}$, $L_{\text{tot}} = 50 \text{fb}^{-1}/\text{yr}$;
- $\sqrt{s} \lesssim 4$ TeV, $\mathcal{L} \sim \text{few} \times 10^{35} \text{cm}^{-2} \text{s}^{-1}$, $L_{\text{tot}} = 200 \text{fb}^{-1}/\text{yr}$.

A $\mu^+\mu^-$ collider has some natural advantages as compared to an e^+e^- collider, including some that are crucial for Higgs boson physics:

- there is essentially no beamstrahlung;
- there is substantially reduced bremsstrahlung;^a
- there is no final focus problem (storage rings are used to build up the effective instantaneous luminosity);
- as required for detailed Higgs studies in the s -channel, excellent beam energy resolution of $R = 0.01\%$ and the ability to set E_{beam} to 1 part in 10^6 using spin

*To appear in the proceedings of the Ringberg workshop on “The Higgs Puzzle”, ed. B. Kniehl, Dec. 8-13, 1996.

^aStill, bremsstrahlung depletes the central Gaussian peak in \sqrt{s} by $\sim 40\%$ — this effect is included in the computations.¹

precession techniques are both possible if the necessary technology is built into the machine;

- $\sqrt{s}_{\text{max}} > 500$ GeV can probably be reached more easily; and
- there is hope that the cost might be lower.

The negatives regarding a muon collider include:

- the design is immature, and five years of research and development projects are needed before a full-fledged proposal would be possible — in particular, cooling tests are required to see if multistage cooling will be sufficiently efficient;
- the exact nature of the detector backgrounds, and how to manage them, is still under investigation — certainly the detector will be more expensive due to higher shielding and segmentation requirements;
- significant polarization probably implies significant loss in \mathcal{L} — for s -channel Higgs production, $S/\sqrt{B} \propto \sqrt{L_{\text{tot}}} \sqrt{(1+P^2)^2/(1-P^2)}$ implies that $[P = 0, L_{\text{tot}}]$ is equivalent to $[P = 0.8, L_{\text{tot}}/10]$;
- it is not possible to have a $\gamma\gamma$ collider facility.




Fig. 1. Feynman diagram for s -channel production of a Higgs boson.

A muon collider can explore Higgs physics in two modes. Away from the s -channel Higgs pole, *e.g.* at the maximum available energy \sqrt{s}_{max} , a $\mu^+\mu^-$ collider can discover and study Higgs bosons in exactly the same ways as an e^+e^- collider with the same \sqrt{s} and \mathcal{L} (barring unmanageable detector backgrounds at the muon collider). On the s -channel pole at $\sqrt{s} \sim m_h$, $\mu^+\mu^- \rightarrow h$ collisions (see Fig. 1) imply unique capabilities for:

- precision studies of $(b\bar{b}h)^2 : (WW h)^2 : (ZZ h)^2$ coupling-squared ratios and of Γ_h^{tot} for a SM-like Higgs (either h_{SM} or h^0) with $m_h \lesssim 2m_W$; and
- discovery and study of the heavier H^0, A^0 of the MSSM up to \sqrt{s}_{max} .

For s -channel Higgs physics, the size of the s -channel cross section, $\bar{\sigma}_h$, is crucial. We obtain $\bar{\sigma}_h$ by convoluting the standard Breit-Wigner shape for the Higgs with a Gaussian distribution of width $\sigma_{\sqrt{s}}$ centered at $\sqrt{s} = m_h$; $\bar{\sigma}_h$ is given by $\bar{\sigma}_h \sim 4\pi m_h^{-2} BF(h \rightarrow \mu^+ \mu^-)$ if $\sigma_{\sqrt{s}} \ll \Gamma_h^{\text{tot}}$ and by $\bar{\sigma}_h \sim 2\pi^2 m_h^{-2} \Gamma(h \rightarrow \mu^+ \mu^-) / (\sqrt{2\pi} \sigma_{\sqrt{s}})$ if $\sigma_{\sqrt{s}} \gg \Gamma_h^{\text{tot}}$. To get near maximal $\bar{\sigma}_h$ and to have sensitivity to Γ_h^{tot} via scanning in \sqrt{s} it is important that $\sigma_{\sqrt{s}}$ be no larger than $2 - 3 \times \Gamma_h^{\text{tot}}$. Very small Γ_h^{tot} is not uncommon. Fig. 2 shows that $\Gamma_h^{\text{tot}} < 1 - 10$ MeV is typical of the h_{SM} for $m_{h_{SM}} \lesssim 140$ GeV and of *all* the MSSM Higgs bosons if $\tan\beta \lesssim 2$ and $m_{A^0} \lesssim 2m_W$. Using the parameterization $\sigma_{\sqrt{s}} \simeq 7 \text{ MeV} \left(\frac{R}{0.01\%}\right) \left(\frac{\sqrt{s}}{100 \text{ GeV}}\right)$ for $\sigma_{\sqrt{s}}$ in terms of the beam energy resolution, R , we see that very excellent resolution $R \sim 0.01\%$ is required for $\sigma_{\sqrt{s}} < 2 - 3 \times \Gamma_h^{\text{tot}}$ in the above cases.




Fig. 2. Total width versus mass of the SM and MSSM Higgs bosons for $m_t = 175$ GeV. In the case of the MSSM, we have plotted results for $\tan\beta = 2$ and 20, taking $m_{\tilde{t}} = 1$ TeV and including two-loop/RGE-improved Higgs mass corrections and neglecting squark mixing; SUSY decay channels are assumed to be absent.

A final important note is that high \mathcal{L} is needed at all \sqrt{s} values where a Higgs boson is discovered or might exist. This will possibly imply that several final storage rings (which fortunately are relatively cheap), designed to maintain near-optimal \mathcal{L} over a span of \sqrt{s} values, will need to be constructed

2. s -channel studies of a Standard-Model-like Higgs boson

Most probably, one would first discover the h at the LHC (e.g. in the $h \rightarrow \gamma\gamma$ or $h \rightarrow 4\ell$ discovery modes), or at the NLC (e.g. in the $Z^* \rightarrow Zh$ production mode) and

then set up the muon collider for running at $\sqrt{s} \sim m_h$. At the LHC or NLC, m_h will be quite well determined. For $m_h \lesssim 2m_W$ one finds: ⁴ $\Delta m_h \sim 100 \text{ MeV} \left(\frac{600 \text{ fb}^{-1}}{L}\right)^{1/2}$ at the LHC, using $h \rightarrow \gamma\gamma, 4\ell$ decays and reconstructing the resonance peak in $m_{\gamma\gamma}, m_{4\ell}$; $\Delta m_h \sim 100 \text{ MeV} \left(\frac{50 \text{ fb}^{-1}}{L}\right)^{1/2}$ for NLC $\sqrt{s} = 500 \text{ GeV}$ running and reconstructing the $m_{b\bar{b}}$ resonance peak in the Zh ($h \rightarrow b\bar{b}$) mode; and $\Delta m_h < 100 \text{ MeV}$ at the NLC, assuming $L = 50 \text{ fb}^{-1}$ devoted to a Zh threshold study at $\sqrt{s} = m_Z + m_h + 0.5 \text{ GeV}$. If there is no LHC or NLC, one would accumulate $L \sim 1 \text{ fb}^{-1}$ at $\sqrt{s} = 500 \text{ GeV}$ at the muon collider in order to observe $\mu^+\mu^- \rightarrow Zh$ and determine m_h to within $\Delta m_h \sim 1 \text{ GeV}$, and then turn to $\sqrt{s} \sim m_h$ running.

For a SM-like Higgs with $m_h \gtrsim 2m_W$, Γ_h^{tot} is large (as a result of $h \rightarrow WW, ZZ$ decays), see Fig. 2, $\bar{\sigma}_h \propto BF(h \rightarrow \mu^+\mu^-)$ is tiny, and $\mu^+\mu^- \rightarrow h$ will not be useful. ¹ But, if $m_h \lesssim 2m_W$ then Γ_h^{tot} is very small, $\sigma_{\sqrt{s}} \sim 2 - 3 \times \Gamma_h^{\text{tot}}$ is typical for $R = 0.01\%$, $\bar{\sigma}_h \propto \Gamma(h \rightarrow \mu^+\mu^-)/\sigma_{\sqrt{s}}$ will be much larger, and the $\mu^+\mu^-$ collider becomes a Higgs factory. There are several reasons to suppose that the SM-like Higgs will have mass $\lesssim 2m_W$: a) precision electroweak measurements currently favor (but only weakly) a relatively light SM-like Higgs boson; b) light Higgs masses are preferred in the SM if the SM couplings are required to remain perturbative up to the GUT scale; and c) in the minimal supersymmetric model (MSSM), the light SM-like Higgs has $m_{h^0} \lesssim 130 \text{ GeV}$. ³

If a SM-like h with $m_h \leq 2m_W$ has been discovered, the FMC final ring would be optimized for $\sqrt{s} \sim m_h$ and a scan of the region of size $2\Delta m_h$ in the vicinity of the known m_h would be performed in order to pin down m_h more precisely (roughly within $\pm\sigma_{\sqrt{s}}/2$). ^b In the “typical case” of $m_h \sim 110 \text{ GeV}$, $\sigma_{\sqrt{s}} \sim 8 \text{ MeV}$ (for $R \sim 0.01\%$) and $2\Delta m_h \sim 200 \text{ MeV}$ (see above), we would require $\gtrsim 25$ scan points with $\sim 0.01 \text{ fb}^{-1}$ per scan point (the luminosity for a 5σ signal if $\sqrt{s} = m_h$), corresponding to a total luminosity of $L_{\text{tot}} \sim 0.25 \text{ fb}^{-1}$. (One must reset \sqrt{s} with $\Delta\sqrt{s} < \sigma_{\sqrt{s}}$ hourly.) Centering on $\sqrt{s} \simeq m_h$ does not require much luminosity for such Higgs masses. The worst mass is $m_h \simeq m_Z$. For the $R = 0.01\%$ value of $\sigma_{\sqrt{s}} \sim 6.5 \text{ MeV}$, $L_{\text{tot}} \sim 25 \text{ fb}^{-1}$ would be needed to center on $\sqrt{s} \simeq m_h$ by scanning. A smaller Δm_h , such as could be obtained using “super” tracking or other improvements at the NLC, ⁴ would be very helpful for avoiding wasting luminosity on the centering process in this case.

Once we have determined m_h to within $\sim \pm\sigma_{\sqrt{s}}/2$, we would perform a three point scan of the Higgs resonance peak employing luminosity distributed as follows: L_1 at $\sqrt{s} \simeq m_h$; $2.5L_1$ at $\sqrt{s} \simeq m_h + 2\sigma_{\sqrt{s}}$; $2.5L_1$ at $\sqrt{s} \simeq m_h - 2\sigma_{\sqrt{s}}$. The total luminosity for the scan is then $L = 6L_1$. This scan would simultaneously yield measurements of Γ_h^{tot} and of the $\mu^+\mu^- \rightarrow h \rightarrow b\bar{b}, WW^*$ and ZZ^* rates. Assuming that $L = 200 \text{ fb}^{-1}$ is devoted over a period of several years to this three point scan (presumably beginning with $\sqrt{s} \simeq m_h$), one obtains accuracies for the event rates as given in Table 1 and errors for Γ_h^{tot} and coupling-squared ratios as given in Table 2.

^bDetails of the strategy for this scan ¹ will not be discussed here; the scan is most efficient for small R (as desirable in any case) in which case \sqrt{s} must be reset rapidly with high precision.

Table 1. Errors for $\sigma(\mu^+\mu^- \rightarrow h_{SM})BF(h_{SM} \rightarrow b\bar{b}, WW^*, ZZ^*)$, for $R = 0.01\%$ and $L_{\text{scan}} = 200 \text{ fb}^{-1}$ (equivalent to $L_{\sqrt{s}=m_h} = 50 \text{ fb}^{-1}$).

Quantity	Errors				
$m_{h_{SM}}(\text{GeV})$	80	90	100	110	120
$\sigma(\mu\mu \rightarrow h_{SM})BF(b\bar{b})$	$\pm 0.2\%$	$\pm 1.6\%$	$\pm 0.4\%$	$\pm 0.3\%$	$\pm 0.3\%$
$\sigma(\mu\mu \rightarrow h_{SM})BF(WW^*)$	—	—	$\pm 3.5\%$	$\pm 1.5\%$	$\pm 0.9\%$
$\sigma(\mu\mu \rightarrow h_{SM})BF(ZZ^*)$	—	—	—	$\pm 34\%$	$\pm 6.2\%$
$m_{h_{SM}}(\text{GeV})$	130	140	150	160	170
$\sigma(\mu\mu \rightarrow h_{SM})BF(b\bar{b})$	$\pm 0.3\%$	$\pm 0.5\%$	$\pm 1.1\%$	$\pm 59\%$	—
$\sigma(\mu\mu \rightarrow h_{SM})BF(WW^*)$	$\pm 0.7\%$	$\pm 0.5\%$	$\pm 0.5\%$	$\pm 1.1\%$	$\pm 9.4\%$
$\sigma(\mu\mu \rightarrow h_{SM})BF(ZZ^*)$	$\pm 2.8\%$	$\pm 2.0\%$	$\pm 2.1\%$	$\pm 22\%$	$\pm 34\%$
$m_{h_{SM}}(\text{GeV})$	180	190	200	210	220
$\sigma(\mu\mu \rightarrow h_{SM})BF(WW^*)$	$\pm 18\%$	$\pm 38\%$	$\pm 58\%$	$\pm 79\%$	—
$\sigma(\mu\mu \rightarrow h_{SM})BF(ZZ^*)$	$\pm 25\%$	$\pm 27\%$	$\pm 35\%$	$\pm 45\%$	$\pm 56\%$

Table 2. Errors for coupling-squared ratios and $\Gamma_{h_{SM}}^{\text{tot}}$ for s -channel Higgs production at the FMC, assuming $L = 6L_1 = 200 \text{ fb}^{-1}$ total scan luminosity and $R = 0.01\%$.

Quantity	Errors			
$m_{h_{SM}}(\text{GeV})$	80	m_Z	100	110
$(WW^*h_{SM})^2/(b\bar{b}h_{SM})^2$	—	—	$\pm 3.5\%$	$\pm 1.6\%$
$(ZZ^*h_{SM})^2/(b\bar{b}h_{SM})^2$	—	—	—	$\pm 34\%$
$(ZZ^*h_{SM})^2/(WW^*h_{SM})^2$	—	—	—	$\pm 34\%$
$\Gamma_{h_{SM}}^{\text{tot}}$	$\pm 2.6\%$	$\pm 32\%$	$\pm 8.3\%$	$\pm 4.2\%$
$m_{h_{SM}}(\text{GeV})$	120	130	140	150
$(WW^*h_{SM})^2/(b\bar{b}h_{SM})^2$	$\pm 1\%$	$\pm 0.7\%$	$\pm 0.7\%$	$\pm 1\%$
$(ZZ^*h_{SM})^2/(b\bar{b}h_{SM})^2$	$\pm 6\%$	$\pm 3\%$	$\pm 2\%$	$\pm 2\%$
$(ZZ^*h_{SM})^2/(WW^*h_{SM})^2$	$\pm 6\%$	$\pm 3\%$	$\pm 2\%$	$\pm 2\%$
$\Gamma_{h_{SM}}^{\text{tot}}$	$\pm 3.6\%$	$\pm 3.6\%$	$\pm 4.1\%$	$\pm 6.5\%$

Very good errors in the $c\bar{c}$ and $\tau^+\tau^-$ channels might also be possible, depending on the detector. For example, $c\bar{c}$ isolation requires topological tagging in which one must distinguish primary, secondary, and tertiary vertices for a b -jet vs. primary and secondary only for a c -jet. The ability to do so depends on how close to the beam the first layer of the vertex detector can be placed.

What do the errors of Table 2 mean in terms of our ability to discriminate between the SM h_{SM} and the MSSM h^0 ? Plots ^{3,4} of the $WW^*/b\bar{b}$ coupling-squared ratio as computed in the MSSM divided by that computed in the SM show: a) there is almost no dependence of the ratio upon the squark mixing scenario; b) the ratio is essentially independent of $\tan\beta$ in the allowed portion of the standard $(m_{A^0}, \tan\beta)$ parameter space; and c) roughly -50% (-20%) deviations from the SM result are predicted for $m_{A^0} \sim 250 \text{ GeV}$ ($m_{A^0} \sim 400 \text{ GeV}$). By the time the muon collider is in operation, theoretical systematic errors (primarily from uncertainty in the running mass, $m_b(m_h)$)

in the ratio should be in the $\pm 5\%$ to $\pm 10\%$ range. Table 2 shows that statistical errors will be much smaller over most of the relevant m_h range. Thus, the $WW^*/b\bar{b}$ event rate ratio will distinguish between the SM and the MSSM at the $\geq 2\sigma$ level for $m_{A^0} \lesssim 400$ GeV. For $m_{A^0} < 400$ GeV, a rough determination of m_{A^0} will be possible. (This will be important for the H^0, A^0 discussion in the next section.) Similar results apply for the $ZZ^*/b\bar{b}$ ratio.

In contrast to the event rate ratios, deviations in Γ_h^{tot} in the MSSM vs. the SM as a function of $(m_{A^0}, \tan \beta)$ are very dependent upon the squark mixing scenario, the amount of SUSY decays present, and so forth; large deviations in Γ_h^{tot} from SM expectations are the rule, but do not pin down either m_{A^0} or $\tan \beta$. However, if one ⁴ uses the very accurate direct determination of Γ_h^{tot} from the FMC scan in combination with other measurements performed with $L = 200 \text{ fb}^{-1}$ and $\sqrt{s} = 500$ GeV at the NLC ^c a large variety of very important coupling-squared magnitudes can be extracted. The ratio of the MSSM prediction to the SM prediction for a squared coupling is always very squark-mixing independent and gives new opportunities for determining m_{A^0} . As an example, there are four ways to determine $\Gamma(h \rightarrow \mu^+ \mu^-)$ by combining NLC and FMC data: ^d $\Gamma(h \rightarrow \mu^+ \mu^-) =$

$$\begin{aligned} 1) & \frac{[\Gamma(h \rightarrow \mu^+ \mu^-) BF(h \rightarrow b\bar{b})]_{\text{FMC}}}{BF(h \rightarrow b\bar{b})_{\text{NLC}}}; & 2) & \frac{[\Gamma(h \rightarrow \mu^+ \mu^-) BF(h \rightarrow WW^*)]_{\text{FMC}}}{BF(h \rightarrow WW^*)_{\text{NLC}}}; \\ 3) & \frac{[\Gamma(h \rightarrow \mu^+ \mu^-) BF(h \rightarrow ZZ^*) \Gamma_{h_{SM}}^{\text{tot}}]_{\text{FMC}}}{\Gamma(h \rightarrow ZZ^*)_{\text{NLC}}}; & 4) & \frac{[\Gamma(h \rightarrow \mu^+ \mu^-) BF(h \rightarrow WW^*) \Gamma_{h_{SM}}^{\text{tot}}]_{\text{FMC}}}{\Gamma(h \rightarrow WW^*)_{\text{NLC}}}. \end{aligned}$$

In the above, $\Gamma_h^{\text{tot}}|_{\text{FMC}}$ refers to the scan measurement at the FMC. Resulting errors for $h = h_{SM}$ are tabulated in Table 3, labelled $(\mu^+ \mu^- h_{SM})^2|_{\text{NLC+FMC}}$. The predicted $(\mu^+ \mu^- h_{SM})^2|_{\text{MSSM}}/(\mu^+ \mu^- h_{SM})^2|_{\text{SM}}$ coupling-squared ratio is independent of $\tan \beta$: one finds values of 1.5, 1.2, 1.15 for $m_{A^0} = 300, 475, 600$ GeV, respectively (for theoretically allowed $\tan \beta$ values at the given m_{A^0}). Since there is no systematic error due to uncertainty in the muon mass and since experimental systematics should be well below 10%, the above numbers imply that the expected $\lesssim \pm 5\%$ statistical error probes out to very high m_{A^0} .

Table 3 gives a number of other quantities that are determined with remarkable precision by combining $\sqrt{s} = 500$ GeV NLC data and s -channel FMC data [and, in the case of $(\gamma\gamma h_{SM})^2$, including the (NLC+LHC) determination of $BF(h_{SM} \rightarrow \gamma\gamma)$]. ⁴ The advantages of having high L data from all three machines are enormous. In particular, if there is a SM-like Higgs bosons with $m_h \lesssim 2m_W$, it would be much more beneficial to have both an e^+e^- collider operating at full energy, $\sqrt{s} \sim 500$ GeV, and a $\mu^+ \mu^-$ collider

^cThe same measurements are also possible at an FMC, but we use NLC notation in what follows.

^dNote that since $\sigma_{\sqrt{s}} \sim 2 - 3 \times \Gamma_h^{\text{tot}}$, the measured $\mu^+ \mu^- \rightarrow h \rightarrow X$ rate is more or less proportional to $\Gamma(h \rightarrow \mu^+ \mu^-)/\sigma_{\sqrt{s}}$ so that $\Gamma(h \rightarrow \mu^+ \mu^-)$ can be computed given the known $\sigma_{\sqrt{s}}$; small corrections from the influence of Γ_h^{tot} can be made using the measured Γ_h^{tot} .

Table 3. Errors for combining NLC ($L = 200 \text{ fb}^{-1}$) data, the NLC + LHC ($L = 600 \text{ fb}^{-1}$, ATLAS+CMS) $BF(h_{SM} \rightarrow \gamma\gamma)$ determination, $\gamma\gamma$ collider ($L = 50 \text{ fb}^{-1}$) data and FMC s -channel ($L = 200 \text{ fb}^{-1}$) data.

Quantity	Errors			
$m_{h_{SM}}(\text{GeV})$	80	100	110	120
$(b\bar{b}h_{SM})^2 _{\text{NLC+FMC}}$	$\pm 6\%$	$\pm 9\%$	$\pm 7\%$	$\pm 6\%$
$(c\bar{c}h_{SM})^2 _{\text{NLC+FMC}}$	$\pm 9\%$	$\pm 11\%$	$\pm 10\%$	$\pm 9\%$
$(\mu^+\mu^-h_{SM})^2 _{\text{NLC+FMC}}$	$\pm 5\%$	$\pm 5\%$	$\pm 4\%$	$\pm 4\%$
$(\gamma\gamma h_{SM})^2 _{\text{FMC}}$	$\pm 15\%$	$\pm 16\%$	$\pm 14\%$	$\pm 13\%$
$(\gamma\gamma h_{SM})^2 _{\text{NLC+FMC}}$	$\pm 9\%$	$\pm 10\%$	$\pm 9\%$	$\pm 9\%$
$m_{h_{SM}}(\text{GeV})$	130	140	150	170
$(b\bar{b}h_{SM})^2 _{\text{NLC+FMC}}$	$\pm 7\%$	$\pm 7\%$	$\pm 10\%$	$\pm 23\%$
$(c\bar{c}h_{SM})^2 _{\text{NLC+FMC}}$	$\pm 10\%$?		
$(\mu^+\mu^-h_{SM})^2 _{\text{NLC+FMC}}$	$\pm 3\%$	$\pm 3\%$	$\pm 4\%$	$\pm 10\%$
$(WW^*h_{SM})^2 _{\text{FMC}}$	$\pm 16\%$	$\pm 9\%$	$\pm 9\%$	—
$(WW^*h_{SM})^2 _{\text{NLC+FMC}}$	$\pm 5\%$	$\pm 4\%$	$\pm 6\%$	$\pm 10\%$
$(\gamma\gamma h_{SM})^2 _{\text{FMC}}$	$\pm 14\%$	$\pm 18\%$	$\pm 36\%$	—
$(\gamma\gamma h_{SM})^2 _{\text{NLC+FMC}}$	$\pm 10\%$	$\pm 13\%$	$\pm 23\%$	—

operating at $\sqrt{s} \sim m_h$, as opposed to two NLC's.

3. The Heavy MSSM Higgs Bosons

Colliders other than the FMC offer various mechanisms to directly search for the A^0, H^0 , but have significant limitations:

- There are regions in $(m_{A^0}, \tan\beta)$ parameter space at moderate $\tan\beta$, $m_{A^0} \gtrsim 200 \text{ GeV}$ in which the H^0, A^0 cannot be detected at the LHC.
- At the NLC one can use the mode $e^+e^- \rightarrow Z^* \rightarrow H^0 A^0$, but it is limited to $m_{H^0} \sim m_{A^0} \lesssim \sqrt{s}/2$.
- A $\gamma\gamma$ collider could probe heavy Higgs up to masses of $m_{H^0} \sim m_{A^0} \sim 0.8\sqrt{s}$, but this would quite likely require $L > 100 \text{ fb}^{-1}$, especially if the Higgs bosons are at the upper end of the $\gamma\gamma$ collider energy spectrum.⁸

In contrast, there is an excellent chance of being able to detect the H^0, A^0 at a $\mu^+\mu^-$ collider provided only that m_{A^0} is smaller than the maximal \sqrt{s} available. This could prove to be very important given that GUT MSSM models usually predict $m_{A^0} \gtrsim 200 \text{ GeV}$.

A detailed study of s -channel production of the H^0, A^0 has been made.¹ The signals become viable when $\tan\beta > 1$ (as favored by GUT models) since the $\mu^+\mu^-H^0$ and $\mu^+\mu^-A^0$ couplings are proportional to $\tan\beta$. In particular, even though $\Gamma_{H^0}^{\text{tot}}, \Gamma_{A^0}^{\text{tot}}$ are big (see Fig. 2) at high $\tan\beta$, due to large $b\bar{b}$ decay widths, $BF(H^0, A^0 \rightarrow \mu^+\mu^-)$ approaches a constant value that is large enough to imply substantial $\bar{\sigma}_{H^0}, \bar{\sigma}_{A^0}$. The optimal strategy

for H^0, A^0 detection and study depends upon the circumstances. First, it could be that the H^0 and/or A^0 will already have been discovered at the LHC. With $L = 300 \text{ fb}^{-1}$ (ATLAS+CMS) of integrated luminosity, this would be the case if $\tan \beta \lesssim 3$ or $\tan \beta$ is above an m_{A^0} -dependent lower bound (*e.g.* $\tan \beta \gtrsim 10$ for $m_{A^0} \sim 400 \text{ GeV}$).^e Even if the H^0, A^0 have not been detected, strong constraints on m_{A^0} are possible if precision measurements of the properties of the h^0 (such as the $b\bar{b}/WW^*$ and $c\bar{c}/b\bar{b}$ event rate ratios and the $(\mu^+\mu^-h^0)^2$ coupling-squared, as discussed earlier) are made via s -channel production at the FMC or in $\sqrt{s} = 500 \text{ GeV}$ running at the NLC⁴ or by combining these two types of data. By limiting the \sqrt{s} scan for the H^0 and A^0 in the s -channel to the $m_{A^0} \sim m_{H^0}$ mass region preferred by h^0 measurements, we would greatly reduce the luminosity needed to find the A^0 and H^0 via an s -channel scan as compared to that required if m_{A^0} is not constrained.

With such pre-knowledge of m_{A^0} , it will be possible to detect and perform detailed studies of the H^0, A^0 for all $\tan \beta \geq 1$ provided only that $m_{A^0} \lesssim \sqrt{s}_{\text{max}}$.^f If $\tan \beta \lesssim 3$ and $m_{H^0}, m_{A^0} \lesssim 2m_t$, then excellent resolution, $R \sim 0.01\%$, will be necessary for detection since the A^0 and H^0 are relatively narrow (see Fig. 2). For higher $\tan \beta$ values, $R \sim 0.1\%$ is adequate for H^0, A^0 detection, but $R \sim 0.01\%$ would be required in order to separate the rather overlapping H^0 and A^0 peaks (as a function of \sqrt{s}) from one another.

Even without pre-knowledge of m_{A^0} , there would be an excellent chance for discovery of the A^0, H^0 Higgs bosons in the s -channel at a $\mu^+\mu^-$ collider if they have not already been observed at the LHC. This is because non-observation at the LHC implies that $\tan \beta \gtrsim 3$ while it is precisely for $\tan \beta \gtrsim 2.5 - 3$ that detection of the A^0, H^0 is possible¹ in the mass range from 200 to 500 GeV via an s -channel scan in $\mu^+\mu^-$ collisions. (The lower $\tan \beta$ reach given assumes that $L_{\text{tot}} = 200 \text{ fb}^{-1}$ is devoted to the scan. A detailed strategy as to how much luminosity to devote to different \sqrt{s} scan settings in the 200 – 500 GeV range must be employed.¹) That the LHC and the FMC are complementary in this respect is a very crucial point. Together, the LHC and FMC guarantee discovery of the A^0, H^0 after 3 to 4 years of high luminosity operation each, provided $m_{A^0} \lesssim 500 \text{ GeV}$. Once m_{A^0}, m_{H^0} are known, very precise measurements of some of the crucial properties of the H^0, A^0 (including a scan determination of their total widths) become possible.¹

In the event that the NLC has not been constructed, it could be that the first mode of operation of the FMC would be to optimize for and accumulate luminosity at, say, $\sqrt{s} = 500 \text{ GeV}$. In this case, there is still a high probability for detecting the H^0, A^0 if they have not been observed at the LHC. First, if $m_{A^0} \sim m_{H^0} \lesssim \sqrt{s}/2 \sim 250 \text{ GeV}$ then $\mu^+\mu^- \rightarrow H^0 A^0$ (and $H^+ H^-$) pair production will be observed. Second, although reduced in magnitude compared to an electron collider, there is a long low-energy bremsstrahlung

^eFor $\tan \beta \lesssim 3$, one makes use of modes such as $H^0 \rightarrow h^0 h^0 \rightarrow b\bar{b}\gamma\gamma$ and $H^0 \rightarrow ZZ^{(*)} \rightarrow 4\ell$, when $m_{H^0} \lesssim 2m_t$, or $H^0, A^0 \rightarrow t\bar{t}$, when $m_{H^0}, m_{A^0} \gtrsim 2m_t$. At high $\tan \beta$, the enhanced production rates for $b\bar{b}H^0, b\bar{b}A^0$ with $H^0, A^0 \rightarrow \tau^+\tau^-$ are employed.

^fWe assume that a final ring optimized for maximal luminosity at $\sqrt{s} \sim m_{A^0}$ would be constructed.

tail at a muon collider that provides a self-scan over the full range of \sqrt{s} values below the nominal operating energy. Observation of A^0, H^0 s -channel peaks in the $b\bar{b}$ mass ($m_{b\bar{b}}$) distribution created by this bremsstrahlung tail may be possible. The region of the $(m_{A^0}, \tan\beta)$ parameter space plane for which a peak is observable depends strongly on the $m_{b\bar{b}}$ resolution. For excellent $m_{b\bar{b}}$ resolution of order ± 5 GeV and integrated luminosity of $L = 200 \text{ fb}^{-1}$ at $\sqrt{s} = 500$ GeV, the A^0, H^0 peak(s) are observable for $\tan\beta \gtrsim 4 - 5$ if $500 \text{ GeV} \geq m_{A^0} \geq 250 \text{ GeV}$.⁹

Finally, if neither the LHC nor a FMC scan of the ≤ 500 GeV region has discovered the H^0, A^0 , but supersymmetric particles and the h^0 have been observed, we would believe that the H^0, A^0 must exist but have $m_{A^0} \sim m_{H^0} \geq 500$ GeV. Analyses of the SUSY spectrum in the GUT context and precision h^0 studies might have yielded some prejudice for the probable m_{A^0} , and an extension of the FMC energy up to the appropriate $\sqrt{s} \sim m_{A^0}$ for s -channel discovery of the H^0, A^0 could be considered. But, a machine with much higher \sqrt{s} , such as the earlier-mentioned $\sqrt{s} = 4$ TeV, might be most worthwhile. It has been shown⁵ that such an energy with appropriately matched luminosity would allow discovery of $\mu^+\mu^- \rightarrow A^0 H^0$ and $H^+ H^-$ pair production, via the $b\bar{b}$ or $t\bar{t}$ decay channels of the H^0, A^0 and $t\bar{b}, \bar{t}b$ decay channels of the H^+, H^- , up to masses very close to $m_{A^0} \sim m_{H^0} \sim m_{H^\pm} \sim 2$ TeV, even if SUSY decays of the H^0, A^0, H^\pm are substantial. (This mass range certainly includes that expected in any supersymmetric model that provides a solution to the naturalness and hierarchy problems.) Detailed studies of the H^0, A^0, H^\pm would be possible once they were discovered.^{5,6}

4. Conclusions and Discussion

In this report, we have briefly reviewed the capabilities of a muon collider to explore Higgs physics in the Standard Model and its minimal supersymmetric extension via direct s -channel Higgs production. If there is a light ($m_h \lesssim 2m_W$) SM-like Higgs boson, an s -channel scan of the Higgs resonance peak provides direct measurements of a number of its important properties (in particular Γ_h^{tot}) that cannot be duplicated using e^+e^- (or $\mu^+\mu^-$) collisions at high \sqrt{s} . Thus, if a light SM-like Higgs boson has already been observed at the LHC or in early operation at the first NLC, it would be much more useful to follow the first NLC with an FMC, rather than a second NLC. In the context of the minimal supersymmetric extension of the SM, the FMC takes on added value. At the FMC, an appropriately designed s -channel scan will allow H^0, A^0 discovery in the $200 \lesssim m_{A^0} \lesssim 500$ GeV range for all $\tan\beta \gtrsim 3$. Thus, if $H^0 A^0$ pair production is not seen in $\sqrt{s} = 500$ GeV running at the NLC (or FMC), implying $m_{A^0} \gtrsim 200 - 240$ GeV, nor at the LHC, implying $\tan\beta \gtrsim 3$ (and below an m_{A^0} -dependent upper limit) if $m_{A^0} \gtrsim 200$ GeV, the H^0, A^0 will be observed at the FMC (provided $m_{A^0} \lesssim 500$ GeV). Once discovered (at any collider), the FMC s -channel production mode allows (for any

⁹Required $\tan\beta$ values increase dramatically as one moves into the $m_{A^0} \sim m_Z$ zone, but this region is covered by $H^0 A^0$ pair production.

$\tan\beta > 1$) a detailed study of some of their key properties and a scan determination of their total widths. As reviewed elsewhere,^{1,4,3,7} a muon collider will be at least as valuable if the Higgs sector is still more exotic than the constrained two-doublet MSSM sector. Particularly noteworthy is the ability of $\mu^-\mu^-$ collisions to probe the $\mu^-\mu^-$ coupling of a doubly-charged Higgs boson^h (as present in many Higgs triplet models) down to an extraordinarily small coupling magnitude. Overall, if there are elementary Higgs bosons, a muon collider will almost certainly be mandated purely on the basis of its ability to explore the Higgs sector via s -channel factory-like production of Higgs bosons.

5. Acknowledgements

I would like to acknowledge the many contributions of my collaborators, V. Barger, M. Berger, and T. Han, to the muon collider Higgs physics results. This work was supported in part by the Department of Energy and by the Davis Institute for High Energy Physics.

6. References

1. V. Barger, M. Berger, J.F. Gunion and T. Han, UCD-96-6 (January, 1996), to appear in Phys. Rep.
2. See, for example, R.B. Palmer, A. Sessler, and A. Tollestrup, BNL-63245 and BNL-63865.
3. See J.F. Gunion, A. Stange, and S. Willenbrock, *Weakly-Coupled Higgs Bosons*, preprint UCD-95-28 (1995), to be published in *Electroweak Physics and Beyond the Standard Model*, World Scientific Publishing Co., eds. T. Barklow, S. Dawson, H. Haber, and J. Siegrist, and references therein.
4. J.F. Gunion, L. Poggioli and R. Van Kooten, to appear in the Proceedings of the 1996 DPF/DPB Summer Study on New Directions for High-Energy Physics (denoted below by Snowmass96), and references therein.
5. J.F. Gunion and J. Kelly, hep/ph-9610421, to appear in Snowmass96.
6. J. Feng and T. Moroi, hep/ph-9612333.
7. J.F. Gunion, to appear in *Perspectives on Higgs Physics*, 2nd edition, ed. G. Kane, World Scientific Publishing.
8. J.F. Gunion and H.E. Haber, Phys. Rev. **D48**, 5109 (1993).
9. J.F. Gunion, Int. J. Mod. Phys. **A11**, 1551 (1996).
10. J.F. Gunion, C. Loomis, and K.T. Pitts, hep/ph-9610237, to appear in Snowmass96.

^hSuch a Higgs is very likely to be seen at the LHC if it has mass below 1 TeV.¹⁰