

UM {P {97/19

To appear in the American Journal of Physics

Classical antiparticles

John P. Costella, Bruce H. J. McKellar,^y and Andrew A. Rawlinson^z
 School of Physics, The University of Melbourne, Parkville, Victoria 3052, Australia

(29 March 1997)

Abstract

We review how antiparticles may be introduced in classical relativistic mechanics, and emphasize that many of their paradoxical properties can be more transparently understood in the classical than in the quantum domain.

I. Introduction

Recently,¹ we reviewed briefly the physics and early history of the Foldy-Wouthuysen transformation,^{2;3} emphasizing that the transformed representation is the only one in which a classical limit of the Dirac equation can be meaningfully extracted, in terms of particles and antiparticles. But few textbooks actually describe how antiparticles can be dealt with in classical mechanics. Discussions of antiparticles usually begin with the "negative energy problem": the inevitable introduction, in relativistic mechanics, of what appears to be a "spurious" set of mirror eigenstates of negative energy; their reinterpretation by Dirac as "holes" in a filled Fermi sea of vacuum electrons; and their further reformulation, in quantum field theory, as completely valid eigenstates in their own right. But this introduction is altogether too late: while its appearance in a course on relativistic quantum mechanics reflects accurately the historical development of the theory of antiparticles, it can tend to hide completely the fact that it is relativistic mechanics itself that makes possible the phenomenon of antiparticle motion | quantum mechanics is by no means a prerequisite.

Arguably, a thorough preliminary understanding of the classical theory of antiparticles better equips the student for tackling the same issues when they arise in relation to the Dirac equation. It is this topic that we shall review in this paper.

II. The proper time

Consider a structureless point particle. Classically, its kinematical state at any time t consists simply of the three components of its three-position $\mathbf{z}(t)$. We assume that $\mathbf{z}(t)$ is a continuous function of t that is sufficiently differentiable for our purposes. In special relativity, we form the four-position z of the particle:

$$z = (t; \mathbf{z});$$

where we shall always use units in which $c = 1$. The continuous function $z(t)$ specifies the path of the particle in Minkowski spacetime its worldline. To parameterize its "length", in a Lorentz-invariant way, we consider an infinitesimal differential element of the path,

$$dz(t) = (dt; dz(t));$$

where $dz(t)$ is the infinitesimal change in position $z(t)$ in the infinitesimal time interval from t to $t + dt$. We now consider the Lorentz-invariant quantity

$$d^2(t) = dz(t)dz(t) = dt^2 - dz^2(t); \quad (1)$$

where we employ a $(+; -; -; -)$ metric. What we would like to do is define a quantity $d(t)$ that would provide a measure of "length" along the worldline. But the Lorentz-invariant expression (1) involves not d , but rather the square of d . Thus, d can only be defined up to a sign:

$$d = \sqrt{\pm dz dz} : \quad (2)$$

To investigate the meaning of this ambiguity in the sense of d , let us consider the special case in which the particle is instantaneously at rest, with respect to our own inertial coordinate system:

$$dz = 0:$$

In this case, we find

$$d = dt:$$

The solution $d = dt$ for a particle at rest is the one usually presented in introductory texts on special relativity: such a d is obviously equal to the passage of time as measured in the instantaneous rest frame of the particle. For a particle undergoing arbitrary relativistic motion, we assume that the particle itself possesses its own "cumulative time" or "age", which we term the proper time, that can be calculated by summing up all of the d along its worldline:

$$(E) = \int_{E_0}^E d; \quad (E)$$

where (E) is the proper time at event E on the worldline, and where the event E_0 on the worldline defines the (arbitrary) origin of (t) . Since the worldline of any classical particle passes through each constant- t hyperplane once and only once, we can replace the events E and E_0 by their corresponding coordinate times t and t_0 , and hence determine (t) as a function of t :

$$(t) = \int_{t_0}^t dt^0 \frac{d}{dt^0} = \int_{t_0}^t dt^0 \frac{1}{(t^0)}; \quad (3)$$

where we have made use of Eq. (1):

$$\frac{d}{dt}(t) = \sqrt{1 - \frac{v^2(t)}{c^2}} = \frac{1}{\gamma(t)}:$$

The standard textbook result (3) shows that when the speed v of the particle is much smaller than the speed of light, the factor $\gamma(t)$ is close to unity, and the passage of proper time is indistinguishable from that of coordinate time; but if the particle's motion is such that its speed rises to an appreciable fraction of the speed of light, the factor $\gamma(t)$ rises above unity, and the particle "ages" more slowly. In all cases, however, the particle gets older: special relativity only seems to modify the rate; it warps our view of the world, but it does not throw it into reverse.

III. Classical antiparticles

Let us now consider the other solution in Eq. (2) for a particle at rest, namely,

$$d = -dt; \quad (4)$$

Even to a student possessing a good knowledge of special relativity, Eq. (4) does not look familiar at all. It seems to imply that a particle at rest with respect to our Lorentz coordinate system might somehow believe that time evolves in the opposite direction to what we do! For example, if we determine that some spacetime event E^0 is definitely earlier than another event E^0 (i.e., E^0 lies within the backward lightcone of E^0), then a particle whose own "proper time" obeys (4) would insist, to the contrary, that E^0 is definitely later than E^0 (i.e., from the particle's point of view, E^0 lies within the forward lightcone of E^0).

The problem is that, by the principles of relativity, such a particle is just as valid an observer of the universe as we are: it agrees with us that the speed of light is unity in all inertial frames. We have no physically acceptable justification for dismissing its counter-intuitive view of the world. We must conclude that both of the solutions (2) are equally valid definitions of the passage of proper time. (This is analogous to the fact⁴ that both the retarded and the advanced Lienard-Wiechert potentials for a point charge are equally valid solutions of Maxwell's equations.)

Armed with a thorough knowledge of relativistic quantum mechanics and quantum field theory, Stueckelberg⁵ and Feynman^{6;7;8} made the following realization: a particle for which d evolves in the opposite sense to the dt in our particular Lorentz frame of reference is simply in antiparticle motion with respect to us. Of course, there are no classical forces that can change "particle motion" into "antiparticle motion" — the two regimes are as disjoint as the interiors of the forward and backward lightcones; but, even classically, this does not bar the possibility that a particle might have always been in antiparticle motion.

Following this argument to its logical conclusion, it could be noted that there is a similar ambiguity of sign when parameterizing path lengths in Euclidean space, since there the invariant interval is also squared:

$$d\ell^2 = dz^2;$$

and hence we could equally well measure length one way along the path, or in the opposite way. But we are already used to the idea that, at a fundamental level, traveling to the left is no more difficult than traveling to the right. The crucial difference in Minkowski space is precisely the fact that classical forces do not reverse the sense in which "time is traversed"; our intuition with Galilean mechanics is rooted firmly in the belief that everyone agrees on

the direction that time is traveling. Relativistically boosting to another frame of reference "warps" the rate at which clocks tick, but it does not reverse it; in contrast, time-reversal is a discrete symmetry, and cannot be brought into contact with "intuitive" physics by a continuous transformation.

IV. C, P, and T

Another way of recognizing the possibility of the existence of antiparticle motion, in any relativistically complete theory of mechanics, is to consider the fundamental symmetries of the Lorentz group — namely, those transformations under which the interval (1) is invariant. Intuitive, introductory constructions of the proper time generally make use of everyday objects, such as people, trains, measuring rods, clocks, and so on. In thinking about such everyday objects — even in relativistic terms — we usually only consider proper Lorentz transformations (boosts and rotations) — or, at most, orthochronous ones (proper transformations with or without the parity transformation). But the interval (1) is also invariant under non-orthochronous Lorentz transformations — those involving the time-reversal operation — and it is precisely such transformations that convert what appears to be "normal" particle motion into "antiparticle" motion.

Let us make this argument more concrete. Consider the parity operation, which in classical physics simply changes the sign of the spatial coordinates in a given Lorentz frame:

$$P : x \rightarrow -x :$$

The time-reversal operation does likewise for the time coordinate:

$$T : t \rightarrow -t :$$

Under the combined operations of P and T , all four components of dz are reversed in sign:

$$PT : dz \rightarrow -dz : \quad (5)$$

The three-velocity, being a ratio of the spatial part dz to the temporal part dz^0 , is therefore unchanged:

$$PT : v = \frac{dz}{dt} \rightarrow v :$$

Let us now try to define proper time so that the sign of d is always taken in the same sense as dt . Consider a free particle, that has a three-velocity v in a given Lorentz frame of reference; and let us define

$$d = + \frac{dt}{\gamma}; \quad (6)$$

where we always define γ as the positive square-root:

$$(v) \quad \gamma = \frac{1}{\sqrt{1 - v^2}} :$$

The choice of sign (6) lets us label the worldline with values of τ in the "standard" way, such that the τ values increase in the direction of increasing coordinate time t .

If we now apply the operation PT to the above Lorentz frame, we are placed in a new, equally valid Lorentz frame, in which all directions of space and time have been reversed. But this operation does not affect our markings on the particle's worldline, since the proper time is a property of the particle itself. Thus, with respect to the new Lorentz frame, τ increases in the direction of decreasing coordinate time t :

$$d\tau = -\frac{dt}{\gamma}.$$

Thus, if we insist on the invariance of classical mechanics under the complete Lorentz group (as we do in all other forms of relativistic mechanics), we find that for every possible solution of the equations of motion with the choice of the positive sign in Eq. (2), there exists an equally possible solution in which the negative sign is chosen. By the Stueckelberg-Feynman interpretation, this transformation is the classical particle-antiparticle (or "charge-conjugation") transformation, so let us label it as such:

$$C : \tau \rightarrow -\tau. \quad (7)$$

Let us denote by η the choice of sign in Eq. (2): $\eta = +1$ if the particle is in "normal particle" motion with respect to our own Lorentz frame (i.e., $d\tau = dt > 0$); whereas $\eta = -1$ if the particle is in "antiparticle" motion (i.e., $d\tau = dt < 0$). (The symbol η is sometimes used in introductory texts for the ratio v/c , but in natural units it is simpler to just use the intuitive symbol v for this latter quantity. It is to maintain consistency with the results of the Foldy-Wouthuysen transformation^{1,3} that we use the symbol η for the classical particle-antiparticle number.)

With the definition (7), classical mechanics possesses all three discrete symmetry operations C , P , and T required for a relativistically invariant system of mechanics. Classically, all of these operations commute, and each of them individually squares to unity. We expect that the equations of motion of classical physics will be invariant under the combined operation CPT .

V. Antiparticles in the real world

Let us now show that the classical C operation yields "antiparticles" in the everyday sense of the word, i.e., that the antiparticle of an electron is a positron, and so on. To do so, it suffices to give our classical point particle two characteristics: an electric charge q , and a mass m . We assume that the quantities q and m are Lorentz scalars, and are unchanged under any of the operations C , P , or T , as defined above. (We shall show how the usual interpretation of C as changing the sign of the "effective" charge is to be understood shortly.)

Let us first consider the electromagnetic interaction. This is a vector interaction, which couples to the electromagnetic vector current density of the point charge,⁴

$$j^\mu(x) = q \int_{-\infty}^{\infty} d\tau \, \dot{x}^\mu(\tau) \delta^4(x - z(\tau)) u(\tau); \quad (8)$$

where $u^\mu(\tau)$ is the four-velocity of the particle:

$$u^\mu = \frac{dz^\mu}{d\tau} = (\gamma; \gamma \mathbf{v}) : \quad (9)$$

The delta function and τ -integration in (8) are necessary to obtain a current density from the trajectory of the point particle (the density of a point particle being infinite on its worldline and zero outside); but the essential properties of $j^\mu(\mathbf{x})$ are contained in the classical electromagnetic current vector,

$$J^\mu = q u^\mu : \quad (10)$$

From Eq. (9) we see that u^μ is both C-odd and PT-odd, since dz^μ is C-even and PT-odd, whereas $d\tau$ is by definition C-odd and PT-even. Thus, since q is assumed to be unchanged by C or PT as we have defined them, J^μ is also C-odd and PT-odd. (The density $j^\mu(\mathbf{x})$ clearly possesses the same symmetries as J^μ , since the delta function in (8) is effectively an even function of its argument, and under the C operation the sign of $d\tau$ is reversed, but so too are the limits of integration.) The former property is of particular importance for us:

$$C : J^\mu \rightarrow -J^\mu : \quad (11)$$

The result (11) tells us that, as far the electromagnetic interaction is concerned, antiparticle ($\eta = -1$) motion of a charged particle appears the same as an "equivalent normal particle", with $\eta = +1$, with the same three-velocity $\mathbf{v}$ as the original particle, but with the opposite "effective" charge, since

$$J^\mu = q \frac{dz^\mu}{d\tau} = (-q) \frac{dz^\mu}{d(-\tau)} :$$

For example, a particle of charge q , at rest, but in antiparticle motion, generates a static Coulomb field that is equivalent to that from a "normal" particle at rest of charge $-q$. It is in this sense of replacing antiparticle motion by an "equivalent normal particle" with an opposite "effective" charge that the classical particle-antiparticle operation C is a "charge conjugation" operation.

Let us now determine the mass of the "equivalent normal particle" corresponding to antiparticle motion. A priori, it may not be clear how we can make such a determination. However, if we believe Einstein's theory of general relativity (which we do here), then the equivalence principle tells us that "inertial" mass and "gravitational" mass are one and the same thing: they both represent the coupling of matter to spacetime. Thus, we can simply carry out the same analysis as we performed above for the electromagnetic interaction, but now with regard to the gravitational interaction.

Gravitation is a tensor interaction, which couples to the mechanical stress-energy tensor current density of our point particle:

$$t_{\mu\nu}(\mathbf{x}) = m \int_{-1}^1 d\tau \int d^3x \delta^4(\mathbf{x} - \mathbf{z}(\tau)) u_\mu(\tau) u_\nu(\tau) : \quad (12)$$

Again, the delta function and integration over τ and $\mathbf{x}$ are required for the purposes of converting a pointlike trajectory into a density; the essential properties of $t_{\mu\nu}(\mathbf{x})$ are contained in the classical mechanical current tensor,

$$T_{\mu\nu} = m u_\mu u_\nu : \quad (13)$$

In this case, we find that T is C-even, since it contains two factors of u :

$$C : T \rightarrow -T :$$

Thus, as far as the gravitational interaction is concerned, a particle of mass m in antiparticle motion ($\epsilon = -1$) is indistinguishable from the same particle of mass m in the corresponding particle motion ($\epsilon = +1$). For example, the gravitational field of a star made of antimatter is the same as that of an identical star in which the antimatter is replaced by normal matter; a collection of such stars would all attract each other gravitationally. By the equivalence principle, this invariance of the mass of the "equivalent normal particle" under the C operation is true in full generality.

The above examples have concentrated on the fields generated by particles in antiparticle motion, but the same conclusions can be drawn from the equations of motion for the particles under the influence of given external fields. In the gravitational case, the mass m actually drops out of the equations of motion (again, by the equivalence principle), and the equation of motion is simply the geodesic equation,⁹

$$\frac{du}{d\tau} = -\Gamma_{\mu\nu}^{\lambda} u^{\mu} u^{\nu} : \quad (14)$$

If the particle is in antiparticle motion, we can again replace it by an "equivalent normal particle": since

$$\frac{du}{d\tau} = -\Gamma_{\mu\nu}^{\lambda} \frac{dz^{\mu}}{d\tau} \frac{dz^{\nu}}{d\tau} = -\Gamma_{\mu\nu}^{\lambda} \left(\frac{dz^{\mu}}{d\tau} \right)^2 \quad (15)$$

and

$$u^{\lambda} \frac{dz^{\mu}}{d\tau} = \frac{dz^{\lambda}}{d\tau} \frac{dz^{\mu}}{d\tau} ; \quad (16)$$

we can rewrite the geodesic equation (14) in the form

$$\frac{d^2 z^{\lambda}}{d\tau^2} = -\Gamma_{\mu\nu}^{\lambda} \frac{dz^{\mu}}{d\tau} \frac{dz^{\nu}}{d\tau} ;$$

which shows that the "equivalent normal particle" acts in the same way as any other particle.

In the electromagnetic case, the equation of motion is the Lorentz force law, which in relativistic form is

$$\frac{du}{d\tau} = \frac{q}{m} F_{\mu\nu} u^{\nu} ; \quad (17)$$

where $F_{\mu\nu}$ represents an external electromagnetic field. Again, if the particle is in antiparticle motion, we can use the properties (15) and (16) to rewrite Eq. (17) in the form

$$\frac{d^2 z^{\lambda}}{d\tau^2} = \frac{q}{m} F_{\mu\nu} \frac{dz^{\mu}}{d\tau} \frac{dz^{\nu}}{d\tau} :$$

Hence the "equivalent normal particle" has the sign of its effective charge-to-mass ratio negated compared to that of the actual particle; and since we have already determined that

its mass is unchanged, we conclude that it is its "effective charge" that changes sign, in agreement with our analysis above.

V I. The "negative energy problem"

Finally, let us discuss a subtlety that is a frequent source of confusion for the student: the existence and interpretation of an energy-momentum four-vector for antiparticles. The subject arises most naturally when we consider the Lagrangian and Hamiltonian formulations of mechanics (which are of course of vital importance in the construction of a quantum mechanical description); but in introductory courses and textbooks it is often presented in a somewhat confusing and contradictory manner.

Let us continue to consider our classical point particle of mass m and charge q . Relativistically, the construction of a manifestly covariant set of generalized coordinates is somewhat delicate:^{4,10} we would like to treat all four components z^μ of the position four-vector of the particle in an equal fashion, with the "time" parameter preferentially given by the Lorentz-invariant proper time τ . But the existence of the definition (1) tells us that only three of the components of z^μ are actually independent of τ ; and the Lagrangian and Hamiltonian formulations of mechanics are greatly complicated if all of the generalized coordinates are not actually independent.¹⁰

One way to circumvent these problems is to take the "time" parameter of the Lagrangian or Hamiltonian formalism to be the "time", where

$$d\tau = \frac{d\tau}{\gamma} : \quad (18)$$

The quantity γ is a Lorentz scalar, since both dt and m are Lorentz scalars; and since m is even under C, P , and T , the quantity γ has the same symmetry properties as t ; in particular,

$$C : \quad t \rightarrow -t : \quad$$

To show that the simple redefinition (18) solves the independence problem, one need simply note that the mass (rest-energy) $m(\tau)$ of a general system need not be a constant of the motion. Thus, while one of the four components of z^μ is still differentially dependent on τ through Eq. (1), all four components of z^μ are, in general, independent of τ , since the m in Eq. (18) may vary as a function of τ . A useful bonus of this approach is that all equations in the "time" formalism can be written in such a way that the quantities m and γ never explicitly appear; the formalism may then be applied equally well to massless particles (for which γ exists, but for which $m = 0$ and γ is undefined).

A somewhat simplified version of this approach is not to actually use the "time" at all, but rather to simply "pretend" that τ is not in fact constrained by Eq. (1) until after the equations of motion have been obtained. The results are the same, so let us follow this latter, simplified approach. A suitable Lagrangian can, for example, be chosen to be⁵

$$L = \frac{1}{2} m u^\mu u_\mu + q u^\mu A_\mu ; \quad (19)$$

where A_μ is the electromagnetic four-potential:

$$A_\mu = (V; \mathbf{A}) :$$

Using Eqs. (10) and (13), we can recognize the Lagrangian (19) as simply a straightforward coupling of the tensor and vector currents of the particle to their corresponding gauge fields:

$$L = \frac{1}{2} T_{\mu\nu} g^{\mu\nu} + J_\mu A^\mu :$$

The canonical momentum components p_μ conjugate to the generalized degrees of freedom z^μ are obtained from (19) in the standard way:

$$p_\mu = \frac{\partial L}{\partial u^\mu} = m u_\mu + q A_\mu : \quad (20)$$

The Euler-Lagrange equations of motion then yield the Lorentz force law (17).¹¹

A corresponding Hamiltonian formulation of this same classical theory may be constructed in two different ways. On the one hand, a "manifestly-covariant Hamiltonian" H can be constructed by the usual Legendre transformation:

$$H = p_\mu u^\mu - L = \frac{1}{2m} (p_\mu - q A_\mu)(p^\mu - q A^\mu) - \frac{(p_\mu - q A_\mu)^2}{2m};$$

from which Hamilton's equations (with respect to the "time" parameter t) again yield the Lorentz force law (17), as well as the equation (20) relating the canonical momentum and velocity four-vectors.

On the other hand, the more conventional way to construct a Hamiltonian formulation of this system — that yields a somewhat simpler transition to the quantum theory — is to recognize that the canonical energy p^0 can be interpreted as the Hamiltonian H of the system, with respect to the coordinate time t . From Eq. (20), we have

$$(p_\mu - q A_\mu)^2 = m^2 (u_\mu u^\mu) = m^2;$$

where in this last expression we can "stop pretending" that $u_\mu u^\mu$ is not identically equal to unity, because it is p_μ , not u_μ , that plays a fundamental role in the Hamiltonian formulation of mechanics. Thus

$$(H - q' A^0)^2 - (p_i - q A_i)^2 = m^2;$$

and hence

$$H = q' A^0 + \frac{q}{\sqrt{m^2 + (p_i - q A_i)^2}}; \quad (21)$$

where $\epsilon = \pm 1$ encapsulates the choice of sign in taking the square root. Hamilton's equations again yield expressions equivalent to (17) and (20).

Let us now consider the simple case of a free particle, which will be sufficient for our purposes. From Eq. (20), we have in this case

$$p_\mu = m u_\mu; \quad (22)$$

equivalently, from Eq. (21), we have

$$H = \frac{q}{\sqrt{m^2 + p^2}}; \quad (23)$$

Eqs. (22) and (23) emphasize the fact that, in antiparticle motion, the canonical momentum four-vector p has negative energy; or, in other words, that p is odd under C :

$$C : p \rightarrow -p :$$

Consider, now, a particle and its corresponding antiparticle, both at rest. For the former, we have

$$p = (m; 0); \quad (24)$$

whereas for the latter we have

$$p = (-m; 0); \quad (25)$$

"Surely," a common argument goes, "does this not tell us that antiparticle motion is really a negative mass solution? Does it not further tell us that the total energy of this pair of particles is zero?" These two statements directly contradict our finding above that the mass of a particle is unchanged, whether it be in particle or antiparticle motion; and we of course know that the total energy of a neutral particle-antiparticle pair at rest is indeed $2m$, not zero. On the other hand, we know that, quantum mechanically, the canonical energy really is negative for an antiparticle solution: for example, the eigenstates for $p = 0$ have a time-dependence of the form $e^{-im t}$ (where we use units in which $\hbar = 1$), which, using the Einstein relation $E = i\hbar \partial_t$, implies that $E = -m$. So what are we to believe?

This "paradox" is usually presented in discussions of relativistic quantum mechanics leading to Dirac's "holes", and so on—but it is fundamentally a feature of relativistic mechanics itself, whether it be of the classical or quantum flavor. Let us now dispense with it once and for all.

The fallacy above is the assumption that the canonical momentum four-vector has anything at all to do with the "total mass" of a system. It does not. In trying to "add together" the four-momenta of the two particles, we are making the implicit assumption that we are in some sense computing the four-momentum generalization of the "total energy" or "total rest mass" of a system. But we have already noted that for a system of gravitational sources, it is the sum of the mechanical stress-energy tensor densities $t_{ij}(x)$ that determines the overall gravitational field generated by the system. From this gravitational field, one can define a "total gravitational mass" of the system (because for our purposes the strengths of these gravitational fields are assumed to be negligible compared to the other forces present, so that special relativity is a good approximation). But by the equivalence principle, this "gravitational" mass is simply the mass of the system; and no definition of "mass" giving a different result can be compatible with the equivalence principle. Now, we know that T is unchanged under the C -operation; in particular, for both the particle and the antiparticle above, we have

$$\begin{aligned} T^{00} &= m; \\ T^{0i} &= T^{i0} = T^{ij} = 0; \end{aligned}$$

and so for the system as a whole we have $T^{00} = 2m$. Thus, the total mass of the particle-antiparticle pair at rest is indeed the commonsense result of $2m$, not zero.

Let us review these issues in more detail. The Hamiltonian H is the zero-component of the canonical momentum four-vector p , and is thus reasonably called the "canonical energy". In quantum mechanics, the canonical momentum four-vector relates directly to frequency and wavelength:

$$p = \hbar \omega :$$

Thus, particle (antiparticle) motion for a free particle corresponds to positive (negative) frequencies. But this has nothing at all to do with those mechanical (or "kinematical") properties of the particle, that are physically observable. Indeed, in the presence of interactions, even the sign of the canonical energy (frequency) loses its relevance completely: for example, for the electromagnetic interaction of a classical point charge, we have

$$p = m u + qA :$$

It will be immediately noted that p is not gauge-invariant, and the value of p^0 can be given any arbitrary value (positive or negative) simply by redefining the zero point of the scalar potential. Thus p cannot possibly, of itself, determine any physically observable property of the particle, such as its mass, or its mechanical stress-energy tensor.

On the other hand, we know that it is possible to define some sort of four-vector that represents the mechanical energy-momentum of a system — after all, we have been adding energies and momenta together for centuries. To define such a vector, we need simply integrate the mechanical stress-energy tensor density $t_{\mu\nu}(x)$ over all three-space, in some given Lorentz frame. We can write this covariantly in the form

$$(t) = \int d^3x t_{\mu\nu}(t; \mathbf{x}); \quad (26)$$

where an element of "three-space" has been written covariantly as d^3x : in the given Lorentz frame, in which it is simply all three-space at a constant time t , we have

$$d^3x = n_\mu d^3x^\mu;$$

where n_μ is the timelike four-normal to the hyperplane, which in this frame has coordinates

$$n_\mu = (1; 0):$$

Let us compute (t) in this Lorentz frame: from Eqs. (12) and (26), we have

$$(t) = m \int_{-1}^1 d^3x \int_1^Z dt \delta^{(4)}[x - z(t)] u^\mu(t) u_\mu(t);$$

where we write (t) on the understanding that the given t is the proper time of the particle at the corresponding coordinate time t . The integration over all three-space can be performed immediately, with the three-part of the delta function simply yielding unity:

$$(t) = m \int_{-1}^1 dt \delta[t - z^0(t)] u^\mu(t) u_\mu(t):$$

We can now perform the integration over $\mathcal{V}$ by noting the standard identity for a delta function of a function:

$$[\delta(f(x))] = \sum_z \frac{\delta(x - z)}{|f'(z)|};$$

where z are the zeroes of $f(x)$. In this case we have

$$f(x) = t - z^0(x);$$

whence

$$\frac{df}{dx} = -\dot{z}^0;$$

and hence

$$= m u \frac{u^0}{\dot{z}^0};$$

where all quantities are assumed evaluated at the given value of t (or x). We now recognize the last factor as the particle (antiparticle) number:

$$\frac{u^0}{\dot{z}^0} = 1;$$

and hence

$$= m u = (m; m \cdot v): \quad (27)$$

We see that the extra factor of $\dot{z}^0$ "cancels" the oddness of u under C , so that the four-vector T is like T itself even under the C operation. It is the mechanical momentum, not the canonical momentum, that should be added together to compute the mechanical four-momentum of a system of particles. (Similarly, one must be careful to never confuse the mechanical stress-energy tensor density $t(x)$ with the canonical stress-energy tensor density $\theta(x)$ of field theory:¹² the former can be derived by functional differentiation of the Lagrangian with respect to the metric tensor g , and is therefore always symmetric;^{9;13} the latter is defined in field theory with respect to the canonical momentum density, and in general possesses no particular symmetry.)

We can perform a similar integration of the electromagnetic current density vector $j(x)$, to obtain the "effective flux of charge" Q passing through a given spacelike hypersurface:

$$Q(t) = \int_{\Sigma} d^3x j(t; x):$$

By a similar analysis to that above, we simply find

$$Q = q;$$

so that Q can be understood as the "effective" charge of the particle, if it had been in "normal" particle motion. (The choice of what is "particle" motion and what is "antiparticle" motion is of course implicitly contained in the direction of the timelike four-normal.)

Quantity	Symbol	Rank	C	P T	Value
Mass	m	0	+	+	Free parameter
Electric charge	q	0	+	+	Free parameter
Proper time		0		+	$d = dt = (t)$
Theta time		0		+	$d = d = m ()$
Particle number		0		+	$u^0 = j u^0 j = u^0 = = 1$
Lagrangian	L	0	+	+	$T g = 2 + J A$
Hamiltonian	H	0	+	+	$p u L = (p q A)^2 = 2m$
Effective charge	Q	0		+	q
Position	z	1	+		$z (),$ state vector
Velocity	u	1			$dz = d = (; v)$
Canonical momentum	p	1			$@L=@u = m u + qA$
Electromagnetic potential	A	1			$@ @ A @ @ A = j$
Electromagnetic current	J	1			qu
Mechanical momentum		1	+		$m u = (m ; m v)$
Electromagnetic field	F	2		+	$@ A @ A$
Mechanical stress (energy)	T	2	+	+	$m u u$
Mechanical angular momentum	M	3	+		$z T z T$

Table 1: Lorentz-covariant quantities for a classical point charge.

V II. C onclusions

The Stueckelberg-Feynman picture of antiparticles being simply particles moving backwards in proper time" can be seen to be an integral and important part of relativistic classical mechanics, which only requires minor additions to standard texts on special relativity. Furthermore, the historical misconceptions of the "negative energy problem" in relativistic quantum mechanics can be avoided by a thorough understanding of the difference between the canonical momentum p of Lagrangian theory, and the "mechanical" momentum that dictates the kinematical and gravitational properties of an object.

Finally, we summarize the properties of the various important Lorentz-covariant quantities for a classical point charge in Table 1.

A cknow ledgm ents

Helpful discussions with S.A.W outhuysen, J.W .G.W ignall, A.G.K lein, G.I.O pat, and M .J.T homson are gratefully acknowledged. This work was supported in part by the Australian Research Council. We were saddened to learn of the death of Prof. W outhuysen during the completion of this work.

R eferences

- [†]pc@physics.unimelb.edu.au; <http://www.ph.unimelb.edu.au/~pc>.
^Ymckellar@physics.unimelb.edu.au.
^Zarawlin@physics.unimelb.edu.au.

- ¹ J.P. Costella and B.H.J. McKellar, "The Foldy-Wouthuysen transformation," *Am. J. Phys.* 63, 1119{1121 (1995).
- ² T.D. Newton and E.P. Wigner, "Localized states for elementary systems," *Rev. Mod. Phys.* 21, 400{406 (1949).
- ³ L.L. Foldy and S.A. Wouthuysen, "On the Dirac theory of spin $1/2$ particles and its non-relativistic limit," *Phys. Rev.* 78, 29{36 (1950).
- ⁴ J.D. Jackson, *Classical Electrodynamics*, 2nd ed. (Wiley, New York, 1975).
- ⁵ E.C.G. Stueckelberg, "La mécanique du point matériel en théorie de la relativité et en théorie des quanta," *Helv. Phys. Act.* 15, 23{37 (1942).
- ⁶ R.P. Feynman, "A relativistic cut-off for classical electrodynamics," *Phys. Rev.* 74, 939{946 (1948).
- ⁷ R.P. Feynman, "The theory of positrons," *Phys. Rev.* 76, 749{759 (1949).
- ⁸ R.P. Feynman, "The reason for antiparticles," in *Elementary Particles and the Laws of Physics: The 1986 Dirac Memorial Lectures* (Cambridge Univ. Press, Cambridge, 1987).
- ⁹ C.W. Misner, K.S. Thorne, and J.A. Wheeler, *Gravitation* (Freeman, New York, 1970).
- ¹⁰ H. Goldstein, *Classical Mechanics*, 2nd ed. (Addison-Wesley, Massachusetts, 1980).
- ¹¹ J.P. Costella, Ph.D. thesis, The University of Melbourne (1994), unpublished; available from the author's home page, listed above.
- ¹² C. Itzykson and J.-B. Zuber, *Quantum Field Theory* (McGraw-Hill, New York, 1980).
- ¹³ F. Belinfante, "On the current and the density of the electric charge, the energy, the linear momentum, and the angular momentum of arbitrary fields," *Physica* 7, 449{474 (1940).