

CERN-TH/98-123
 FERMILAB-PUB-98/101-T
 IC/98/34
 hep-ph/9804254

Resolving a Discrete Ambiguity in the CKM Angle θ_1 through $B_{u,d} \rightarrow J/\psi K$ and $B_s \rightarrow J/\psi D$ decays

Amol S. Dighe

The Abdus Salam International Centre for Theoretical Physics
 34100 Trieste, Italy

Isard Dunietz

Theoretical Physics Division, Fermi National Accelerator Laboratory
 Batavia, IL 60510, USA

Robert Fleischer

Theory Division, CERN, CH-1211 Geneva 23, Switzerland

Abstract

It is well known that $\sin(2\theta_1)$, where θ_1 is one of the angles of the unitarity triangle of the CKM matrix, can be determined in a theoretically clean way by measuring mixing-induced CP violation in the decay $B_d \rightarrow J/\psi K_S$. Another clean extraction of this CKM angle is provided by the time-dependent angular distribution for the decay products of $B_d \rightarrow J/\psi (l^+ l^-) K^0 (l^+ l^- K_S)$, where we have more observables at our disposal than in the case of $B_d \rightarrow J/\psi K_S$, so that in addition to $\sin(2\theta_1)$ also $\cos(2\theta_1)$ can be probed in a direct way. Unfortunately a sign ambiguity remains in $\cos(2\theta_1)$. If it could be resolved, a discrete ambiguity in the extraction of the CKM angle θ_1 could be resolved as well, which would allow a more incisive test of the CKM model of CP violation. This note shows that detailed time-dependent studies of $B_{u,d} \rightarrow J/\psi K$ and $B_s \rightarrow J/\psi D$ decay processes can determine the sign of $\cos(2\theta_1)$, thereby removing the corresponding ambiguity in the extraction of the CKM angle θ_1 .

The conventional methods for determining the angles α , β , and γ of the usual unitarity triangle [1] of the Cabibbo-Kobayashi-Maskawa matrix (CKM matrix) [2] leave several discrete ambiguities [3]. This is also the case for the "gold-plated" mode $B_d \rightarrow J/\psi K_S$. The mixing-induced CP asymmetry arising in this channel allows only a theoretically clean determination of $\sin(2\beta)$, so that a discrete four-fold ambiguity for the extracted value of $\beta \in [0; 360]$ remains. In the recent literature, several strategies were proposed to resolve ambiguities of this kind [4].

Another clean probe of the CKM angle β is provided by the observables of the angular distributions for the decay products of $B_d \rightarrow J/\psi (1^3 1^-) K^0 (\bar{1}^0 K_S)$ modes [5][7]. Such observables can in general be expressed in terms of decay amplitudes as

$$\dot{A}_f(t)^2; \quad \text{Re}[A_{\bar{e}}(t)A_f(t)]; \quad \text{Im}[A_{\bar{e}}(t)A_f(t)]; \quad (1)$$

where f and \bar{f} are labels for specific final-state configurations. The full three-angle distributions for tagged $B_d(t) \rightarrow J/\psi K^0 (\bar{1}^0 K_S)$ decays are given in [6][8]. Throughout this note, by tagging we mean making the distinction of initially, i.e. at $t=0$, present unmixed $B_{d;s}^0$ and $\bar{B}_{d;s}^0$ mesons. Weighting functions have been derived to extract the corresponding observables in an efficient way from experimental data [6, 7]. The time evolution of the interference terms in (1), i.e. of the real and imaginary parts of bilinear combinations of certain decay amplitudes, allows the determination [7] of $\sin(\alpha_1 - \alpha_2)$ and $\cos(\alpha_1 - \alpha_2)$, where α_1 and α_2 are CP-conserving strong phases, and of $\sin(2\beta)$ and

$$\cos(\alpha_1 - \alpha_2) \cos(2\beta): \quad (2)$$

The CP-conserving observables $\dot{A}_f(t=0)$, $\sin(\alpha_1 - \alpha_2)$ and $\cos(\alpha_1 - \alpha_2)$ can be determined to a higher accuracy from the much larger data samples arising for $B_{u;d} \rightarrow J/\psi K$ transitions, and untagged B_d decays into $J/\psi K^0 (\bar{1}^0 K^+)$ and $J/\psi \bar{K}^0 (\bar{1}^0 K^-)$ states [6, 7]. At first sight, one may think that $\sin(\alpha_1 - \alpha_2)$ and $\cos(\alpha_1 - \alpha_2)$ extracted this way from the $B_{u;d} \rightarrow J/\psi K$ angular distributions will allow the determination of $\cos(2\beta)$ with the help of the terms given in (2). A closer look shows, however, that this is unfortunately not the case, since we do not have sufficient information to fix the signs of $\cos(\alpha_1 - \alpha_2)$, thereby leaving a sign ambiguity for $\cos(2\beta)$.

The purpose of this letter is to point out that this ambiguity can be resolved with the help of tagged, time-dependent studies of $B_s \rightarrow J/\psi$ decays. The angular distributions are given in [7, 8], and weighting functions to extract the observables from experimental data can be found in [7]. An important feature of these observables is that they allow the determination of a CP-violating weak phase β_s [7, 9], which takes a very small value, of 0.03 , within the Standard Model, and represents a sensitive probe for new-physics contributions to B_s^0 (\bar{B}_s^0) mixing. Provided there is a sizeable mass difference between the mass eigenstates B_s^H and B_s^L , this phase can even be extracted from untagged B_s data samples [10], where the rapid m_s oscillations cancel [11].

Another important feature is the fact that the tagged, time-dependent $B_s(t) \rightarrow J/\psi (1^3 1^-) (\bar{1}^0 K^+ K^-)$ observables corresponding to the "Im" terms in (1) provide

sufficient information to determine $b_{1,2}$ unambiguously, where the strong phases $b_{1,2}$ are the flavour SU (3) counterparts of $\delta_{1,2}$. In the strict SU (3) limit, we have $b_{1,2} = \delta_{1,2}$. The time evolution of these observables takes the following form [7]:

$$e^{-\Gamma t} \sin(b_k - m_s t) + \frac{1}{2} e^{-\Gamma_H t} e^{-\Gamma_L t} \cos(b_k) ; \quad (3)$$

where terms of $O(\epsilon^2)$ have been neglected, Γ_H, Γ_L denote the decay widths of the B_s mass eigenstates, $\Gamma = (\Gamma_H + \Gamma_L)/2$, and $k = 1, 2$. Consequently, the strong phases $b_{1,2}$ can be determined unambiguously by resolving the rapid $m_s t$ oscillations in (3). Comparing the resulting values for $\sin(b_{1,2})$ and $\cos(b_1 - b_2)$ with their unhatted analogues, which can be determined from the $B_{u,d} \rightarrow J/\psi K$ observables, we obtain valuable information on SU (3) breaking. In order to fix the sign of $\cos(2\delta)$ with the help of (2), we just need the sign of $\cos(\delta_1)$ or $\cos(\delta_2)$, which is provided by the sign of $\cos(b_k)$ determined from the $B_s(t) \rightarrow J/\psi$ observables.

The SU (3) flavour symmetry should work reasonably well to determine this sign, unless $|\sin(b_k)|$ is close to 1, implying b_k close to 90 or 270, where $\cos(b_k)$ flips its sign. However, for such values of b_k , the $\cos(b_k)\cos(2\delta)$ terms (2) appearing in the $B_d \rightarrow J/\psi (1^+1^-) K^0 (1^0 K_S)$ angular distribution { which are essential for our strategy } will anyway be highly suppressed, so that it is doubtful that the sign ambiguity can be resolved in this case. It is of course not yet clear whether future experiments will encounter such an unfortunate situation. Within the framework of "factorization", we have $\delta_{1,2} \approx 0; 180^\circ$, i.e. $|\cos(\delta_{1,2})| = 1$. Since $B_{u,d} \rightarrow J/\psi K$ decays and their B_s counterpart $B_s \rightarrow J/\psi$ are colour-suppressed modes, "factorization" is not expected to be a good approximation in this case. Consequently, the actual values of $\delta_{1,2}$ may deviate significantly from these trivial values.

Angular distribution measurements for $B \rightarrow J/\psi K$ modes have already been reported [12], and others may soon be made public [13]. The modes considered here are very appealing, because of the ability to trigger on the J/ψ meson. The prospects are bright for resolving the rapid $m_s t$ oscillations in $B_s(t) \rightarrow J/\psi$ decays at planned experiments at the Tevatron and the LHC. Thus, in a not too distant future, the determination of b_k and the resolution of the ambiguity (related to the sign of $\cos(2\delta)$) in the extraction of the CKM angle θ from $B_d \rightarrow J/\psi K^0 (1^0 K_S)$ decays may become feasible. Let us note that there remains a two-fold ambiguity for θ in this approach, since we cannot decide whether θ lies within the intervals $[0; 180]$ or $[180; 360]$. In each interval, θ is, however, fixed unambiguously. Consequently, the original four-fold ambiguity arising in the extraction of θ from $\sin(2\delta)$ can be reduced to just a two-fold ambiguity. Usually it is argued that θ_K , which measures indirect CP violation in the kaon system, implies the former range [4].

While the $B_d \rightarrow J/\psi K^0 (1^0 K_S)$ mode is very accessible at B factories operating at the $(4S)$ resonance, detectors at hadron accelerators should study the feasibility of the 0^- reconstruction. The $B_d \rightarrow J/\psi 0^-, J/\psi 1^-$ modes could be added to $B_d \rightarrow J/\psi K^0 (1^0 K_S)$ in order to resolve the ambiguity. If penguin amplitudes are neglected, the time evolution of these decay modes also depends on the CKM angle θ and, in the limit

of the $SU(3)$ flavour symmetry, their strong phases are equal to those of their $SU(3)$ counterparts.

In summary, traditional methods allow tests of the CKM picture of CP violation only up to discrete ambiguities. The resolution of these ambiguities would make such CKM tests significantly more powerful. In this letter, making use of the many observables that are available from angular correlations, we have proposed an approach to resolve a discrete ambiguity in the determination of the CKM angle β that may be simpler than strategies advocated earlier [4]. More generally, angular-correlation methods can also be formulated to remove discrete CKM ambiguities in $B \rightarrow 2\pi$ and from colour-allowed processes [14].

This work was supported in part by the Department of Energy, Contract No. DE-AC02-76CH03000.

References

- [1] L.L. Chau and W.-Y. Keung, Phys. Rev. Lett. 53 (1984) 1802; J.D. Bjorken, private communication; C. Jarlskog and R. Stora, Phys. Lett. B 208 (1988) 268.
- [2] N. Cabibbo, Phys. Rev. Lett. 10 (1963) 531; M. Kobayashi and T. Maskawa, Progr. Theor. Phys. 49 (1973) 652.
- [3] For a recent review, see, for instance, A.J. Buras and R. Fleischer, University of Karlsruhe preprint TTP 97-15 (1997) [hep-ph/9704376], to appear in Heavy Flavours II, eds. A.J. Buras and M. Lindner (World Scientific, Singapore, 1998).
- [4] Y. Grossman, Y. Nir and M.P. Worah, Phys. Lett. B 407 (1997) 307; Y. Grossman and H.R. Quinn, Phys. Rev. D 56 (1997) 7259; Y. Grossman, B. Kayser, Y. Nir, Phys. Lett. B 415 (1997) 90; J. Charles, A. Le Yaouanc, L. Oliver, O. Pene and J.-C. Raynal, LPTHE-Orsay 97/70 (1997) [hep-ph/9801363]; L. Wolfenstein, Carnegie Mellon University preprint CMU-HEP-98-01 (1998) [hep-ph/9801386]; B. Kayser, talk given at the Moriond Workshop on Electroweak Interactions and Unified Theories, Les Arcs, France, March 1997 [hep-ph/9709382], to appear in the Proceedings.
- [5] B. Kayser, M. Kuroda, R.D. Peccei and A.J. Sanda, Phys. Lett. B 237 (1990) 508.
- [6] I. Dunietz, H.R. Quinn, A. Snyder, W. Toki and H.J. Lipkin, Phys. Rev. D 43 (1991) 2193.
- [7] A.S. Dighe, I. Dunietz and R. Fleischer, preprint CERN-TH/98-85, FERMILAB-PUB-98/093-T (1998).
- [8] A.S. Dighe, I. Dunietz, H.J. Lipkin and J.L. Rosner, Phys. Lett. B 369 (1996) 144.

- [9] I. Dunietz, in Proceedings of the Workshop on B Physics at Hadron Accelerators, Snowmass, Colorado, eds. P. McBride and C. Shekhar Mishra, Fermilab-CONF-93/267 (Batavia, 1993), p. 83.
- [10] R. Fleischer and I. Dunietz, Phys. Rev. D 55 (1997) 259.
- [11] I. Dunietz, Phys. Rev. D 52 (1995) 3048.
- [12] C. P. Jessop et al. (CLEO collaboration), Phys. Rev. Lett. 79 (1997) 4533.
- [13] M. Schmidt and S. Pappas (CDF collaboration), private communication.
- [14] I. Dunietz, in progress.