

UR-1523

April 1998

A Simple Phenomenological Parameterization of Supersymmetry without R-Parity

Mike Bisset, Otto C. W. Kong, Cosmin Macesanu, and Lynne H. Orr

Department of Physics and Astronomy,

University of Rochester, Rochester NY 14627-0171

Abstract

We present a parameterization of the supersymmetric standard model without R-parity that permits efficient phenomenological analyses of the full model without a priori assumptions. Under the parameterization, which is characterized by a single vacuum expectation value for the scalar components of the $Y = 1/2$ superfields, the expressions for tree-level mass matrices are quite simple. They do not involve the trilinear R-parity violating couplings; however, the bilinear μ_i terms do enter and cannot be set to zero without additional assumptions. We set up a framework for doing phenomenology and show some illustrative results for fermion mass matrices and related bounds on parameters. We find in particular that large values of $\tan\beta$ can suppress R-parity violating effects, substantially weakening experimental constraints.

Typeset using REVTeX

E-mails: bisset@urhep.fas.rochester.edu ; kong@pas.rochester.edu ;

mcos@pas.rochester.edu ; orr@urhep.fas.rochester.edu .

I. INTRODUCTION

A softly broken supersymmetry (SUSY) around or below the scale of a TeV is no doubt the most popular extension of the Standard Model. Most SUSY studies concentrate on a "minimal" version of such a model which contains two electroweak symmetry breaking Higgs doublets and an ad hoc discrete symmetry, called R-parity, which essentially distinguishes particles from superparticles. The phenomenological role of R-parity is to forbid B- or L-number violating couplings for which there are important experimental bounds, for instance those from superparticle mediated proton decays. However, strict R-parity conservation is not required to satisfy these bounds. Furthermore, allowing R-parity to be broken, either spontaneously through sneutrino VEV's or explicitly in the Lagrangian, gives rise to interesting phenomenology. This has become a subject of much interest recently (for some recent reviews and references, see [1,2]). Because of the large number of possible R-parity violating (RPV) couplings, most studies impose assumptions to restrict the analyses to a particular subset of RPV couplings. We wish to adopt a purely phenomenological point of view that allows an analysis of all the RPV couplings without a priori assumptions. In this letter we present our perspective on the parametrization of the general RPV Lagrangian and show some initial results. Here we concentrate on tree level results for mass matrices in the fermion sector, especially in the large $\tan\beta$ regime; a more detailed study will appear elsewhere.

II. PARAMETRIZATION OF R-PARITY VIOLATION

The most general renormalizable superpotential for the supersymmetric standard model without R-parity can be written as

$$W = \frac{1}{2} \mu_{ab} \hat{L}^a \hat{H}_u^b + h_{ik}^u \hat{Q}_i^a \hat{H}_u^b \hat{U}_k^c + \frac{1}{2} \mu_{ik}^0 \hat{Q}_i^a \hat{L}^b \hat{D}_k^c + \frac{1}{2} \mu_{kij}^1 \hat{L}^a \hat{L}^b \hat{E}_k^c + \frac{1}{2} \mu_{ijk}^0 \hat{D}_i^c \hat{D}_j^c \hat{U}_k^c ; \quad (1)$$

where (a;b) are SU(2) indices, (i;j;k) are family (flavor) indices, and (;) are (extended) flavor indices from 0 to 3 with $\hat{L}$'s denoting the four doublet superfields with $Y = -1/2$.

and ϵ_{ij} are antisymmetric in the first two indices as required by SU (2) and SU (3) product rules respectively. In the limit where $\epsilon_{ijk} = 0$; $\epsilon_{ijk} = 0$ and ϵ_{ij} all vanish, one recovers the expression for the R-parity preserving minimal supersymmetric standard model (MSSM), with $\hat{L}_0$ identified as $\hat{H}_d$. In that case, the two Higgses acquire vacuum expectation values (VEV's) and the lepton and quark Yukawa couplings are given by $h_{ijk}^e = 2\epsilon_{ijk}$ ($= 2\epsilon_{ijk}$), $h_{ijk}^d = 0$, and h_{ijk}^u , respectively. In the general case, the full expression for W together with all admissible soft SUSY breaking terms should be used to construct the scalar potential and solve for the vacuum. The solution is then expected to involve VEV's for all ve neutral scalars, i.e. for both the Higgses and sneutrinos in the usual terminology.

It is not necessary, however, to retain all ve VEV's in parametrizing the model, because the freedom associated with the choice of flavor basis creates redundancy amongst the parameters. For example, in the MSSM, the two 3×3 complex mass matrices $\frac{Y_u}{2}h_{ijk}^u$ and $\frac{Y_d}{2}h_{ijk}^d$ correspond only to ten real parameters describing the six mass eigenvalues and the CKM matrix. And although there are models attempting to construct the full high-energy mass matrices [3] in particular flavor bases, so far as low-energy phenomenology is concerned the different bases cannot be distinguished. A fruitful strategy in the MSSM is therefore to choose a flavor basis that parametrizes the two Yukawa matrices h^u and h^d with exactly ten parameters, namely assuming one mass matrix to be given by the diagonal eigenvalues and the other a multiple of the CKM matrix and the other diagonal eigenvalue matrix. Similarly, in parametrizing the general supersymmetric standard model without R-parity, U (3) flavor rotations for $\hat{Q}_i$; $\hat{U}_i^c$; $\hat{D}_i^c$ and $\hat{E}_i^c$ as well as a U (4) rotation for $\hat{L}$ can be exploited.

The above observation is not new. The popular parametrization exploiting the flavor rotations has all 's except ν_0 as well as two of the sneutrino VEV's ($\tilde{\nu}_i$'s) set to zero (rotated away) [4]. (In this context sneutrinos refer to the scalar components of the three $\hat{L}_i$ superfields as defined in the basis where the $\tilde{\nu}_i$'s are zero.) Note that the lepton Yukawa matrix h_{ijk}^e cannot then be taken as diagonalized, since the full $\hat{L}$ basis is already fixed. This (single-) parametrization was introduced originally in studies of RPV effects on the leptons

under the assumption that the trilinear RPV couplings are zero [4]. Extending its usage to the most general RPV scenario proves difficult. For instance, the chargino-charged-lepton mass matrix in a generic basis is given by

$$M_c = \begin{pmatrix} 0 & \frac{g_2 v_u}{2} & 0 & 0 & 0 \\ \frac{g_2 v_d}{2} & 0 & h_{ik}^e \frac{\tilde{v}_i}{2} & 0 & 0 \\ \frac{g_2 v_i}{2} & h_{ik}^e \frac{\tilde{v}_i}{2} + 2 m_{ijk} \frac{\tilde{v}_i}{2} & 0 & 0 & 0 \end{pmatrix} : \begin{matrix} C \\ C \\ C \\ C \\ A \end{matrix} \quad (2)$$

It is easy to see that under the single- $\tilde{v}$ parametrization a set of $\tilde{v}$ -couplings associated with the nonzero sneutrino VEV still remain, making the analysis of the mass eigenstates quite complicated. Analogously, $\tilde{v}$ -couplings will enter the down-quark mass matrix along with the nonzero sneutrino VEV. It has also been pointed out that rotating away the $\tilde{v}_i$'s does not simplify the analysis of the scalar potential because the RPV soft mass terms also contribute to the same quadratic terms [5].

We wish to find a relatively simple parametrization not requiring a priori assumptions. We propose here to use the $U(4)$ rotation to set all sneutrino VEV's ($\tilde{v}_i^D, \hat{\tilde{L}}_i^E$) to zero, leaving a single VEV for $\hat{\tilde{L}}_0^E$. We keep all of the $\tilde{v}$ while the rest of the leptonic flavor rotations are used to set h_{ik}^e diagonal. In our new basis, the (tree-level) mass matrices for all the fermions do not involve any trilinear RPV couplings¹. In particular, M_c is given by

$$M_c = \begin{pmatrix} 0 & \frac{g_2 v_u}{2} & 0 & 0 & 0 \\ \frac{g_2 v_d}{2} & 0 & 0 & 0 & 0 \\ 0 & m_1 & 0 & 0 & 0 \\ 0 & m_2 & 0 & m_2 & 0 \\ 0 & m_3 & 0 & 0 & m_3 \end{pmatrix} ; \begin{matrix} C \\ C \\ C \\ C \\ A \end{matrix} \quad (3)$$

where $m_i = h_{ii}^e \frac{\tilde{v}_i}{2}$. The quark mass matrix for each sector involves only one VEV and therefore assumes the same form as in the MSSM. The neutralino-neutrino mass matrix is then given by

¹The scalar mass matrices are also much simplified as there are only two non-zero VEV's, v_u and v_d . Further details will be discussed in [6].

some higher scale. In that case, the extra super field(s) have to be integrated out to recover the supersymmetric standard model. Such models are naturally formulated in a mixed parametrization [2], i.e. with both $\tilde{L}_i$'s and $\hat{L}_i^D \hat{L}_i^E$'s.

There are also discussions in the literature which consider the MSSM with only a few RPV-terms added while the "neutrino VEV's" are assumed to be zero. One should be particularly careful in interpreting the meaning of the assumptions in such cases. As implied in the above discussion, imposing such assumptions at the Lagrangian level is not a (flavor) basis or parametrization independent procedure. Combining such assumptions with a specific choice of basis is a potential source of confusion and sometimes inconsistency. In our opinion, a clear interpretation is provided by beginning with the Lagrangian in the single-VEV parametrization we proposed above. A given model is then identified by specifying which RPV-terms are not admitted. An important related point is that so long as the $\tilde{L}_i$'s are not all assumed to be zero, the $\tilde{L}_i$'s are not exactly identifiable as the physical charged leptons, nor are the charginos and neutralinos, for instance, the same states as in the R-parity conserving MSSM. The vanishing of the $\tilde{L}_i$'s, if taken, would be an assumption. All in all, while specific RPV models may be more naturally formulated in a particular parametrization, the single-VEV parametrization, we believe, provides a particularly efficient framework for a model independent study of constraints on and phenomenology of R-parity violation.

III. ANALYSIS OF NEUTRAL FERMION MASS MATRIX

The generation of non-zero neutrino mass(es) is one of the most prominent features of R-parity violation. As a result the experimental neutrino mass bound has been used to put constraints on various RPV-couplings [8]. Two neutrino eigenstates are left massless at the tree level, while the third one gains a mass through the RPV-couplings ($\tilde{L}_i$'s) to the higgsino, as can be seen from Eq.(4). Note that the massive eigenstate is in general a mixture of all three neutrino states. In fact one can use a simple rotation to decouple the massless states. The remaining 5×5 mass matrix is then given by

$$M_N^{(5)} = \begin{pmatrix} 0 & 0 & \frac{g_1 v_u}{2} & \frac{g_1 v_d}{2} & 0 \\ 0 & M_2 & \frac{g_2 v_u}{2} & \frac{g_2 v_d}{2} & 0 \\ \frac{g_1 v_u}{2} & \frac{g_2 v_u}{2} & 0 & 0 & 0 \\ \frac{g_1 v_d}{2} & \frac{g_2 v_d}{2} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}; \quad (5)$$

where

$$m_5 = \frac{1}{2} \sqrt{\frac{g_1^2 v_u^2 + g_2^2 v_d^2}{M_2^2}}; \quad (6)$$

and the corresponding massless neutrino state is given by

$$j_5 = \frac{1}{5} j_1 + \frac{2}{5} j_2 + \frac{3}{5} j_3; \quad (7)$$

The common strategy to obtain the neutrino mass corresponds to assuming a small m_5 where $M_N^{(5)}$ adopts a "seesaw" structure. This gives

$$m_5 = \frac{\det M_N^{(5)}}{\det M_{4 \times 4}} = \frac{1}{2} \frac{v^2 \cos^2 \theta (x g_2^2 + g_1^2)}{[2xM_2 + (x g_2^2 + g_1^2) v^2 \sin^2 \theta]}; \quad (8)$$

where we have substituted $v_d = v \cos \theta$, $v_u = v \sin \theta$, and $M_1 = xM_2$. Note that for large $\tan \theta$, $\cos \theta$ is a suppression factor. For example, at $\tan \theta = 45$, saturation of the machine bound of 24 MeV [9] for m_5 allows a m_5 value as large as the chargino mass scale M_2 and the higgsino mass mixing parameter, μ . Now, large m_5 values are beyond the validity of the "seesaw" analysis; however, an alternative perturbative analysis can be performed treating the EW-symmetry breaking terms in $M_N^{(5)}$ as a perturbation. The first order part can then be diagonalized exactly without any assumptions about the magnitude of m_5 . The resulting zero eigenvalue is lifted by the perturbation to give

$$m_5 = \frac{1}{4} \frac{v^2 \cos^2 \theta (x g_2^2 + g_1^2)}{(x g_2^2 + g_1^2) x M_2}; \quad (9)$$

with the eigenvector in the original basis given by

$$\begin{pmatrix} \frac{g_1 v \cos \theta}{2xM_2} \\ \frac{g_2 v \cos \theta}{2M_2} \\ 0 \\ 0 \\ 0 \end{pmatrix}; \quad (10)$$

One can rewrite Eq.(9) to obtain a bound on m_5 :

$$m_5^2 < \frac{4x_0^2 M_2 m_{5(\text{bound})}}{v^2 \cos^2(\alpha g_2^2 + g_1^2) - 4x M_2 m_{5(\text{bound})}} : \quad (11)$$

As M_2 increases, the denominator above drops to zero, beyond which there is no bound on m_5 . For $\tan\beta = 45$, this happens at $M_2 = 210 \text{ GeV}$. We note that in order to use the machine mass bound of 24 MeV for $m_{5(\text{bound})}$, we have assumed $\mu_1 = \mu_2 = 0$, i.e. $m_5 = m_3$ ($\mu_1 : \mu_2 : \mu_3 = 0 : 0 : 1$).

The perturbative result in Eq.(11) is borne out by exact numerical results from diagonalizing the neutral fermion mass matrix, as illustrated in Figure 1, for $\tan\beta = 45$; the corresponding result for $\tan\beta = 2$ is also shown for comparison. The difference between m_5 bounds for the two cases is striking. In both cases we see that the neutrino mass bound on m_5 ($= m_3$ here) is tighter for low values of $|j_0|$ and weakens as $|j_0|$ is increased. For $\tan\beta = 45$, values of m_5 in the hundreds of GeV are completely consistent with the 24 MeV bound for viable regions of the $(M_2; j_0)$ parameter space. And again, for M_2 large enough, we get no limit on m_5 at all.

There are potentially much stronger bounds on neutrino masses from cosmological considerations which however depend on the decay modes and other assumptions so that a neutrino mass above an MeV is not definitely ruled out [10]. There are also other experimental constraints on neutrino masses and mixings. In this first paper, our purpose is simply to illustrate the advantage of performing the analysis in the single-VEV parametrization as well as to point out the interesting suppression of RPV effects in the large $\tan\beta$ regime.

IV. CONSTRAINTS FROM CHARGINO-CHARGED LEPTON MASS MATRIX

How are the μ_i 's otherwise constrained particularly in the large $\tan\beta$ regime? Couplings of the charged leptons to the Z^0 are well measured and can constrain the μ_i 's through the chargino-charged lepton mass matrix. In Ref. [11], various Z^0 -couplings constraints on R-parity violation are studied assuming the trilinear RPV couplings vanish. We follow basically

the same strategy (but without the latter assumption) and our parametrization allows the constraints to be cast explicitly in terms of the magnitudes of the μ_i . For the large $\tan\beta$ regime, we find a weakening of the constraints as a result of $\cos\beta$ suppression factor(s) even stronger than that illustrated above for the neutrino case.

We begin with a perturbative approach as in the neutral fermion sector. After we diagonalize its upper 2×2 (chargino) block, M_c (Eq.(3)) consists of two diagonal blocks of fully diagonal sub-matrices and a lower 3×2 off-diagonal block containing the μ_i parameters. The latter is taken as a perturbation to the diagonal matrix. It is then a simple exercise to obtain the matrix elements [12] :

$$U_L^Y(i+2;1) = \frac{\mu_i \bar{M}_2 \cos\beta}{M_2^2 - 2M_W^2 \sin^2\beta} ;$$

$$U_L^Y(i+2;2) = \frac{\mu_i M_2}{M_2^2 - 2M_W^2 \sin^2\beta} ; \quad (12)$$

$$U_R^Y(i+2;1) = \frac{m_i \bar{M}_2 \cos\beta (M_2 \sin\beta + M_0 \cos\beta)}{(M_2^2 - 2M_W^2 \sin^2\beta)^2} ;$$

$$U_R^Y(i+2;2) = \frac{m_i (M_2^2 + 2M_W^2 \cos^2\beta)}{(M_2^2 - 2M_W^2 \sin^2\beta)^2} ; \quad (13)$$

with

$$U_L^Y M_c U_R = \text{diag}(M_{\chi_1}, M_{\chi_2}, m_1, m_2, m_3, g) ; \quad (14)$$

where index i refers to one of the three leptonic states. These matrix elements are the ones needed for studying leptonic physics. They characterize the gaugino and higgsino contents of each leptonic mass eigenstate.

The Z^0 -boson coupling to the mass eigenstates is given by

$$L_{int}^Z = \frac{g_Z}{2\cos\beta} Z_\mu \left[A_{ij}^L \frac{1}{2} \gamma^\mu + A_{ij}^R \frac{1+\gamma^5}{2} \right] \psi_j ; \quad (15)$$

Using the results above, together with unitarity of the U_L^Y and U_R^Y matrices, we have, for large $\tan\beta$ where formulae simplify,

$$A_{ij}^L = \frac{2\mu_i \mu_j M_W^2 \cos^2\beta}{M_2^2} + \delta_{ij} (1 - 2\sin^2\beta) ;$$

$$A_{ij}^R = \frac{\mu_i \mu_j (M_2^2 + 4M_W^2)}{M_2^2} \delta_{ij} 2\sin^2\beta ; \quad (16)$$

Note that while Eq.(15) is valid for all ve charged fermion states, the above given formul for the couplings $\tilde{A}_{ij}^{L,R}$'s are only for the three leptonic states, to which we limit the present discussion. In accordance with the perturbational approach, the results contain \sm all" ratios such as $\frac{j}{0}$. The deviations from standard universality in $\tilde{A}_{ij}^R$ are typically ignored, being suppressed by two factors of the $\frac{m_i}{0}$ mass ratios. But the corresponding deviations in $\tilde{A}_{ij}^L$ have a $\cos^2$ suppression, which for $\tan = 45$, for example, gives a factor of 10^{-3} . The important point to note here, in the large $\tan$ regime, is that $\tilde{A}_{33}^R$ or $\tilde{A}_{23}^R$ could well be significant in comparison with the $\tilde{A}_{ij}^L$'s.

With the $\tilde{A}_{ij}^{L,R}$ formul, it is then straightforward to check the constraints different processes involving universality violation or FCNC [11] impose on the $\frac{j}{0}$ ratios. Specifically, we check coupling universality and left-right asymmetry, as well as tree level $Z^0 \rightarrow i \bar{i}$ couplings through branching ratios of various $Z^0 \rightarrow 2\gamma$, $\rightarrow 3\gamma$ and $\rightarrow 3\pi$ processes. When a constraint allows the ratios to be larger than unity, it essentially goes away, from the present perturbative perspective. This happens for all of these constraints for a sufficiently large M_2 except the one from $\rightarrow e^- e^+ e^-$, which has a much stronger experimental bound. For $\tan = 45$, $M_2 > 15-35 \text{ GeV}$ eliminates all but the latter process from which we obtain

$$\frac{j_{12}}{0} < 4.7 \cdot 10^{-7} M_2^2 : \quad (17)$$

With an admissible gaugino mass M_2 of the order 100 GeV , this only surviving constraint gives a numerical bound on $\frac{j_{12}}{0} \cdot 10^{-3}$ only (the signs of the j_i do not affect any of the results presented here). Furthermore, the same behavior appears when the $Z^0 \rightarrow \gamma \gamma$ width constraint is considered.

In the perturbative calculation, the m_i 's are approximated by the m_i 's, the physical charged lepton masses. In the exact computations, we numerically integrate from $j_i = 0$ (for which the m_i are exactly the m_i) to the final j_i values. This is necessary to find an acceptable set of m_i 's that yield the correct physical charged lepton masses for a given set of j_i 's. We also then find the chargino masses, which now depend on the j_i 's. For example, the minimum j_i values required to give both chargino masses above 90 GeV for $\tan = 45$

and $\mu_1 : \mu_2 : \mu_3 = 0 : 1 : 1$ are shown in Figure 2.

The numerical results also bear out the perturbative analysis of the couplings (in lepton and Z^0 decays) mentioned above. Bounds from $\text{B.R.}(\tau \rightarrow e e^+ e^-) < 1.0^{12}$ [9] are shown in Figure 3. Here we plot contours of μ_5 assuming a ratio $\mu_1 : \mu_2 : \mu_3 = 1 : 1 : 0$. As with the neutrino mass bound, the limit on μ_5 is more strict for low values of j_0 . The substantial weakening of the constraint for large $\tan \beta$ is well illustrated in the figure. Numerical studies of the other processes mentioned above give no restriction on the maximum allowed values of μ_5 for $j_0, M_2 > 10 \text{ GeV}$ (again for $\tan \beta = 45$), as indicated by the perturbational results.

V. SUMMARY

In summary, we have presented and illustrated with examples the merits of the single-VEV parametrization of supersymmetry without R-parity (see also [7]), which can be used in phenomenological studies without requiring specific model-dependent assumptions. Our analysis as outlined above also indicates a strong suppression of RPV effects from the bilinear μ_i terms in the large $\tan \beta$ regime. We note that the trilinear RPV couplings play no role in the analysis discussed here, though they are expected to have an important role in the other aspects of the model. This is, however, exactly what makes a comprehensive analysis of the full model feasible under the parametrization. Further details of such an analysis will be reported in future work.

ACKNOWLEDGMENTS

We thank K.S. Babu, E. Nardi and S. Pakvasa for helpful discussions. This work was supported in part by the U.S. Department of Energy, under grant DE-FG 02-91ER 40685 and by the U.S. National Science Foundation, under grant PHY-9600155.

REFERENCES

- [1] G . Bhattacharyya, Nucl. Phys. Proc. Suppl. 52A (1997) 83; H . Dreiner, hep-ph/9707435, to be published in *Perspectives on Supersymmetry*, ed. by G . Kane, World Scientific.
- [2] J W F . Valle, FTUV-98-18, hep-ph/9802292.
- [3] See, for example, P H . Frampton and O C W . Kong, Phys. Rev. D 53 (1995) R2293; Phys. Rev. Lett. 77 (1996) 1699, and O C W . Kong, UNC-CH Ph.D . Dissertation (1997), for a comprehensive review and more references.
- [4] R . Barbieri, D E . Brahm , L J . Hall, and S D H . Hsu, Phys. Lett. B 238 (1990) 86; L J . Hall and M . Suzuki, Nucl. Phys. B 231 (1984) 419.
- [5] A . Joshipura and M . Nowakowski, Phys. Rev. D 51 (1995) 5271.
- [6] M . Bisset, O C W . Kong, C . Macesanu, and L H . Orr, in preparation.
- [7] See also E . Nardi, Phys. Rev. D 55 (1997) 5772, where a similar parametrization to what we advocate here (single-VEV) has also been used to obtain the fermion mass matrices (without setting h_{ik}^e diagonal). There R-parity violation is studied under the assumptions of universality and $\hat{L}^D - \hat{L}^E$ alignment at some high energy scale and the analysis focuses on the neutrino mass induced through renormalization-group runnings.
- [8] See, for example, R . Hemping, Nucl. Phys. B 478 (1996) 3; H P . Nilles and N . Polonsky, *ibid* B 484 (1997) 33; R . Adhikari and G . Domancovic, IMSC/98/02/07, hep-ph/9802390.
- [9] R M . Barnett et al. (Particle Data Group), Phys. Rev. D 54 (1996) 1.
- [10] K S . Babu, T M . Gould, and I Z . Rothstein, Phys. Lett. B 321 (1994) 140; A D . Dolgov, S . Pastor, and J W F . Valle, Phys. Lett. B 383 (1996) 193; K . Kainulainen, hep-ph/9608215. M . Kawasaki, P . Keman, H . S . Kang R J . Scherrer, G . Steigman, and T P . Walker, Nucl. Phys. B 419 (1994), 105.

[11] M . Nowakowski and A . P ilafts is, Nucl. Phys. B 461 (1996) 19.

[12] See, for example, G . C . Branco and L . Lavoura, Nucl. Phys. B 278 (1986) 738 and Ref.

[11].

Figure Captions

Figure 1:

Maximum allowed values of m_5 (in GeV) consistent with $m_{\tilde{\chi}_1^0} < 24 \text{ MeV}$ ($m_1 : m_2 : m_3 = 0 : 0 : 1$). $M_1 = x M_2$, with $x = \frac{5}{3} \tan^2 \theta_w$ assumed from gaugino unification ($M_Z = 91.19 \text{ GeV}$; $\sin^2 \theta_w = 0.23$). The region above or outside of a given contour is excluded for m_5 's above the indicated value.

Figure 2:

Minimum values of m_5 (in GeV) required to give both chargino masses above 90 GeV ($m_1 : m_2 : m_3 = 0 : 1 : 1$). The area above or outside of a given contour has both chargino masses $> 90 \text{ GeV}$ for m_5 's above the indicated value.

Figure 3:

Maximum allowed values of m_5 (in GeV) consistent with $\text{BR}(\tilde{\chi}_1^0 \rightarrow e^+ e^-) < 1.0 \times 10^{-12}$ ($m_1 : m_2 : m_3 = 1 : 1 : 0$). The region above or outside of a given contour is excluded for m_5 's above the indicated value.

FIGURES

FIG .1. Maximum allowed values of μ_5 (in GeV) consistent with $m_{\tilde{\chi}_1^0} < 24$ MeV
 $(\mu_1 : \mu_2 : \mu_3 = 0 : 0 : 1)$. $M_1 = xM_2$, with $x = \frac{5}{3}\tan^2 \theta_w$ assumed from gaugino unification
 $(M_Z = 91.19 \text{ GeV}; \sin^2 \theta_w = 0.23)$. The region above or outside of a given contour is excluded for
 μ_5 's above the indicated value.




FIG .2. Minimum values of μ_0 (in GeV) required to give both chargino masses above 90 GeV ($\mu_1 : \mu_2 : \mu_3 = 0 : 1 : 1$). The area above or outside of a given contour has both chargino masses > 90 GeV for μ_0 's above the indicated value.




FIG. 3. Maximum allowed values of μ_5 (in GeV) consistent with $\text{BR.}(\tau \rightarrow e e^+ e^-) < 10^{-12}$

($\mu_1 : \mu_2 : \mu_3 = 1 : 1 : 0$). The region above or outside of a given contour is excluded for μ_5 's above the indicated value.

