Tools for Tunneling

UriSarid

Department of Physics, University of Notre Dame, Notre Dame, IN 46556 USA

(April 16, 1998)

If the universe is trapped and cooled in a metastable false vacuum state, that state will eventually decay by bubble nucleation and expansion. For example, many extensions of the standard model incorporate new scalar elds ~ whose potential has a local minimum at ~ = 0 but a global minimum elsewhere, to which the vacuum will eventually tunnel. I calculate the lifetime of the false vacuum, and the eld pro le of the bubble after tunneling, for any potential that is approximately a polynomial of degree 4 near the false vacuum. E sentially exact results are given for a single eld; for multiple elds a strict lower bound is placed on the tunneling rate.

PACS num bers: 12.60 Jv, 03.65 Sq, 98.80 Hw

UND-HEP-97-US01 hep-ph/9804308

Consider a quantum eld theory of som e scalar elds (grouped for convenience into a vector) whose potential U (~) has local m in in a at the origin ~ = 0 and at some other $\sim = \sim_1$. If U (\sim_1) < U (0) then the vacuum at the origin is a false and metastable one. If the universe is in this false vacuum at some early epoch, and the tem perature is su ciently low { as I will assume for the rem ainder of this paper { then quantum uctuations will eventually initiate a phase transition from the supercooled initial state to the true vacuum through bubble nucleation and expansion [1]. A very elegant and tractable approach to studying this process was introduced by Coleman [2]. The bubble's most likely eld con guration, called the bounce, is the one which extrem izes the Euclidean action, and the lifetim e of the false vacuum is proportional to the exponential of this extrem al action. The bounce also determines the bubble's form in Minkowski space, after its form ation and outside the lightcone of its center.

In this work I calculate the bounce and its Euclidean action for a wide class of models, namely all those for which the potential is at least approximately a polynom ial of degree 4 near the false vacuum . The results are essentially exact for a single eld ; with more elds the action and therefore the false-vacuum lifetim e I calculate are strict upper bounds, while the one-dim ensional bounce given here may only give a qualitative picture of the true eld con guration. One class of applications for this work is to extensions of the standard m odel containing elds ~ whose vacuum expectation values (VEVs) in the present epoch are phenom enologically required to vanish, for example if they are electrically charged. Thus if a lower minimum of U develops away from the origin, the origin must not only remain a local minimum, but must also be su ciently long-lived that we would probably still be inhabiting it today. For this type of application the lifetim e is of prim ary im portance. It is hoped that, within other contexts, the shape of the bounce will be useful in calculating various other properties of the decay, for example the degree of supercooling, the thickness of the transition region and the distribution of energies

within its volume. Of course, for many of these contexts the e ects of gravity [3] and the expansion of the universe, and possibly of nite temperature [4], must be included, but the techniques developed here may readily be extended.

Most previous work on false vacuum decay either assum es a thin-wall approxim ation to derive analytical results, or specializes to a particularm odel and num erically extrem izes the action. But the form er approach is often only applicable when the lifetime is extremely long and hence not very interesting, while the latter can be intricate and tim e-consum ing, requiring a new computation for each model and often also ingenious methods of extrem ization [5]. The results presented here yield either a bound on the action or an essentially exact answer for a very wide class of potentials, with any choice of param eters and without any computer reanalysis. There is nevertheless som e overlap with previous authors. For the case of a single scalar eld , rescaling was used (as I do below) to exactly study two lim iting-case potentials [6,7] and a general quartic potential [8,9], though most of the latter results are given only graphically and for a limited range of parameters; in the high-tem perature lim it the action was calculated and presented com pletely in Ref. [10]. For the case of multiple elds, the reduction to a single eld was employed, for example, in Refs. [8,11]. These were particular realizations of som e of the sam e techniques used in the present work, and, as further em phasized in Ref. [12], illustrate the usefulness of this approach.

Consider any potential U in any number of scalar elds ~ with a hom ogeneous false vacuum at ~ = 0 and with U (0) = 0. In the sem iclassical approximation [2], the probability of bubble form ation per unit time and volume is =V ' $m^4 \exp(S_E)$. The prefactor is dimensionally the fourth power of the typical mass scale in the theory and thus can be readily estimated. I will concentrate exclusively on the argument of the exponential, which is the Euclidean action for the bounce solution. However, care should be taken to account for any large numerical prefactors, for instance [13] group-theoretical factors re-

lated to vacuum degeneracy in models with spontaneous symmetry breaking.

The requirem ent that the false vacuum has probably not decayed in our past light-cone reads: $(=V)L^4$ 1, where L is roughly the present size and age of the vis-10⁴⁵TeV ¹ ible universe. Num erically, L 10¹⁰yr so $S_E > 400 + 4 \ln (m = TeV)$. The bounce is an O (4)symmetric [14] solution of the Euclidean equations of motion which approaches the false-vacuum eld con guration $\sim = 0$ at Euclidean time = 1. To calculate it, one must extrem ize the action with respect to all possible non-trivial paths through eld space which satisfy the boundary conditions. This requires som e careful and som ew hat tim e-intensive num erical techniques [5], and the results are di cult to generalize. To simplify this task, consider a single, straight-line path: $\sim = *$ where v is a constant. For example, v can be xed to point tow ards the true vacuum , or tow ards any other direction where tunneling is possible. (The action can also later be minimized with respect to v, or to small perturbations thereon [10]). In any case, the extrem um of the true action S_E must be no larger than the extremum of the action $\mathbf{S}_{\mathrm{E}}~$ for ~ restricted to lie along v, by a theorem ofColeman [15]. Any model not obeying

$$S_{E} > 400 + 4 \ln (m = TeV)$$
 : (1)

will have an even smaller $S_{\rm E}$ and hence its false vacuum would have decayed by today. Constraining a model based on Eq. (1) isn't as stringent as contraining it based on the full $S_{\rm E}$, but it is certainly much simpler, can be studied in general as I do below, and is often su cient to rule out large regions of parameter space. For instance, D asgupta [12] has shown for a particular supersymmetric model that most of the parameter range excluded by an exhaustive $S_{\rm E}$ constraint; see also Ref. [8].

Thus we consider a potential function of a single eld , in particular:

U () = M
$$_{2}^{2}$$
 M $_{3}^{3}$ + 4 : (2)

This form encompasses all renorm alizable potentials, but is also a good approximation form any nonrenormalizable ones which can be expanded as a quartic polynomial near the false vacuum. I assume M $_2^2 > 0$ to ensure at least m etastability of the origin, and M $_3 > 0$ without loss of generality. By Euclidean spherical symmetry, the action is

$$S_E = 2 \frac{2}{0} \frac{3}{1} \frac{1}{2} z @ @ + U () d$$
 (3)

where z is some constant factor resulting from $p \frac{e^{\frac{1}{2}}}{x^2 + 2}$. Using a dimensionless coordinate $x = M_2 = z^{1-2}$) and

eld y (x) = M $_3$ =M $_2^2$ (), I and S = $z^2M \,_2^2$ =M $_3^2 \, \stackrel{1}{S}_{\rm E}$, where



FIG.1. The rescaled Euclidean action $\$_{\rm E}$ as a function of , with asymptotes as dashed lines.

$$\mathfrak{B}_{E} = 2 \frac{2}{0} \frac{x^{3}}{1} \frac{1}{2} \frac{dy}{dx}^{2} + \mathfrak{B}(y) dx \qquad (4)$$

Thus the general problem has been reduced to extrem – izing an action $\overset{1}{B}_{E}$ depending on a single parameter . Tunneling is possible for any $\frac{1}{4} > > 1$; when < 0 some other mechanism must eventually stabilize the potential, but will not in uence the following if the true potential is well-approximated by $\overset{1}{D}$ at all 0 < y y_{0} , where y_{0} is the escape point de ned below.

Extrem izing $\mathcal{D}_{\rm E}$ amounts [2] to solving the equations of motion

$$\frac{d^2y}{dx^2} + \frac{3}{x}\frac{dy}{dx} = \frac{0}{0}(\frac{10}{2}); \quad \frac{dy}{dx} = 0; \quad y(1) = 0 \quad (6)$$

for a particle located at position y as a function of time x 0 and subject to a potential $\overset{1}{b}$ [y (x)] and a timedependent viscous frictional force. The particle starts at some y (x = 0) y₀ > 0 and slides down the potential $\overset{1}{b}$ m onotonically towards the origin, with y₀ chosen so y (x ! 1) = 0.

The above equations are numerically integrated throughout the allowed range for . The resulting action, from Eq. (4) or equivalently from $\vartheta_{\rm E} = 2 \frac{2^{\rm R}}{0} x^3 \vartheta$ [y(x)]dx, is shown in Fig.1. To better than 1% accuracy this action is the the sem iem pirical expressions

where the + () subscript indicates > 0 (< 0), $\mathcal{B}_{E \text{ ;thick}}$ ' 45:4 (see also [6]), $\mathcal{B}_{E \text{ ;thin}} = 2^{-2} = [12(1 - 4)^3]$ and $\mathcal{B}_{E \text{ ;F}} = 2^{-2} = 3$ ensure the correct action in the thick



FIG.2. The (mescaled) bounce y(x) for 0:05 < x < 20 and various values: (A) 5, (B) 2, (C) 1, (D) 0:5, (E) 0:25, (F) 0:1, (G) 0:01, (H) 0:01, (I) 0:1, (J) 0:175, (K) 0:2, (L) 0:22, (M) 0:23, (N) 0:235.



FIG.3. The (rescaled) bounce is well-approximated by $y_t(x) = y_0 = [1 + x e^{\frac{p}{2}(x R)}]$; the \initial" value y_0 and the t parameters R and are plotted in (a), (b), and (c), respectively.

wall, thin walland Fubiniasym ptotic lim its, respectively, as discussed below .

The bounce solutions to the equations of motion are plotted in Fig. 2 for a various $\$. Perhaps surprisingly, they all t quite accurately the simple form

$$y(x) ' y_{t}(x) = 1 + x e^{p \frac{1}{2}(x R)}$$
 (9)

where $y_0 = y (x = 0)$, while and R are empirically chosen as functions of to optimize the t. Fig.3 shows y_0 and the best-t parameters and R as functions of . Note that when is positive but signi cantly below its maximal value, say < 0.15, the shape of the bubble and especially its action are very dienent from the thin-wall results. In particular, the eld tunnels out to a value y_0 far from the true vacuum value $\frac{3}{8}$ $\frac{1}{1}$ 1+ $(1 \quad \frac{32}{9})^{1=2}$ shown as the long-dashed gray curve in Fig.3a. The t parameters are them selves well-approximated by

У0 ; +	'	$y_{0;thin} + 1:18(1 4)^2 1:40(1 4)^3$	(10)
У0;	′	$y_{0,thick} + 10:4 \ln (1 + j) = 1:67 \ln (1 + ^2)$	(11)
+	′	2 [2 7:2 11:4 ² 0:72 ln (1 4)] ¹	(12)
	'	_F 1=[2+4:9jj 1:3j ¹ j ³]	(13)
R +	'	$R_{thin} + 0.91 = (1 \ 4)^{1=2} \ 1.41 + 6.4^{1:2}$	(14)
R	'	0.21 1:13 ln (1 + 11j $\frac{1}{2}$:15)	(15)

Here $y_{0,thin} = p^2 + 4(1 - 4)$, $y_{0,thick} ' 5:78$, $_F = 2$, and $R_{thin} = [2(1 - 4_p)]^1$ are the asymptotic values described below. The 2x in the exponential makes y_t tend towards zero at the correct rate as x ! 1, and the various expressions (10)-(15) give an excellent t wherever y(x) is signi cantly di erent from zero: $jy_t(x) = y(x) = y_0 < 1$ % for all x.

Several lim iting cases are of interest. The best-known is the thin-wall regime [2], when ! 1=4 and the two vacua approach degeneracy. To leading order in 4 the equations of $m \text{ otion}_{D} \underline{m}$ ay be solved analyt-1 ically: $y_{thin} = y_{0;thin} = f1 + exp[\frac{1}{2}(x)]$ R_{thin})]g. Note that, while in general the solution is better t by Eq. (9) with nite , in the thin-wall limit becomes entirely immaterial and y t ! ythin. The opposite limit, in a sense, is when ! 0, which I call (as did Ref. [16]) the thick-wall regime. The values thick ' 1:5 and R_{thick} ' 0.21 provide a better t over the entire range of x than the exact solution of R ef. [16] to the linearized equations of motion. A more subtle limit is ! 1. discussed in some detail in Refs. [6,7]. Then the cubic term in b becomes irrelevant, so $b_{\rm E}$! $jj^1 b_{\rm E,F}$ $2 \frac{2}{0} \frac{R_1}{R_1} x^3 \left[\frac{1}{2} (dy_F = dx)^2 + y_F^2\right]$ where $\$_{\rm E}$; y₄]dx and $j^{\frac{1}{2}}$ y. This action can only be extrem ized asym p-VF totically, Consider the action of the fam ily of functions $y_{\rm F}$; = $\frac{1}{2} = (2 + x^2)$. When ! 0, the quadratic term s m ay be neglected, in which case the $y_{\rm F}$; are exact solutions (known as Fubini instantons [17]) with a

-independent action $\frac{2}{3}$ 2 . Thus when the complete action $\$_{E\,;F}$ is computed for y_F ; with ! 0, its variation also tends to 0, so y_F ; becomes an increasingly acceptable sem iclassical tunneling solution with an asymptotic extrem al action $\$_{E\,;F}$ = $\frac{2}{3}$ 2 . Returning to the original problem, I expect $\$_E$! jj¹ $(\frac{2}{3}$ $^2)$ and y ! jj¹⁼² y_F ; ! 0, which is largely what I nd. (For exponentially large j jone expects R ! $2^{1=2}$ ln j j but the

tting function of Eq. (15) was chosen to obtain a simple and adequate t, not to generate the correct asymptotic behavior.)

The above results allow an easy calculation of the tunneling action, and also of the bubble's eld con guration for all t 0 and $j_{\mathbf{x}}j_{\mathbf{y}}$ tby using [2] the bounce solution with argument = $\frac{1}{\mathbf{x}^2} \neq \frac{1}{\mathbf{c}}$. (Within the lightcone, the

eld typically oscillates about the true vacuum; for this post-tunneling evolution, the equations of motion must be solved anew with imaginary .)

One sample application of these results is [18] in the m in im al m odel of supersym m etric gauge m ediation, in which the large Higgs VEV hierarchy can generate a new globalm in im um in the potential of the scalar superpartners of the tau lepton. The potential in fact involves three elds, two of which are electrically charged and so should not acquire a VEV. Using the above m ethods, and restricting to a straight path connecting the false and true vacua, a signi cant range of that model's param eters is ruled out on the grounds that tunneling would be too fast. A small span of parameters remains for which a true multidim ensional analysis would be needed to test whether the lifetim e would be long enough. Note, how ever, that no vacuum stability study can establish that a param eter value is de nitely allow ed: even if it produces a long-lived false vacuum, the evolution of the early universem ight put the initial state of ~ near the true vacuum and the model would be unacceptable. Vacuum stability can only rule out param eter values, and the present analysis does just that.

Som ewhat stronger bounds m ay result not only from tunneling via other paths through a multidim ensional

eld space, but also from therm al uctuations in the early, hot universe (for a related discussion, see Ref. [19] and references therein). However, such bounds are in general very model-dependent and, in the latter case, may be subject to more uncertainties (see, e.g., Ref. [20]).

The results of this work provide signi cant, robust and easy to determ ine bounds on a wide class of models in which the acceptable vacuum which we presumably inhabit can be destabilized by quantum tunneling. It is true that for such models the shape of the bounce, also determ ined in this work, is of only academ ic interest to us, since we will not survive the tunneling. The analysis of this shape can be useful, however, in a di erent class of models: ones in which the true vacuum is the one we live in, while the phase transition occurred long ago. Then the pro le of the bubble determ ines various properties of the early universe, such as how and where the latent heat was deposited, what was the spectrum of the particles produced, and what rem nants were left behind. To properly study such a scenario m ay require the inclusion of gravity, the tim e-dependent evolution of the universe, and possibly nonzero tem perature. The results of such studies will quantitatively be quite di erent from the present work, but hopefully m uch of the m ethodology introduced here will still be useful.

Enlightening discussions with S. Coleman, I. Dasgupta, S. Kusenko, A. Linde and especially R. Rattazzi, who collaborated in an early stage of this work, are gratefully acknow ledged.

E lectronic address: sarid@ particle phys.nd.edu

- M. B. Voloshin, IB. Kobzarev and L.B. Okun, Sov. J. Nucl. Phys. 20 (1974) 644.
- [2] S.Colem an, Phys. Rev. D 15 (1977) 2929; C.Callan and S.Colem an, Phys. Rev. D 16 (1977) 1762.
- [3] S. Colem an and F. De Luccia, Phys. Rev. D 21 (1980)
 3305; S. Parke, Phys. Lett. B 121 (1983) 313.
- [4] A D. Linde, Phys. Lett. B 100 (1981) 37 and Ref. [6].
- [5] A.Kusenko, Phys.Lett.B 358 (1995) 51.See also Ref. [8] below.
- [6] A D. Linde, Nucl. Phys. B 216 (1983) 421, and Particle Physics and In ationary Cosmology (Harwood, Chur, Switzerland, 1990).
- [7] A D. Linde, Nucl. Phys. B 372 (1992) 421.
- [8] M. Claudson, L.J. Hall and I. Hinchlie, Nucl. Phys. B 228 (1983) 501.
- [9] T.C. Shen, Phys. Rev. D 37 (1988) 3537; K. Enqvist, J. Ignatius, K. Kajantie and K. Rummukainen, Phys. Rev. D 45 (1992) 3415.
- [10] M. Dine, R.G. Leigh, P. Huet, A. Linde and D. Linde, Phys. Rev. D 46 (1992) 550.
- [11] I.D asgupta, B A.D obrescu and L.R andall, Nucl. Phys. B 483 (1997) 95; I.D asgupta, R.R adem acher and P. Suranyi, hep-ph/9804229.
- [12] I.Dasgupta, Phys. Lett. B 394 (1997) 116.
- [13] A.Kusenko, Phys. Lett. B 358 (1995) 47.
- [14] S.Colem an, V.G laser and A.M artin, Commun.M ath. Phys.58 (1978) 211.
- [15] S.Colem an, private com munication, based on the correspondence between the bounce and the minimal barrierpenetration path as established in S.Coleman, Nucl. Phys. B 298 (1988) 178 and Ref. [2]. See also Ref. [12].
- [16] A. Kusenko, P. Langacker and G. Segre, Phys. Rev. D 54 (1996) 5824.
- [17] S.Fubini, Nuovo Cimento 34A (1976) 521.
- [18] R.Rattazziand U.Sarid, Nucl.Phys.B 501 (1997) 297.
 [19] M.Sher, Phys.Lett.B 317 (1993) 159; Addendum, ibid. B 331 (1994) 448.
- [20] T.Falk et al, Phys. Lett. B 396 (1997) 50.