

FTUV/97-47
 IFIC/97-63
 UG-FT-77/97
 hep-ph/9804462
 April 1998

PERTURBATIVE QUARK MASS CORRECTIONS TO THE TAU HADRONIC WIDTH

Antonio Pich^a and Joaquim Paredes^b

^a Departament de Física Teòrica, IFIC, Universitat de València | CSIC
 Dr. Moliner 50, E-46100 Burjassot (Valencia), Spain.

^b Departamento de Física Teórica y del Cosmos, Universidad de Granada,
 Campus de Fuente Nueva, E-18002 Granada, Spain.

Abstract

The perturbative quark mass corrections to the τ hadronic width are analysed to $O(\frac{3}{s}m_q^2)$, using the presently available theoretical information. The behaviour of the perturbative series is investigated in order to assess the associated uncertainties. The implications for the determination of the strange quark mass from τ decay data are discussed.

1. Introduction

The inclusive character of the total hadronic width renders possible an accurate calculation of the ratio [1,5]

$$R = \frac{[\sum_{\text{hadrons}} \Gamma]}{[\sum_{\text{leptons}} \Gamma]}; \quad (1)$$

using standard field theory methods. The result turns out to be very sensitive to the value of $\alpha_s(M^2)$. Moreover, the uncertainties in the theoretical calculation are quite small and dominated by the perturbative errors. This has been used to perform a very precise determination of the QCD coupling at low energies [5].

Quark masses play a rather minor rôle in R . Owing to the tiny values of m_u and m_d , their associated corrections are very small [3] ($< 0.1\%$). The strange quark contribution to the total hadronic width is suppressed by the Cabibbo factor V_{us}^2 , which puts the induced m_s correction also at the per cent level. However, if one analyses separately the semi-inclusive decay width of the τ into Cabibbo-suppressed modes (i.e. final states with an odd number of kaons), the relatively large value of m_s induces an important effect of a size similar to the massless perturbative correction and of opposite sign [3]. The corresponding R_{ss} prediction is then very sensitive to the strange quark mass and could be used to extract information on this important, and nowadays controversial, parameter. A very preliminary value of m_s , extracted from the ALEPH decay data, has been already presented in recent workshops [6,7].

The determinations of light quark masses are usually obtained from analyses of the divergences of the vector and axial-vector current two-point function correlators or related observables [8-13]. These correlators are proportional to quark masses and, therefore, are very sensitive to their numerical values. Unfortunately, one needs phenomenological information on the associated scalar and pseudo-scalar spectral functions, which are not well known at present. The obvious advantage of a possible determination of m_s analysing quark mass effects in τ decays is that the experimental error can be systematically reduced in foreseen facilities like tau-charm or B factories. There is then some hope to achieve a precise determination of m_s from such analyses.

Recently the $O(\alpha_s^3)$ corrections to the $J = 0$ quark correlators have been calculated [14], and have been found to be rather large. The influence of these $O(\alpha_s^3)$ corrections on the determination of quark masses and the uncertainties coming from the truncation of the QCD perturbative series depend very much on the observable. One can see for instance that the QCD perturbative series behaves geometrically to $O(\alpha_s^3)$ for the divergence of pseudo-scalar (scalar) currents if resummed perturbatively in terms of $\alpha_s(s)$ [12,13]. This convergence improves [13] using other resummations like the Principle of Minimal Sensitivity (PMS) [15] or the one advocated in Ref. [4].

The hadronic decay width has also a $J = 0$ contribution, which, as we shall see, behaves rather badly. However, the largest quark mass correction originates in a piece of the left-handed current correlation function, involving the $J = 0 + 1$ combination, which shows a much better perturbative convergence.

The purpose of this paper is to study the perturbative behaviour of the corrections to R which are proportional to m_q^2 , in order to assess the associated uncertainties. These are the leading theoretical uncertainties in the m_s determination. The $O(m_q^2)$ contributions were already studied in Ref. [3]. In Ref. [16] the contributions of $O(m_q^2)$ to the relevant correlators were worked out. More recently, some partial information on $O(m_q^2)$ corrections has become available [14,17]. A much detailed analysis of all contributions up to dimension four will be presented elsewhere.

2. Theoretical Framework

The theoretical analysis of R involves the two-point correlation functions for the vector $V_{ij} = \bar{\psi}_j \gamma_\mu \psi_i$ and axial-vector $A_{ij} = \bar{\psi}_j \gamma_\mu \gamma_5 \psi_i$ colour-singlet quark currents ($i, j = u, d, s$):

$$\chi_{ijV}^{(J)}(q) = i \int \frac{d^4x}{(2\pi)^4} e^{iqx} \langle 0 | T(V_{ij}(x) V_{ij}^\dagger(0)^Y) | 0 \rangle \quad (2)$$

$$\chi_{ijA}^{(J)}(q) = i \int \frac{d^4x}{(2\pi)^4} e^{iqx} \langle 0 | T(A_{ij}(x) A_{ij}^\dagger(0)^Y) | 0 \rangle \quad (3)$$

They have the Lorentz decompositions

$$\chi_{ijV/A}^{(J)}(q) = (g_{\mu\nu} - \frac{q_\mu q_\nu}{q^2}) \chi_{ijV/A}^{(J)\perp}(q^2) + \frac{q_\mu q_\nu}{q^2} \chi_{ijV/A}^{(J)\parallel}(q^2); \quad (4)$$

where the superscript (J) in the transverse and longitudinal components denotes the corresponding angular momentum $J = 1$ (T) and $J = 0$ (L) in the hadronic rest frame.

The imaginary parts of the two-point functions $\chi_{ijV/A}^{(J)}(q^2)$ are proportional to the spectral functions for hadrons with the corresponding quantum numbers. The semihadronic decay rate of the B can be written as an integral of these spectral functions over the invariant mass s of the final state hadrons:

$$R = 12 \int_0^{M^2} \frac{ds}{M^2} \left(1 - \frac{s}{M^2} \right)^2 \left(1 + 2 \frac{s}{M^2} \right) \text{Im} \chi_{udV}^{(1)}(s) + \text{Im} \chi_{udA}^{(0)}(s) : \quad (5)$$

The appropriate combinations of correlators are

$$\chi^{(J)}(s) = \chi_{udV}^{(J)}(s) + \chi_{udA}^{(J)}(s) + \chi_{usV}^{(J)}(s) + \chi_{usA}^{(J)}(s) : \quad (6)$$

We can decompose the predictions for R into contributions associated with specific quark currents:

$$R = R_V + R_A + R_S : \quad (7)$$

R_{ijV} and R_{ijA} correspond to the contributions from the first two terms in Eq. (6), while R_{ijS} contains the remaining Cabibbo-suppressed contributions.

Exploiting the analytic properties of the correlators $\langle J \rangle(s)$, Eq. (5) can be expressed as a contour integral in the complex s plane running counter-clockwise around the circle $|s| = M^2$:

$$R = \frac{1}{i} \oint_{|s|=M^2} \frac{ds}{s} \left(1 - \frac{s}{M^2} \right)^3 \left(1 + \frac{s}{M^2} \right) \left(D^{L+T}(s) + 4D^L(s) \right); \quad (8)$$

We have used integration by parts to rewrite R in terms of the logarithmic derivative of the relevant correlators,

$$D^{L+T}(s) = s \frac{d}{ds} h^{(0+1)}(s); \quad D^L(s) = \frac{s}{M^2} \frac{d}{ds} h^{(0)}(s); \quad (9)$$

which satisfy homogeneous renormalization group equations.

Using the Operator Product Expansion to organise the perturbative and non-perturbative contributions to the correlators into a systematic expansion [18] in powers of $1/s$, the total ratio R can be expressed as an expansion in powers of $1/M^2$, with coefficients that depend only logarithmically on M^2 [3]:

$$R = 3 \langle V_{ud}^2 \rangle + \langle V_{us}^2 \rangle S_{EW} \left(1 + \frac{0}{EW} + \langle^{(0)} \rangle + \sum_{D=2,4,\dots}^X \cos^2 c_{ud}^{(D)} + \sin^2 c_{us}^{(D)} \right); \quad (10)$$

where $\langle^{(D)}_{ij} \rangle = (\langle^{(D)}_{ijV} \rangle + \langle^{(D)}_{ijA} \rangle)/2$ is the average of the vector and axial-vector correlations of dimension D , S_{EW} and $\frac{0}{EW}$ contain the known [19,20] electroweak corrections, and $\sin^2 c_{us} = \langle V_{us}^2 \rangle / (\langle V_{ud}^2 \rangle + \langle V_{us}^2 \rangle)$.

The dimension-zero contribution is the purely perturbative correction neglecting quark masses, which, owing to chiral symmetry, is identical for the vector and axial-vector correlators. It is fully generated by the Adler function $D^{L+T}(s)$, because $D^L(s)$ vanishes in the chiral limit. The correction $\langle^{(0)} \rangle$ has been investigated in great detail in Ref. [4]. We will follow a similar procedure to analyse the perturbative quark-mass corrections of dimension two.

3. Dimension-Two Corrections

For the sake of simplicity, let us take here $m_u = m_d = 0$. The arguments we shall put forward don't depend on it. In this limit, the vector and axial-vector correlators get the same quark-mass corrections, i.e. $D_{usA}^J(s) = D_{usV}^J(s)$ ($J = L + T; L$). The dimension-two contributions can be written in the form:

$$D_{us}^{L+T}(s) \Big|_{D=2} = \frac{3}{4} \frac{m_s^2(s)}{s} \sum_{n=0}^X \alpha_n^{L+T} a^n(s); \quad (11)$$

$$D_{us}^L(s) \Big|_{D=2} = \frac{3}{8} \frac{m_s^2(s)}{M^2} \sum_{n=0}^X \alpha_n^L a^n(s); \quad (12)$$

where $a = \mu_s = \mu$, μ is an arbitrary scale factor (of order unity) and the coefficients $\bar{\alpha}_n^J(\mu)$ are constrained by the homogeneous renormalization group equations satisfied by the corresponding functions $D_{us}^J(s)$:

$$\frac{d}{d} \bar{\alpha}_n^J(\mu) = \sum_{k=1}^{X^n} [2 - k - (n - k) \gamma_k] \bar{\alpha}_{n-k}^J(\mu); \quad (13)$$

for $n = 1$ and

$$\frac{d}{d} \bar{\alpha}_0^J(\mu) = 0 \quad (14)$$

i.e.

$$\begin{aligned} \bar{\alpha}_0^J(\mu) &= \bar{\alpha}_0^J; \\ \bar{\alpha}_1^J(\mu) &= \bar{\alpha}_1^J + 2 \gamma_1 \bar{\alpha}_0^J \log \mu; \\ \bar{\alpha}_2^J(\mu) &= \bar{\alpha}_2^J + 2 \gamma_2 \bar{\alpha}_0^J + (2 - \gamma_1 - \gamma_1) \bar{\alpha}_1^J \log \mu + \gamma_1 (2 - \gamma_1 - \gamma_1) \bar{\alpha}_0^J \log^2 \mu; \\ \bar{\alpha}_3^J(\mu) &= \bar{\alpha}_3^J + 2 \gamma_3 \bar{\alpha}_0^J + (2 - \gamma_2 - \gamma_2) \bar{\alpha}_1^J + 2 (\gamma_1 - \gamma_1) \bar{\alpha}_2^J \log \mu \\ &\quad + (\gamma_1 - \gamma_2 + 2 \gamma_2 (2 - \gamma_1 - \gamma_1)) \bar{\alpha}_0^J + (\gamma_1 - \gamma_1) (2 - \gamma_1 - \gamma_1) \bar{\alpha}_1^J \log^2 \mu \\ &\quad + \frac{2}{3} \gamma_1 (\gamma_1 - \gamma_1) (2 - \gamma_1 - \gamma_1) \bar{\alpha}_0^J \log^3 \mu; \\ \bar{\alpha}_4^J(\mu) &= \bar{\alpha}_4^J + \end{aligned} \quad (15)$$

The factors γ_k and γ_k are the expansion coefficients of the QCD β and γ functions,

$$\frac{da}{d} = \beta(a) a; \quad \beta(a) = - \sum_{k=1}^X \beta_k a^k; \quad (16)$$

$$\frac{dm}{d} = \gamma(a) m; \quad \gamma(a) = \sum_{k=1}^X \gamma_k a^k; \quad (17)$$

which are known to four loops [21{23]. The coefficients $\bar{\alpha}_n^J$ and $\bar{\alpha}_n^J(1)$ are only known to order μ_s^2 for $J = L + T$ and μ_s^3 for $J = L$ [8,14,17,24{28].

For three flavours and in the $\overline{MS}$ scheme, one has:

$$\gamma_1 = \frac{9}{2}; \quad \gamma_2 = -8; \quad \gamma_3 = \frac{3863}{192};$$

$$\gamma_4 = \frac{140599}{2304} - \frac{445}{16} \gamma_3 - 94.456079; \quad (18)$$

$$\gamma_1 = 2; \quad \gamma_2 = \frac{91}{12}; \quad \gamma_3 = \frac{8885}{288} - 5 \gamma_3 - 24.840410;$$

$$\gamma_4 = \frac{2977517}{20736} - \frac{9295}{216} \gamma_3 + \frac{135}{8} \gamma_4 - \frac{125}{6} \gamma_5 - 88.525817; \quad (19)$$

$$\begin{aligned}
d_0^{L+T} &= 1; & d_1^{L+T} &= \frac{13}{3}; & d_2^{L+T} &= \frac{21541}{432} + \frac{323}{54} \left(\frac{520}{27} \right)^3 - 37.083047; \\
d_0^L &= 1; & d_1^L &= \frac{17}{3}; & d_2^L &= \frac{9631}{144} - \frac{35}{2} \left(\frac{520}{27} \right)^3 - 45.845949; \\
d_3^L &= \frac{4748953}{5184} - \frac{91519}{216} \left(\frac{520}{27} \right)^3 + \frac{5}{2} \left(\frac{520}{27} \right)^4 + \frac{715}{12} \left(\frac{520}{27} \right)^5 - 465.846304;
\end{aligned} \quad (20)$$

Notice the rather bad perturbative behaviour of the $D = 2$ corrections to the correlation functions D_{us}^J . Remember that $a(M^2) \sim 0.11$.

Inserting the expansions (11) and (12) in Eq. (8), the $D = 2$ corrections to R_{us} can be expressed as

$$R_{us}^{(2)} = 8 \frac{m_s^2(M^2)}{M^2} [a(M^2)]; \quad [a] = \frac{1}{4} \left(\frac{520}{27} \right)^3 [a]^{L+T} + [a]^L; \quad (21)$$

where

$$[a(M^2)] = \sum_{n=0}^{\infty} d_n^J \left(\frac{520}{27} \right)^n [a]^{(n)}; \quad (22)$$

and the contour integrations are contained in the functions

$$B_{L+T}^{(n)}(a) = \frac{1}{4} \int_{-1}^1 \frac{dx}{x^2} (1+x)(1-x)^3 \frac{m^2(M^2 x)}{m^2(M^2)} a^n(M^2 x); \quad (23)$$

$$B_L^{(n)}(a) = \frac{1}{2} \int_{-1}^1 \frac{dx}{x} (1-x)^3 \frac{m^2(M^2 x)}{m^2(M^2)} a^n(M^2 x); \quad (24)$$

Since the quark mass ratio is flavour independent, the integrals $B_J^{(n)}(a)$ regulate also the small corrections proportional to m_u and m_d , which we are neglecting. These functions depend only on $i, j, a_s(M^2) = \alpha_s(M^2)$ and $\log \frac{M^2}{\mu^2}$. Moreover, they satisfy the following homogeneous renormalization group equations:

$$\frac{d}{d \log \frac{M^2}{\mu^2}} B_J^{(n)}(a) = \sum_{k=1}^{\infty} (n_k - 2_k) B_J^{(n+k)}(a); \quad (25)$$

4. Perturbative α_s Expansion

The usual perturbative approach expands the $B_J^{(n)}(a)$ functions in powers of α_s . This gives,

$$\begin{aligned}
B_J^{(0)}(a) &= 1 + \frac{h}{2} \log \frac{M^2}{\mu^2} + H_1^J a \\
&+ \frac{2}{2} \log^2 \frac{M^2}{\mu^2} + \frac{1}{2} (2h_1 + h_2) \log \frac{M^2}{\mu^2} + \frac{1}{2} H_2^J a^2 \\
&+ \frac{1}{2} \log^3 \frac{M^2}{\mu^2} + \frac{1}{2} (3h_1 + 3h_2 + h_3) \log^2 \frac{M^2}{\mu^2} + \frac{1}{2} (2h_1 h_2 + h_4) \log \frac{M^2}{\mu^2} + \frac{1}{2} H_3^J a^3 \\
&+ \frac{1}{2} \log^4 \frac{M^2}{\mu^2} + \frac{1}{2} (4h_1 + 4h_2 + 4h_3 + h_4) \log^3 \frac{M^2}{\mu^2} + \frac{1}{2} (3h_1^2 + 3h_1 h_2 + 3h_2^2 + h_5) \log^2 \frac{M^2}{\mu^2} \\
&+ \frac{1}{2} (2h_1 h_3 + 2h_2 h_3 + h_6) \log \frac{M^2}{\mu^2} + \frac{1}{2} H_4^J a^4 + \dots
\end{aligned}$$

$$\begin{aligned}
& 2 \log^3 + (2 \log^2 + \frac{2}{3} \log) + \frac{2}{3} \log^3 \\
& + \frac{1}{2} \log^2 + \frac{1}{4} \log + \frac{1}{12} \log^3 + \frac{1}{2} \log^2 + \frac{1}{4} \log + \frac{1}{12} \log^3 \\
& + \frac{1}{12} (2 \log^2 + \frac{2}{3} \log) + \frac{1}{12} \log^3 + \frac{1}{12} \log^3
\end{aligned} \tag{26}$$

$$\begin{aligned}
B_J^{(1)}(a) &= a \left[2 \log + \frac{1}{2} \log^2 + \frac{1}{4} \log^3 + \frac{1}{12} \log^4 \right] \\
&+ \frac{1}{2} \log^2 + \frac{1}{4} \log + \frac{1}{12} \log^3 + \frac{1}{12} \log^4
\end{aligned} \tag{27}$$

$$B_J^{(2)}(a) = a^2 \left[2 \log + \frac{1}{2} \log^2 + \frac{1}{4} \log^3 + \frac{1}{12} \log^4 \right] + \frac{1}{12} \log^4 \tag{28}$$

$$B_J^{(3)}(a) = a^3 \tag{29}$$

where

$$H_n^{L+T} = \frac{1}{4} \int_0^1 \frac{dx}{x^2} (1+x)(1-x)^3 \log^n(x); \tag{30}$$

$$H_n^L = \frac{1}{2} \int_0^1 \frac{dx}{x} (1-x)^3 \log^n(x); \tag{31}$$

To $O(\epsilon^4)$, the needed integrals are

$$\begin{aligned}
H_0^{L+T} &= 1; & H_1^{L+T} &= \frac{1}{6}; & H_2^{L+T} &= \frac{25}{18} - \frac{2}{3}; \\
H_3^{L+T} &= \frac{85}{36} - \frac{2}{6}; & H_4^{L+T} &= \frac{721}{54} - \frac{25}{9} + \frac{4}{5}; \\
H_0^L &= 1; & H_1^L &= \frac{11}{6}; & H_2^L &= \frac{85}{18} - \frac{2}{3}; \\
H_3^L &= \frac{575}{36} + \frac{11}{6}; & H_4^L &= \frac{3661}{54} - \frac{85}{9} + \frac{4}{5};
\end{aligned} \tag{32}$$

$$\begin{aligned}
H_3^L &= \frac{575}{36} + \frac{11}{6}; & H_4^L &= \frac{3661}{54} - \frac{85}{9} + \frac{4}{5};
\end{aligned} \tag{33}$$

The perturbative expansions $\alpha^n[a]$ then take the form

$$\alpha^n[a(M^2)] = \sum_{n=0}^{\infty} \alpha_n^J(\epsilon) + \tilde{\alpha}_n^J(\epsilon) a^n; \tag{34}$$

where the coefficients $R_n^J(\cdot)$ depend on $d_{m < n}^J(\cdot)$, $m < n$ and $m = n$; thus, they are known up to $O(a^3)$ and $O(a^4)$ for $J = L + T$ and $J = L$ respectively. For $n = 1$, one has $[h_n^J, R_n^J(1)]$:

$$h_0^{L+T} = 0; \quad h_1^{L+T} = \frac{1}{3}; \quad h_2^{L+T} = \frac{113}{72} \frac{17}{12} a^2 - 12.412495;$$

$$h_3^{L+T} = \frac{114517}{2592} \frac{4391}{144} a^2 - \frac{3659}{648} a^3 + \frac{1690}{81} a^5 - 241.926329; \quad (35)$$

$$h_4^{L+T} = \frac{26864009}{13824} \frac{3110783}{5184} a^2 - \frac{1073}{1920} a^4 + \frac{3051761}{15552} a^3 - \frac{123745}{2592} a^2 a^3 \\ + \frac{171845}{243} a^5 + \frac{29575}{162} a^2 a^5 - \frac{35}{24} d_3^{L+T} - 3229.101787 - \frac{35}{24} d_3^{L+T};$$

$$h_0^L = 0; \quad h_1^L = \frac{11}{3}; \quad h_2^L = \frac{625}{8} \frac{17}{12} a^2 - 64.143060;$$

$$h_3^L = \frac{1435691}{864} \frac{7927}{144} a^2 - \frac{5225}{24} a^3 - 856.673579; \quad (36)$$

$$h_4^L = \frac{693706385}{20736} \frac{2295071}{1728} a^2 - \frac{10877}{17280} a^4 - \frac{1429525}{144} a^3 + \frac{5595}{32} a^2 a^3 + \frac{264275}{288} a^5 \\ - 11377.111254;$$

The contour integration generates rather large numerical factors, which show an opposite behaviour for the transverse and longitudinal pieces. In the $L+T$ expansion the h_n^{L+T} contributions cancel to some extent with the original correlation function coefficients d_n^{L+T} ,

$$^{L+T} [a(M^2)] = 1 + 4a(M^2) + \frac{22219}{432} \frac{17}{12} a^2 + \frac{323}{54} a^3 - \frac{520}{27} a^5 - a(M^2)^2 \\ + d_3^{L+T} + \frac{114517}{2592} \frac{4391}{144} a^2 - \frac{3659}{648} a^3 + \frac{1690}{81} a^5 - a(M^2)^3 + \\ = 1 + 4a(M^2) + 24.671 a(M^2)^2 + d_3^{L+T} - 241.926 a(M^2)^3 + \quad (37)$$

However, both d_n^L and h_n^L contributions are large and positive, which gives rise to a badly behaved expansion for L :

$$^L [a(M^2)] = 1 + \frac{28}{3} a(M^2) + \frac{20881}{144} \frac{17}{12} a^2 - \frac{35}{2} a^3 - a(M^2)^2 \\ + \frac{13363099}{5184} \frac{7927}{144} a^2 - \frac{4}{36} \frac{17318}{27} a^3 + \frac{715}{12} a^5 - a(M^2)^3 + \\ = 1 + 9.333 a(M^2) + 109.989 a(M^2)^2 + 1322.520 a(M^2)^3 + \quad (38)$$

The bold{guess estimate $d_4^L = d_3^L$ ($d_3^L = d_2^L$) = 4733 would result in a huge $O(a^4)$ coefficient $d_4^L + h_4^L = 16110$.

Since d_3^{L+T} has a larger weight on the total contribution to $D_{us}^{(2)}$, the final combination of the transverse and longitudinal pieces has a better behaviour^a:

$$\begin{aligned}
[a(M^2)] &= 1 + \frac{16}{3} a(M^2) + \frac{10775}{144} \frac{17}{12} a^2 + \frac{1}{9} a^3 + \frac{130}{9} a^5 + a(M^2)^2 \\
&+ \frac{3}{4} d_3^{L+T} + \frac{14050201}{20736} \frac{5275}{144} a^2 + \frac{4}{144} a^4 + \frac{47401}{288} a^3 + \frac{13195}{432} a^5 + a(M^2)^3 \\
&+ \\
&= 1 + 5.333 a(M^2) + 46.000 a(M^2)^2 + 149.185 + \frac{3}{4} d_3^{L+T} a(M^2)^3 +
\end{aligned} \tag{39}$$

Nevertheless, the convergence of this perturbative series is very poor for the range of the strong coupling relevant in B decays, $a(M^2) = 0.11$. With $d_3^{L+T} = d_2^{L+T}$ ($d_2^{L+T} = d_1^{L+T}$) = 317, the $O(a^3)$ correction would be of the same size as the $O(a)$ and $O(a^2)$ contributions.

5. Resummation of Running Effects along the Integration Contour

At the moment, we can do very little about the apparent growth of the d_n^J coefficients, specially for $J = L$. We clearly need a deeper understanding of the perturbative $D_{us}^J(s)_{b=2}$ expansions. However, we can try to control better the large contributions contained in the h_n^J factors.

The integration along the circle $x = e^i$ gives rise to a long running of the quark mass and the QCD coupling. The expansion of $m^2(2M^2x) a^n(2M^2x)$ in powers of a generates imaginary logarithms $\log^n(x) = i^n (\dots)^n$, which are large in some parts of the integration range. The radius of convergence of such expansion is actually quite small [4]. However, there is no need to perform this ill{defined power expansion.

Using in Eqs. (23) and (24) the exact solution form (s) and $a(s)$ obtained from the renormalization group equations, the $B_J^{(n)}(a)$ integrals can be calculated to all orders in s , apart from the unknown $n > 4$ and $n > 4$ contributions, which are likely to be small. Thus, a more appropriate approach is to directly use the expansions (22), in terms of the original d_n^J coefficients, and to fully keep the known four{loop information on the functions $B_J^{(n)}(a)$.

Tables 1 and 2 show the exact results for $B_{L+T}^{(n)}(a)$ and $B_L^{(n)}(a)$ ($n = 0; 1; 2; 3$) with $\beta = 1$ obtained at different orders in the s and a expansions, together with

^a The $O(a^2)$ correction agrees with the numerical result recently reported in Ref. [29], which is larger than the value originally quoted in Ref. [16]. This larger $O(a^2)$ correction has been also confirmed by K.Chetyrkin and A.Kwiatkowski [30].

the numerical values of $B_J(a)$, for $a = 0:1$ ($a = 1$). For comparison the numbers coming from the truncated perturbative expressions at $O(a^3)$ are also given.

Table 1: Exact results for $B_{L+T}^{(n)}(a)$ ($n = 0;1;2;3$) obtained at the k loop ($k = 1;2;3;4$) approximation ($j > k = j > k = 0$), together with the numerical value of $B_{L+T}^{(n)}(a) = \sum_{n=0}^3 d_n^{L+T} B_{L+T}^{(n)}(a)$, for $a = 0:1$ and $a = 1$. For comparison the numbers coming from the truncated expressions at $O(a^3)$ are also given.

Loops	$B_{L+T}^{(0)}(a)$	$B_{L+T}^{(1)}(a)$	$B_{L+T}^{(2)}(a)$	$B_{L+T}^{(3)}(a)$	$B_{L+T}^{(n)}(a)$
1	0.89032	0.06965	0.00452	0.000186	1.360
2	0.81719	0.05666	0.00278	0.000008	1.166
3	0.79143	0.05296	0.00236	0.000048	1.108
4	0.78237	0.05168	0.00222	0.000060	1.089
$O(a^3)$	0.79363	0.06473	0.00892	0.001000	1.405

Table 2: Exact results for $B_L^{(n)}(a)$ ($n = 0;1;2;3$) obtained at the k loop ($k = 1;2;3;4$) approximation ($j > k = j > k = 0$), together with the numerical value of $B_L^{(n)}(a) = \sum_{n=0}^3 d_n^L B_L^{(n)}(a)$, for $a = 0:1$ and $a = 1$. For comparison the numbers coming from the truncated expressions at $O(a^3)$ are also given.

Loops	$B_L^{(0)}(a)$	$B_L^{(1)}(a)$	$B_L^{(2)}(a)$	$B_L^{(3)}(a)$	$B_L^{(n)}(a)$
1	1.39908	0.18473	0.02255	0.002588	4.686
2	1.54013	0.20247	0.02421	0.002692	5.052
3	1.57853	0.20617	0.02444	0.002690	5.120
4	1.58910	0.20706	0.02446	0.002681	5.133
$O(a^3)$	1.64446	0.21894	0.02192	0.001000	4.356

These numerical results show a reasonable convergence of the $B_J^{(n)}(a)$ integrals, as higher order k and k contributions are taken into account. Increasing the number of loops one gets a small decrease (increase) of the transverse (longitudinal) contribution. It is also clear that the truncated $O(a^3)$ expressions overestimate (underestimate) B_{L+T} (B_L). Taking the full four-loop information into account, we get the following perturbative behaviour:

$$B_{L+T}[0:1] = 0.7824 + 0.2239 + 0.0823 - 0.0000601 d_3^{L+T} + \quad (40)$$

$$B_L[0:1] = 1.5891 + 1.1733 + 1.1214 + 1.2489 + \quad (41)$$

The $L+T$ series converges very well. Owing to the negative running contributions the B_{L+T} series behaves better than the original perturbative expansion of $D_{us}^{L+T}(s)_{j=2}$. Unfortunately, the longitudinal series is much more problematic.

The bad perturbative behaviour of $D_{us}^L(s) \big|_{\mu=2}$ gets reinforced by the running effects, giving rise to a badly defined series.

The combined α_s expansion,

$$[0:1] = 0.9840 + 0.4613 + 0.3421 + 0.3122 - 0.000045 d_3^{L+T} + \dots \quad (42)$$

looks acceptable for the first terms because d_3^{L+T} is weighted by a larger factor. In fact, this series behaves better than the one in Eq. (39), obtained with the usual perturbative truncation of the contour integrals. Nevertheless, after the third term the series appears to be dominated by the longitudinal contribution, and the bad perturbative behaviour becomes again manifest.

Using the full four-loop result we have certainly gained in convergence for the $L+T$ series [compare the fourth term in the series (37) and (40) for $a = 0.1$], which is otherwise the one we don't know the $O(a^3)$ coefficient. We can take advantage that the $O(a^3)$ correction to $[a]$ is almost completely given by the known L contribution. Using $d_3^{L+T} - d_2^{L+T} (d_2^{L+T} = d_1^{L+T}) \approx 317$, the fourth term in (42) becomes 0.298, i.e. a 5% reduction only. Taking the size of the $O(a^3)$ contribution to L as an educated estimate of the perturbative uncertainty, we finally get

$$[0:1] = 2.1 \pm 0.3 \quad (43)$$

6. Renormalization {Scale Dependence

The expansion (22) depends order by order on μ and this dependence cancels out only when we sum the infinite series. In practice, we only know a few first terms of the series (three for $L+T$ $[a]$ and four for L $[a]$); so we should worry how much the predictions depend on our previous choice $\mu = 1$. Obviously, μ should be close to one in order to avoid large logarithms; but variations within a reasonable range, let us say from 0.75 to 2, should not affect too much the final results. Smaller values of μ would put the QCD coupling in the non-perturbative regime and are therefore not acceptable.

Figures 1, 2 and 3 show the sensitivity to the selection of renormalization scale of the final predictions for $L+T$, L , and T , respectively, for $(M^2) = 0.1$.

The behaviour of $L+T$ is quite good. The predicted value remains very stable in the whole range $\mu \in [0.75; 2]$, showing that the perturbative series is very reliable. Below $\mu = 1$, the perturbative expansion breaks down, as expected, because the coupling a is already outside the radius of convergence of the series.

The longitudinal series, on the other side, has a quite wild dependence on the renormalization scale. Changing μ from 1 to 2, amounts to a reduction of L of about 65%. Thus, the theoretical uncertainty is very large in this case.

The μ dependence of the complete expansion $[a]$, reflects obviously the behaviour of its two components. The larger weight of $L+T$ keeps the result still acceptable, within the range of μ considered, but the sizeable L contribution

spoils the stability and generates a monotonic decrease of the prediction for increasing values of α_s . Taking this variation into account, the theoretical error in Eq. (43) should be increased to about 0.6, i.e. a 30% uncertainty in the final prediction.

6. Discussion

The bad perturbative behaviour of the longitudinal contribution does not allow to make an accurate determination of the strange quark mass from R_{us} . Nevertheless, taking

$$|D_1| = 2.1 \pm 0.6; \quad (44)$$

$m_s(M^2)$ could be still obtained with a theoretical uncertainty of about 15%, which is not so bad.

Notice that it is the phase space integration of the original correlation functions the responsible for the different behaviour of the longitudinal and transverse components. Therefore, the perturbative convergence could probably be improved through an appropriate use of weight factors in Eqs. (5) and (8). This requires an accurate measurement of the final hadrons mass distribution in the decay, which so far has only been performed for the dominant Cabibbo-allowed modes [31]. The measurement of $R_{\text{us}}(s)$ could be feasible at the forthcoming flavour factories, where a very good kaon identification is foreseen.

From the theoretical point of view, the analysis of weighted moments of the final hadrons mass distribution proceeds in a completely analogous way [32]. A detailed study will be presented in a forthcoming publication.

Acknowledgements

We would like to thank Kostja Chetyrkin for informing us about the existence of a misprint in Ref. [16]. We have benefited from many discussions (and program exchanges) with Matthias Jamnitsch and Arcadi Santamaría, concerning the numerical implementation of running effects at higher orders. Useful discussions with Michel Davier, Andreas Hocker and Eduardo de Rafael are also acknowledged. This work has been supported in part by the European Union TMR Network EURODAPHNE | Contract No. ERBFMRX-CT98-0169 (DG 12 { M IHT})| and by CICYT, Spain, under Grants No. AEN-96/1672 and AEN-96/1718.

References

- [1] E. Braaten, Phys. Rev. Lett. 60 (1988) 1606; Phys. Rev. D 39 (1989) 1458.
- [2] S. Narison and A. Pich, Phys. Lett. B 211 (1988) 183.
- [3] E. Braaten, S. Narison, and A. Pich, Nucl. Phys. B 373 (1992) 581.
- [4] F. Le Diberder and A. Pich, Phys. Lett. B 286 (1992) 147.
- [5] A. Pich, QCD Tests from Tau Decays, Proc. of the 20th Johns Hopkins Workshop on Current Problems in Particle Theory | Non-Perturbative Particle Theory and Experimental Tests | (Heidelberg, June 1996), eds. M. Jamín et al (World Scientific, Singapore, 1997), p. 247.
- [6] S. Chen, Nucl. Phys. B (Proc. Suppl.) 64 (1998) | .
- [7] M. Davier, Nucl. Phys. B (Proc. Suppl.) 55C (1997) 395.
- [8] C. Becchi et al, Z. Phys. C 8 (1981) 335.
- [9] C. A. Dominguez and E. de Rafael, Ann. Phys., NY 174 (1987) 372.
- [10] M. Jamín and M. Münz, Z. Phys. C 66 (1995) 633.
- [11] J. Bijnens, J. Prades, and E. de Rafael, Phys. Lett. B 348 (1995) 226.
- [12] M. Jamín, Nucl. Phys. B (Proc. Suppl.) 64 (1998) 250.
- [13] J. Prades, Nucl. Phys. B (Proc. Suppl.) 64 (1998) 253.
- [14] K. G. Chetyrkin, Phys. Lett. B 390 (1997) 309.
- [15] P. M. Stevenson, Phys. Rev. D 23 (1981) 2916.
- [16] K. G. Chetyrkin and A. Kwiatkowski, Z. Phys. C 59 (1993) 525.
- [17] K. G. Chetyrkin and J. H. Kühn, Phys. Lett. B 406 (1997) 102.
- [18] M. A. Shifman, A. L. Vainshtein, and V. I. Zakharov, Nucl. Phys. B 147 (1979) 385; 448; 519.
- [19] W. J. Marciano and A. Sirlin, Phys. Rev. Lett. 61 (1988) 1815; 56 (1986) 22.
- [20] E. Braaten and C. S. Li, Phys. Rev. D 42 (1990) 3888.
- [21] T. van Ritbergen, J. A. M. Vermaseren, and S. A. Larin, Phys. Lett. B 400 (1997) 379; B 405 (1997) 327.

- [22] K.G. Chetyrkin, B.A. Kniehl, and M. Steinhauser, Phys. Rev. Lett. 79 (1997) 2184.
- [23] K.G. Chetyrkin, Phys. Lett. B 404 (1997) 161.
- [24] L.R. Surguladze, Sov. J. Nucl. Phys. 50 (1989) 372.
- [25] S.G. Gorishny et al, Mod. Phys. Lett. A 5 (1990) 2703.
- [26] S.G. Gorishny et al, Phys. Rev. D 43 (1991) 1633.
- [27] S.G. Gorishny, A.L. Kataev, and S.L. Larin, Nuovo Cimento 92 (1986) 119.
- [28] W. Bernreuther and W. Wetzel, Z. Phys. C 11 (1981) 113.
- [29] K. Maltman, Problem with Extracting m_s from Flavor Breaking in Hadronic Decays, York preprint YU-PP-K/M-98-1, hep-ph/9804298.
- [30] K.G. Chetyrkin, private communication.
- [31] R. Barate et al (ALEPH Coll.), Measurement of the Spectral Functions of Axial-Vector Hadronic Decays and Determination of $m_s(M^2)$, preprint CERN-EP-98-012
- [32] F. Le Diberder and A. Pich, Phys. Lett. B 289 (1992) 165.

Figure Captions

Figure 1.-Variation of $\alpha_{L+T} [0:1]$ with the renormalization {scale factor μ , to four loops.

Figure 2.-Variation of $\alpha_L [0:1]$ with the renormalization {scale factor μ , to four loops.

Figure 3.-Variation of $\alpha [0:1]$ with the renormalization {scale factor μ , to four loops.




Figure 1: Variation of $\Delta_{L+T}[0.1]$ with the renormalization scale factor ξ , to four loops.




Figure 2: Variation of $\Delta_L[0.1]$ with the renormalization scale factor ξ , to four loops.




Figure 3: Variation of $\Delta[0.1]$ with the renormalization scale factor ξ , to four loops.