

Gauge mediated supersymmetry breaking and neutrino anomalies

Anjan S. Joshipura and Sudhir K. Vempati

*Theoretical Physics Group, Physical Research Laboratory,
Navarangpura, Ahmedabad, 380 009, India.*

Abstract

Supersymmetric standard model with softly broken lepton symmetry provides a suitable framework to accommodate the solar and atmospheric neutrino anomalies. This model contains a natural explanation for large mixing and hierarchical masses without fine tuning of the parameters. Neutrino spectrum is particularly constrained in the minimal messenger model (MMM) of gauge mediated SUSY breaking, since all SUSY breaking effects are controlled in MMM by a single parameter. We study the structure of neutrino masses and mixing both in MMM and in simple extensions of it in the context of solar and atmospheric neutrino anomalies.

I. INTRODUCTION

The hints in favour of a non-zero neutrino mass are greatly strengthened by the recent high statistics results of the atmospheric ν_μ deficit at the Superkamioka [1]. These results are claimed to be the evidence of a non-zero neutrino (mass)² difference $\Delta_A \sim (.07 \text{ eV})^2$ and large mixing between ν_μ and ν_τ or a sterile state. Additional hint for one more mass scale comes from the observed deficit in the solar neutrinos. These results require [2] the solar mass scale Δ_S of $10^{-5} - 10^{-6} \text{ eV}^2$ (MSW conversion [3]) or 10^{-10} eV^2 (vacuum oscillations). While the former can be consistent with the small mixing of ν_e , the latter explanation requires [2] one more large mixing. On the basis of these results, the neutrino spectrum seems to be characterized by hierarchical masses and one or two large mixings. The above neutrino spectrum is quite different from the quark spectrum suggesting a characteristically different origin for the neutrino masses and mixing.

The standard seesaw model [4] based on simple $SO(10)$ theory links respectively the ν -masses and leptonic mixing to the up quark masses and the KM matrix. It leads to $\frac{\Delta_S}{\Delta_A} \sim \left(\frac{m_c}{m_t}\right)^4 \sim 10^{-8}$ if simplifying assumptions are made on the structure of the right handed neutrino masses [4]. With $\Delta_A \sim 5 \times 10^{-3} \text{ eV}^2$ one obtains $\Delta_S \sim 5 \times 10^{-11} \text{ eV}^2$ in the range required for the vacuum solution to the solar anomalies. The MSW and the atmospheric mass scales cannot be easily reconciled in this simple picture although additional contribution from a left handed triplet [5] field can be used to do so. Moreover, large mixing angles though not impossible [6] are not natural in the seesaw picture.

We wish to discuss an alternative scheme for neutrino masses based on supersymmetry. The scheme utilizes soft lepton number violation and soft supersymmetry (SUSY) breaking terms generated by gauge mediated interactions. As we will see, this scheme is quite predictive and can provide natural and simultaneous understanding of the hierarchical masses and large mixings in the leptonic sector hinted on experimental grounds.

The Minimal Supersymmetric Standard Model (MSSM) naturally contains lepton number violation if the conventional R symmetry is not imposed on it. This violation is manifested through hard (trilinear) or soft (bilinear) operators [7]. The soft terms are parameterized in terms of three dimensionful co-efficients ϵ_i :

$$W = \epsilon_i L'_i H_2 \tag{1}$$

We will assume that eq (1) is the only source of lepton number violation. This assumption is theoretically well motivated. Spontaneous breaking of lepton number [8] could normally result in such a term. Alternatively, one could imagine a generalized Peccei-Quinn symmetry whose spontaneous breaking leads to μ and ϵ_i at the weak scale through dim 5 operators [9,10]. Moreover, it is possible to choose the PQ charges of different fields in such a way that the generation of effective trilinear operators is enormously suppressed.

The presence of ϵ_i leads to neutrino masses and mixing among them. In fact, these three parameters control three neutrino masses and three mixing angles. This is easily seen from the fact that the limit $\epsilon_i \rightarrow 0$ corresponds to no lepton number violation, zero neutrino masses and trivial mixing in the leptonic sector. In practice, the neutrino masses also depend upon parameters responsible for the SUSY breaking. But this breaking can be characterized by only one parameter in a minimal version of the gauge mediated breaking [11] of SUSY. This scenario can therefore provide a constrained framework for the description of the neutrino masses and mixing in which four input parameters determine three masses and three mixing angles.

The neutrino spectrum resulting from eq.(1) has been extensively studied in the context

of MSSM with supergravity induced SUSY breaking [10,12]. The following features make it a very attractive scenario for the description of the neutrino spectrum:

(i) Neutrinos obtain masses [7] through two sources: through mixing with gauginos induced by sneutrino vacuum expectation value (vev) and through coupling of neutrinos to squarks and sleptons. Both these contributions are suppressed by the b and τ Yukawa couplings in the supergravity framework as well as in the gauge mediated models. This leads to suppressed (\sim MeV) neutrino masses even when ϵ_i are large (\sim 100 GeV).

(ii) The sneutrino vev makes only one combination of neutrinos massive. The other combinations pick up masses from the loop diagrams [13]. In particular, one of the neutrinos remains massless when Yukawa couplings of the first two generations are neglected [10]. This generates clear hierarchy in masses of all the three neutrinos without fine tuning or without imposing any horizontal symmetries.

(iii) Mixing among neutrinos is essentially controlled [14] by ratios of ϵ_i which can be large when these parameters are not hierarchical. One can therefore simultaneously obtain large mixing and hierarchal masses.

The above features (ii) and (iii) can allow simultaneous solution of the solar and atmospheric neutrino anomalies.

We discuss the details of this in the following section. Section (2) summarizes the analytic structure of neutrino masses and mixing one obtains in the presence of eq.(1). In the next section, we summarize salient features of the gauge mediated models and introduce the minimal model in this category. Section (4) contains detailed prediction of this model and phenomenological discussions on the solar and the atmospheric neutrino problem. We introduce an extended version of the minimal model which is capable of solving the solar and atmospheric neutrino anomalies simultaneously in Section (5). We end with a discussion in the last section.

II. NEUTRINO MASSES AND SUSY

The structure of neutrino masses crucially depends upon the nature of SUSY breaking soft terms. It is now recognized [10,12] that neutrino masses are calculable in terms of basic parameters if soft terms associated with the leptonic and one of the Higgs doublet (H'_1) superfields are identical at some high scale Λ . This happens in the minimal supergravity model as well as in gauge mediated models. The latter class of models contain fewer parameters and thus are more predictive. We will specialize to this case in the next section. The detailed analytic discussion of the structure of neutrino masses in the present context was

given in [10]. Here we briefly recall some salient features to be used in this paper. Consider the full superpotential of the MSSM [15] including the lepton number violating term (1):

$$W = \lambda_i L'_i E_i^c H'_1 + \mu' H'_1 H_2 + \epsilon_i L'_i H_2 + \lambda_i^D Q_i D_i^c H'_1 + \lambda_{ij}^U Q_i U_j^c H_2, \quad (2)$$

where, $i, j = 1, 2, 3$. We have made a specific choice of the basis such that $L'_i(Q_i)$ denote the mass eigenstates of the charged lepton (down quarks) in the absence of the ϵ_i terms. Following [7] we now redefine the leptonic and the Higgs fields in such a way that the superpotential (2) does not contain bilinear ϵ_i -dependent terms. This happens in the following unprimed basis:

$$\begin{aligned} H_1 &= c_3 H'_1 + s_3(s_2(c_1 L'_2 + s_1 L'_1) + c_2 L'_3), \\ L_1 &= -s_1 L'_2 + c_1 L'_1, \\ L_2 &= c_2(c_1 L'_2 + s_1 L'_1) - s_2 L'_3, \\ L_3 &= -s_3 H'_1 + c_3(s_2(c_1 L'_2 + s_1 L'_1) + c_2 L'_3). \end{aligned} \quad (3)$$

Where,

$$\begin{aligned} \epsilon_1 &= \mu s_1 s_2 s_3, & \epsilon_2 &= \mu c_1 s_2 s_3, \\ \epsilon_3 &= \mu s_3 c_2, & \mu' &= \mu c_3, \end{aligned} \quad (4)$$

and $\mu \equiv (\mu'^2 + \epsilon_1^2 + \epsilon_2^2 + \epsilon_3^2)^{1/2}$.

The superpotential in the new basis contains trilinear lepton number violating terms. Moreover, the originally diagonal charged lepton mass matrix M_l now acquires [10] non-diagonal parts [16]:

$$\frac{M_l}{\langle H_1 \rangle} = \begin{pmatrix} \lambda_1 c_1 c_3 & -\lambda_2 s_1 c_3 & 0 \\ \lambda_1 s_1 c_2 c_3 & \lambda_2 c_1 c_2 c_3 & -\lambda_3 s_2 c_3 \\ \lambda_1 s_1 s_2 & \lambda_2 c_1 s_2 & \lambda_3 c_2 \end{pmatrix} \quad (5)$$

The mass basis for the charged leptons in the presence of non-zero ϵ_i are denoted by $\alpha = e, \mu, \tau$ and are defined as:

$$L_i = (O_L^T)_{i\alpha} L_\alpha, \quad e_i^c = (O_R^T)_{i\alpha} e_\alpha^c, \quad (6)$$

where,

$$O_L M_l O_R^T = \text{diagonal}. \quad (7)$$

Note that the parameters λ_i which denote charged lepton masses when $\epsilon_i = 0$ still need to be hierarchical if M_l in (5) is to reproduce the charged lepton masses. One can determine [10] O_L by assuming $\lambda_1 \ll \lambda_2 \ll \lambda_3$ and neglecting λ_1 :

$$O_L^T \approx N_1 \begin{pmatrix} c_1 & -s_1 N_2 & 0 \\ s_1 c_2 & c_1 c_2 N_2^{-1} & -s_2 c_3 N_1^{-1} N_2^{-1} \\ s_1 s_2 c_3 & c_1 s_2 c_3 N_2^{-1} & c_2 N_1^{-1} N_2^{-1} \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos \theta_{23} & -\sin \theta_{23} \\ 0 & \sin \theta_{23} & \cos \theta_{23} \end{pmatrix}. \quad (8)$$

Where,

$$\begin{aligned} N_1 &= (1 - s_1^2 s_2^2 s_3^2)^{(-1/2)}, \\ N_2 &= (1 - s_2^2 s_3^2)^{(1/2)}, \\ \theta_{23} &\approx N_1 c_1 c_2 s_2 \left(\frac{s_3 m_\mu}{c_3 m_\tau} \right)^2. \end{aligned} \quad (9)$$

θ_{23} is small even for $s_{1,2,3} \sim O(1)$. We shall therefore neglect it.

The trilinear terms generated in the superpotential due to rotation (3) assume the following form [10] in the physical mass eigenstate basis of charged leptons:

$$W = -\frac{\tan \theta_3}{\langle H_1 \rangle} [(O_L^T)_{3\alpha} L_\alpha] (m_\beta^l L_\beta e_\beta^c + m_i^D Q_i d_i^c). \quad (10)$$

The above trilinear terms generated due to the basis rotation lead to neutrino masses at 1-loop [13]. The other contribution to neutrino mass is generated by the soft SUSY breaking terms in a manner discussed below.

A. Sneutrino vevs and neutrino masses

The soft supersymmetry breaking part of the scalar potential contains linear terms in sneutrino fields and lead to their vev. These follow from the following soft terms written in the primed basis:

$$\begin{aligned} V_{soft} &= m_{H_1^2} |H_1^{0\prime}|^2 + m_{H_2^2} |H_2^0|^2 + m_{\tilde{\nu}_i^{\prime 2}} |\tilde{\nu}_i^{\prime}|^2 \\ &- [H_2^0 (\mu' B_\mu H_1^{0\prime} + \epsilon_i B_i \tilde{\nu}_i^{\prime}) + H.c.]. \end{aligned} \quad (11)$$

It is always possible to choose a minimum with zero sneutrino vev if $B_\mu = B_i$ and $m_{H_1^2} = m_{\tilde{\nu}_i^{\prime 2}}$. But, these equalities are generally not satisfied at the weak scale even if they are imposed at a high scale. This is due to the presence of Yukawa couplings which

distinguish between Higgs and leptons. If one neglects the Yukawa couplings of the first two generations then the soft mass parameters related to the first two leptonic generations evolve in the same way. Thus, one would have the following non-zero differences among low energy parameters:

$$\Delta m_{L,H} \equiv (m_{\tilde{\nu}'_3}{}^2 - m_{\tilde{\nu}'_2, H_1}^2), \quad \Delta B_{L,H} \equiv (B_3 - B_{2,\mu}). \quad (12)$$

These differences determine the sneutrino vev. The latter are obtained by minimizing the scalar potential expressed in the redefined basis of eqs.(3). One finds:

$$\begin{aligned} V = & (m_{H_1}^2 + \mu^2)|H_1^0|^2 + (m_{H_2}^2 + \mu^2)|H_2^0|^2 + m_{\tilde{\nu}_3}^2|\tilde{\nu}_3|^2 + m_{\tilde{\nu}'_2}^2|\tilde{\nu}_2|^2 + m_{\tilde{\nu}'_1}^2|\tilde{\nu}_1|^2 \\ & - \left[\mu H_2^0 \left(B H_1^0 + c_3 s_3 \tilde{\nu}_3 (\Delta B_H - s_2^2 \Delta B_L) - c_2 s_2 s_3 \tilde{\nu}_2 \Delta B_L \right) + H.c. \right] \\ & + \left[-c_2 s_2 \Delta m_L (s_3 H_1^0 + c_3 \tilde{\nu}_3) \tilde{\nu}_2^* + s_3 c_3 \tilde{\nu}_3 H_1^{0*} (\Delta m_H - s_2^2 \Delta m_L) + H.c. \right] \\ & + \frac{1}{8} (g^2 + g'^2) (|H_1^0|^2 + |\tilde{\nu}_1|^2 + |\tilde{\nu}_2|^2 + |\tilde{\nu}_3|^2 - |H_2^0|^2)^2. \end{aligned} \quad (13)$$

Where,

$$\begin{aligned} m_{H_1}^2 &= m_{H'_1}^2 + s_3^2 \Delta m_H - s_2^2 s_3^2 \Delta m_L, \\ m_{\tilde{\nu}_3}^2 &= m_{\tilde{\nu}'_3}^2 - s_3^2 \Delta m_H - s_2^2 c_3^2 \Delta m_L, \\ m_{\tilde{\nu}'_2}^2 &= m_{\tilde{\nu}'_2}^2 + s_2^2 \Delta m_L, \\ B &= B_\mu + s_3^2 \Delta B_H - s_2^2 s_3^2 \Delta B_L. \end{aligned} \quad (14)$$

The additional terms do not significantly effect the vev for the standard Higgs since the sneutrino vevs and $\Delta m_{H,L}$, $\Delta B_{H,L}$ are suppressed compared to m_{SUSY}^2 . This remains true even for $s_3 \sim O(1)$. In addition, we will be considering $s_3 \ll 1$ to account for the atmospheric neutrino scale. Hence, MSSM results are hardly altered by additional terms. The effect of these terms is to induce vevs for the sneutrinos:

$$\begin{aligned} \langle \tilde{\nu}_2 \rangle &\approx \frac{c_2 s_2 s_3}{(m_{\tilde{\nu}'_2}^2 + D)} (v_1 \Delta m_L - \mu v_2 \Delta B_L), \\ \langle \tilde{\nu}_3 \rangle &\approx \frac{c_3 s_3}{(m_{\tilde{\nu}'_3}^2 + D)} (v_1 (-\Delta m_H + s_2^2 \Delta m_L) - \mu v_2 (-\Delta B_H + s_2^2 \Delta B_L)). \end{aligned} \quad (15)$$

We have neglected terms higher order in $\Delta m_{L,H}$, $\Delta B_{H,L}$ while writing the above equations. Note that one of the sneutrino field ($\equiv \tilde{\nu}_1$) does not acquire a vev in this basis. This vev would arise if Yukawa couplings of the first two generations neglected here are turned on.

The sneutrino vevs are zero at the boundary scale Λ corresponding to the universal masses. Their weak scale values are determined by solving the relevant RG equations involving the above differences:

$$\begin{aligned}
\frac{d \Delta m_H}{d t} &= 3Y_b(t)(m_{H'_1}^2 + m_{\tilde{b}}^2 + m_{\tilde{b}^c}^2 + A_b^2) , \\
\frac{d \Delta m_L}{d t} &= -Y_\tau(t)(m_{H'_1}^2 + m_{\tilde{\tau}}^2 + m_{\tilde{\tau}^c}^2 + A_\tau^2) , \\
\frac{d \Delta B_H}{d t} &= 3Y_b(t)A_b(t) , \\
\frac{d \Delta B_L}{d t} &= -Y_\tau(t)A_\tau .
\end{aligned} \tag{16}$$

Where, $Y_f \equiv \frac{\lambda_f^2}{(4\pi)^2}$, $m_{\tilde{f}}$ is the mass of the sfermion concerned, A_f are the trilinear soft SUSY breaking terms and $t = 2 \ln(M_{GUT}/Q)$.

The tree level mass matrix generated due to these vev can be written in the physical basis ν_α as:

$$M_0 = m_0 O_L \begin{pmatrix} 0 & 0 & 0 \\ 0 & s_\phi^2 & s_\phi c_\phi \\ 0 & s_\phi c_\phi & c_\phi^2 \end{pmatrix} O_L^T . \tag{17}$$

Where,

$$\tan \phi = \frac{\langle \tilde{\nu}_2 \rangle}{\langle \tilde{\nu}_3 \rangle} . \tag{18}$$

O_L is defined by eqs.(7) and,

$$m_0 = \frac{\mu(cg^2 + g'^2)(\langle \tilde{\nu}_2 \rangle^2 + \langle \tilde{\nu}_3 \rangle^2)}{2(-c\mu M + 2M_W^2 c_\beta s_\beta (c + \tan^2 \theta_W))} . \tag{19}$$

B. 1-loop mass

The trilinear interactions in eq. (10) lead to diagrams involving squarks and sleptons in the loop and generate the neutrino masses at the 1-loop level [13]. These contributions depend upon the masses as well as mixing between the left and the right handed squarks as well as the sleptons. These are however fixed in terms of the basic parameters of the MSSM. In the present case, the trilinear couplings are not independent and are controlled by the fermion masses. As a consequence, the dominant contribution arises when the b-squark or τ slepton are exchanged in the loop. We shall retain only this contribution.

Let us define:

$$\begin{aligned}
\tilde{b} &= \tilde{b}_1 \cos \phi_b + \tilde{b}_2 \sin \phi_b , \\
\tilde{b}^{c\dagger} &= \tilde{b}_2 \cos \phi_b - \tilde{b}_1 \sin \phi_b .
\end{aligned} \tag{20}$$

Where, $\tilde{b}_{1,2}$ are the mass eigenstates with masses M_{b_1,b_2} respectively. The mixing angles ϕ_τ and masses M_{τ_1,τ_2} are defined analogously in case of the tau slepton.

The exchange of b-squark produces the following mass matrix for the neutrinos:

$$(M_{1b})_{\alpha\beta} = m_{1b}(O_L)_{\alpha 3}(O_L)_{\beta 3}. \quad (21)$$

Due to the antisymmetry of the leptonic couplings in eq. (10), the exchange of the τ slepton leads to the following contribution:

$$M_{1\tau} = m_{1\tau} \begin{pmatrix} O_{L13}^2 & O_{L13}O_{L23} & 0 \\ O_{L13}O_{L23} & O_{L23}^2 & 0 \\ 0 & 0 & 0 \end{pmatrix}. \quad (22)$$

The mixing induced by these contributions is completely fixed by the matrix O_L while the overall scale of both these contributions is set by,

$$m_{1b,1\tau} = N_c \frac{m_{b,\tau}^3}{16\pi^2 v_1^2} \tan^2 \theta_3^2 \sin \phi_{b,\tau} \cos \phi_{b,\tau} \ln \left(\frac{M_{b_2,\tau_2}^2}{M_{b_1,\tau_1}^2} \right). \quad (23)$$

Where, $N_c = 3, 1$ for the \tilde{b} and $\tilde{\tau}$ contribution respectively. The total mass matrix including the 1-loop corrections is given by,

$$\mathcal{M}_\nu = M_0 + M_{1b} + M_{1\tau}. \quad (24)$$

We stress that the above \mathcal{M}_ν is in the physical basis with diagonal charged lepton masses. This matrix assumes particularly simple form when rotated by the matrix O_L :

$$O_L^T \mathcal{M}_\nu O_L \approx \begin{pmatrix} 0 & 0 & 0 \\ 0 & m_0 s_\phi^2 + m_{1\tau} N_2^{-4} c_2^2 s_2^2 c_3^2 & m_0 s_\phi c_\phi + m_{1\tau} N_2^{-4} c_2 s_2^3 c_3^3 \\ 0 & m_0 s_\phi c_\phi + m_{1\tau} N_2^{-4} c_2 s_2^3 c_3^3 & m_0 c_\phi^2 + m_{1b} + m_{1\tau} N_2^{-4} s_2^4 c_3^4 \end{pmatrix} \quad (25)$$

This explicitly shows that one of the neutrinos is massless in our approximation of neglecting Yukawa couplings of the first two generations.

The full mixing matrix analogous to the KM matrix is given by,

$$U = O_L O_\nu^T. \quad (26)$$

Where, O_ν is the matrix diagonalizing the RHS of eq.(25). As we will show, the mixing angle appearing in O_ν is small due to hierarchy in neutrino masses while, the O_L can contain large

mixing. Hence, the neutrino masses are determined by the matrix (25) and mixing among neutrinos is essentially fixed by eq.(8). We shall use these equations in specific case of the gauge mediated models in the next section.

The above formalism shows that the neutrino masses are greatly suppressed compared to the typical SUSY breaking scale if $\Delta m_{H,L}, \Delta B_{H,L}$ vanish at some scale Λ . The weak scale values of sneutrino vev and hence neutrino masses follow from evolution of these parameters. It is clear from eq.(16) that the b and τ Yukawa couplings control the evolution of sneutrino vev. Similarly, the 1-loop masses following from eq. (23) are also controlled by the same couplings. As a result, all the effects of lepton number violating parameters ϵ_i can be rotated away from the full Lagrangian when the down quark and the charged lepton Yukawa couplings vanish. Neutrino masses also vanish in this limit.

III. GAUGE MEDIATED MODELS AND NEUTRINO MASSES

The suppression in neutrino masses mentioned above crucially depends upon vanishing of $\Delta m_{H,L}, \Delta B_{H,L}$ at some scale. This happens in two of the most popular scenarios of supersymmetry breaking namely, supergravity induced breaking in its minimal form and the gauge mediated breaking of SUSY [11,17,18]. We now discuss neutrino masses in the latter context.

The basic approach adopted in most gauge mediated models of SUSY breaking is to assume that a singlet sector is responsible for such breaking. Effect of this is felt by the standard fields through a messenger sector which is a gauge non-singlet. The minimal version of this sector contains a pair of fields $\Phi, \bar{\Phi}$ transforming as $5+\bar{5}$ representation of the $SU(5)$ group. Their coupling to a SUSY breaking field S introduces a supersymmetric mass scale $X \equiv \lambda \langle S \rangle$ as well as a SUSY breaking (mass)² differences of order F_S . Models with minimal messenger sector are thus characterized by two parameters $\Lambda \equiv \frac{F_S}{X}$ and $x \equiv \frac{\Lambda}{X}$ with $x \sim O(1)$ on natural grounds.

All the soft parameters related to MSSM fields are fixed at Λ in terms of Λ, x and the gauge couplings. The masses of the squarks and sleptons and the gauginos are given in this case by:

$$\begin{aligned} m_i^2(X) &= 2\Lambda^2 \left\{ C_3 \tilde{\alpha}_3^2(X) + C_2 \tilde{\alpha}_2^2(X) + \frac{3}{5} Y^2 \tilde{\alpha}_1^2(X) \right\} f(x), \\ M_j(X) &= \tilde{\alpha}_j(X) \Lambda g(x). \end{aligned} \tag{27}$$

m_i^2 represents the scalar masses with i running over all the scalars, whereas, M_j represents

the gaugino masses with j representing the three gauge couplings. The functions $f(x)$ and $g(x)$ have been derived in [19]. Here,

$$\tilde{\alpha}_j(X) = \frac{\alpha_j(X)}{(4\pi)}; \quad (28)$$

$C_3 = 4/3, 0$ for triplets and singlets of $SU(3)_C$, $C_2 = 3/4, 0$ for doublets and singlets of $SU(2)_L$ and $Y = Q - T_3$ is the hypercharge.

In this paper, we will consider two different versions of the model. The popular [11,17,20,21] minimal messenger model (MMM) which is further characterized by the assumption of the vanishing bilinear (B) and trilinear (A) soft mass parameters at Λ . This is attractive in view of the very restricted structure it offers. But as we will show it turns out to be too restrictive if one wants to solve the solar and atmospheric neutrino problems simultaneously. We shall thus consider an alternative version on phenomenological grounds in which the boundary conditions (27) are still imposed but the value of B at Λ is not taken to be zero.

In MMM, all the soft parameters in the low energy theory are essentially determined by one parameter Λ since dependence of the boundary condition eq.(27) on x is very mild. In particular, the value of the B parameter at the weak scale gets fixed through its running. This in turn determines both μ as well as $\tan\beta$ through the following equations:

$$\begin{aligned} \mu^2 &= \frac{m_{H_1}^2 - m_{H_2}^2 \tan^2\beta}{\tan^2\beta - 1} - \frac{1}{2}M_Z^2, \\ \text{Sin}2\beta &= \frac{2B\mu}{m_{H_2}^2 + m_{H_1}^2 + 2\mu^2}. \end{aligned} \quad (29)$$

The presence of the ϵ_i induces corrections to these equations, but they are very small as discussed below eq.(14). The eq.(29) therefore holds to a very good approximation.

In spite of the restricted structure, it is possible to self consistently solve the above equations [17,20,21] and implement breaking of the $SU(2) \times U(1)$ symmetry at low energy. Vanishing of the soft B parameter at Λ makes the analysis of this breaking little more involved than in the case of the supergravity induced breaking. One needs to include two loop corrections to the evolution of the B parameter and also needs to use fully one loop corrected effective potential. Details of this analysis are presented in [17,20,21]. We follow the treatment given in [21]. The smallness of the B at the weak scale results in this scheme in relatively large value of $\tan\beta$ and its sign fixes the sign of μ to be positive. The full 1-loop corrected potential was employed in the analysis of [21] but it was found that working with RG improved tree level potential also gives similar results provided one evolves soft parameters of the supersymmetric partners up to a scale $Q_0^2 \equiv (m_Q^2(X)m_{\bar{U}}^2(X))^{\frac{1}{2}}$. We prefer

to follow this approach and use the RG improved tree level potential of eq.(13) in order to determine the low energy parameters at the minimum. We have however included two loop corrections to the RG equations [22] for B and $\Delta B_{L,H}$ in determining their values at the weak scale. Use of RG improved tree level potential allows us to analytically understand the structure of neutrino masses and mixing in a transparent way.

The three mass parameters $m_{0,1b,1\tau}$ introduced earlier control the neutrino masses. m_0 is determined by solving RG equations (16) along with similar ones for parameters occurring in them. We have numerically solved them imposing eq.(27) as boundary conditions at Λ . We evolved these equations self consistently up to the scale Q_0 defined above. The m_0 determined in this manner depends upon μ as well as $\tan\beta$ both of which are fixed in terms of Λ .

The loop contributions are fixed in terms of the squark and slepton masses and mixings defined in eq. (20). These are determined from the standard 2×2 matrices involving left and right squarks and slepton mixing. The elements in these matrices are also completely fixed in terms of Λ . All the three parameters $m_0, m_{1b}, m_{1\tau}$ depend upon an overall scale s_3 of the R breaking. For small s_3^2 they roughly scale as s_3^2 . The ratios $m_0/s_3^2, m_{1b}/s_3^2, m_{1\tau}/s_3^2$ are thus determined by Λ alone [23]. We have displayed in Fig. 1 variations of $\frac{m_0}{\text{GeV}s_3^2}, \frac{m_{1b}}{m_0}$ and $\frac{m_{1\tau}}{m_0}$ with Λ . One clearly sees hierarchy in the loop and sneutrino vev induced contributions. This hierarchy gets reflected in the neutrino masses and one obtains hierarchical neutrino masses independent of the overall strength of the R violating parameter s_3 . The mass ratio and hence the hierarchy among neutrino masses are seen to be less sensitive to Λ . The m_0 roughly scales linearly with Λ . But since the over all scale of m_0 is set by s_3^2 which is also unknown a change in Λ is equivalent to changing s_3 . Thus, we may use one specific value of Λ and neutrino mass spectrum is then completely fixed by three angles $s_{1,2,3}$ or equivalently by the three R violating parameters ϵ_i .

The $\Delta m_{H,L}, \Delta B_{H,L}$ entering m_0 are determined from the RG equations (16) and are fixed in terms of Λ . For example,

$$\begin{aligned} \mu &\sim 397.0 \text{ GeV} , & \tan\beta &\sim 46.39 , \\ \Delta m_H &\sim 192661.23 \text{ GeV}^2 , & \Delta m_L &\sim -2392.35 \text{ GeV}^2 , \\ \Delta B_H &\sim -14.07 \text{ GeV} , & \Delta B_L &\sim 0.12 \text{ GeV} , \end{aligned} \tag{30}$$

when $\Lambda = 100 \text{ TeV}$. The suppression in $\Delta m_L, \Delta B_L$ is due to color factors and larger squark masses compared to the slepton masses in the model. It follows that the ratio $\tan\phi$ of the sneutrino vev, eq. (18) gets considerably suppressed even when the angle s_2 is large. We

show in Fig 2. the value of s_ϕ^2 as function of s_2 for $\Lambda = 100 \text{ TeV}$. Note that this ratio is independent of the values of the other R violating parameters when s_3 is small.

The small value of s_ϕ leads to very simple expression for neutrino masses. The neutrino mass matrix in eq.(25) is almost diagonal and one finds:

$$\begin{aligned} m_{\nu_3} &\sim m_0 + m_{1b} , \\ \frac{m_{\nu_2}}{m_{\nu_3}} &\sim c_2^2 s_2^2 \frac{m_{1\tau}}{(m_0 + m_{1b})} \sim c_2^2 s_2^2 (7.1 \times 10^{-3} - 5.6 \times 10^{-3}) , \\ \theta_{23}^\nu &\sim \frac{m_{1\tau} c_2 s_2^3}{(m_0 + m_{1b})} \sim \tan \theta_2 \frac{m_{\nu_2}}{m_{\nu_3}} . \end{aligned} \quad (31)$$

The masses are fixed in terms of $m_{0,1b,1\tau}$ which are determined in terms of Λ and s_3 . The mass ratio is fixed in terms of s_2 . The range indicated on the RHS in above equation corresponds to variation in Λ from (51 TeV- 150 TeV) and θ_{23}^ν represents the angle diagonalizing the matrix in eq.(25).

IV. NEUTRINO MASSES: PHENOMENOLOGY

As discussed in the last section, the model considered here implies hierarchical masses and large mixing without any fine tuning of the parameters. We now try to see if the predicted spectrum can be used to simultaneously reconcile both the solar and the atmospheric neutrino anomalies. The model is quite constrained. Three neutrino masses and three mixing angles get completely determined in the model in terms of four parameters namely, Λ and three R violating angles $s_{1,2,3}$. In particular, the angle s_1 characterizing the electron number violation does not enter the muon and tau neutrino masses, see eq.(25). The mixing between neutrinos is largely fixed by the matrix O_L with a small correction coming from the angle θ_{23}^ν in eq.(31). Thus one has approximately,

$$\begin{aligned} \nu_e &\approx N_1(c_1\nu_1 - s_1c_2\nu_2 + s_1s_2c_3\nu_3) \\ \nu_\mu &\approx N_1(-s_1N_2\nu_1 + c_1c_2N_2^{-1}\nu_2 + c_1s_2c_3N_2^{-1}\nu_3) \\ \nu_\tau &\approx N_1^{-1}N_2^{-1}(-s_2c_3\nu_2s_1 + c_2\nu_3) \end{aligned} \quad (32)$$

Note that s_1 (s_2) determines $\nu_e - \nu_\mu$ ($\nu_\mu - \nu_\tau$) mixing. We must thus require s_2 to be large in order to account for the atmospheric muon neutrino deficit. The s_1 should be small for the small angle MSW solution and large for the vacuum oscillation solution to the solar neutrino problem. As we now demonstrate these constraints are too tight and one does not obtain parameter space in case of the MMM allowing simultaneous solution for both these problems.

A. MSW and atmospheric neutrino problem in MMM

The angle s_1 can be appropriately chosen to fix the required mixing for the small angle MSW conversion. The angle s_3 which determines the overall scale of neutrino masses is also required to be small. In such a case, the survival probability for the atmospheric ν_μ assumes two generation form and one can take the restrictions on relevant parameters from the standard analysis [24]. We have determined the effective $\nu_\mu - \nu_\tau$ mixing and neutrino masses following from eq.(26) by the procedure outlined in the last section. We show this mixing in Fig.(3a). In Fig.(3b), we show the masses for two values of $\Lambda = 70 \text{ TeV}, 150 \text{ TeV}$. As seen from Fig.(3a), the $s_2 = 0.3 - 0.75$ leads to the required $\sin^2 2\theta_{\mu\tau} = 0.8 - 1$. Fig.(3b) displays the contours corresponding to $\Delta_S \sim (3. - 12.) \cdot 10^{-6} \text{ eV}^2$ and $\Delta_A = (0.3 - 3.) \cdot 10^{-3} \text{ eV}^2$ in the $s_2 - s_3$ plane. It is seen that hierarchy among neutrino masses obtained in the required region is stronger than needed for a simultaneous solution of the solar and atmospheric neutrino problems and there is no overlapping region in the $s_2 - s_3$ plane for a combined solution. It is of course possible to solve each of this problem separately and get the required amount of mixing as well.

B. Vacuum oscillations and atmospheric neutrino problem in MMM

Unlike in the case of the MSW interpretation, the model can nicely account for the hierarchies required for solving the solar and atmospheric neutrino problems through vacuum oscillations. This is displayed in Fig.(4a) where we show contours corresponding to $\Delta_S = (0.5 - 3) \cdot 10^{-10} \text{ eV}^2$ and $\Delta_A = (0.3 - 3) \cdot 10^{-3} \text{ eV}^2$ in the $s_2 - s_3$ plane. Unlike in case of the MSW conversion, here there is a large overlap region in $s_2 - s_3$ plane which leads to the required values for $\Delta_{S,A}$. Despite this one unfortunately cannot explain both the problems simultaneously in a phenomenologically consistent way. This is due to the very restricted mixing structure displayed in eqs.(26). The vacuum oscillation probability in the present case is given by,

$$P_e = 1 - 4U_{e1}^2 U_{e2}^2 \sin^2 \left(\frac{\Delta_S t}{4E} \right) - 2U_{e3}^2 (1 - U_{e3}^2), \quad (33)$$

where the last term comes from the averaged oscillations corresponding to the atmospheric neutrino scale. Likewise, the muon neutrino survival probability which determines the atmospheric neutrino flux is given by,

$$P_\mu = 1 - 4U_{\mu 2}^2 U_{\mu 3}^2 \sin^2 \left(\frac{\Delta_A t}{4E} \right). \quad (34)$$

The amplitude of oscillations is controlled by two effective angles:

$$\sin^2 2\theta_S = 4U_{e1}^2 U_{e2}^2, \quad \sin^2 2\theta_A = 4U_{\mu 2}^2 U_{\mu 3}^2. \quad (35)$$

The matrix U appearing above is given by eq.(26). Restrictions on these angles required for a combined solution of the solar and atmospheric anomaly are worked out in [25] for different values of U_{e3} . Independent of the chosen values for U_{e3} one requires,

$$\sin^2 2\theta_S = 0.5 - 1, \quad \sin^2 2\theta_A = 0.8 - 1. \quad (36)$$

It is possible to choose these angles independently and satisfy above equations in a generic three generation case. In our case, the mixings are also determined in terms of $s_{1,2}$ through eq.(8). We have plotted the contours corresponding to restrictions in eq.(36) in Fig. 4b. It is seen that there is no region in $s_1 - s_2$ plane for which the solar and vacuum mixing angles can be simultaneously large ruling out the possibility of reconciling atmospheric anomaly with vacuum solution in the case of the MMM.

V. NON-MINIMAL MODEL AND NEUTRINO ANOMALIES

We had restricted our analysis so far to the MMM which is characterized by eq.(27) and the vanishing of the B and A parameters at Λ . Apart from predictivity, there are no strong theoretical arguments in favour of this minimal choice. One could consider variations of the MMM which in general result in introduction of additional low energy parameters. A class of non-minimal models could contain more complicated messenger sector which would influence boundary conditions in eq. (27). Alternatively, one may keep the same messenger sector but introduce some direct coupling between messenger and matter fields. This could result in non-zero B values at Λ . In fact, B gets generated [26,27] in models which try to understand origin of μ term in gauge mediated scenario [28]. B may be generated in the absence of messenger-matter coupling if MSSM itself is extended.

We shall not consider any specific model here, but would adopt a purely phenomenological attitude to point out possible ways which can allow simultaneous understanding of the solar and atmospheric neutrino anomalies. It turns out that prediction of m_0 is quite sensitive to the sign of μ term which is fixed to be positive in MMM. This follows from eq.(15) which shows that two contributions to $\langle \tilde{\nu} \rangle$ add or cancel depending on the sign of μ . We may thus consider a slightly less restrictive form of the MMM in which we regard the value and sign of B_μ as independent parameters to be determined phenomenologically. We still assume that mechanism responsible for generation of B parameters does not distinguish between leptonic

and the Higgs doublet H_1 and hence B_μ and B_i coincide at the scale Λ . Due to this, sneutrino vevs are still characterized by the differences in eq.(12) and hence are suppressed. The boundary conditions on soft masses are still assumed to be given by eq.(27). This particular scenario is now characterized by parameters Λ and B_μ . As follows from the minimum equation, (29), one may regard the value of $\tan\beta$ and sign of μ as independent parameters instead of B_μ . The magnitude of μ is determined in terms of these parameters by eq.(29). It is now possible to simultaneously account for the atmospheric neutrino deficit and have the MSW conversion for the solar neutrinos. This is depicted in Fig. (5) in which we show contours (in $s_2 - s_3$ plane) corresponding to $\Delta_S = 3 - 12 \cdot 10^{-6} \text{ eV}^2$, $\Delta_A = 0.3 - 3 \cdot 10^{-3} \text{ eV}^2$ for negative μ and two representative values of $\tan\beta = 40, 50$. The magnitude of μ gets fixed by eq. (29) to 379.28 GeV and 382.03 GeV in the respective cases. It is seen that now there is a considerable overlap where two mass scales arise simultaneously. As mentioned before, these masses are independent of the value of s_1 which can be chosen in the required range namely,

$$s_1 = 0.0225 - 0.071$$

to allow MSW conversion. The angle θ_A relevant for the atmospheric anomaly coincides roughly with s_2 and as follows from Fig.(5), one can simultaneously account for mixing as well as masses needed to solve the atmospheric and the solar neutrino problems.

VI. DISCUSSION

The supersymmetric standard model contains natural source of lepton number violation and hence of neutrino masses. The resulting neutrino mass pattern is quite constrained if source of lepton number violation is provided by soft bilinear operators and if the SUSY breaking is introduced through gauge mediated interactions. This scenario has the virtue that one can obtain hierarchical masses and large mixing in the neutrino sector. The hierarchy in masses results from hierarchy in the two different sources of neutrino masses while large mixing can be linked to ratio of R violating parameters ϵ_i . Overall scale of neutrino mass is set by s_3 and by the Yukawa couplings of the b and τ . Neutrino masses are thus naturally suppressed and hierarchical. One however needs to assume relatively suppressed R violation, i.e. $s_3 \sim 10^{-3}$ in order to obtain the mass scale relevant for the atmospheric neutrino anomaly. This requires that $\epsilon_2 \sim \epsilon_3 \sim 1 \text{ GeV}$ when $\mu \sim 1 \text{ TeV}$.

In case of the minimal gauge mediated model, the three neutrino masses and three mixing angles are controlled by four parameters Λ and $s_{1,2,3}$. This proves to be quite constraining

and does not allow one to obtain simultaneous solution of the solar and atmospheric neutrino anomalies. However, a non-minimal version which allows negative μ parameter is capable of accommodating the MSW effect and atmospheric neutrino anomaly. The number of parameters needed are still less than in the models based on the minimal supergravity scenario.

Although we concentrated on the bilinear terms a similar situation is obtained if neutrino masses are generated from the trilinear lepton number violating terms also. Here also, neutrinos obtain their masses from the loop as well as sneutrino vev induced contribution [29]. Unlike the bilinear terms, the number of possible trilinear terms is quite large and one needs to fine tune some of these or impose additional discrete symmetries in order to get a consistent picture [30].

We have mainly concentrated here on generating the atmospheric and the solar scales. The LSND result or the presence of hot neutrino dark matter needs an additional scale. It should be possible to introduce this through a sterile neutrino. Supersymmetry with gauge mediated interaction may be ideally suited to do so as it may explain both the lightness as well as the very existence of a sterile state [31].

Acknowledgments: We have greatly benefited from numerous discussions we had with K. S. Babu, G. Rajasekaran, Probir Roy and J. W. F. Valle.

- [1] Y.Fakuda et. al, hep-ex/9803006; hep-ex/9805006; hep-ex/9807003; T.Kajita, in Proc. of the XV111th Conf. on Neutrino Physics and Astrophysics, Takayama, Japan (1998).
- [2] Updated analysis including the recent Superkamioka results is presented in J. N. Bahcall, P. L. Krastev and A. Yu. Smirnov, hep-ph/9807216.
- [3] L. Wolfenstien *Phys. Rev. D* **17** (1978) 2369; S.P.Mikheyev and A.Yu.Smirnov *Yad. Fiz.* **42** (1985) 1441 [*Sov. J. Nucl. Phys.* **42** (1985) 913].
- [4] R.N.Mohapatra and P.B.Pal, Massive Neutrinos in Physics and Astrophysics, World Scientific lecture Notes in physics Vol.41, 1991.
- [5] D. O. Caldwell and R. N. Mohapatra, *Phys. Rev. D* **48** (1993) 3259; A. S. Joshipura, *Z.Phys.C* **64** (1994) 31; B. Brahmachari and R. N. Mohapatra, *Phys. Rev. D* **58** (1998) 15003.

- [6] A.Yu. Smirnov, *Phys. Rev. D* **48** (1993) 3264.
- [7] L.J.Hall and M.Suzuki, *Nucl. Phys. B* **231** (1984) 419.
- [8] A. Masiero and J.W.F. Valle, *Phys. Lett. B* **251** (1990) 273.
- [9] K.Tamvakis, *Phys. Lett. B* **383** (1996) 307.
- [10] K.S.Babu and A.S.Joshi-pura, Talks at Trieste Conf., Oct. 1996 and Oct. 1997.
- [11] See, for a review and original references, G.Giudice and R.Rattazi, hep-ph/9801271.
- [12] R. Hempfling, *Nucl. Phys. B* **478** (1997) 3; H.P.Nilles and N.Polonsky, *Nucl. Phys. B* **484** (1997) 33; A.Yu.Smirnov and F.Vissani, *Nucl. Phys. B* **460** (1996) 3427; E.Nardi *Phys. Rev. D* **55** (1997) 5772; B. de Carlos and P.L. White, *Phys. Rev. D* **54** (1996) 3427; T.Banks et. al, *Phys. Rev. D* **52** (1995) 5319; M.A.Diaz, J.C.Romao and J.W.F. Valle, *Nucl. Phys. B* **524** (1998) 23; E.J.Chun, S.K.Kang, C.W.Kim and U.W. Lee, hep-ph/9807327.
- [13] K.S.Babu and R.N.Mohapatra, *Phys. Rev. D* **42** (1990) 3778; R.Barbieri et. al, *Phys. Lett. B* **252** (1990) 251; E.Roulet and D.Tommasini, *Phys. Lett. B* **256** (1991) 218.
- [14] A. S. Joshipura and M. Nowakowski, *Phys. Rev. D* **51** (1995) 2421.
- [15] We use the notation $H_1' H_2 \equiv \epsilon_{ab} H_1'^a H_2^b$ with $\epsilon_{12} = -\epsilon_{21} = 1$.
- [16] Note that we have neglected here a sub-dominant contribution to \mathcal{M}_l arising due to sneutrino vevs.
- [17] K.S.Babu, C.Kolda and F.Wilczek, *Phys. Rev. Lett.* **77** (1996) 3070.
- [18] S.Dimopoulos, S.Thomas and J.D.Wells, *Nucl. Phys. B* **488** (1997) 39; J.A.Bagger, K.Matchev, D.M.Pierce and R.Zhang, *Phys. Rev. D* **55** (1997) 3188.
- [19] S.P.Martin, *Phys. Rev. D* **55** (1997) 3177; S.Dimopoulos, G.F.Giudice and A.Pomarol, *Phys. Lett. B* **389** (1996) 37.
- [20] R.Rattazzi and U.Sarid *Nucl. Phys. B* **501** (1997) 297.
- [21] F. Borzumati, hep-ph/9702307.

- [22] S.P.Martin and M.T.Vaughn, *Phys. Rev. D* **50** (1994) 2282.
- [23] m_0 also depends on s_2 very mildly through the vev $\langle \tilde{\nu}_2 \rangle$ which is greatly suppressed compared to $\langle \tilde{\nu}_3 \rangle$.
- [24] R. Foot, R.R. Volkas, O.Yasuda, *Phys. Rev. D* **58** (1998) 13006 (hep-ph/9801431); hep-ph/9802287, M.C. Gonzalez-Garcia, H.Nunokawa, O.L.G. Peres, T.Stanev and J.W.F. Valle, hep-ph/9801368.
- [25] P.Osland and G.Vigdel, hep-ph/9806339.
- [26] M.Dine, Y.Nir and Y.Shirman, *Phys. Rev. D* **55** (1997) 1501, T.Gherghetta, G.Jungman and E.Poppitz, hep-ph/9511317.
- [27] G.Dvali, G.F.Giudice and A.Pomarol, *Nucl. Phys. B* **478** (1996) 31.
- [28] A.de Gouvêa, A. Friedland and H. Murayama, *Phys. Rev. D* **57** (1998) 5676. This paper contains summary of the attempts to understand the μ problem.
- [29] A.S.Joshi, V.Ravindran and S.K.Vempati, hep-ph/9706482.
- [30] M.Drees, S.Pakvasa, X.Tata and T. ter Veldhuis, *Phys. Rev. D* **57** (1998) 5335.
- [31] E.J.Chun, A.S.Joshi and A.Yu. Smirnov, *Phys. Rev. D* **54** (1996) 4654; *Phys. Lett. B* **357** (1995) 608; K. Benakli and A.Yu. Smirnov, *Phys. Rev. Lett.* **79** (1997) 4314; A.S.Joshi and A. Yu.Smirnov, hep-ph/9806376.

Fig 1




Fig. 1. The variations of $\frac{m_0}{(\text{GeV } s_3^2)}$, $\frac{m_{1b}}{m_0}$, $\frac{m_{1\tau}}{m_0}$ are shown here with respect to Λ . m_0 mildly depends upon s_2 and the displayed curve is for $s_2 = 0.8$.

Fig 2




Fig. 2. The function s_ϕ^2 is plotted here with respect to s_2 .

Fig 3a




Fig. 3a. The effective ν_μ - ν_τ mixing angle is plotted here with respect to s_2^2 for $\Lambda = 100$ TeV in the case of minimal messenger model.




Fig. 3b. Contours of Δm^2 are plotted in MMM case, for $\Lambda = 70$ TeV (continuous lines) and $\Lambda = 150$ TeV (dash-dot). For Δ_A , the upper (lower) lines correspond to $3 \times 10^{-3} \text{ eV}^2$ ($0.3 \times 10^{-3} \text{ eV}^2$). For Δ_S , the upper(lower) lines correspond to $12 \times 10^{-6} \text{ eV}^2$ ($3 \times 10^{-6} \text{ eV}^2$).




Fig. 4a. Contours of Δm^2 in MMM are plotted for $\Lambda = 100$ TeV. For Δ_A , the upper (lower) line corresponds to $3 \times 10^{-3} \text{ eV}^2$ ($0.3 \times 10^{-3} \text{ eV}^2$). For Δ_S , the upper (lower) line corresponds to $3 \times 10^{-10} \text{ eV}^2$ ($0.5 \times 10^{-10} \text{ eV}^2$).




Fig. 4b. The effective $\text{Sin}^2 2\theta_S$ and $\text{Sin}^2 2\theta_A$ are plotted in the case of the minimal messenger model. The inner lines represent contours for 0.9 in both the cases whereas, the outer lines correspond to contours for 0.5 (0.8) for $\text{Sin}^2 2\theta_S$ ($\text{Sin}^2 2\theta_A$).

Fig5




Fig. 5. Contours of Δm^2 are plotted in **Non- MMM** case with $\mu < 0$, for $\tan \beta = 50$ (continuous lines) and $\tan \beta = 40$ (dash-dot) with $\Lambda = 100 \text{ TeV}$. For Δ_A , the upper (lower) lines correspond to $3 \times 10^{-3} \text{ eV}^2$ ($0.3 \times 10^{-3} \text{ eV}^2$). For Δ_S , the upper (lower) lines correspond to $12 \times 10^{-6} \text{ eV}^2$ ($3 \times 10^{-6} \text{ eV}^2$).