Current tensor with heavy photon for double hard photon emission by longitudinally polarized electron

M Konchatnij, N P M erenkov

K harkov Institute of P hysics and Technology

Abstract

The electron current tensor for the scattering of heavy photon on longitudinally polarized electron accompanied with two hard real photon is considered. The contribution of collinear and sem icollinear kinem atics is computed. The obtained result allows to calculate the corresponding contribution into second order radiative correction to D IS or electron {positron annihilation cross{sections with the next{to{leading accuracy.

1.The recent polarised experiments on deep inelastic scattering [1,2] cover the kinematical region y' 0:9, where the electrom agnetic corrections to the cross{section are extremely large. The rst orderQED correction have been computed in [3,4], and it is of the order the B orn cross{section in this region. That is why the calculation of the second orderQED correction becames very important for interpretation of these experiments. The rst step in such calculation was done in [5] where one{loop corrected C on pton tensor with a heavy photon had been considered. That is one of the contribution into polarized electron current tensor which appears in the second order of perturbation theory. A nother contributions arise due to double hard photon em ission and pair production.

Here we calculate the contribution into the polarized electron current tensor caused by two hard photon emission. We investigate double collinear kinematics and semicollinear one. This allow sus to compute the corresponding second order radiative correction to dierent observables with the next{to{leading accuracy in the same manner as it was done, for example, for small{ angle Bhabha scattering [6] and tagged photon cross{sections at D IS [7] and electron{positron annihilation [8].

In the Born approximation the electron current tensor with longitudinally polarized electron has a form

 $L^{B} = Q + iE$; $Q = 4(p_{1}p_{2})g + p_{1}p_{2} + p_{1}p_{2}$; (1) E = 4 $p_{1}p_{2}$;

where $p_1(p_2)$ is the 4{m om entum of the initial (nal) electron, and = 1(1) if the initial electron is polarized along (against) its 3{m om entum direction.

In the case of single collinear photon emission the corresponding contribution into electron current tensor conserves the born structure for radiation along the scattered electron m om entum direction

$$L^{(1)f} = \frac{1}{2} \frac{1 + (1 + y)^2}{y} \hat{E}_0 \frac{2(1 + y)}{y} dy L^B ; y = \frac{1}{m_2}; \hat{E}_0 = \ln \frac{m_2^2}{m_2^2};$$
(2)

and gets an additional part (which is proportional to $i \in D$) for radiation along the initial electron momentum direction [9]

$$L^{(1)i} = \frac{1}{2} - \frac{1 + (1 - x)^2}{x} L_0 - \frac{2(1 - x)}{x} L^B - 2xiE - dx; \quad x = \frac{!}{!}; \quad L_0 = \ln \frac{\frac{1}{10}}{m^2}: \quad (3)$$

In Eqs.(2) and (3) ! is the photon energy, $"_1("_2)$ is the energy of initial (nal) electron, m is the electron mass, and parameter $_0$ de nes the angular phase space of hard colinear photon. Index i(f) labels initial (nal) electron state.

Looking at Eq.(3) we see that the additional part does not contribute in m ain logarithm ic approximation and has not infrared divergence. Another words, the born structure of the electron current tensor in the case of longitudinally polarized electron is disturbed only in the next{to{bading approximation due to radiation just by initial polarized electron.

In general, the contribution into current tensor L due to emission of n collinear photons can be written as follows:

$$L^{(n)} = \frac{1}{2^{2}} \left[L^{(n)} L^{B} + K^{(n)} i E \right]_{i=1}^{Y^{n}} \frac{d^{3}k_{i}}{!_{i}}; \qquad (4)$$

where the quantity K ⁽ⁿ⁾ equals to zero if (and only if) all n collinear photons are emitted along the scattered (unpolarized) electron momentum direction. The set term in the right side of Eq.(4) had been obtained in [10] with the next{to{leading accuracy. Our goal is to nd also the second one with the same accuracy.

2 W e use the covariant m ethod of calculation and start from the general expression for polarized current tensor which arises due to two hard photon em ission

$$L^{(2)} = \frac{1}{4} - \frac{{}^{2} d^{3} k_{1} d^{3} k_{2}}{!_{1} !_{2}} Sp(p_{2} + m)Q \quad (p_{1} + m)(1 - {}_{5}P^{2})(Q)^{+};$$
(5)

where P is the polarization 4 {vector of initial electron. The quantity Q reads

$$Q = \frac{\hat{r} + m}{2m^2} \frac{\hat{p}_1 + \hat{k}_1 + m}{2p_1k_1} + \frac{\hat{p}_2 + \hat{k}_2 + m}{2p_2k_2} \frac{\hat{p}_1 + \hat{k}_1 + m}{2p_1k_1} + \frac{\hat{p}_2 = \hat{k}_2 + m}{2p_2k_2} \frac{\hat{r} + m}{2p_1k_1} + (1 + 2); = p_1 + k_1 + k_2; \quad (6)$$

For the important case of the longitudinally polarized electron in the frame of the choosen accuracy we can write the polarization vector in the form

$$P = \frac{1}{m} p_1 \frac{m^2 k}{p_1 k} ; \qquad (7)$$

where is the doubled electron helicity, and $4\{\text{vector } k \text{ has a components } ("_1; p_1); k^2 = m^2:$ It is easy to see that

 $P^{2} = 1 + 0 \text{ (m}^{4} = "^{4}) \text{ ; } P p_{1} = 0 \text{ :}$

Note that for calculation in the leading approximation we can neglect with the second term in the right side of Eq.(7) as it was done in [5].

There are four collinear regions in the case of double photon em ission: $(k_1; k_2 \ k \ p_1); (k_1; k_2 \ k \ p_2); (k_1 \ k \ p_1; k_2 \ k \ p_2)$ and $(k_1 \ k \ p_2; k_2 \ k \ p_1)$: The strightforward calculation in the region

 $(K_1; K_2 \text{ k } p_1)$ when both hard collinear photons are emitted by the initial (state polarized electron gives

$$\frac{2}{d_{1}^{2}} x_{2} + \frac{2y(1 x_{1})}{x_{2}} \qquad \frac{4y}{d_{1}^{2}} + \frac{(1 y)(2y + x_{1}x_{2})}{x_{1}x_{2}d_{12}} + \frac{4y}{d_{1}^{2}} + \frac{1}{1} + \frac{1}{2} + (1 \ 2); \quad (8)$$

W hen writing Eqs.(8),(9) we used the following notations

$$2p_1k_{1,2} = m^2_{1,2}$$
; $m^2 = m^2d$; $x_{1,2} = \frac{!_{1,2}}{!_1}$:

In the region $(\tilde{k}_1; \tilde{k}_2 \ k \ p_2)$ when both hard collinear photons are emitted by the nal{state unpolarized electron we have

$$K_{ff}^{(2)} = 0$$
; $I_{ff}^{(2)} = I_{ii}^{(2)} (x_{1,2} ! y_{1,2} ; 1,2 ! 1,2 ; d! ; y! = 1 + y_1 + y_2)$; (10)

where

$$y_{1,2} = \frac{!_{1,2}}{!'_2}$$
; $2p_2k_{1,2} = m^2_{1,2}$; $m^2 = m^2$:

In accordance with the quasireal electron method [9] we can express the electron current tensor in the region $(K_1 \ k \ p_1; K_2 \ k \ p_2)$ as a production of the radiation probability of the collinear photon with the energy $!_2$ by the scattered electron (which is the coe cient at L^B in the right side of Eq.(2) with $y = y_2$) and electron current tensor due to single photon emission by the initial electron as given by Eq.(3) with $x = x_1$. Therefore, the contribution of the regions $(K_1 \ k \ p_1; K_2 \ k \ p_2)$ and $(K_2 \ k \ p_1; K_1 \ k \ p_2)$ reads

$$L^{(2)if} = \frac{2}{2} \frac{{}^{2h}1 + (1 + y_2)^2}{y_2} \hat{E}_0 \frac{2(1 + y_2)}{y_2} \frac{inh_1 + (1 - x_1)^2}{x_1} L_0 \frac{2(1 - x_1)^i}{x_1} L^B$$
(11)
$$2x_1 i E \stackrel{o}{dy_2 dx_1} + (1 \$ 2) :$$

In order to derive the corresponding contributions in the regions $(k_1; k_2 \ k_{p_1})$ and $(k_1; k_2 \ k_{p_2})$ we have to perform the angular integration in Eq.(4) using Eqs. (8) and (9). Moreover, we can also integrate over the energy fraction x_1 (y_1) in the region $(k_1; k_2 \ k_{p_1})$ ($(k_1; k_2 \ k_{p_2})$) at xed value of the quantity $x_1 + x_2 = 1$ y ($y_1 + y_2 = 1$) because of 4{m omentum of the heavy photon which interacts with hadronic part of the amplitude depends on 1 y (1) in this case.

The expressions (8) and (9) for $I_{ii}^{(2)}$ and K $_{ii}^{(2)}$ are suitable for the calculations with a power accuracy (up to term s of the order m $^2=$ ¹/₁). But here we restrict ourselves with the logarithm ic accuracy and therefore can om itt term s proportional to 1=d $_{1}$ $_{2}$; 1=d² $_{1}$; 1=d² $_{1}^{2}$ and 1=d² $_{1}$ $_{2}$

in the right sides of Eqs.(8) and (9). In this approximation the integration of the quantity $I_{ii}^{(2)}$ leads to (see[10])

$$\frac{d^{3}k_{1}d^{3}k_{2}}{!_{1}!_{2}}\frac{I_{ii}^{(2)}}{m^{4}} = \frac{2}{2}\frac{1}{2}L_{0}^{2}A(y;) + L_{0}B(y;) dy; \qquad (12)$$

$$A = 4 \frac{1+y^2}{1-y} \ln \frac{1-y}{1-y} + (1+y) \ln y - 2(1-y);$$
(13)

$$B = 3(1 \quad y) + \frac{3 + y^2}{2(1 \quad y)} \ln^2 y \quad \frac{2(1 + y)^2}{1 \quad y} \ln \frac{1 \quad y}{1 \quad y} ; \qquad (14)$$

where <<1 is the infrared cut for the energy fraction of each photon. Analogously, the integration of the quantity K $_{\rm ii}^{(2)}$ reads

$$\frac{d^{3}k_{1}d^{3}k_{2}}{!_{1}!_{2}}\frac{K_{ii}^{(2)}}{m^{4}} = {}^{2}L_{0}C(y;)dy; C = 2(1 \ y) 2 \ \ln y \ 2\ln\frac{1 \ y}{m^{4}}dy:$$
(15)

By using the Eqs.(12),(15) together with Eq.(4) we obtain

$$L^{(2)ii} = \frac{2h}{2} \frac{1}{2} L_0^2 A(y;) + L_0 B(y;) L^B + C(y;) L_0 i E dy$$
(16)

for the contribution of the region $(\tilde{k}_1; \tilde{k}_2 \ k \ p_1)$ into the current tensor of longitudinally polarized electron. In some applications the quantyity y kept xed (for example, for calculation of the tagged photon cross{sections}). In this case we can write $\ln((1 \ y) =)$ instead of $\ln((1 \ y) =)$ in expressions for the quantities A and B.

The corresponding contribution of the region $(k_1; k_2 \mid p_2)$ can be written as follows

$$L^{(2)ff} = \frac{2^{h}}{2} \left(\frac{1}{2} \hat{E}_{0}^{2} \hat{R}(;)^{0} \right) + \hat{E}_{0} \hat{E}(;)^{0} L^{B} d; \quad ; \quad 0 = \frac{"}{"_{2}}; \quad (17)$$

where

$$R^{e} = 4 \frac{1+2}{1} \ln \frac{1}{2} \ln \frac{$$

$$B^{2} = 3(1) + \frac{3+2}{2(1)}\ln^{2} - 2\frac{(1+2)^{2}}{1}\ln\frac{1}{2} = 0 \qquad (19)$$

Note that the quantities A^{c} and B^{c} can be reconstructed from the quantities A and B by the rule

 $\hat{R}(; {}^{0}) = A(; {}^{0}); \hat{B}(;) = B(; {}^{0}):$

As we saw above (Eq.(3)) the additional part to the Born structure of polarized electron current tensor due to single collinear photon emission has neither colinear (does not contain large logarithm) nor infrared (is nite in the limit x go to zero) singularities. But these singularities appear in the corresponding contribution due to double collinear photon emission (Eqs.(11),(16)). Nevertheless, the additional part never conribute in the leading approximation.

The infrared parameter must eliminate in any physical application if photons are unobserved. Such elimination takes place because of contributions due to double virtual and soft photon emission as well as virtual and soft correction to single hard photon emission. The last contribution have been consider recenty [5] inside approximation $m^2 = 0$ which describes the large{angle photon radiation. If we put $m^2 = 0$ in our calculations than we will leave only with the bom{like structure in Eqs.(3),(11) and (16). Moreover, quantities B and B in Eqs.(16) and

(17) will be changed in this approximation. We see consequently, that it needs to keep nite the electron mass to be correct inside next{to{leading approximation in any physical application with unobserved photons (for example in classical deep inelastic scattering). We conclude therefore that the results of work [5] have to be improved for such kind applications.

3 Let us consider double hard photon emission in semicollinear regions $\tilde{k}_1 \ k \ p_1$ or p_2 and \tilde{k}_2 is arbitrary. In this situation we can use the quasireal electron method for longitudinally polarized initial electron [9]. In accordance with this method the contribution of the region $\tilde{k}_1 \ k \ p_2$ into electron current tensor is defined by its born {like structure L as follows

$$L \quad (k_1 \ k \ p_2) = \frac{2}{8^{-3}} \frac{d^3 k_2}{!_2} \frac{d y_1}{1 + y_1}^h \frac{1 + (1 + y_1)^2}{y_1} E_0 \quad \frac{2(1 + y_1)^i}{y_1} L \quad (p_1; p_2(1 + y_1); k_2); \quad (20)$$

where for large (angle em ission tensor L we can use the approximation $m^2 = 0$ [3,5,11]

$$L (p_{1}; p_{2}; k_{2}) = 4 (B + i E); \qquad (21)$$

$$B = \frac{1}{st} [(s + u)^{2} + (t + u)^{2}]g + \frac{4q^{2}}{st} (p_{1} p_{2} + p_{1} p_{2});$$

$$E = \frac{2}{st} [(u + t)p_{1} q + (u + s)p_{2} q]; g = g \frac{q q}{q^{2}};$$

$$= p \frac{(pq)q}{q^{2}}; u = 2p_{1}p_{2}; s = 2p_{2}k_{2}; t = 2p_{1}k_{2}; q = p_{2} + k_{2} p_{1}:$$

As above, the emission of collinear photon by the initial electron disturbs the Born structure of the electron current tensor just in the same manner as it done in Eq.(11)

$$L (\tilde{k}_{1} k p_{1}) = \frac{2}{8^{-3}} \frac{d^{3}k_{2}}{!_{2}} \frac{dx_{1}}{1 x_{1}} \left[\frac{1 + (1 x_{1})^{2}}{x_{1}} L_{0} - \frac{2(1 x_{1})}{x_{1}} \right] L (p_{1} (1 x_{1}); p_{2}; k_{2})$$

$$2x_{1} i E (p_{1} (1 x_{1}); p_{2}; k_{2})$$
(22)

Form ulae (20) and (22) are derived by us independently on quasireal electron m ethod starting from the general expression for current tensor as given by Eqs.(5), (6) and (7).

W hen calculating the radiative corrections to polarized D IS cross{section we have to integrate over all phase space of photons. At this case the angular cut parameter $_0$ is unphysical and must vanish in the sum of contributions of double collinear and sem icollinear regions. At taken accuracy this fact goes to cancellation term softhe type $L_0 \ln \frac{2}{0}$, and that can be verifyied by separation of $\ln \frac{2}{0}$ at integration of L ($p_1 (1 x_1); p_2; k_2$) in the lim it $k_2 k p_1$:

$$\frac{d^{3}k_{2}}{!_{2}}L \quad (p_{1}(1 \quad x_{1}); p_{2}; k_{2} \quad x_{2}p_{1}) = 2 \ln \frac{2}{0}dx_{2}\frac{y^{2} + (1 \quad x_{1})^{2}}{(1 \quad x_{1})x_{2}}L^{B} :$$
(23)

Taking into account that at $x = x_1 + x_2 = 1$ y

p

^Z dx₁dx₂
$$\frac{[1 + (1 x_1)^2][y^2 + (1 x_1)^2]}{x_1 x_2 (1 x_1)^2} = A (y;)dy;$$
 (24)

we carry conviction that the term s of the type $L_0 \ln \frac{2}{0}$ indeed vanish in the sum of contributions due to collinear kinem atics and sem icollinear one. A nalogous cancellation take place of course for radiation in the nal state.

Note in conclusion that the electron current tensor has an universal character. It can be used for calculation of cross{sections in di erent processes including most interesting D IS and e^+e {annihilation into hadrons. To obtain the corresponding cross{sections we have to multiply the electron current tensor by the hadron one. Just hadron tensor carry important inform ation about hadronic structure and fragmentation functions [12], and the study of radiative corrections to electron current tensor is necessary for interpretation experimental data in terms of these hadronic functions.

Authors thank A B A rbuzov and I.V. A kushevich for discussion. This work supported in part (N P M.) by INTAS grant 93{1867 ext and U kranian DFFD by grant N 24/379.

References

- 1. SM C., D. Adam set al, PhysRev. D 56 (1997) 5330.
- 2. HERMES, K. Achersta et al, PhysLett. B 404 (1997) 383.
- 3. T.V.Kukhto and N.M. Shumeiko, NuclPhysB 219 (1983) 412.
- 4. IV. Akushevich and N.M. Shumeiko, J.P. hys. G 20 (1994) 513.
- 5. IV. Akushevich, A B. Arbuzov and E A. Kuraev, PhysLett. B 432 (1998) 222.
- 6. A B. Arbuzov et al, NuclPhys. B 485 (1997) 457; PhysLettB 399 (1997) 312; N P. Merenkov, JETP 112 (1997) 400.
- 7. H. Anlauf, A. B. Arbuzov, E. A. Kuraev and N. P. Merenkov, hep {ph/9711333.
- 8. A B. Arbuzov, E A. Kuraev, N P. Merenkov and L. Trentadue, (to be published in JHEP).
- 9. V N. Baier, V S. Fadin V A. Khoze, NuclPhys. B 65 (1973) 381; V N Baier, V S. Fadin V A. Khoze and E A. Kuraev, PhysRep. 78 (1981) 293.
- 10. N P.M erenkov, Sov J Nucl Phys. 48 (1988) 1073.
- 11. EA.Kuraev, NP.Merenkov and VS.Fadin, YadFiz. 45 (1987) 782.
- 12. X. Ji, PhysRev. D 49 (1994) 114; R.L. Ja e and X. Ji, PhysRevLett. 67 (1991) 552.