Universal P ion Freeze-out Phase-Space D ensity

D. Ferenc, B. Tom asik, U. Heinz, Inst. f. Theor. Physik, Universitat Regensburg

G.Bertsch has indicated [1] a possibility to measure the pion freeze-out phase-space density and thereby test the bcalthern alequilibrium in a pion source. In case of therm al equilibrium at temperature T, identical pions of energy E would follow the Bose-E instein distribution

$$f = \frac{1}{e^{\frac{E}{T}} - 1}$$
 (1)

An average of this function over di erent phase space regions is the quantity to be measured. When the p_T - spectrum is parameterized by an exponential with $T_{\rm e}~(y)$ being the inverse slope parameter, averaging over the spatial coordinates yields

$$hfi(p_{T};y) = \frac{\frac{p}{2} - \frac{p}{\frac{dir(p_{T};y)}{E_{p}T_{eff}^{2}(y)}} \exp \frac{p_{T}}{T_{eff}(y)} \frac{dn}{dy}(y)}{R_{s}(p_{T};y) - \frac{p}{R_{o}^{2}(p_{T};y)R_{1}^{2}(p_{T};y)} - \frac{R_{o1}^{4}(p_{T};y)}{R_{o1}^{4}(p_{T};y)}};$$
(2)

This equation comprises inform ation from essentially two di erent classes of experim ental results: the single particle momentum spectra (dn =dy; T_{eff}), and the two pion Bose-E instein correlations (R_s ; R_o ; R_1 ; R_{ol} ; dir). We have calculated hfi(p_T ;y) for the S-S, S-Cu, S-Ag, S-Au, S-Pb and Pb-Pb data from the experiments NA 35 [2], NA 49 [3], NA 44 [4], and for the -p data from the NA 22 experiment 5] at CERN-SPS.From the results for hfi as a function of p_T , presented in Fig. 1, one may conclude:

1. universal phase-space density

All the nuclear collision data from the SPS in Fig.1 are indistinguishably similar, in spite of a factor of 10 difference in multiplicity density.

2. agreem ent with Bose-Einstein distribution

U sing simultaneously pion spectra and pion correlations NA 49 has disentangled therm alm otion from collective expansion [3]. Taking the NA 49 result for the local freezeout temperature T = 120 MeV as the only parameter in Eq. (1) one indeed nds good agreement with the data. This is consistent with the therm al nature of the pion source, and inconsistent with the presence of a hypothetic pion condensate.

3. radial ow

Looking in more detail, one nds that the data indicate a som ewhat slower decrease with increasing p_T than the Bose-E instein curve. This is most likely due to radial collective expansion which adds extra transverse momentum to particles, i.e. the local hf i values appear in the measurement at a p_T that is higher than the local h p_T i in the source reference frame. A detailed study is under way.

4. rapidity dependence

A certain departure from the universal scaling is seen for the data at rapidities close to the projectile rapidity, both at AGS and SPS; moreover, the two results are consistent.



Figure 1: Phase-space density as a function of p_T for di erent data sets. Heavy-ion data from SPS are indistinguishably sim ilar, although they span over an order of m agnitude range in multiplicity. A Bose-E instein function (Eq. (1)) is superim posed with the three choices of the local freeze-out tem perature: 80 MeV, 120 MeV and 180 MeV.

5. high tem perature decoupling in p collisions In contrast to freeze-out in nuclear collisions which takes place in two steps (chem ical at T ' 170-180 M eV and therm al at T ' 120 M eV), pion production in -p collisions [5] is essentially immediate, without the second evolution stage, and therefore freeze-out tem peratures of around 180 M eV should be expected. The data [5] are indeed consistent with this expectation, as seen in Fig.1.

References

- G.F.Bertsch, Phys. Rev. Lett. 72 (1994) 2349, and 77 (1996) 789e.
- [2] NA 35, T.A ber et al., Z.Phys. C 66 (1995) 77, Phys. Rev. Lett. 74 (1995) 1303.
- [3] NA 49, H. Appelshauser, Ph D. Thesis, Univ. Frankfurt, 1997. H. Appelshauser et al., Eur. Phys. J. C 2 (1998) 661.
- [4] NA 44, H. Beker et al, Phys. Rev. Lett. 74 (1995)
 3340. I.G. Bearden et al, Phys. Rev. C 58 (1998)
 1656.
- [5] N M . A gababyan et al., Z . P hys. C 71 (1996) 405.
- [6] E 877, J. Barrette et al, Phys. Rev. Lett. 78 (1997) 2961.