

FCNC transitions $c \rightarrow u$ and $s \rightarrow d$ in $B_c \rightarrow B_u$ and $B_s \rightarrow B_d$ decays

S. Fažfer^{a,b} and S. Prelovsek^a

a) J. Stefan Institute, Jamova 39, 1001 Ljubljana, Slovenia

b) Department of Physics, University of Ljubljana, 1000 Ljubljana, Slovenia

P. Singer

Department of Physics, Technion - Israel Institute of Technology, Haifa 32000, Israel

ABSTRACT

We propose the $B_c \rightarrow B_u$ decay as the most suitable probe for the flavour changing neutral transition $c \rightarrow u$. We estimate the short and long distance contributions to this decay within the standard model and we find them to be comparable; this is in contrast to radiative decays of D mesons, that are completely dominated by the long distance contributions. Since the $c \rightarrow u$ transition is very sensitive to the physics beyond the standard model, the standard model prediction $\text{Br}(B_c \rightarrow B_u) \sim 10^{-8}$ obtained here opens a new window for future experiments. The detection of $B_c \rightarrow B_u$ decay at branching ratio well above 10^{-8} would signal new physics. In addition we study the $s \rightarrow d$ transition in $B_s \rightarrow B_d$ decay and we find it to be dominated by the long distance contribution. We use the Isgur-Scora-Grinstein-Wise (ISGW) constituent quark model for the calculation of these decays.

1. INTRODUCTION

Flavour changing neutral current (FCNC) transitions occur in the standard model only at the loop level. Hence, they are very rare in the standard model and they present a suitable probe for new physics. The FCNC transitions in the down-quark sector are relatively frequent due to the large mass of the top quark running in the loop and the transition $b \rightarrow s$ has indeed been observed [1]. The FCNC transitions in the up-quark sector are especially rare in the standard model due to the small masses of the intermediate down-like quarks that run in the loop. For these transitions, the standard model represents a small background for the possible contributions arising from some new physics. At present, only upper experimental limits on the FCNC transitions in the up-quark sector are available [2].

In the present paper we study the transition $c \rightarrow u$, which is the most probable FCNC transition in the up-quark sector within the standard model. To probe the $c \rightarrow u$ transition we propose the radiative beauty-conserving decay $B_c \rightarrow B_u \gamma$; the B_c meson has been detected recently at Fermilab [3]. We estimate the short distance (SD) and long distance (LD) contributions to $B_c \rightarrow B_u$ decay within the standard model and find them to be comparable, which allows us in principle to probe $c \rightarrow u$ transition in this decay. This is in contrast to previously discussed D meson decays, where the dynamics is completely dominated by the LD contributions [4-7] and it is impossible to extract the short distance $c \rightarrow u$ contribution from the experiment.

In addition we study $s \rightarrow d$ transition in $B_s \rightarrow B_d$ decay. The dynamics of $B_s \rightarrow B_d$ is very similar to the dynamics of $B_c \rightarrow B_u$ due to the spectator b in both decays. In contrast to $B_c \rightarrow B_u$, the decay $B_s \rightarrow B_d$ is found to be dominated by LD contributions and consequently the signal of new physics is not expected in this decay.

In Sec. 2 the $B_c \rightarrow B_u$ decay is studied: we define SD and LD contributions, present the model and the results. The same reasoning is applied to $B_s \rightarrow B_d$ decay in Sec. 3. We conclude with a summary in Sec. 4.

2. $B_c \rightarrow B_u$ DECAY AND $c \rightarrow u$ TRANSITION

2a. The short distance contribution

The SD contribution in $B_c \rightarrow B_u$ decay is driven by FCNC $c \rightarrow u$ transition and b is a spectator. The $c \rightarrow u$ transition is strongly GIM suppressed at one-loop, QCD logarithms enhance the amplitude by two orders of magnitude [8], while the complete 2-loop QCD corrections further increase the amplitude by two orders of magnitude [8]. The Lagrangian that induces the $c \rightarrow u$ transition is given by

$$L_{SD}^{c \rightarrow u} = \frac{G_F}{2} \frac{e}{4} V_{cs} V_{us} C_7^{c \rightarrow u} (\gamma) \bar{u} \left[m_c \frac{1 + \gamma_5}{2} + m_u \frac{1 - \gamma_5}{2} \right] c F \quad : \quad (1)$$

The appropriate scale for $C_7^{c \rightarrow u} (\gamma)$ in $B_c \rightarrow B_u$ decay is $\mu = m_c$ (and not $\mu = m_b$), since b is merely a spectator in the SD process. The 2-loop QCD calculation was performed in [8], giving $C_7^{c \rightarrow u} (m_c) = 0.0068 - 0.020i$.

The corresponding amplitude for $B_c \rightarrow B_u (q; \gamma)$ decay is proportional to $\langle \gamma | \bar{u} \gamma_\mu (1 - \gamma_5) c | B_c \rangle$ taken at $q^2 = 0$, which can be expressed in terms of the form factors $F_1(0)$ and $F_2(0)$ [9]:

$$\begin{aligned} \langle \gamma | \bar{u} \gamma_\mu (1 - \gamma_5) c | B_c \rangle_{q^2=0} &= i \langle \gamma | \bar{u} \gamma_\mu (1 - \gamma_5) c | B_c \rangle_{q^2=0} F_1(0); \\ \langle \gamma | \bar{u} \gamma_\mu (1 - \gamma_5) c | B_u \rangle_{q^2=0} &= [(m_{B_c}^2 - m_{B_u}^2) \gamma_\mu - 2(\not{q} \cdot \not{p}) \gamma_\mu] F_2(0); \end{aligned} \quad (2)$$

The form factors defined in Eqs. (2) will be calculated using the ISGW model [10] later on in this section.

2b. The long distance contributions

The long distance contributions will be calculated by using the nonleptonic weak La-

grangian, which on the quark level can be written as [5]

$$L^{\text{eff}} (c = 1) = \frac{G_F}{2} V_{uq_i} V_{cq_j} [a_1 (uq_i)_{V-A} (q_j c)_{V-A}; + a_2 (uc)_{V-A} (q_j q_i)_{V-A};]; \quad (3)$$

where $(1\ 2)_{V-A} = 1 - \gamma_5$, q_i, q_j are the down quarks d, s, b and a_1, a_2 include the QCD corrections [11].

Quite generally, the LD contributions to $B_c \rightarrow B_u$ decay can be separated into two classes related to the two terms of (3), as performed previously [4] for $D \rightarrow V$ decays. The class (I), called also the VMD contribution, is related to the a_2 term (3) and corresponds to the processes $c \rightarrow uq_i q_i$ followed by $q_i q_i \rightarrow \gamma$, while b is the spectator in $B_c \rightarrow B_u$ decay. At the hadron level the $q_i q_i \rightarrow \gamma$ transition is expressed using the vector meson dominance (VMD) and the corresponding diagram is depicted in Fig. (1a). The class (II), called also the pole contribution, is related to the a_1 term (3) and corresponds to the process $cb \rightarrow ub$ with the photon attached to incoming or outgoing quark lines. Selecting the lowest contributing states, the pole contributions are depicted in Fig. (1b).

We turn now to the estimation of these two classes of contributions and we start with the VMD contribution (class (I)) represented by Fig. (1a). The underlying quark processes are $c \rightarrow us(d)$ with $ss; dd$ hadronizing into vector mesons ρ, ω which then turn to a photon, while b remains a spectator. We neglect the contribution of $bb \rightarrow \gamma$ in view of the large mass of b . The relevant part of the Lagrangian, after using the relations among CKM matrix elements, is

$$L_{(I)}^{\text{eff}} = \frac{G_F}{2} a_2 (V_{cs} V_{us}) u (1 - \gamma_5) c [s (1 - \gamma_5) s - d (1 - \gamma_5) d]; \quad (4)$$

The appropriate scale for $a_2(\mu)$ in $B_c \rightarrow B_u$ decay is $\mu = m_c$, since b is again merely a spectator in VMD contribution. Thus, we may use $a_2(m_c) = 0.5$, as obtained in the successful phenomenological fit to D meson decays [11]. Defining $h_V(q; \mu) \bar{V} \gamma_j V = g_V(q^2)$ and using the factorization approximation, the effective Lagrangian that induces the VMD contribution is given by

$$L_{\text{VMD}}^{c \rightarrow u}(\mu) = \frac{G_F}{2} e a_2(m_c) V_{cs} V_{us} C_{\text{VMD}}^0 u (1 - \gamma_5) c; \quad (5)$$

where

$$C_{\text{VMD}}^0 = \frac{g^2(0)}{2m^2} = \frac{g_\rho^2(0)}{6m_\rho^2} = \frac{g_\omega^2(0)}{3m_\omega^2} = (1.2 \pm 1.2) \cdot 10^3 \text{ GeV}^2 \quad (6)$$

is obtained by assuming $g_V(m_V) = g_V(0)$, with the mean value and the error in (6) calculated from the experimental data on $(V \rightarrow e^+ e^-)$ [2]. Note here the remarkable GIM cancellation carried over to the hadronic level.

Lagrangian (5) implies that the VMD amplitude for $B_c \rightarrow B_u(q; \mu)$ is proportional to $h_{B_u \gamma}(\mu) (1 - \gamma_5) c \beta_c$ taken at $q^2 = 0$. For the hadronic matrix elements, one defines appropriate form factors for the vector and axial transitions as follows [9]:

$$h_{B_u}(p^0; 0) \gamma_\mu (1 - \gamma_5) c \beta_c(p) = \frac{2i}{m_{B_c} + m_{B_u}} [p^0 p_V(q^2) + (m_{B_c} + m_{B_u}) A_1(q^2) - \frac{q_\mu}{m_{B_c} + m_{B_u}} (p + p^0) A_2(q^2) - 2m_{B_u} \frac{q_\mu}{q^2} [A_3(q^2) - A_0(q^2)]]; \quad (7)$$

The requirements of the finite matrix elements at $q^2 = 0$ [11] and of gauge invariance lead to the relations among the various form factors [6], which imply $A_0(0) = A_3(0) = 0$ and $A_2(0) = (m_{B_c} + m_{B_u})(m_{B_c} - m_{B_u})A_1(0)$. The same relations are obtained by using the prescription that the photon couples only to the transverse polarization of the current [6, 12]. Accordingly, the VMD amplitude will be expressed in terms of two form factors only, $V(0)$ and $A_1(0)$.

At this point, we remark that the form factors F_1, F_2, V and A_1 , needed for the SD and VMD amplitudes cannot be safely related using the Isgur-Wise relations [13], since the masses of b and c quarks composing B_c meson do not permit the b quark to be at rest. Therefore we shall determine the corresponding form factors at $q^2 = 0$ independently, using the ISGW model [10].

We now turn to the discussion of the LD contributions of class (II), the pole contribution, where the quark process $cb \rightarrow ub$ is driven by

$$L_{(II)}^{eff} = \frac{G_F}{2} a_1(\gamma) V_{cb} V_{ub} \bar{u} (1 - \gamma_5) b b (1 - \gamma_5) c \quad (8)$$

and the photon line is attached to any of four quark lines. In terms of hadronic degrees of freedom this diagram is given in Fig. (1b), where the white box represents the action of the Lagrangian (8) (we have neglected the contribution of the scalar and axial poles). Considering the scale for $a_1(\gamma)$ in $cb \rightarrow ub$, it is difficult to decide between $\mu = m_c$ or $\mu = m_b$, since b is not spectator in the pole contribution. As the difference between $a_1(m_c) = 1.2$ and $a_1(m_b) = 1.1$ [11] and is not essential, we take $a_1(m_b) = 1.1$. Note that the pole contribution is relatively small due to the factor $V_{cb} V_{ub}$ in (8). In D meson decays, the corresponding factor $V_{cs} V_{us}$ is much bigger, which makes the pole contribution dominant over the SD and VMD ones [4, 5, 6]. The different CKM matrix elements in the pole contributions of B_c and D decays is a major factor in establishing the $B_c \rightarrow B_u$ decay as more suitable for the investigation of $c \rightarrow u$ than the D decays.

To evaluate the amplitude for the pole diagrams given in Fig. (1b) we define

$$\begin{aligned} \langle 0 | \bar{A} | p \rangle &= f_p p \\ \langle 0 | V | p \rangle &= g_V \\ A(B_c(p) \rightarrow B_u(p^0; 0) | \gamma) &= \frac{e}{p} \langle 0 | p^0 \rangle; \end{aligned} \quad (9)$$

where $f_{B_c}, f_{B_u}, g_{B_c}$ and g_{B_u} will be determined using ISGW model.

2c. The amplitude

Using the above Lagrangians and form factor decomposition of Eqs. (2), (7), (9), the total amplitude for $B_c \rightarrow B_u$ containing SD and LD contributions can be expressed as

$$A(B_c(p) \rightarrow B_u(p^0; 0) | \gamma) = i^0 [A_{PV}(p, p, g, p, q) + iA_C(p^0, p, l); \quad (10)$$

where

$$A_{PV} = \frac{G_F}{2} e V_{cs} V_{ud} \frac{C_7^{c \rightarrow u}(m_c)}{2} (m_c - m_u) F_2(0) + 2a_2(m_c) C_{VMD}^0 \frac{A_1(0)}{m_{B_c} - m_{B_u}}; \quad ;$$

$$\begin{aligned}
A_{PC} = & \frac{G_F}{\sqrt{2}} e V_{cs} V_{ud} \frac{G_7^{c!u} (m_c)}{4} (m_c + m_u) F_1(0) + 2a_2 (m_c) C_{VMD}^0 \frac{V(0)}{m_{B_c} + m_{B_u}} \\
& + V_{cb} V_{ub} a_1 (m_b) \frac{g_{B_c} g_{B_u}}{m_{B_c}^2 m_{B_u}^2} + \frac{B_u m_{B_c}^2 f_{B_c} f_{B_u}}{m_{B_c}^2 m_{B_u}^2} : \quad (11)
\end{aligned}$$

The first term in Eqs. (11) comes from SD contribution, the second term from VMD contribution and the third term from the pole contribution. The decay width is then given by

$$= \frac{1}{4} \frac{m_{B_c}^2 m_{B_u}^2}{2m_{B_c}} (\mathcal{A}_{PV}^2 + \mathcal{A}_{PC}^2) : \quad (12)$$

2d. The model

To account for the nonperturbative dynamics within the mesons we use the nonrelativistic constituent ISGW quark model [10]. This model is considered to be reliable for a state composed of two heavy quarks, which makes it suitable for treating B_c ; in addition the velocity of B_u in the rest frame of B_c is to a fair approximation nonrelativistic. In the ISGW model the constituent quarks of mass M move under the influence of the effective potential $V(r) = -4_s = (3r) + c + br$, $c = 0.81 \text{ GeV}$, $b = 0.18 \text{ GeV}^2$ [14]. Instead of the accurate solutions of the Schrodinger equation, the variational solutions

$$\langle r \rangle = \frac{3}{4} \frac{3}{2} e^{-\frac{2r^2}{2}} \quad \text{or} \quad \langle k \rangle = \frac{3}{4} \frac{3}{2} e^{-\frac{k^2}{2}} \quad \text{for S state}$$

are used, where α is employed as the variational parameter. The meson state composed of constituent quarks q_1 and q_2 is given by

$$\mathcal{M}(p)_{i} = \int_{C_1; s_1; s_2}^X \frac{1}{\sqrt{3}} \frac{S}{(2)^3} d\mathbf{k}(\mathbf{k}) \frac{M_1}{E_1} \frac{M_2}{E_2} f_{s_2; s_1} (p_1 p_2) b_1^y(p_1; s_1; C) d_2^y(p_2; s_2; C) \mathcal{J}_i; \quad (13)$$

where \mathbf{k} is the momentum of the constituents in the meson rest frame, C denotes the colour, while $f_{s_2; s_1} = (\text{"# + \#"} = \frac{1}{\sqrt{2}}$ for pseudoscalar and $f_{s_2; s_1} = (\text{"# \#"} = \frac{1}{\sqrt{2}}$; " " ; # # for vector mesons. Using the normalization of the spinors as in [15], we obtain in the nonrelativistic limit

$$\begin{aligned}
V(q^2) &= \frac{m_{B_c} + m_{B_u}}{2} F_3(q^2) \frac{1}{M_u} \frac{M_b (M_c + M_u) \frac{2}{B_c}}{M_c M_u m_{B_u} (\frac{2}{B_c} + \frac{2}{B_u})} ; \\
A_1(q^2) &= F_2(q^2) = \frac{2m_{B_c}}{m_{B_c} + m_{B_u}} F_3(q^2) ; \\
F_1(q^2) &= 2F_3(q^2) \left[1 + (m_{B_c} - m_{B_u}) \frac{1}{2M_u} \frac{M_b (M_c + M_u) \frac{2}{B_c}}{2M_c M_u m_{B_u} (\frac{2}{B_c} + \frac{2}{B_u})} \right] ; \\
g_{B_c} &= \frac{S}{m_{B_c}} \frac{2}{\frac{2}{B_c} + \frac{2}{B_c}} \frac{3}{2} \frac{2}{3M_c} \frac{1}{3M_b} ; \\
f_{B_c} &= \frac{2}{\frac{3}{4}} \frac{p}{3} \frac{3}{2} \frac{1}{m_{B_c}} ;
\end{aligned}$$

$$g_{B_c} = m_{B_c} \frac{2^{\frac{3}{2}} \sqrt{3}}{4^{\frac{3}{2}} m_{B_c}} \quad (14)$$

and analogously for f_{B_u} , f_{B_u} and g_{B_u} . Here

$$F_3(q^2) = \frac{m_{B_u}}{m_{B_c}} \frac{2}{\frac{2}{B_c} + \frac{2}{B_u}} \exp^{3=2} \frac{M_b^2 [(m_{B_c} - m_{B_u})^2 - q^2]}{2m_{B_c}m_{B_u} \left(\frac{2}{B_c} + \frac{2}{B_u} \right)} ;$$

where $\frac{2}{B_c} = 0.7$ [10]. The results for $V(q^2)$ and $A_1(q^2)$ reproduce the results of [10], while $F_1(q^2)$ and $F_2(q^2)$ represent, to our knowledge, the new results within ISGW model. Using parameters [14] and meson masses given in Table 1 and the constituent quark masses $M_u = 0.33$ GeV; $M_c = 1.82$ GeV and $M_b = 5.2$ GeV [14] we get

$$f_{B_u} = 0.18 \text{ GeV} ; g_{B_u} = 0.86 \text{ GeV}^2 ; \frac{2}{B_u} = 1.81 \text{ GeV}^{-1}$$

$$f_{B_c} = 0.51 \text{ GeV} ; g_{B_c} = 2.41 \text{ GeV}^2 ; \frac{2}{B_c} = 0.28 \text{ GeV}^{-1} ;$$

while the form factors evaluated at $q^2 = 0$ are given in Table 2.

2e. The results for $B_c \rightarrow B_u$

We use the central value of the current quark masses $m_u = 0.0035$ GeV, $m_c = 1.25$ GeV from [2] and $V_{cb} = 0.04$, $V_{ub} = 0.0035$. The SD, VM D and pole contributions to amplitudes A_{PC} and A_{PV} needed to compute the amplitude (10) and the decay rate (12) are given in Table 3, where the error is due only to the uncertainty in parameter $C_{VM D}^0$ (6). In Table 4 we present the total branching ratio and separately also the SD and LD part of the branching ratios for $B_c \rightarrow B_u$ decay, where we have taken $\tau(B_c) = 0.46^{+0.18}_{-0.16}$ ps as measured by CDF Collaboration recently [3]. Note that SD and LD contributions give branching ratios of comparable size 10^{-8} , which in principle allows to probe the $c \rightarrow u$ transition in $B_c \rightarrow B_u$ decay. Experimental detection of $B_c \rightarrow B_u$ decay at the branching ratio well above 10^{-8} would clearly indicate a signal for new physics. The measurement of this decay would probe different scenarios of physics beyond the standard model: the non-minimal supersymmetric model [16] and the standard model with four generations [17], for example, predict $Br(c \rightarrow u)$ up to 10^{-5} , which would enhance $Br(B_c \rightarrow B_u)$ up to 10^{-6} . In D meson decays ($cq \rightarrow uq$), on the other hand, the branching ratios are of order 10^{-6} even within the standard model [5, 6, 8]: they are driven mainly by the long distance pole contributions, which overshadow the $c \rightarrow u$ transition (predicted at the branching ratio 10^{-9} in the standard model) and possible signals of new physics.

3. $B_s \rightarrow B_d$ DECAY AND $s \rightarrow d$ TRANSITION

The calculation of SD and LD contributions to $B_s \rightarrow B_d$ decay is analogous to what was presented in the previous section. Due to the small difference $m_{B_s} - m_{B_d}$, the final B_d meson is almost at rest in the rest frame of B_s .

The SD Lagrangian is

$$L_{SD}^{s \rightarrow d} = \frac{G_F}{2} \frac{e}{4} V_{cs} V_{us} C_7^{s \rightarrow d} (m_s - m_d) \left[\bar{m}_s \frac{1 + \gamma_5}{2} + m_d \frac{1 - \gamma_5}{2} \right] s \bar{F} ; \quad (15)$$

where $C_7^{s!d}(m_s) = 0.23$ [18].

The Lagrangian, from which the long distance VMD contribution is calculated, is given by

$$L_{VMD}^{s!d} = \frac{G_F}{2} a_2(m_s) V_{ud} V_{us}^* d (1-\gamma_5) s [u (1-\gamma_5) u - c (1-\gamma_5) c]; \quad (16)$$

where $|a_2(m_s)| \approx 0.5$ [18] and

$$C_{VMD} = \frac{g^2(0)}{2m^2} + \frac{g_1^2(0)}{6m_1^2} \times \frac{2g_i^2(0)}{3m_i^2}; \quad (17)$$

From $\exp(-\Gamma \tau) < 3.7 \cdot 10^9 \text{ eV}$ the upper limit $|C_{VMD}| < 0.01 \text{ GeV}^2$ has been obtained [18].

The pole contribution is absent in $B_s \rightarrow B_d$ decay since the decay $sb \rightarrow db$ involves four quarks of the same charge.

We use the constituent quark masses $M_d = 0.33 \text{ GeV}$, $M_s = 0.55 \text{ GeV}$ [14], the central value of the current quark masses $m_d = 0.006 \text{ GeV}$, $m_s = 0.115 \text{ GeV}$ [2] and the masses and parameters of the mesons given in Table 1. Using the formulas of the previous section and discarding the pole amplitudes, the resulting amplitudes $A_{PC,PV}$ and the branching ratios are given in Tables 3 and 4, respectively, where the upper limits are due to $|C_{VMD}| < 0.01 \text{ GeV}^2$. The upper limit $\text{Br}(B_s \rightarrow B_d) < 1.8 \cdot 10^7$ is dominated by the LD contribution. The same conclusion with a smaller upper bound was obtained in [18] assuming the simple free quark decay, which is a reasonable assumption for $s \rightarrow d$ in $B_s \rightarrow B_d$ decay. However, the calculation of the VMD amplitude in [18] was based on the formalism presented in [19], which is not reliable for $s \rightarrow d$ transition.

4. SUMMARY

We have studied flavour changing neutral transitions $c \rightarrow u$ and $s \rightarrow d$ in $B_c \rightarrow B_u$ and $B_s \rightarrow B_d$ decays, respectively. The predicted short and long distance contributions to these decays within the standard model are presented in Table 4.

We predict $\text{Br}(B_s \rightarrow B_d) < 1.8 \cdot 10^7$, which is dominated by the long distance contribution.

The short distance part (driven by $c \rightarrow u$) and the long distance part of the branching ratio for $B_c \rightarrow B_u$ decay are found to be of comparable size; they are both of order 10^8 . This makes $B_c \rightarrow B_u$ decay the most suitable decay to probe the $c \rightarrow u$ transition. Since $c \rightarrow u$ transition is very sensitive to the physics beyond the standard model, it would be very desirable to compare the standard model prediction of $\text{Br}(B_c \rightarrow B_u) = (8.5^{+5.8}_{-2.5}) \cdot 10^9$ presented here to the experimental data in the future. The detection of $B_c \rightarrow B_u$ decay at a branching ratio well above 10^8 would signal new physics. In comparison to $B_c \rightarrow B_u$ decay, the D meson decays are far less suitable for probing $c \rightarrow u$ transition, since they are almost completely dominated by the long distance effects.

Finally, we wish to stress that $B_c \rightarrow B_u$ and $B_s \rightarrow B_d$ decays are characterized by a very clear signature: their detection requires the observation of a $B_{u=d}$ decay in coincidence with two photons. The $B_c \rightarrow B_u$ transition involves the emission of a high energy

(985 MeV) and of a low energy (45 MeV) photon in the respective centers of mass of B_c , B_u , while in the decay $B_s \rightarrow B_d$ two photons of nearly equal energy (45 MeV) are emitted.

ACKNOWLEDGMENTS

The research of S.P. and S.F. was supported in part by the Ministry of Science of the Republic of Slovenia. The research of P.S. was supported in part by the Fund for Promotion of Research at the Technion.

Figure caption

Fig. 1: Long distance contributions in $B_c \rightarrow B_u$ decay. a) VMD contribution; the black box denotes the action of the Lagrangian (5). b) pole contribution; the white box denotes the action of the Lagrangian (8).

	B_c	B_c	B_u	B_u	B_s	B_d
m	6.40 [3]	6.42 [14]	5.28 [2]	5.325 [2]	5.37 [2]	5.325 [2]
	0.92	0.75	0.43	0.40	0.54	0.40

Table 1: Parameters (taken from [14]) and masses of pseudoscalar and vector mesons in GeV.

	$A_1(0)$	$V(0)$	$F_1(0)$	$F_2(0)$
$B_c \rightarrow B_u$	0.24	1.3	0.48	0.24
$B_s \rightarrow B_d$	0.90	11	1.8	0.90

Table 2: Form factors at $q^2 = 0$ calculated using ISGW model [10].

	A_{PV}^{SD}	A_{PV}^{VMD}	A_{PV}^{pole}	A_{PC}^{SD}	A_{PC}^{VMD}	A_{PC}^{pole}
$B_c \rightarrow B_u$	$5.7 + 17i$	$14 - 14i$	0	$5.7 + 17i$	$7.3 - 7.3i$	21
$B_s \rightarrow B_d$	62	$< 1 - 10i$	0	70	$< 5.6 - 10i$	0

Table 3: The amplitudes $A_{PV,PC}$ defined in (10) for SD, VMD and pole contributions in units of 10^{11} GeV^{-1} as predicted by ISGW model. The error-bars in $B_c \rightarrow B_u$ are due to the uncertainty in $C_{VMD}^0 = (1.2 - 1.2) \cdot 10^3 \text{ GeV}^2$ (6), while the upper bounds for $B_s \rightarrow B_d$ are due to $|C_{VMD}| < 0.01 \text{ GeV}^2$.

	$B r^{SD}$	$B r^{LD}$	$B r^{tot}$
$B_c \rightarrow B_u$	$4.7 \cdot 10^9$	$(7.5^{+7.7}_{-4.3}) \cdot 10^9$	$(8.5^{+5.8}_{-2.5}) \cdot 10^9$
$B_s \rightarrow B_d$	$1.4 \cdot 10^{11}$	$< 2.0 \cdot 10^7$	$< 2.0 \cdot 10^7$

Table 4: SD, LD and total branching ratios as predicted by ISGW model. The error bars in $B_c \rightarrow B_u$ are due to the uncertainty in $C_{VM D}^0 = (1.2 \pm 1.2) \cdot 10^3 \text{ GeV}^2$ (6), while the upper bounds for $B_s \rightarrow B_d$ are due to $|C_{VM D}| < 0.01 \text{ GeV}^2$.

References

- [1] R. Ammar et al. (CLEO Collab.), Phys. Rev. Lett. 71, 674 (1993); M. S. Alam et al. (CLEO Collab.), Phys. Rev. Lett. 74, 2885 (1995).
- [2] Particle Data Group, Eur. Phys. J. C 3, 1 (1998).
- [3] CDF Collaboration, Phys. Rev. Lett. 81, 2432-2437 (1998).
- [4] G. Burdman, E. Golowich, J.L. Hewett and S. Pakvasa, Phys. Rev. D 52, 6383 (1995).
- [5] S. Faﬀer and P. Singer, Phys. Rev. D 56, 4302 (1997).
- [6] S. Faﬀer, S. Prelovsek and P. Singer, Eur. Phys. J. C 6, 471 (1999).
- [7] S. Faﬀer, S. Prelovsek and P. Singer, Phys. Rev. D 58 094038 (1998).
- [8] G. Greub, T. Hurth, M. Misiak and D. Wyler, Phys. Lett. B 382, 415 (1996).
- [9] J.M. Soares, Phys. Rev. D 54, 6837 (1996).
- [10] N. Isgur, D. Scora, B. Grinstein and M. B. Wise, Phys. Rev. D 39, 799 (1989).
- [11] M. Bauer, B. Stech and M. Wirbel, Z. Phys. C 34, 103 (1987).
- [12] E. Golowich and S. Pakvasa, Phys. Rev. D 51, 1215 (1995).
- [13] N. Isgur and M. B. Wise, Phys. Rev. D 42, 2388 (1990).
- [14] D. Scora and N. Isgur, Phys. Rev. D 52, 2783 (1995).
- [15] C. Itzykson and J.B. Zuber, Quantum field theory, McGraw-Hill, New York (1985).
- [16] I. Bigi, F. Gabbiani and A. Masiero, Z. Phys. C 48, 633 (1990).
- [17] K.S. Babu, X.G. He, X.Q. Li and S. Pakvasa, Phys. Lett. B 205, 540 (1988).
- [18] G. Eilam, A. Ioannissian, R.R. Mendel and P. Singer, Phys. Rev. D 53, 3629 (1996).
- [19] N.G. Deshpande, X.G. He and J. Trampetic, Phys. Lett. B 367, 362 (1996).




Fig. 1