

New Insight into the Photoproduction of D -meson in QCD

A.V. Berezhnoy

Nuclear Physics Institute of Moscow State University,

Moscow, Russia

and

V.V. Kiselev, A.K. Likhoded

State Research Center of Russia "Institute of High Energy Physics",

Protvino, Moscow region, 142284, Russia

Abstract

In the framework of the model motivated by perturbative calculations in the fourth $O(\alpha_s^3)$ -order, the estimates for the p -cross-section of the D -meson production in the ZEUS experiment are performed. We factorize the hadronization of $(c\bar{q})$ -state hardly produced in perturbative QCD, which allows us to take into account the higher twist terms in the powers of $1/p_T$ at $p_T \sim m_c$ and to correctly reproduce the c -quark fragmentation dominant at high $p_T \sim m_c$ to the given order of α_s . We find a good agreement with the experimental data on the photoproduction, if the color-octet $(c\bar{q})$ -state is taken into account, which yields $\sigma_{(8)} \approx 0.33 - 0.49 \text{ GeV}^2$.

The measurement of the charm production by the ZEUS and H1 collaborations at HERA in the interaction of quazireal photon with the proton [1, 2] rises the problem on the data interpretation in the framework of QCD. The description of process for the heavy quark production in the perturbative QCD (pQCD) is completely justified due to the hardness of partonic subprocesses. At present, there are two types of calculations based on the results in the next-to-leading order over α_s , i.e. in $O(\alpha_s^2)$: for the production of massive and massless quarks, correspondingly [3, 4].

In the both approaches the fragmentation model for the c -quark hadronization into D was used at $p_T \sim m_c$. It was applied in the following form:

$$\frac{d^2 D}{dz dp_T} = \frac{d\hat{\sigma}_{cc}(k_T; \mu)}{dk_T} \frac{D_{c \rightarrow D}(z; \mu)}{z}; \quad (1)$$

$k_T = \frac{p_T}{z}$

where $D_{c \rightarrow D}(z; \mu)$ is a fragmentation function normalized by a probability of the c -quark fragmentation into the D -meson, $W(c \rightarrow D)$, which is empirically determined from the e^+e^- -annihilation data [5], where $W(c \rightarrow D) = 0.22 - 0.014 - 0.14$, and μ is the factorization scale for the perturbative partonic cross section $d\hat{\sigma} = dk_T$. It is clear that such the approach does not allow one to describe the experimental data in the region

of $p_T \gg m_c$, where the fragmentation is not in the saturation regime, yet. The fact is valid even for the case of calculations with the massive quarks, wherein the pQCD can be correctly applied in the region of $p_T \gg m_c$.

The most important assumption of this method is that the factorization takes place for the probability of isolated c-quark hadronization and its hard production. This circumstance automatically leads to the mentioned restriction: $p_T \gg m_c$.

In this paper, we discuss a model of the charm meson production with the hadronization of $(c\bar{q})$ -system into the D-meson instead of the hadronization of isolated c-quark. So, it is quite reasonable to apply this approach in the region of $p_T \gg m_c$, too, wherein the pQCD can be used. In the production of $(c\bar{q})$ -state, it is necessary to take into account the production of the additional quark pair, so that, in the leading order of α_s , the $O(\alpha_s^3)$ calculation must be used. Among the full set of the tree level diagrams on Fig. 1, we can see several diagrams (i.e. 16 and 19) of heavy quark fragmentation, which dominates at $p_T \gg m_c$. At high transverse momenta, they reproduce the results of fragmentation model with the following fragmentation function [6]:

$$D_{c \rightarrow D}(z) = \frac{8^2 h_0^{\text{eff}}}{27m_c^3} \frac{rz(1-z)^2}{(1-r)z^6} (2 - 2(3-2r)z + 3(3-2r+4r^2)z^2 - 2(1-r)(4-r+2r^2)z^3 + (1-r)^2(3-2r+2r^2)z^4); \quad (2)$$

where $r = m_q/(m_q + m_c)$ and

$$h_0^{\text{eff}} = h_{(1)} = \frac{1}{12M} \left(g + \frac{p \cdot p}{M^2} h_D(p) j(c \rightarrow q)(q \rightarrow c) \right) j(p);$$

which corresponds to the squared value of the quarkonium wave function at the origin for two quarks in the framework of nonrelativistic potential model: $h_{(1)} = |j_R(0)|^2$. We have introduced the effective mass of light quark m_q , which determines both the normalization and the form of fragmentation function. It is worth to stress that, in the higher orders of QCD, the hard part of processes does not depend on m_q . However, the problem of infrared divergences always presents in these cases. Usually, it can be removed by a redefinition of initial and final states. For the initial states one takes into account the QCD-evolution of intrinsic quark-gluon sea. One can introduce a regularization parameter (τ in the NLO-calculations in [3, 4]) to avoid the difficulties connected to the nonperturbative hadronic sea in the final state. In the model under consideration, $m_q = m = 2$ is used as an infrared regulator. In this case the gluonic virtualities are greater than $k_g^2 = q^2 + m^2$. So, the fact allows us to believe in the reliability of pQCD calculation. The D-meson production in e^+e^- -annihilation has been used to fix the model parameters. We find that the form and normalization of fragmentation function with the account for the QCD-evolution is in rather a good agreement with the LEP data, if the following values of parameters are chosen as below:

$$\begin{aligned} \alpha_s &= 0.35; \\ m_q &= 0.3 \text{ GeV}; \\ m_c &= 1.5 \text{ GeV}; \\ h_0^{\text{eff}} &= 0.25 \text{ GeV}^3; \\ W(c \rightarrow D) &= 0.22; \end{aligned} \quad (3)$$

In e^+e^- annihilation the enforcement of color-singlet (α)-state takes place in comparison with the octet contribution due to the color amplitude structure, so that

$$hO^{eff}i = hO_{(1)}i + \frac{1}{8}hO_{(8)}i; \quad (4)$$

where

$$hO_{(8)}i = \frac{1}{8M} \left[g + \frac{p \cdot p}{M^2} hD(p)j(c^a q)(q^b c)D(p)i \frac{ab}{8} \right];$$

It is qualitatively clear that the relative contribution of the octet four quark operator $hO_{(8)}i = hO_{(1)}i$ is determined by the relative velocity v of the quark motion in the hadron. For example, in the hard production of heavy quarkonium the octet fraction is strongly suppressed due to the nonrelativistic expansion in $v \ll 1$ [7]. For the meson, containing the light relativistic quark, we can expect that $hO_{(8)}i = hO_{(1)}i \approx 1$, since $v \approx 1$. Supposing this arrangement in e^+e^- annihilation, we see that $hO_{(1)}i$ is dominant in $hO^{eff}i$. The normalization of octet operator $hO_{(8)}i$ remains the free parameter. It can be determined from a process with another color structure of the production amplitude, for example, from the photoproduction. In this case, the essential contribution due to diagrams of nonfragmentation type (recombination) takes place at $p_T \approx m_c$. We have to emphasize that, in this kinematical region, the octet contribution is not suppressed in comparison with the singlet one.

The calculation results are shown in Fig. 2 in comparison with the experimental data of ZEUS [1] in the following kinematical region: $p_T > 2 \text{ GeV}$, $1.5 < \eta < 1.5$, $130 \text{ GeV} < W < 280 \text{ GeV}$, $Q^2 < 1 \text{ GeV}^2$, where W is the total energy of p interaction, Q^2 is the photonic virtuality, and η is the pseudorapidity of D -meson.

We see that the value of the D -meson production cross section cannot be explained by the singlet mechanism only. The relative value of octet contribution $hO_{(8)}i = hO_{(1)}i = 1.3$ allows us to describe the spectra of D -mesons. We would like to emphasize that the singlet term, say, at $p_T > 4 \text{ GeV}$ is very close to the predictions in the model for the production of isolated c -quark [3, 4].

Following a request by the ZEUS Collaboration we have calculated the differential cross sections for the D -meson production, as predicted by the model under consideration and shown in Fig. 3, at low W for the following kinematical region: $p_T > 2 \text{ GeV}$, $1 < \eta < 1.5$, $80 \text{ GeV} < W < 120 \text{ GeV}$, $Q^2 < 0.01 \text{ GeV}^2$. The same values of $hO_{(1)}i$ and $hO_{(8)}i$ have been used in the both calculations.

Using the experimental data by the ZEUS collaboration, only, we cannot exclude the possibility of other ratio between the singlet and octet operators, if the octet fraction is not suppressed in the fragmentation regime in e^+e^- annihilation. Then, for example, at $hO_{(1)}i \approx 0.14 \text{ GeV}^3$ and $hO_{(8)}i = hO_{(1)}i \approx 3.5$ we find a good description of the experimental data (see Fig. 4). However, the low value of singlet operator in combination with the high fraction of octet contribution results in quite a slow variation of the octet term.

Further, in the calculations we have neglected the differences between the spectra of color-octet vector and pseudoscalar states, whose contributions have been effectively summed up in the corresponding operator. Moreover, we have not considered a possible influence by the octet P -wave states, which seem to be very unstable for the heavy-light system.

In conclusion, we have presented the model for the charmed meson production in photon-proton collisions with the hadronization of $(c\bar{q})$ -state in contrast to the usual restriction by the isolated c -quark. It allows us to make justified predictions in the range of $p_T \sim m_c$, which improves the results of simple c -quark fragmentation regime dominant at high transverse momenta. In addition, the model includes the contributions of color-singlet and octet $(c\bar{q})$ -states, which depend on the color structure of production amplitude. So, in e^+e^- -annihilation the octet fraction is suppressed, while in the photoproduction we have found that the singlet contribution underestimates the data, and the octet operator can be measured so that, in the conservative evaluation, we extract

$$f_{D(s)} \approx 0.33 \pm 0.49 \text{ GeV}^3.$$

To improve the estimation, we plan to analyze the model predictions for the electroproduction, where the photon can be essentially virtual, in comparison with the current data.

This work is in part supported by the Russian Foundation for Basic Research, grants 96-02-18216 and 96-15-96575.

The authors would like to express the gratitude to Prof. A. W. Agner and members of DESY Theory Group for their kind hospitality during the visit to DESY, where this work was elaborated, as well as to Prof. S. S. Gershtein for discussions. The authors thank Prof. G. Kramer for conversation and remarks, which essentially improved the presentation of paper.

We thank Leonid Gladilin (ZEUS) for useful discussions, explanations on the data sample and help in our research.

References

- [1] J. Breitweg et al., ZEUS Collab., *Eur. Phys. J. C* **6**, 67 (1999).
- [2] C. Adloff et al., H1 Collab., preprint DESY 98-204, hep-ex/9812023 (1998).
- [3] S. Frixione, M. L. Mangano, P. Nason, G. Ridolfi, *Phys. Lett. B* **348**, 633 (1995);
S. Frixione, P. Nason, G. Ridolfi, *Nucl. Phys. B* **454**, 3 (1995).
- [4] B. A. Kniehl, G. Kramer, M. Spira, *Z. Phys. C* **76**, 689 (1997);
J. Binnewies, B. A. Kniehl, G. Kramer, *Z. Phys. C* **76**, 677 (1997);
J. Binnewies, B. A. Kniehl, G. Kramer, *Phys. Rev. D* **58**, 014014 (1998);
M. Cacciari, M. Greco, *Z. Phys. C* **69**, 459 (1996);
M. Cacciari, M. Greco, S. Rolli, A. Tanzini, *Phys. Rev. D* **55**, 2736 (1997);
M. Cacciari, M. Greco, *Phys. Rev. D* **55**, 7134 (1997);
- [5] K. Akerstedt et al., OPAL Collab., *Eur. Phys. J. C* **1**, 439 (1998).
- [6] C.-H. Chang, Y.-Q. Chen, *Phys. Rev. D* **46**, 3845 (1992), **D 50**, 6013 (E) (1994);
E. Braaten, K. Cheung, T. C. Yuan, *Phys. Rev. D* **48**, 4230 (1993);
V. V. Kiselev, A. K. Likhoded, M. V. Shevlyagin, *Z. Phys. C* **63**, 77 (1994);

T C Yuan, Phys. Rev. D 50, 5664 (1994);
K Cheung, T C Yuan, Phys. Rev. D 53, 3591 (1996).

- [7] G T Bodwin, E Braaten, G P Lepage, Phys. Rev. D 51, 1125 (1995);
T Mannel, G A Schuler, Z. Phys. C 67, 159 (1995);
E Braaten, S Fleming, T C Yuan, Ann. Rev. Nucl. Part. Sci. 46, 197 (1996).




Fig. 1. The leading order diagrams for the (αq) -state production in g interactions.




Fig.2. The distributions of D-meson photoproduction for $\langle \sigma_8 \rangle = \langle \sigma_1 \rangle = 1:3$ in comparison with the ZEUS data at $130 GeV < W < 280 GeV$ and $Q^2 < 1 GeV^2$. The color-singlet term is shown by the dashed histogram, the octet contribution is given by the dotted histogram, and the solid histogram presents the sum of two terms.




Fig.3. The distributions of D-meson photoproduction for $\langle x_8 \rangle = \langle x_1 \rangle = 1/3$ at $80 \text{ GeV} < W < 120 \text{ GeV}$ and $Q^2 < 0.01 \text{ GeV}^2$. The notations are the same as in Fig.2.




Fig.4. The distributions of D-meson photoproduction for $\langle x_8 \rangle = \langle x_1 \rangle = 3.5$ in comparison with the ZEUS data at $130 \text{ GeV} < W < 280 \text{ GeV}$ and $Q^2 < 1 \text{ GeV}^2$. The notations are the same as in Fig. 2.




Fig.5. The distributions of D-meson photoproduction for $\langle \theta_8 \rangle = \langle \theta_1 \rangle = 3.5$ at $80 \text{ GeV} < W < 120 \text{ GeV}$ and $Q^2 < 0.01 \text{ GeV}^2$. The notations are the same as in Fig.2.