#### RECENT ADVANCES IN ODDERON PHYSICS

Basarab NICOLESCU

LPNHE<sup>1</sup> - LPTPE, Université Pierre et Marie Curie, 4, Place Jussieu, 75252 Paris Cedex 05, France E-mail: nicolesc@ipno.in2p3.fr

Talk at International Conference on Elastic and Diffractive Scattering (VIIIth EDS Blois Workshop), Protvino, Russia, June 28 - July 2, 1999 (to be published in the Proceedings of this Conference)

Abstract

We present in this talk the phenomenological and theoretical advances in Odderon physics which occured since the last EDS Blois Workshop (Seoul, 1997).

LPNHE 99-06

November 1999

 $<sup>^1 \</sup>mathrm{Unit\acute{e}}$  de Recherche des Universités Paris 6 et Paris 7, Associée au CNRS

## 1 Introduction

The elastic and diffractive scattering is controlled by two singularities located near J = 1 in the complex J-plane : the Pomeron[1], in the even-under-crossing amplitude  $F_+$  and the Odderon[2], in the odd-under-crossing amplitude  $F_-$ .

The Odderon idea was longtime forced to stay in the Purgatory, because it contradicted the belief, founded on the dominant approach of the 70's - the Regge-pole model, that  $F_{-}$  is dominated by singularities located near J = 1/2( $\rho$  and  $\omega$  Regge poles and their cuts). In spite of its rediscovery in QCD in the 80's[3], and of its solid theoretical foundation in the framework of asymptotic theorems[2],[4] and derivative relations[5], the Odderon continued to be considered as an heretical concept. As late as in October 1990, here, in Dubna, André Martin did not hesitate to associate the words "revolution" and "Odderon" [6]. The complete theoretical legitimacy of the Odderon came only in the last few years from the calculation of the Odderon intercept in QCD. In the first part of my talk I will discuss the last results in this field.

From the phenomenological point of view, the interest clearly shifted from the somewhat biased study of Odderon effects in  $\bar{p}p$  and pp scattering towards HERA Odderon physics. The surprisingly rich activity in Odderon phenomenology in the last two years will be described, before drawing conclusions, in the second part of the talk.

## 2 Calculation of the Odderon intercept in QCD

The most important recent result is, beyond any doubt, the discovery by Janik and Wosiek[7] and, independently, by Lipatov[8], of an exact solution for the Odderon intercept in LLA.

In QCD the Odderon is a C-odd state of 3 reggeized gluons which interact pairwise with a well-defined potential (Fig. 1). The problem is to find an operator  $\hat{q}_3$ ,

$$\hat{q}_3^2 = -r_{12}^2 r_{23}^2 r_{31}^2 p_1^2 p_2^2 p_3^2, \tag{1}$$

which commutes with the Odderon hamiltonian H,

$$\hat{q}_3^2, H] = 0$$
 (2)

and has a much simpler form than H.

The exact solution of the problem is formulated in terms of the eigenequation

$$\hat{q}_3 f = q_3 f \tag{3}$$

where

$$f = \left(\frac{\rho_{12}\rho_{13}\rho_{23}}{\rho_{10}^2\rho_{20}^2\rho_{30}^2}\right)^{\mu}\Phi(z), \tag{4}$$

with  $\mu = h/3$ , h being the conformal weight. The function  $\Phi$  satisfies a third order linear equation

$$a(z)\frac{d^{3}}{dz^{3}}\Phi(z) + b(z)\frac{d^{2}}{dz^{2}}\Phi(z) + c(z)\frac{d}{dz}\Phi(z) + d(z)\Phi(z) = 0,$$
(5)

where

$$a(z) = z^3 (1-z)^3,$$
(6)

$$b(z) = 2z^{2}(1-z)^{2}(1-2z),$$
(7)

$$c(z) = z(z-1) \left( z(z-1)(3\mu+2)(\mu-1) + 3\mu^2 - \mu \right),$$
(8)

$$d(z) = \mu^2 (1-\mu)(z+1)(z-2)(2z-1) - iq_3 z(1-z),$$
(9)

$$z = \frac{\rho_{12}\rho_{30}}{\rho_{10}\rho_{32}}, \rho_k = x_k + iy_k(k = 1, 2, 3), \rho_{ij} = \rho_i - \rho_j.$$
(10)

In Ref. 7, the numerical solution

$$q_3 = -0.20526\,i\tag{11}$$

was found, corresponding to the Odderon energy

$$\epsilon = 0.16478. \tag{12}$$

The relation between the Odderon energy  $\epsilon$  and the Odderon intercept  $\alpha_O(0)$  is given by the equation

$$\alpha_O(0) = 1 - (9\alpha_s/2\pi)\epsilon. \tag{13}$$

For realistic values of  $\alpha_s$  ( $\alpha_s \simeq 0.19$ ) one gets

$$\alpha_O(0) = 0.94. \tag{14}$$

In collaboration with M.A. Braun and P. Gauron we recently performed a direct calculation of the lower bound for the Odderon intercept[9] in the framework of the variational approach we formulated earlier in collaboration with L. Lipatov[10]. In this variational approach the Odderon energy is defined as

$$\epsilon = E/D \tag{15}$$

In Eq.(15) D is a normalization constant and the energy functional is given by

$$E = \sum_{n=-\infty}^{\infty} \int_{-\infty}^{\infty} d\nu \epsilon_n(\nu) |\alpha_n(\nu)|^2, \qquad (16)$$

where

$$\epsilon_n(\nu) = 2Re[\psi(\frac{1+|n|}{2}+i\nu)-\psi(1)], \qquad (17)$$

$$\alpha_{n}(\nu) = \int_{0}^{\infty} dr r^{-2-2i\nu} \int_{0}^{2\pi} d\phi e^{-in\phi} \left( i\nu + \frac{n+1}{2} + re^{i\phi}(h-i\nu - \frac{n-1}{2}) \right) (i\nu - \frac{n-1}{2}) \left( -\tilde{h} + i\nu - \frac{n-1}{2} \right) Z(r,\phi),$$
(18)

$$h = 1/2 + n/2 - i\nu, \dot{h} = 1/2 - n/2 + i\nu, -\infty < \nu < \infty, n = \dots, -1, 0, 1, \dots,$$
(19)

$$Z(z, z^*) = |z(1-z)|^{2h/3} \Psi(z, z^*).$$
(20)

The function  $\Psi$  in Eq. (20) is invariant under the transformations  $z \to 1-z$ and  $z \to 1/z$  (Bose symmetry in the 3 gluons).

The following trial functions  $\Psi$  were used in Ref. 9 :

$$\Psi = \sum_{k=1}^{N-N_1} c_k a^{k/2-1/6} + \sum_{k=1}^{N_1} d_k a^{k-1/6} \ln a, \qquad (21)$$

where

$$a = \frac{r^2 r_1^2}{(1+r^2)(1+r_1^2)(r^2+r_1^2)}.$$
(22)

The result is

$$\epsilon = 0.22269,\tag{23}$$

a value which has to be compared with Eq. (12). The corresponding Odderon intercept is

$$\alpha_O(0) = 0.96. \tag{24}$$

By comparing the values (24) and (14), one sees that there is only a 2% difference between the "exact" result and the variational one.

We draw from this section the conclusion that the Odderon intercept is very close to 1, i.e. is much higher than the 1/2 ( $\rho,\omega$ ) intercept. We therefore expect important Odderon effects at high energy.

The LLA result shows that the gap  $\alpha_O(0) - 1$  is surprisingly small and therefore very sensitive to higher order corrections. The crucial problem of knowing if  $\alpha_O(0)$  is bigger than 1, smaller than 1 or just equal to one, is therefore still an open problem. The last case in the one I favor. The best way of solving this problem is, in my opinion, the study of the non-perturbative Odderon. Promising results on this line were already obtained[11].

Let me add, before closing this section, that very recently, G.P. Korchemsky and J. Wosiek obtained a new representation for the Odderon wave function[12], which is in agreement with the results of Refs. 7 and 8 and which allows identification of a new quantum number - *triality* - associated with the Odderon.

An intriguing problem is if there is or not a Pomeron-Odderon exchangedegeneracy, like in the non-leading reggeon  $\rho - \omega - f - A_2$  sector. The study of the *C*-even state of 3 reggeized gluons is crucial in this context[13].

# 3 The Odderon phenomenology

### 3.1 What is the problem with the Odderon ?

A quarter of century after its birth the Odderon has still an uncertain existential status. Of course, there are some interesting experimental indications of its existence :

- the difference between pp and  $\bar{p}p$  differential cross-sections in the dipshoulder region at ISR energies[14];
- the unusual shape of the polarisation in  $\pi^- p \to \pi^o n$  at low energies[15] (indicating a  $\rho$ -type Odderon, distinct from the  $\omega$ -type Odderon as required in LLA);
- the extraction of the semi-theoretical  $\rho$ -parameter from the dN/dt UA4/2  $\bar{p}p$  data in the presence of oscillations at very small t and high energies or of a more complicated phase of the forward scattering amplitude[16].

However, these experimental indications are either isolated or controversial. The real problem with the Odderon is the paradoxical scarcity of the highenergy data in hadron-hadron scattering, leading to an excessive focus on the pp and  $\bar{p}p$  scattering. In other words we fit high-energy parameters by using mainly low-energy data and, even worse, we draw conclusions about the Odderon based only on pp and  $\bar{p}p$  scattering. The folklore about the "suppression" of the Odderon has its source in these facts. The recent shift in attention from pp and  $\bar{p}p$  scattering is, in this context, very positive.

### 3.2 HERA-Odderon phenomenology

The most active team in the Odderon phenomenology in the last two years is, of course, the Heidelberg group (H.G. Dosch, O. Nachtmann, E.R. Berger) and its associates (A. Donnachie, P.V. Landshoff, W. Kilian, M. Rueter). In a series of papers[17],[18], they produced impressive results with a solid theoretical ground.

For example, the pseudoscalar meson production

$$e^{\pm}p \to e^{\pm}pPS$$
, where  $PS = \pi^o, \ \eta, \ \eta', \ \eta_c$  (25)

at HERA ( $\sqrt{s} = 300.6 \text{ GeV}$ ) constitutes a direct probe for the Odderon. The Odderon is here in competition with the photon only (see Fig. 2) : the Odderon contribution is not obscured by the huge Pomeron contribution as in the hadron-hadron reactions.

By taking, as a toy model, the Odderon as a Regge pole located near J = 1,

$$\eta_O \beta(t) \left(\frac{s}{s_0}\right)^{\alpha_O(t)-1} \xi(t) \tag{26}$$

where  $\alpha_O(0) \simeq 1$ ,  $\xi(t)$  is the signature fractor,  $\beta|t|$ -the Odderon residue and  $\eta_O = \pm 1$  (due to the absence of a positivity property for the Odderon contribution), one gets important Odderon effects in a variety of observables. I give just one example in Fig. 3 : the  $p_{\perp}$  distribution for pion production in the photoproduction region. One can see from Fig. 3 (where  $c_O$  is proportional to  $\eta_O$ ) that dramatic effects are induced by the presence of the Odderon. Moreover, one can see that the effects for the case  $\eta_O = +1$  are drastically different from the ones for  $\eta_O = -1$ : the HERA data can give important indications on the sign of  $\Delta \sigma$  - the difference between hadron-hadron and hadron-antihadron total cross-sections.

An important theoretical ingredient in several works of the Heidelberg group[18] is the Stochastic Vacuum Model (SVM)[19] which established a very interesting connection between the topological Y-shape of the baryons and the coupling of the Odderon. If the angle between two sheets of the baryon is very small (i.e. the baryon has a diquark-quark structure) the Regge-pole Odderon is suppressed. An interesting process is the photoproduction of pions with single dissociation (breakup of the target proton), because this process is independent of the particular structure of the baryon. Its cross-section is  $\simeq 300$  nb, i.e. 50 times larger for the corresponding one in the elastic photoproduction.

Of course, taking the Odderon as a Regge-pole could be too simplistic : for example, perturbative QCD indicates a more complicate singularity structure in the complex J-plane. Moreover, we showed longtime ago[20] that the Reggepole Odderon induces an overall shift of the low-energy data for the real parts which are already very well described by the Pomeron and the secondary Regge poles. Therefore, the Regge-pole Odderon is the worst case to be considered as a possible Odderon singularity. However, as a toy model, it can be still used as an illustration of typical Odderon effects, present even if the Odderon coupling is very much suppressed as compared with the Pomeron one.

A very interesting Odderon effect was recently studied by S.J. Brodsky et al.[9] : the assymptry in the fractional energy of charm versus anticharm jets in the diffractive photoproduction  $\gamma p \rightarrow c\bar{c}Y$  at HERA. This assymmetry is very sensitive to the Pomeron-Odderon interference (Fig. 4) : it measures the Odderon amplitude linearly. Namely

$$A(t, M_x^2, z_c) \simeq \epsilon \frac{\sin[(\pi/2)(\alpha_O - \alpha_P)]}{\cos((\pi/2)\alpha_O)} \left(\frac{s_{\gamma p}}{M_x^2}\right)^{\alpha_O - \alpha_P}$$
(27)

where  $z_c = E_{c(\bar{c})}/E_{\gamma}$  and  $\epsilon = \pm 1$ . It can be seen from eq.(27) that the sign of the asymmetry is controlled by the gap  $(\alpha_O - 1)$  and the energy dependence

by the gap  $(\alpha_O - \alpha_P)$ . By taking, as a numerical illustration,  $\alpha_P = 1.13$  and  $\alpha_O = 0.95$ , Brodsky et al. get a lower bound for the asymmetry equal to 15%.

Before closing this section, let me mention other interesting phenomenological studies of the Odderon : diffractive C = + neutral meson production from virtual photons[22], exclusive  $\eta_c$  photo- and electroproduction[23], exclusive  $f_2$ leptoproduction[24] and single-spin asymmetries for small-angle pion production in hadron collisions[25].

### 4 Conclusions : the Odderon in the future

As we understood from the talk of S. Weisz at this Conference [26], the TOTEM experiment at LHC will not be very helpful for the Odderon physics : at least in the initial stage of the experiment, priority will be given to just mesuring the pp total cross-section.

Great hopes arise from the talk of S.B. Nurushev[27] concerning the R7 experiment at RHIC[28] with a higher luminosity that the UA4/2 experiment at CERN. Several measurements concern directly the Odderon physics : extraction of the  $\rho$  parameter with a precision  $\delta \rho = 0.01$ ; research for oscilation at very small t ( $|t| \simeq (1 - 4) \cdot 10^{-4} \text{ GeV}^2$ ; the evolution of the dip in  $d\sigma/dt$ ; the polarisation parameters. In connection with this last measurement, E. Leader and T.L. Trueman showed in a very recent paper[29] the high sensitivity of the spin dependence of pp scattering, particularly the parameter  $A_{NN}$ , to the Odderon.

We had a nice surprise learning that the  $\bar{p}p$  option at RHIC is realistic. That would allow the detection of the Odderon through the measurement of the difference  $\Delta\sigma$  of the  $\bar{p}p$  and pp total cross-sections : the Regge-pole model predicts  $\Delta\sigma = 40 \ \mu b$  at RHIC while the maximal-Odderon approach[2] predicts[14]  $\Delta\sigma = -2.4$  mb. The expected precision at RHIC being  $\Delta\sigma = 0.5$  mb, one can clearly detect the presence of the Odderon through  $\Delta\sigma$ .

Let me also stress the importance of the systematic study of the energydependence of  $\sigma_T(s)$  in the RHIC range 50 GeV  $\leq \sqrt{s} \leq 500$  GeV. A huge gap in the high-energy hadron data will be filled.

However, we have not to wait till the next millenium in order to get a long awaited evidence of the Odderon. The results of the  $H_1$  experiment on the pseudoscalar meson production at HERA[30] will be soon available.

## Acknowledgments

It is a great pleasure to thank Prof. Vladimir Petrov and his young and dynamic staff for the remarkable organization of this Conference.

### References

- I.Ya. Pomeranchuk, Zh. Eksp. i Teor. Fiz. 34, 725 (1958); G.F. Chew and S.C. Frautschi, *Phys. Rev. Lett.* 7, 394 (1961).
- [2] L. Lukaszuk and B. Nicolescu, Nuovo Cimento Lett. 8, 405 (1973).
- [3] J. Bartels, Nucl. Phys. B 175, 365 (1980); J. Kwiecinski and M. Praszalowicz, Phys. Lett. B 94, 413 (1980).
- [4] D. Joynson, E. Leader, C. Lopez and B. Nicolescu, Nuovo Cimento 30A, 345 (1975).
- [5] K. Kang and B. Nicolescu, *Phys. Rev.* D **11**, 2461 (1975).
- [6] A. Martin, talk given at CERN-Dubna-Serpukhov Triangle Seminar, Dubna, 1990, CERN-TH. 5630/90 preprint.
- [7] R.A Janik and J. Wosiek, *Phys. Rev. Lett.* 82, 1092 (1999).
- [8] L.N. Lipatov, talk given at IPN Orsay, France, March 1997 (unpublished).
- [9] M.A. Braun, P. Gauron and B. Nicolescu, Nucl. Phys. B 542, 329 (1999).
- [10] P. Gauron, L.N. Lipatov and B. Nicolescu, *Phys. Lett.* B **260**, 407 (1991).
- P. Desgrolard, M. Giffon, E. Martynov and E. Predazzi, hep-ph/9907451;
   E. Predazzi, talk at this Conference.
- [12] G.P. Korchemsky and J. Wosiek, hep-ph/9908304.
- [13] L.N. Lipatov, private communication.
- [14] A. Breakstone et al., Phys. Rev. Lett. 54, 2180 (1985); S. Erhan et al., Phys. Lett. B 152, 131 (1985); A. Donnachie and P.V. Landshoff, Nucl. Phys. B 244, 322 (1984); P. Gauron, E. Leader and B. Nicolescu, Phys. Rev. Lett. 54, 2656 (1985).
- [15] V.P. Apokin *et al.*, Z. Phys. C **15**, 293 (1982) (Prof. S.B. Nurushev was part of this experiment); P. Gauron, E. Leader and B. Nicolescu, Phys. *Rev. Lett.* **52**, 1952 (1984).
- [16] UA4/2 Coll., C. Augier *et al.*, *Phys. Lett.* B **316**, 448 (1993); P. Gauron,
  B. Nicolescu and O.V. Selyugin, *Phys. Lett.* B **397**, 305 (1997); O.V. Selyugin, talk at this Conference; V. Kundrat, talk at this Conference.
- [17] W. Kilian and O. Nachtmann, *Eur. Phys. J.* C 5, 317 (1998); P.V. Landshoff and O. Nachtmann, hep-ph/9808233; E.R. Berger, A. Donnachie, H.G. Dosch, W. Kilian, O. Nachtmann and M. Rueter, *Eur. Phys. J.* C 9, 491 (1999); E.R. Berger, hep-ph/9905519.

- [18] M. Rueter and H.G. Dosch, *Phys. Lett.* B 380, 177 (1996); *Phys. Rev.* D 57, 4097 (1998); E.R. Berger and O. Nachtmann, *Eur. Phys. J.* C 7, 459 (1999); E.R. Berger, *Nucl. Phys. Proc. Suppl.* 74, 96 (1999); M. Rueter, H.G. Dosch and O. Nachtmann, *Phys. Rev.* D 59, 014018 (1999); E.R. Berger, talk at this Conference.
- [19] H.G. Dosch and Y.A. Simonov, *Phys. Lett.* B **205**, 339 (1988).
- [20] P. Gauron, E. Leader and B. Nicolescu, *Phys. Lett.* B 238, 406 (1990); P. Gauron and B. Nicolescu, *Phys. Lett.* B 292, 448 (1992).
- [21] S.J. Brodsky, C. Merino and J. Rathsman, hep-ph/99044280; SLAC-PUB-8272 and SLAC-PUB-8273, October 1999; S.J. Brodsky, hep-ph/9907346.
- [22] R. Engel, D. Yu. Ivanov, R. Kirschner and L. Szymanowski, *Eur. Phys. J.* C 4, 93 (1998).
- [23] J. Czyzewski, J. Kwiecinski, L. Motyka and M. Sadzikowski, *Phys. Lett. B* 398, 400 (1997); B 411, 402 (1997).
- [24] M.G. Ryskin, Eur. Phys. J. C 2, 339 (1998).
- [25] A. Ahmedov, I.V. Akushevich, E.A. Kuraev and P.G. Ratcliffe, hepph/9902418.
- [26] TOTEM Coll., W. Kienzle *et al.*, CERN/LHCC 99-7 ; S. Weisz, talk at this Conference.
- [27] S.B. Nurushev, talk at this Conference.
- [28] R7 Coll., N. Akchurin et al., Proposal, September 1995.
- [29] E. Leader and T.L. Trueman, hep-ph/9908221.
- [30] H1 Coll. S. Tapprogge et al., Proc. of the Int. Conf. on the Structure and the Interactions of the Photon (Photon 97) including the 11th Int. Workshop on Photon-Photon Collisions, ed. F.C. Erne and A. Buijs (World Scientific, Singapore, 1997); S. Tapprogge, talk at the Pomeron-Odderon Workshop, Heidelberg, March 1998; S. Tapprogge, PhD Thesis, Heidelberg University, HD-IHEP-96-19, December 1996.



Fig. 1. The Odderon in QCD as a C-odd state of 3 reggeized gluons.



Fig. 2. The Odderon exchange in  $e^{\pm}p \rightarrow e^{\pm}pPS$ .

Fig. 3. The  $p_{\perp}$  distribution for pion production in the photoproduction region (Ref. 17).



Fig. 4. The Pomeron-Odderon interference in  $\gamma p \rightarrow c \bar{c} p'$  (Ref. 21).