

Symmetric path integrals for stochastic equations with multiplicative noise

Peter A mold

Department of Physics, University of Virginia, Charlottesville, VA 22901
(November 1999)

Abstract

A Langevin equation with multiplicative noise is an equation schematically of the form $dq=dt = F(q) + e(q)$, where $e(q)$ is Gaussian white noise whose amplitude $e(q)$ depends on q itself. I show how to convert such equations into path integrals. The definition of the path integral depends crucially on the convention used for discretizing time, and I specifically derive the correct path integral when the convention used is the natural, time-symmetric one that time derivatives are $(q_t - q_{t-\epsilon})/\epsilon = \dot{q}$ and coordinates are $(q_t + q_{t-\epsilon})/2$. [This is the convention that permits standard manipulations of calculus on the action, like naive integration by parts.] It has sometimes been assumed in the literature that a Stratonovich Langevin equation can be quickly converted to a path integral by treating time as continuous but using the rule $\dot{q}(t=0) = \frac{1}{2}\dot{q}$. I show that this prescription fails when the amplitude $e(q)$ is q -dependent.

I. INTRODUCTION

Let η be Gaussian white noise, which I'll normalize as

$$\langle \eta_i(t) \eta_j(t') \rangle = \delta_{ij} \delta(t - t') \quad (1.1)$$

It's long been known that a Langevin equation of the form

$$\frac{d}{dt} q_i = F_i(q) + \eta_i \quad (1.2)$$

can be alternatively described in terms of a path integral of the form¹

$$P(q^0; q^0; t^0) = \int_{q(t^0)=q^0}^{q(t^0)=q^0} [dq(t)] \exp \left(- \int_{t^0}^{t^0} L(q; \dot{q}) dt \right) \quad (1.3)$$

¹ For a review of background material in notation close to that I use here, see, for example, chapter 4 of ref. [1]. The most substantial difference in notation is that my F is that reference's $\frac{1}{2}f$.

Here, $P(q^0; q^0; t)$ is the probability density that the system will end up at q^0 at time t if it started at q^0 at time zero. However, the exact form of L depends on the convention used in discretizing time when defining the path integral. With a symmetric discretization,²

$$P(q^0; q^0; t) = \lim_{t \rightarrow 0} N \int_{q(0)=q^0}^{q(t)=q^0} \prod_t dq_t \exp \left[- \int_0^t L \left(\frac{q_t + q_{t-\tau}}{2}; \frac{q_t - q_{t-\tau}}{\tau} \right) dt \right]; \quad (1.4)$$

with Lagrangian

$$L(q; \dot{q}) = \frac{1}{2} \dot{q}^2 + F(q) \quad (1.5)$$

Here and throughout, I adopt the notation that indices after a comma represent derivatives: $F_{i,j} = \partial F_i / \partial q_j$ and $F_{i,j,k} = \partial^2 F_i / \partial q_j \partial q_k$. N is the usual overall normalization of the path integral, which I will not bother being explicit about.

What hasn't been properly discussed, to my knowledge, is how to correctly form such a symmetrically-discretized path integral for the case of Langevin equations with multiplicative noise (meaning noise whose amplitude $e(q)$ depends on q). Schematically,

$$\frac{d}{dt} q_i = F_i(q) + e_{ia}(q) \xi_a; \quad (1.6)$$

with as before (1.1). I will assume that the matrix e_{ia} is invertible. There are a wide variety of applications of such equations, but I'll just mention one particular example of interest to me, which motivated this work and for which a path integral formulation is particularly convenient: the calculation of the rate of electroweak baryon number violation in the early universe [3].

By itself, the continuum equation (1.6) suffers a well-known ambiguity. To define the problem more clearly, we must discretize time and take $\tau \rightarrow 0$ at the end of the day. Specifically, I will interpret (1.6) in Stratonovich convention, writing

$$q_t - q_{t-\tau} = \tau F(q) + e(q) \xi; \quad (1.7)$$

$$h_{at} h_{bt}^0 = \tau \delta_{ab} \xi^0; \quad (1.8)$$

where

$$q = \frac{q_t + q_{t-\tau}}{2}; \quad (1.9)$$

[The fact that I've labelled the noise ξ instead of $\xi_{t-\tau}$ in (1.7) is just an inessential choice of convention.] The Stratonovich equation (1.7) is equivalent to the Itô equation

$$q_t - q_{t-\tau} = \tau F_i(q_{t-\tau}) + e(q_{t-\tau}) \xi; \quad (1.10)$$

² For a discussion of what changes if other discretizations are used in this case, see refs. [2]

with

$$F_i = F_i - \frac{1}{2} e_{ia;j} e_{ja} : \quad (1.11)$$

I will give two different methods for deriving the corresponding path integral. The result is

$$P(q^0; q^0; t) = \lim_{t \rightarrow 0} N \int_{q(0)=q^0}^{q(t)=q^0} dq_t \exp \left[- \int_0^t dt \left(\frac{1}{2} \dot{q}_t^2 + L(q_t, \dot{q}_t) \right) \right] ; \quad (1.12)$$

$$L(q; \dot{q}) = \frac{1}{2} (\dot{q} + F)_i g_{ij} (\dot{q} + F)_j - \frac{1}{2} F_{i;i} + \frac{1}{2} e_{ia}^1 e_{ka;k} (\dot{q} + F)_i + \frac{1}{8} e_{ia;j} e_{ja;i} ; \quad (1.13)$$

where

$$g_{ij} = (e^1)_{ia} (e^1)_{ja} : \quad (1.14)$$

This differs from a result previously given by Zinn-Justin³ [1] by the inclusion of the last term in L . Zinn-Justin's derivation was done in continuous time, resolving ambiguities using the prescription $\dot{q}(0) = \frac{1}{2}$, which is known to work in the case where $e(q)$ is constant.

Because of the confusion surrounding these issues, I will show how to derive the result in two different ways. First, I will follow the standard procedure for directly turning Langevin equations into path integrals, but I will be careful to keep time discrete throughout the derivation. The second method will be to start from the Fokker-Planck equation equivalent to the Langevin equations (1.7) and (1.10) and to then turn that Fokker-Planck equation into a path integral, again using standard methods.

II. DIRECT DERIVATION FROM THE LANGEVIN EQUATION

Rewrite the discretized Langevin equation (1.7) as

$$E_n = 0; \quad (2.1)$$

where n is a discrete time index and

$$E_n = e^1(q) [\dot{q}_n - q_{n-1} + \tau F(q)]; \quad (2.2)$$

with $\dot{q}_n = (q_n - q_{n-1})/\tau$. The corresponding path integral is obtained by implementing these equations, for each value of n , as δ -functions, with appropriate Jacobian, integrated over the Gaussian noise distribution:

$$P(q^0; q^0; t) = \lim_{t \rightarrow 0} N \int_{q(0)=q^0}^{q(t)=q^0} dq \exp \left[- \frac{1}{2\tau} \sum_n \dot{q}_n^2 \right] \det \left(\frac{\partial E_n}{\partial q_{n-1}} \right) \det \left(\frac{\partial E_n}{\partial q_n} \right) : \quad (2.3)$$

³ Specifically, eq. (4.79) or ref. [1]. See also ref. [4] and section 17.8 or ref. [1] for a continuum time discussion of formulating the path integral for this problem using ghosts.

The noise integral then gives

$$P(q^0; q^0; t) = \int_Y dq \exp \left[-\frac{1}{2t} E^2 \right] \det_{0a; 0i} \frac{\partial E_{0a}}{\partial q_{0i}} : \quad (2.4)$$

In our case (2.2), the determinant takes the form

$$\det_{0a; 0i} \frac{\partial E_{0a}}{\partial q_{0i}} = \det \begin{pmatrix} \frac{\partial E_1}{\partial q_1} & 0 & 0 & 0 \\ \frac{\partial E_2}{\partial q_1} & \frac{\partial E_2}{\partial q_2} & 0 & 0 \\ 0 & \frac{\partial E_3}{\partial q_2} & \frac{\partial E_3}{\partial q_3} & 0 \\ 0 & 0 & \frac{\partial E_4}{\partial q_3} & \frac{\partial E_4}{\partial q_4} \\ \vdots & \vdots & \vdots & \vdots \end{pmatrix} = \det_{ai} \frac{\partial E_a}{\partial q_i} : \quad (2.5)$$

The registration of the diagonals is determined by the nature of the initial boundary condition, which is that q_0 is fixed. From (2.2), we then have

$$\begin{aligned} \det_{0a; 0i} \frac{\partial E_{0a}}{\partial q_{0i}} &= \det_{ai} (e^1)_{ai} + \frac{1}{2} (e^1)_{ak; i} (q - q_1)_k + t \frac{1}{2} (e^1 F)_{a; i} \\ &= \det_{ij} e^1(q) \det_{ij} + \frac{1}{2} e_{ja} (e^1)_{ak; i} (q - q_1)_k + t \frac{1}{2} e_{ja} (e^1 F)_{a; i} ; \end{aligned} \quad (2.6)$$

where all e 's and F 's should now be understood as evaluated at q . Now rewrite the determinants in the last factor of (2.6) as exponentials in the usual way, using

$$\det(1 + A) = e^{\text{tr} \ln(1+A)} = \exp \left[\text{tr} \left(A + \frac{1}{2} A^2 + \dots \right) \right] : \quad (2.7)$$

To construct a path integral, we need to keep track of the terms in each time step up to and including $O(t)$, but we can ignore corrections that are higher-order in t . For this purpose, the size of $q - q_1$ should be treated as $O(\sqrt{t})$, which is the size for which the q^2 term in the action [the exponent in (1.12)] becomes $O(1)$ per degree of freedom. So, using the expansion (2.7), we get

$$\begin{aligned} \det_{ij} e^1(q) &= \exp \left(\text{constant} + \frac{1}{2} e_{ia} (e^1)_{ak; i} (q - q_1)_k + t \frac{1}{2} e_{ia} (e^1 F)_{a; i} \right. \\ &\quad \left. + \frac{1}{8} e_{ia} (e^1)_{ak; m} e_{m b} (e^1)_{bl; i} (q - q_1)_k (q - q_1)_l + O[(t)^{3/2}] \right) : \end{aligned} \quad (2.8)$$

It is the $q - q_1$ term in this equation, which came from the second-order term in the expansion (2.7), that will generate the difference with the result quoted in ref. [1]. Putting everything together, we get the path integral (1.12) with Lagrangian

$$\begin{aligned} L(q; q) &= \frac{1}{2} (q + F)_i g_{ij} (q + F)_j - \frac{1}{2} F_{ij} + \frac{1}{2} e_{ia}^1 e_{ka; j} (q + F)_i \\ &\quad + \frac{t}{8} e_{ia} (e^1)_{ak; m} e_{m b} (e^1)_{bl; i} q_k q_l : \end{aligned} \quad (2.9)$$

We can simplify this by realizing that the $q_k q_l$ in the last term can be replaced by its leading-order behavior in ϵ . Specifically, recall that $q(t) - q(t_0)$ is order ϵ . So one can go for a large number of discrete time steps $1 \leq N \leq 1/\epsilon$ without any net change in q at leading order in ϵ . Moreover, the force F doesn't have any net effect, at leading order in ϵ , over that number of steps. The upshot is that $q_k q_l$ can be replaced at leading order in ϵ by its average over a large number of steps, ignoring F and treating the background value of $e(q)$ as constant. The Gaussian integral for q in (1.12) and (2.9) then gives the replacement rule

$$q_k q_l \rightarrow \frac{1}{t} (q^2)_{kl} \quad (2.10)$$

at leading order in ϵ . This substitution turns (2.9) into the result (1.13) presented earlier.

III. DERIVATION FROM FOKKER-PLANCK EQUATION

The Stratonovich Langevin equation (1.7) is well-known to be equivalent to the Fokker-Planck equation

$$\dot{P} = \frac{\partial}{\partial q_i} \left(\frac{1}{2} e_{ia} \frac{\partial}{\partial q_j} (e_{ja} P) \right) + F_i P \quad (3.1)$$

where $P = P(q; t)$ is the probability distribution of the system as a function of time. This is just a Euclidean Schrodinger equation, and one can transform Schrodinger equations into path integrals by standard methods. Specifically, rewrite the equation as

$$\dot{P} = -\hat{H} P \quad (3.2)$$

with the Hamiltonian

$$\hat{H} = \frac{1}{2} \hat{p}_i e_{ia}(q) \hat{p}_j e_{ja}(q) - F(q) \quad (3.3)$$

To obtain a path integral with symmetric time discretization, it is convenient to rewrite \hat{H} in terms of Weyl-ordered operators. The Weyl order corresponding to a classical expression $O(p; q; t)$ is defined as the operator \hat{O}_W with

$$\langle p | \hat{O}_W | p^0 \rangle = \int \frac{d^N q}{(2\pi)^N} e^{ip \cdot (q - q^0)} O(p; \frac{1}{2}(q + q^0); t) \quad (3.4)$$

For the sake of completeness, I'll briefly review how to obtain Weyl ordering of operators in simple cases by considering the application of the operators to an arbitrary function $\phi(q)$. For example,

$$\begin{aligned}
[p_i A(q)]_W(q) &= \lim_{Z \rightarrow 0} [p_i A(q)]_W \Big|_{q^0=0} \\
&= \lim_{Z \rightarrow 0} [p_i A(q)]_W \Big|_{q^0=0} \\
&= \lim_{Z \rightarrow 0} e^{ip \cdot (q - q^0)} p_i A \left(\frac{q + q^0}{2} \right) \Big|_{q^0=0} \\
&= i \lim_{q^0 \rightarrow 0} \frac{\partial}{\partial q_i^0} (q - q^0) A \left(\frac{q + q^0}{2} \right) \Big|_{q^0=0} \\
&= i \frac{\partial}{\partial q_i^0} A \left(\frac{q + q^0}{2} \right) \Big|_{q^0=0} \\
&= i \frac{1}{2} \frac{\partial}{\partial q_i} A(q) + A(q) \frac{\partial}{\partial q_i} (q) :
\end{aligned} \tag{3.5}$$

So

$$[p_i A(q)]_W = \frac{1}{2} f_{p_i; A}(q) g : \tag{3.6}$$

One can similarly show that

$$[p_i p_j A(q)]_W = \frac{1}{4} f_{p_i; p_j; A}(q) g g : \tag{3.7}$$

Now write the Hamiltonian (3.3) in terms of Weyl-ordered operators. One finds

$$\hat{p} \cdot F = [p \cdot W F] - \frac{i}{2} F_{i,i} ; \tag{3.8}$$

$$p_i e_{ia} p_j e_{ja} = [p g^1 p]_W - i e_{ia} e_{ja,j} p_i + \frac{1}{4} (g^1)_{ij,i,j} - \frac{1}{2} (e_{ia} e_{ja,j})_{,i} : \tag{3.9}$$

So

$$\hat{H} = [H(p;q)]_W ; \tag{3.10}$$

with

$$H(p;q) = \frac{1}{2} p g^1(q) p - i p_i F_i(q) + \frac{1}{2} e_{ia}(q) e_{ja,j}(q) + u(q) ; \tag{3.11}$$

$$u = \frac{1}{2} F_{i,i} + \frac{1}{8} (g^1)_{ij,i,j} - \frac{1}{4} (e_{ia} e_{ja,j})_{,i} : \tag{3.12}$$

The usual derivation of the path integral then gives

$$P(q^0; q^0; t) = \lim_{t \rightarrow 0} \int_{q(0)=q^0}^{q(t)=q^0} \frac{dp \, dq}{(2\pi)^d} e^{S(p,q)} ; \tag{3.13}$$

$$S(p;q) = \int_0^1 \left[ip \cdot (\dot{q} - \dot{q}_1) + t H \left(p ; \frac{q + q_1}{2} \right) \right] dt ; \tag{3.14}$$

Doing the p integrals with our Hamiltonian (3.11) then reproduces the path integral (1.4) with Lagrangian

$$L(q;\dot{q}) = \frac{1}{2} \left(\dot{q}_i + F_i + \frac{1}{2} e_{ia} e_{ka;k} g_{ij} \dot{q}_j + F_j + \frac{1}{2} e_{jb} e_{lb;l} \right)^2 + u(q); \quad (3.15)$$

Now note that

$$(g^{-1})_{ij;ij} = 2e_{ia}e_{ja;ij} + e_{ia;i}e_{ja;j} + e_{ia;j}e_{ja;i}; \quad (3.16)$$

and so

$$\begin{aligned} -\frac{1}{8} (e_{ia}e_{ka;k})g_{ij} (e_{jb}e_{lb;l}) + u &= -\frac{1}{8} e_{ia;i}e_{ja;j} + \frac{1}{2} F_{i;i} + \frac{1}{8} (g^{-1})_{ij;ij} - \frac{1}{4} (e_{ia}e_{ja;j})_{;i} \\ &= \frac{1}{2} F_{i;i} + \frac{1}{8} e_{ia;j}e_{ja;i}; \end{aligned} \quad (3.17)$$

Combining (3.15) and (3.17) reproduces the Lagrangian (1.13) asserted in the introduction.

ACKNOWLEDGMENTS

I thank Larry Ya e, Dam Son, and Tim Newman for useful conversations. This work was supported by the U.S. Department of Energy under Grant Nos. DEFG 02-97ER 41027.

REFERENCES

- [1] J. Zinn-Justin, *Quantum Field Theory and Critical Phenomena*, 2nd edition (Oxford University Press, 1993).
- [2] F. Langouche, D. Roekaerts, and E. Tirapegui, *Physics* 95A, 252 (1979); H. Kawara, M. Namiki, H. Okamoto, and S. Tanaka, *Prog. Theor. Phys.* 84, 749 (1990); N. Komike, *Prog. Theor. Phys.* 86, 575 (1991).
- [3] P. Arnold and L. Ya e, "Non-perturbative dynamics of hot non-Abelian gauge fields: Beyond leading log," Univ. of Washington preprint UW /PT 99{25 (coming soon to hep-ph); "High temperature color conductivity at next-to-leading log order," Univ. of Washington preprint UW /PT 99{24 (coming soon to hep-ph); P. Arnold, "An effective theory for $1/k$ gT color dynamics in hot non-Abelian plasmas," Univ. of Virginia preprint UVA /Arnold{99{45 (coming soon to hep-ph); "Langevin equations with multiplicative noise: resolution of time discretization ambiguities for equilibrium systems," hep-ph/9912208.
- [4] J. Zinn-Justin, *Nucl. Phys. B* 275 [FS17] 135, (1986).