

# CP Violation in rare semileptonic B decays and Supersymmetry

E. Lunghi

SISSA -ISA S, Via Beirut 2-4, Trieste, Italy and  
INFN, Sezione di Trieste, Trieste, Italy

and

I. Scimemi

Dep. de Física Teórica, Univ. de Valencia,  
Edificio de Institutos de Investigación de Paterna  
Aptdo Correos 2085, E-46071 Valencia, Spain

## Abstract

We study the effect of new flavor changing SUSY phases arising in the squark mass matrix in semileptonic decays  $B \rightarrow X_s \ell^+ \ell^-$ , and  $B \rightarrow K^{(*)} \ell^+ \ell^-$ , ( $\ell = e, \mu$ ). SUSY effects are parametrized using the mass insertion approximation formalism. Constraints on SUSY contributions coming from other processes (e.g.  $b \rightarrow s$ ,  $B \rightarrow K \ell^+ \ell^-$ ) are taken into account. Chargino and gluino contributions to photon and Z-mediated decays are included and non-perturbative corrections are considered. We study the correlation between the CP (and forward/backward) asymmetries and the expected value of the inclusive branching ratio. Several possible scenarios are distinguished and discussed according to the mass insertion that is considered.

# 1 Introduction

Forthcoming B-factories are assumed to provide a great amount of data on many decays of the B mesons which have not been observed so far [1]. A study of these decays both in the SM and in its extensions is compelling.

The transitions  $B \rightarrow X_s^{*+} \gamma$  decays have been deeply studied. The dominant perturbative SM contribution has been evaluated in ref. [2] and later two loop QCD corrections have been provided [3]{ [5]. Recently the two loop SM matching conditions has been worked out in ref. [6]. cc resonances contributions to these results are included in the papers listed in ref. [7]. Long distance corrections can have a different origin according to the value of the dilepton invariant mass one considers.  $O(1-m_b^2)$  corrections have been first calculated in ref. [8] and recently corrected in refs. [9, 10]. Near the peaks, non-perturbative contributions generated by cc resonances by means of resonance-exchange models have been provided in refs. [9, 11, 12, 13]. Far from the resonance region, instead, ref. [14] (see also ref. [15]) estimate cc long-distance effects using a heavy quark expansion in inverse powers of the charm-quark mass ( $O(1-m_c^2)$  corrections).

An analysis of the SUSY contributions has been presented in refs. [16]{ [19] where the authors estimate the contribution of the Minimal Supersymmetric Standard Model (MSSM). In refs. [17, 18] the authors consider first a universal soft supersymmetry breaking sector at the Grand Unification scale (Constrained MSSM) and then partly relax this universality condition. In the latter case they find that there can be a substantial difference between the SM and the SUSY results in the Branching Ratios and in the forward/backward asymmetries. One of the reasons of this enhancement is that the Wilson coefficient  $C_7(M_W)$  (see section 2 for a precise definition) can change sign with respect to the SM in some region of the parameter space while respecting constraints coming from  $b \rightarrow s$ . The recent measurements of  $b \rightarrow s$  [20] have narrowed the window of the possible values of  $C_7(M_W)$ . Hence, it is worthwhile considering  $B \rightarrow X_s^{*+} \gamma$  in a more general SUSY framework than just the Constrained MSSM. In reference [21] the possibility of new-physics effects coming from gluino-mediated FCNC is studied. Possible relevant SUSY contributions coming from neutral Higgs bosons loops in the large  $\tan \beta$  regime are analyzed in ref. [22]. The process  $B \rightarrow X_s^{*+} \gamma$  in the flipped SU(5) SUSY GUT model is considered in ref. [23]. Effects of SUSY phases in models with heavy first and second generation sfermions have been recently discussed by the authors of ref. [24]. They study the effect of the  $\mu$  and  $A_t$  terms in the MSSM Lagrangian on the branching ratio of  $B \rightarrow X_s^{*+} \gamma$  and no flavor changing phase is assumed. Here we consider the impact of possible new phases in the flavor changing part of the squark mass matrix.

A "SUSY model independent" analysis of these decays has been provided in ref. [25] where SUSY effects are parametrized using the Mass Insertion Approximation (MIA) [26]. In this framework one chooses a basis for fermion and sfermion states in which all the couplings of these particles to neutral gauginos are flavor diagonal. Flavor changes in the squark sector are provided by the non-diagonality of the sfermion propagators. The pattern

of flavor change is then given by the ratios

$$(\tilde{f}_{ij})_{AB} = \frac{(m_{ij}^f)_{AB}^2}{M_{sq}^2}; \quad (1)$$

where  $(m_{ij}^f)_{AB}^2$  are the off-diagonal elements of the  $f = u, d$  mass squared matrix that mixes flavor  $i, j$  for both left- and right-handed scalars ( $A, B = \text{Left, Right}$ ) and  $M_{sq}$  is the average squark mass (see e.g. [27]). The sfermion propagators are expanded in terms of the  $\tilde{f}$ 's and the contribution of the first term of this expansion is considered. The genuine SUSY contributions to the Wilson coefficients will be simply proportional to the various  $\tilde{f}$ 's and a keen analysis of the different Feynman diagrams involved will allow us to isolate the few insertions really relevant for a given process. In this way we see that only a small number of the new parameters is involved and a general SUSY analysis is made possible. The hypothesis regarding the smallness of the  $\tilde{f}$ 's and so the reliability of the approximation can then be checked a posteriori. Many of these  $\tilde{f}$ 's are strongly constrained by FCNC effects [27, 28, 29] or by vacuum stability arguments [30]. Nevertheless it may happen that such limits are not strong enough to prevent large contributions to some rare processes. An estimation of branching ratios and asymmetries was usually provided considering the limits on the  $\tilde{f}$ 's existing at the time.

In ref. [31] it is noted that recent CLEO [32] (90% C.L.) upper limit on the branching ratio of

$$BR(B^0 \rightarrow K^0 + \dots) < 4.0 \cdot 10^{-6} \quad (2)$$

gives limits on the MI stronger than the ones of ref. [25] and new constraints on Wilson coefficients are derived.

So far in the SUSY-MIA analyses all  $\tilde{f}$ 's have been considered as real parameters as this is the relevant part in order to better contribute to branching ratios and FB-asymmetries. However eventual phases of these MI can modify other observables which measure the CP violation (CPV) in these decays. In this paper we want to investigate this possibility in the inclusive decay  $B \rightarrow X_s^{*\prime}$  and give an estimate also in the exclusive channels  $B \rightarrow K^{(*)\prime}$  for  $\prime = e$  or  $\mu$ .

For the exclusive channels a comment is in order. While the SM estimate for the inclusive process  $B \rightarrow X_s^{*\prime}$  is quite well established, much more uncertain is the issue of the exclusive modes. Theoretical errors are still large for  $B \rightarrow K^{(*)\prime}$  due especially to the computation of hadronic matrix elements and/or form factors. A number of approaches has been used to determine them and the interested reader can look in ref. [31, 33] and references therein. A generic analysis of possible new physics effect on these exclusive decays is presented in ref. [34]; however in this paper the authors do not make any hypothesis on the source of new physics. SUSY-MIA models can predict deviations from the SM much larger than the actual theoretical errors. From an experimental point of view the presence of new phases in the mass insertions is surely much easier to be observed in the exclusive channels than in the inclusive one. In order to give an estimate of CP violation in the exclusive channels we refer to the computation of the authors of ref. [31]. We find that the amount of CPV is practically the same on the inclusive and the exclusive transitions. The treatment of

the exclusive decay however allow us to take into account the constraint coming from Eq. (2) and thus to update the results of ref. [25].

The measure of CPV gives us important informations on possible SUSY extension of the SM. Either the experimental measure will or will NOT reproduce within the theoretical error (say 20% for the inclusive decay and 30-40% for the exclusive ones) the SM expectation for the BR's and FBA's, the detection of CPV will represent an important test for supersymmetry. Moreover, in ref. [25] it was pointed out that all SUSY effects in  $B \rightarrow X_s \gamma$  can be parametrized in terms of few MIT's, namely  $(\frac{u,d}{23})_{LL}$  and  $(\frac{u,d}{23})_{LR}$ . We have considered both the possibility that all the  $s$  contribute to the various observables and the case in which only the LL sector of the squark mass matrix is flavor changing. In all of these scenarios we have studied how the measures of BR's, FBA's and CPV are correlated. Given a measure of one observable it is so possible to deduce what are the expectations for the others.

A very particular behavior is observed if only LL mass insertions are allowed. If these insertions are real, no big signal has to be expected. However if they acquire an imaginary part they can sizably enhance the values of the BR's.

In all cases where an important increase of the BR's is provided the amount of CPV can be at best 2-3 times the SM [9] value although it can be of the opposite sign.

Instead, a depression of the SM result can be explained only with the help of the LR insertions and higher values of CPV can be expected in this case.

The paper is organized as follows. In order to estimate the CP violation a number of observables can be defined. We have considered CP asymmetries both in the branching ratio and in the FBA's and we have collected our basic definitions in section 2. In section 3 we summarize the existing constraints on mass insertions. Section 4 and 5 discuss the impact of SUSY on CP non-violating and CP violating observables respectively. Finally conclusions are provided.

## 2 Operator basis and observables

The effective Hamiltonian for the decay  $B \rightarrow X \gamma$  ( $X = X_s; K; \bar{K}$ ) in the SM and in the MSSM is given by

$$H_e = \frac{4G_F}{\sqrt{2}} K_{ts} K_{tb} \sum_{i=1}^8 C_i(\mu) Q_i + \frac{\alpha^0}{4} \sum_{i=9}^{\#} \tilde{C}_i(\mu) \tilde{Q}_i ; \quad (3)$$

where

$$\begin{aligned} Q_1 &= s_L \bar{c}_L c_L c_L ; & Q_2 &= s_L \bar{c}_L c_L c_L ; \\ Q_3 &= s_L \bar{b}_L c_L c_L ; & Q_4 &= s_L \bar{b}_L c_L c_L ; \\ Q_5 &= s_L \bar{b}_L c_R c_R ; & Q_6 &= s_L \bar{b}_L c_R c_R ; \end{aligned}$$

$$\begin{aligned}
Q_7 &= \frac{e}{16} m_b s_L \bar{b}_R F ; & Q_8 &= \frac{g_s}{16} m_b s_L T^a \bar{b}_R G^a ; \\
Q_9 &= (s_L \bar{b}_L) l \bar{l} ; & Q_{10} &= (s_L \bar{b}_L) l \gamma_5 l ;
\end{aligned} \tag{4}$$

$K$  is the CKM matrix and  $q_{L(R)} = \frac{(1 \pm \gamma_5)}{2} q$ . We neglect the small contribution proportional to  $K_{us} K_{ub}$ . This is the only SM source of CPV in these decays. Our approximation is thus equivalent to say that SUSY is the only origin of CP violation. This Hamiltonian is known at next-to-leading order both in the SM [4, 5] and in the MSSM [18, 19]. The most general low-energy SUSY Hamiltonian also contains the operators

$$\begin{aligned}
Q_7^0 &= \frac{e}{8} m_b s_R \bar{b}_L F ; \\
Q_9^0 &= (s_R \bar{b}_R) l \bar{l} ; \\
Q_{10}^0 &= (s_R \bar{b}_R) l \gamma_5 l ;
\end{aligned} \tag{5}$$

However the contribution of these operators is negligible and so they are not considered in the final discussion of physical quantities [25, 31]. SUSY contributions to other operators are negligible because they influence our observables at an higher perturbative order.

The definitions of the differential branching ratio and of the forward-backward asymmetries are  $s = (p_+ + p_-)^2 = m_b^2$  for the inclusive process and  $s = (p_+ + p_-)^2 = M_B^2$  in the exclusive ones,  $\theta$  is the angle between the positively charged lepton and the B flight direction in the rest frame of the dilepton system)

$$\text{(s)} \quad \frac{d(\Gamma(B \rightarrow X \ell^+ \ell^-))}{ds} ; \tag{6}$$

$$A_{FB}(\text{s}) = \frac{\int_{-1}^1 d \cos \theta \frac{d^2(\Gamma(B \rightarrow X \ell^+ \ell^-))}{d \cos \theta ds} \text{Sgn}(\cos \theta)}{\int_{-1}^1 d \cos \theta \frac{d^2(\Gamma(B \rightarrow X \ell^+ \ell^-))}{d \cos \theta ds}} ; \tag{7}$$

$$A_{FB}^0(\text{s}) = \frac{\int_{-1}^1 d \cos \theta \frac{d^2(\Gamma(B \rightarrow X \ell^+ \ell^-))}{d \cos \theta ds} \text{Sgn}(\cos \theta)}{\int_{-1}^1 d \cos \theta \frac{d^2(\Gamma(B \rightarrow X_s \ell^+ \ell^-))}{d \cos \theta ds}} \tag{8}$$

where  $X = X_s; K; \bar{K}$ . Note that for the inclusive case it is usually considered the normalized ratio  $R(\text{s}) = \frac{\Gamma(B \rightarrow X_c e)}{\Gamma(B \rightarrow X_c \mu)}$ ; the explicit formulae can be found in ref [1]. For what concerns the exclusive decays a discussion of the non-perturbative methods necessary to compute the form factors is beyond the scope of this paper. We have considered the approach of ref. [31] of Light Cone Sum Rules and the theoretical errors considered in this reference. In this case SUSY contributions can be disentangled only if they are far beyond the form factors theoretical uncertainty.

In order to compute CP violating effects it is necessary to include the absorptive parts of the matrix elements of the various operators. The only contributions come from  $Q_9$ . In order to use only prescription independent quantities it is usually considered the following

effective coefficient

$$\tilde{C}_9^e(s) = \tilde{C}_9^{\text{eff}}(s) + Y_{\text{res}}(s) : \quad (9)$$

$\tilde{C}_9^{\text{eff}}(s)$  includes both the mixing with of the operators  $Q_1 - Q_6$  and  $Q_8$  and the perturbative estimation of the hadronic matrix elements. Its complete definition for the SM and MSSM can be found again in refs. [4, 5, 18].  $Y_{\text{res}}(s)$  provides the Breit-Wigner ansatz for  $\rho$  resonances [7],

$$Y_{\text{res}} = \frac{3}{2} C^{(0)} \sum_{V_i = (1s); \dots; (6s)} k_i \frac{(V_i! \dots) m_{V_i}}{m_{V_i}^2 \text{Im}_{V_i} \text{Im}_{V_i}} \quad (10)$$

with  $C^{(0)} k_i = 0.875$  (for  $i = 1; \dots; 6$ ) [35]. In the literature there are at least other three different parameterizations of the resonant  $\rho$  contribution: the HQET {based approach [3], the KS {approach [12] and the LSW {approach [13]

Our final estimates of these observables include non-perturbative effects ( $O(1-m_b^2)$  [9], HHChPT [10],  $O(1-m_c^2)$  [14] corrections) and we refer to refs. [1, 25] for their complete expressions.

It is possible to construct several CPV observables using the branching ratios and the FBA's. We will consider the following.

The global branching ratio CP asymmetry,

$$A_G^{\text{BR}}(s) = \frac{\int ds \text{CP} [ (s) ]}{\int ds ( (s) + \text{CP} [ (s) ] )} : \quad (11)$$

The local branching ratio CP asymmetry

$$A_L^{\text{BR}}(s) = \frac{\text{CP} [ (s) ]}{(s) + \text{CP} [ (s) ]} : \quad (12)$$

The FBA differences

$$D_{A, A^0} = A_{\text{FB}}(s) (A_{\text{FB}}^0(s)) - \text{CP} [ A_{\text{FB}}(s) (A_{\text{FB}}^0(s)) ] : \quad (13)$$

In this case it is better not to normalize this difference with the sum of the two observables.  $A_{\text{FB}}(s)$  and  $A_{\text{FB}}^0(s)$  can have a different sign in different  $s$  regions and their sum can be nearly zero according to the value of the  $M$ 's.

All the effects coming from the mass insertion approximation can be included in formulae (6-8) writing the coefficients  $C_7, \tilde{C}_9^e(s), \tilde{C}_{10}$  as

$$\begin{aligned} C_7 &= C_7^{\text{SM}} + C_7^{\text{D ia g}} + C_7^{\text{M I}}; \\ \tilde{C}_9^e(s) &= (\tilde{C}_9^e(s))^{\text{SM}} + (\tilde{C}_9^e(s))^{\text{D ia g}} + (\tilde{C}_9^e(s))^{\text{M I}}; \\ \tilde{C}_{10} &= \tilde{C}_{10}^{\text{SM}} + \tilde{C}_{10}^{\text{D ia g}} + \tilde{C}_{10}^{\text{M I}} \end{aligned} \quad (14)$$

where all the contributions are evaluated at the  $M_B$  scale and the various  $C_i^{\text{D ia g}}$  summarize all the contributions coming from graphs including SUSY Higgs bosons and sparticles in

the limit in which we neglect all the mass insertion contributions (they would be the only SUSY diagrams if the scalar mass matrices were diagonalized by the same rotations as those needed by the fermions). The explicit expressions for  $C_i^{\text{diag}}$  can be found in ref. [18]. In the following we will use [18, 25],

$$\begin{aligned} & \lesssim C_7^{\text{diag}}(M_B) = 0.18 \\ & \gtrsim C_9^{\text{diag}}(M_B) = 0.35 \quad : \\ & \gtrsim C_{10}^{\text{diag}}(M_B) = 0.27 \end{aligned} \quad (15)$$

$C_i^{\text{M I}}$  have been estimated in ref. [25] and within the same range of parameters ( $\mu = 160$ ,  $M_{\text{gl}} = M_{\text{sq}} = 250$  GeV,  $M_{\text{t}} = 90$  GeV,  $M_{\text{b}} = 50$  GeV,  $M_{\text{H}} = 100$  GeV,  $\tan\beta = 2$ ) one gets

$$\begin{aligned} \ll C_7^{\text{M I}}(M_B) &= [ 0.19(\frac{d}{23})_{\text{LL}} \quad 33.4(\frac{d}{23})_{\text{LR}} \quad 1.75(\frac{u}{23})_{\text{LL}} \quad 0.25(\frac{u}{23})_{\text{LR}} ] \frac{s(M)}{s(M_B)}^{16=23} \\ \ll C_9^{\text{M I}}(M_B) &= 1.2(\frac{u}{23})_{\text{LL}} + 0.69(\frac{u}{23})_{\text{LR}} \quad 0.51(\frac{d}{23})_{\text{LL}} \\ \ll C_{10}^{\text{M I}}(M_B) &= 1.75(\frac{u}{23})_{\text{LL}} \quad 8.25(\frac{u}{23})_{\text{LR}} \quad : \end{aligned} \quad (16)$$

In this equation  $M$  is the scale at which all gauge bosons, top quark and SUSY particles are integrated out at the same time. The masses of these particles range between say 80 and 250 GeV so that we have chosen  $M$  at an intermediate scale of about the top quark mass. Corrections to this approximation are included in the errors of our estimates.

The sign and the value of the coefficient  $C_7$  has a great importance. In fact the integral of the BR is dominated by the  $\mathcal{J}_7 \mathcal{J}_9 = s$  and  $C_7 C_9$  term for low values of  $s$ . In the SM the interference between  $O_7$  and  $O_9$  is destructive and this behavior can be easily modified in the general class of models we are dealing with.

The big coefficient of  $(\frac{u}{23})_{\text{LR}}$  in  $C_{10}^{\text{M I}}(M_B)$  implies that the total coefficient  $C_{10}(M_B)$  can have a different sign with respect to the SM estimate. As a consequence of this, the sign of F-B asymmetries can be the opposite of the one calculated in the SM.

Finally the values of the physical constants we use are reported in table 1.

### 3 Constraints on mass insertions

In order to give an estimate of CPV it is necessary to take into account all possible constraints on mass insertions.

The most relevant is interested in the determination of the Wilson coefficients  $C_7$ ,  $C_9$  and  $C_{10}$  are  $(\frac{u}{23})_{\text{LL,LR}}^{\text{d,u}}$ .

Vacuum stability arguments regarding the absence in the potential of color and charge breaking minima and of directions unbounded from below [30] give

$$j(\frac{u}{23})_{\text{LR}} j \lesssim m_{\text{t}} \frac{2M_{\text{u}}^2 + 2M_{\text{t}}^2}{M_{\text{sq}}^2} \lesssim 2 \frac{m_{\text{t}}}{M_{\text{sq}}} \quad (17)$$

For  $M_{\text{sq}} = 300$  GeV this is not an effective constraint on the mass insertions.

A constraint on  $(\frac{d_{23}^{d_{11}}}{23})_{LL}$  can come from the possible measure of  $M_{B_s}$ .

In fact the gluino{box contribution to  $M_{B_s}$  [29] is proportional to  $(\frac{d}{23})_{LL}^2$  (see for instance ref. [29]). A possible experimental determination of  $M_{B_s}$ , say

$$M_{B_s} < 30 \text{ ps}^{-1} \quad (18)$$

would imply that

$$\text{Re}(\frac{d}{23})_{LL}^2 < 0.5 \quad (19)$$

for squark masses about 250 GeV. Moreover the LL up- and down-squark soft breaking mass matrices, at scales higher than the EW {breaking one, are related by a Cabibbo-Kobayashi-Maskawa rotation

$$(M_{sq}^d)_{LL}^2 = K^Y (M_{sq}^u)_{LL}^2 K \quad (20)$$

so that the limit (19) would be valid for the up sector too:

$$\text{Re}(\frac{u}{23})_{LL}^2 < 0.5 : \quad (21)$$

Strong constraints come from the measure of  $B \rightarrow X_s$ . The branching ratio of this process depends almost completely on the Wilson coefficients  $C_7$  and  $C_7^0$  which are proportional respectively to  $(\frac{d}{23})_{LR \text{ or } RL}$  and  $(\frac{u}{23})_{LL}$ . The most recent CLEO estimate of the branching ratio for  $B \rightarrow X_s$  is [20]

$$B_{\text{exp}}(B \rightarrow X_s) = (3.15 \pm 0.35 \pm 0.32 \pm 0.26) \cdot 10^{-4} : \quad (22)$$

where the first error is statistical, the second is systematic and the third comes from the model dependence of the signal. The limits given at 95% C.L. are [20]:

$$2.0 \cdot 10^{-4} < B_{\text{exp}}(B \rightarrow X_s) < 4.5 \cdot 10^{-4} : \quad (23)$$

We can define a  $C_7^e(M_B)$  as

$$C_7^e(M_B)^2 = \frac{B_{\text{exp}}(B \rightarrow X_s)}{\frac{K_{ts} K_{tb}^2}{K_{cb}} \frac{6 F}{g(z)}} \quad (24)$$

where  $F$  can be found for instance in ref. [36]. Considering the limits in Eq. (23) we find at 95% C.L.:

$$0.29 < |C_7^e(M_B)| < 0.41 : \quad (25)$$

Actually  $|C_7^e(M_B)|^2 = |C_7(M_B)|^2 + |C_7^0(M_B)|^2$  and the constraint given in Eq. (25) should be shared between the two coefficients. However in order to get the maximum SUSY contribution, we observe that in physical observables  $C_7^0$  does not interfere with  $C_7$ , the  $C_7^0 C_9$  term is suppressed by a factor  $m_s = m_b$  with respect to the  $C_7 C_9$  one and

$C_7^0 C_9^0$  is numerically negligible (in fact  $C_9^0$  is much smaller than  $C_9$ ). For these reasons we choose to lift the constraint of Eq. (25) with  $C_7(M_B)$  alone.

Thus, we can choose the total  $C_7^{\text{eff}}(M_B)$  anywhere inside the allowed region given in Eq. (25) still remaining consistent with the MIA.

The limit we get for  $(\frac{d}{23})_{LR}$  is of order  $10^{-2}$  and this rules out Z-mediated gluino penguins contributions to  $C_9$  and  $C_{10}$ .

For what concerns  $(\frac{u}{23})_{LL}$  we find that the constraint changes significantly according to the sign of  $C_7^{\text{eff}}(M_B)$ . In this case it is important to consider both the positive and negative region as this delta can give a non negligible contribution to  $C_9$  and  $C_{10}$ . The limits depend on the choice of the parameters in the chargino sector; the numerical results given below are computed for  $M_{sq} \simeq 250$  GeV,  $\mu \simeq 160$  GeV,  $m_{\tilde{t}_R} \simeq 90$  GeV,  $M_{\tilde{g}} \simeq 50$  GeV,  $\tan \beta \simeq 2$  (in ref. [25] these are the conditions under which we find the best SUSY contributions).

Constraints on mass insertions are provided by the updated CLEO [32] upper limit on the exclusive decay  $B^0 \rightarrow K^+ \pi^-$  written in Eq. (2), while no new limits are provided by

$$\begin{aligned} \text{BR}(B^+ \rightarrow K^+ \pi^+ \pi^-) &< 5.2 \cdot 10^{-6}; \\ \text{BR}(B^0 \rightarrow K^+ e^- e^+) &< 1.0 \cdot 10^{-5}; \end{aligned} \quad (26)$$

In ref. [31] constraints on  $C_9, C_{10}$  are derived in the hypothesis of real mass insertions while we now consider complex ones. In order to perform our numerical analysis taking into account this new limit we use the form factors of ref. [31]. Although a better understanding of the strong dynamics is compelling, we think that a clear signal of SUSY should overcome the theoretical uncertainties related to the calculation of hadronic matrix elements.

Choosing the input parameters given in [31] in such a way to obtain the loosest limits on the Wilson coefficients, the result we obtain for the integrated  $\text{BR}(B^0 \rightarrow K^+ \pi^-)$  is

$$\begin{aligned} \text{BR}(B^0 \rightarrow K^+ \pi^-) &= 10^{-6} \{ 1.52 + 0.0399 \mathcal{C}_{10}^2 + 0.0399 \mathcal{C}_9^2 + 1.676 \mathcal{C}_7^2 \\ &\quad + 0.365 \text{Re}[\mathcal{C}_{10}] + 0.294 \text{Re}[\mathcal{C}_9] + 0.197 \text{Re}[\mathcal{C}_7] + \\ &\quad + 0.0488 \text{Im}[\mathcal{C}_7] + 0.0168 \text{Im}[\mathcal{C}_9] + 0.273 \text{Re}[\mathcal{C}_9 \mathcal{C}_7] \} \end{aligned} \quad (27)$$

where the coefficients  $C_7, C_9$  and  $C_{10}$  are defined as follow

$$\begin{aligned} C_7 &= C_7^{\text{D ia g}}(M_b) + C_7^{\text{M I}}(M_b); \\ C_9 &= (C_9^{\text{eff}})^{\text{D ia g}}(M_b) + (C_9^{\text{eff}})^{\text{M I}}(M_b); \\ C_{10} &= C_{10}^{\text{D ia g}}(M_b) + C_{10}^{\text{M I}}(M_b); \end{aligned} \quad (28)$$

## 4 CP non-violating observables

In this section we present the numerical results obtained for the maximum enhancement and depression with respect to the SM expectation of the BR and of the FB asymmetries for the inclusive decays  $B \rightarrow X_s \ell^+ \ell^-$  and for the exclusive channel  $B \rightarrow (K; K^*) \ell^+ \ell^-$  both for  $\ell = e$  and  $\ell = \mu$ . For the inclusive case we update the estimates given in ref. [25] due to the new limits on the  $s$  coming from Eq. (2).

The relevant formulae for the above quantities can be found in ref. [25] where non-perturbative  $O(1=m_b^2)$  and  $O(1=m_c^2)$  corrections are provided as well; we refer to [25] for a complete list of references.

A useful way to study the dependence of these observables on the mass insertions is to observe that the integrated BR and non-normalized FB asymmetry are polynomial in the Wilson coefficients. It is thus possible to write the following master formulae:

$$\begin{aligned} (BR_{e;})_{X_s} &= 10^{-6} [(9.79; 6.53) + 0.168 \mathcal{C}_{10}^2 + 0.168 \mathcal{C}_9^2 + (41.5; 13.0) \mathcal{C}_7^2 \\ &\quad + 1.54 \text{Re}[\mathcal{C}_{10}] + 1.10 \text{Re}[\mathcal{C}_9] - (22.0; 2.86) \text{Re}[\mathcal{C}_7] + \\ &\quad + 0.172 \text{Im}[\mathcal{C}_7] + 0.0541 \text{Im}[\mathcal{C}_9] + 1.30 \text{Re}[\mathcal{C}_9 \mathcal{C}_7]]; \end{aligned} \quad (29)$$

$$\begin{aligned} (BR_{e;})_K &= 10^{-6} [0.519 + 0.0133 \mathcal{C}_{10}^2 + 0.0133 \mathcal{C}_9^2 + 0.0373 \mathcal{C}_7^2 \\ &\quad + 0.122 \text{Re}[\mathcal{C}_{10}] + 0.113 \text{Re}[\mathcal{C}_9] + 0.189 \text{Re}[\mathcal{C}_7] + \\ &\quad + 0.00783 \text{Im}[\mathcal{C}_7] + 0.00468 \text{Im}[\mathcal{C}_9] + 0.0444 \text{Re}[\mathcal{C}_9 \mathcal{C}_7]]; \end{aligned} \quad (30)$$

$$\begin{aligned} (BR_{e;})_K &= 10^{-6} [(2.25; 1.83) + 0.0481 \mathcal{C}_{10}^2 + 0.0481 \mathcal{C}_9^2 + (5.90; 2.13) \mathcal{C}_7^2 \\ &\quad + 0.441 \text{Re}[\mathcal{C}_{10}] + 0.351 \text{Re}[\mathcal{C}_9] + (2.31; 0.203) \text{Re}[\mathcal{C}_7] + \\ &\quad + 0.0584 \text{Im}[\mathcal{C}_7] + 0.0196 \text{Im}[\mathcal{C}_9] + 0.338 \text{Re}[\mathcal{C}_9 \mathcal{C}_7]]; \end{aligned} \quad (31)$$

$$\begin{aligned} (A_{FB}^{NN})_{X_s} &= 2.14 - 0.0542 \text{Im}[\mathcal{C}_{10}] - 0.467 \text{Re}[\mathcal{C}_{10}] - 1.98 \text{Re}[\mathcal{C}_7 \mathcal{C}_{10}] \\ &\quad - 0.246 \text{Re}[\mathcal{C}_9 \mathcal{C}_{10}] + 9.09 \text{Re}[\mathcal{C}_7] + 1.13 \text{Re}[\mathcal{C}_9]; \end{aligned} \quad (32)$$

$$\begin{aligned} (A_{FB}^{NN})_K &= 1.21 - 0.0211 \text{Im}[\mathcal{C}_{10}] - 0.263 \text{Re}[\mathcal{C}_{10}] - 0.488 \text{Re}[\mathcal{C}_7 \mathcal{C}_{10}] \\ &\quad - 0.0882 \text{Re}[\mathcal{C}_9 \mathcal{C}_{10}] + 2.24 \text{Re}[\mathcal{C}_7] + 0.404 \text{Re}[\mathcal{C}_9] \end{aligned} \quad (33)$$

where the suffixes  $X_s$ ,  $K$  and  $K^*$  correspond to the decays  $B \rightarrow (X_s; K; K^*) \ell^+ \ell^-$  and the coefficient  $C_i$  are defined as in Eq. (28).

Some remarks on the above equations are necessary.

The integrated normalized global asymmetries are given by

$$A_{FB}^0(s) = A_{FB}^{NN} \frac{N_i}{BR_{(e;)}} \quad (34)$$

where

$$N_{X_s} = \frac{BR_{B \rightarrow X_s \ell^+ \ell^-}}{0.104} \frac{1}{4} \frac{K_{ts}}{K_{cb}} \frac{1}{f(z)k(z)} = 3.30 \cdot 10^7 \quad (35)$$

$$N_{K;K^*} = \frac{1}{1.54 \text{ ps}} \frac{G_F^2}{2^{11}} \frac{1}{5} K_{ts}^2 K_{tb}^2 = 3.70 \cdot 10^7 \quad (36)$$

(the functions  $f$  and  $g$  can be found for instance in refs. [5, 37],  $z = M_c^2/M_b^2$ ).

There is no difference in the integrated non-normalized asymmetry for electrons and muons within an accuracy of 0.1%. The asymmetry for  $B \rightarrow K^+ \ell^-$  vanishes identically.

Taking into account the discussion of sect. 3 and the expressions (16) for the coefficients it is possible to scan the allowed parameter space looking for the extremal values of the observables we are interested in. We present the results of this analysis drawing scatter plots of the two integrated asymmetries as a function of the integrated BRs. In this way the correlation between the expectations of branching ratios and asymmetries is explicitly worked out. We report the plots concerning both the inclusive and exclusive decays.

We consider two possible scenarios.

- A. In Figs. 1{2 we have allowed all  $\mu$ s to vary at the same time. The range of possible values of the asymmetries is quite broad and in order to compare the SUSY result with the SM expectation we have reported in table 2 the central SM values for the same observables.

Even for values of the branching ratio around the SM expectation, a sensible SUSY contribution to the asymmetries is possible. An enhancement of the asymmetry is particularly favored when also a branching ratio increase is provided. In the MIA it is however possible to have a strongly depressed branching ratio with respect to the SM. In this case the possible values of the asymmetries can both be largely positive and largely negative and, in any case, very different from the SM expectations. Note then, the strong effect of the constraint from the exclusive channel  $B \rightarrow K^+ \ell^-$ : it is in fact responsible for the sharp cuts which are present for high values of the BR's and of the asymmetries.

- B. In Figs. 3{4 we have considered what can be expected if only the  $\mu_{LL}$  are different from zero. This more restricted scenario can occur in several models and is quite peculiar. In this case it is still possible a change of sign of  $C_7$  if  $\mu_{LL}$ 's are real and interesting results can in principle occur. Moreover we note that if the  $\mu_{LL}$ s acquire an imaginary part, the BR for the inclusive decay can increase considerably with respect to the SM but a "depression" is hardly possible. In this case the behavior of the FBA's is greatly modified, in fact they are non more allowed to change sign. This is due to the absence of the big  $(\mu_{23}^u)_{LR}$  contribution to  $C_{10}^{MI}$ . Sensitive enhancements of the asymmetries are anyway possible.

In tables 3{4 we present the results and the values of the Wilson coefficients for which the extremal values are obtained. In these tables  $C_i^{SUSY}$  are the total SUSY contribution to the Wilson coefficients at the  $M_B$  scale.

## 5 CP violating observables

The study of the CP-violating asymmetries defined in sect. 2 follows the same guide-lines of the last section. Now, due to the presence of the resonances, we are forced to perform the

analysis of the integrated CP asymmetries into s regions far from resonances.

First we consider the low-s region ( $1 \leq \sqrt{s} \leq 6 \text{ GeV}^2$ ), see fig. 5. Here we have quite high values of the differential branching ratios but the value of the CP asymmetries are just few percents. An analysis presented in ref. [38] compute a BR CP-asymmetry of about  $0.19^{+0.17}_{-0.19}\%$  for the SM in this region. In our computation we have not considered the SM contribution to CPV as we expect that SUSY can overcome it. This is indeed the case on a relevant part of the parameter space.

Then we take into account the high-s region ( $14 \leq \sqrt{s} \leq 23 \text{ GeV}^2$ ), see fig. 5. However for this part a comment is in order. The branching ratio in this zone of the spectrum is small, plagued with resonances (see fig. 5) and making any measurement on it represents an experimental challenge. Also from the theoretical point of view the presence of resonances makes it difficult to perform a rigorous estimate of CP observables. While a better comprehension of the resonant behavior of the BR is necessary, we present here a simple estimate of SUSY effects one can expect according to our present knowledge. CP asymmetry in this part of the spectrum was considered in [39] and found to be potentially large. We have checked it happens also in our framework.

Our results are plotted in figs. 6{9. In the case of exclusive decays we find very similar results so that we have not reported the corresponding graphs. Let us comment on the two possible scenarios of the previous section.

- A . In figs. 6{7 we have allowed all s to vary at the same time. The extremal values obtained in this case are in tables 7{8. We find that the values of the CP-asymmetries are much more constrained in the case of an enhancement of the BR. In the case of a "depression" of the SM result larger values of CP asymmetries are possible. In particular, from the first plot in fig. 6 we note that

the extremal values of  $A_G^{BR}$  can be in absolute value up to 6 times the one predicted in the SM ;

for values of the BR around the SM result, it is possible that the CP asymmetry be up to 2-3 times the one predicted in the SM .

Thus, the measure of CPV in these decays would represent an important issue for SUSY signals. So far a SM computation of  $A_L^{BR}$  and  $D_{A,A}^R$  is not reported to our knowledge in literature, as their measure is experimentally more difficult to be performed. However we think that similar considerations are still valid.

- B . In figs. 8{9 only the contribution of the  $\mu_{LL}$ 's is permitted. In figs. 8{9 for  $(\mu_{23}^{ud})_{LL} = 0$  the CP asymmetries are zero and the value of the BR for the SM + Diagonal contributions is obtained. If an enhancement of the BR is realized the value of the CP asymmetry is about the one obtainable in the SM . In this case a more refined study of the problem is necessary in order to see the interference between the SM and the SUSY contributions.

## 6 Conclusions

A detailed analysis of SUSY contributions to CPV in semileptonic rare B decays has been performed using the mass insertion approximation. We have estimated the amount of CPV that can be expected given the present limits on the  $M_I$ 's. Several observables have been taken into account. We have studied the correlation between the observables and the integrated BR for the inclusive decays. We have discussed the possible scenarios that can be realized according to the particular insertion one considers. The largest deviation with respect to the SM are of course expected when LR mass insertions are present. In the case in which only LL insertions are taken into account a detectable enhancement of the BR can be expected if the insertion has got an imaginary part. However in this case CP asymmetries are of the order of the SM ones and a more involved discussion about the interference of the two contributions has to be performed.

## Acknowledgments

We want to thank Antonio Masiero for fruitful discussions and suggestions.

I.S. wants to thank Della Riccia Foundation (Florence, Italy) for partial support. This work was partially supported by INFN, by the TM R {EEC network "Beyond the Standard Model" (contract number ERBFMRX {CT 96{0090) and by TM R {EEC network "Eurodaphne" (contract number ERBFMX {CT 98{0169).

## References

- [1] See, for instance, The BaBar Physics Book, SLAC-R-504 and references therein.
- [2] B. Grinstein, M. J. Savage and M. B. Wise Nucl. Phys. B 319 (1989) 271.
- [3] G. Buchalla and A. J. Buras, Nucl. Phys. B 400 (1993) 225.
- [4] M. Misiak, Nucl. Phys. B 393 (1993) 23, erratum *ibid.* B 439 (1995) 461;  
A. J. Buras and M. M. M. un z, Phys. Rev. D 52 (1995) 186.
- [5] G. Buchalla, A. J. Buras and M. E. Lautenbacher, Rev. Mod. Phys. 68 (1996) 1125.
- [6] C. Bobeth, M. Misiak, J. Urban, hep-ph/9910220.
- [7] N. G. Deshpande, J. Trampetic and K. Panose, Phys. Rev. D 39 (1989) 1461;  
C. S. Lim, T. M. Orozum i and A. I. Sanda, Phys. Lett. B 218 (1989) 343;  
A. I. Vainshtein et al., Yad. Fiz. 24 (1976) 820 [Sov. J. Nucl. Phys. 24 (1976) 427];  
P. J. O'Donnell and H. K. Tung, Phys. Rev. D 43 (1991) 2076.
- [8] A. F. Falk, M. Luke and M. J. Savage, Phys. Rev. D 49 (1994) 3367.

- [9] A . A li, G . H iller, L . T . H andoko and T . M orozum i, Phys. Rev. D 55 (1997) 4105;  
A . A li and G . H iller, Phys. Rev. D 58 (1998) 071501 and Phys. Rev. D 58 (1998) 074001.
- [10] G . Buchalla and G . Isidori, Nucl. Phys. B 525 (1998) 333.
- [11] A . A li, T . M annel and T . M orozum i, Phys. Rev. B 273 (1991) 505;  
M . R . A hm adi, Phys. Rev. D 53 (1996) 2843;  
C . D . Lu and D . X . Zhang, Phys. Lett. B 397 (1997) 279.
- [12] F . K ruger and L . M . Sehgal, Phys. Lett. B 380 (1996) 199.
- [13] Z . Ligeti, I . W . Stewart and M . B . W ise, Phys. Lett. B 420 (1998) 359.
- [14] G . Buchalla, G . Isidori and S . J . Rey, Nucl. Phys. B 511 (1998) 594.
- [15] J . W . Chen, G . Rupak and M . J . Savage, Phys. Rev. B 410 (1997) 285.
- [16] S . Bertolini, F . Borzum ati, A . M asiero and G . R idol , Nucl. Phys. B 353 (1991) 591;  
A . A li, G . F . G iudice and T . M annel, Z . Phys. C 67 (1995) 417.
- [17] J . Hewett and J . D . W ells, Phys. Rev. D 55 (1997) 5549.
- [18] P . Cho, M . M isiak and D . W yler, Phys. Rev. D 54 (1996) 3329.
- [19] T . G oto, Y . O kada, Y . Shim izu and M . Tanaka, Phys. Rev. D 55 (1997) 4273;  
T . G oto, Y . O kada and Y . Shim izu, Phys. Rev. D 58 (1998) 094006.
- [20] CLEO Collaboration, CONF98-17, ICHEP98, 1011.
- [21] Y . G . K im , P . K o, J . S . Lee, Nucl. Phys. B 544 (1999) 64.
- [22] C . S . Huang, W . Liao and Q . S . Yan, Phys. Rev. D 59 (1999) 011701.
- [23] C . S . Huang and Q . S . Yan, Phys. Lett. B 442 (1998) 209.
- [24] S . Baek and P . K o, hep-ph/9904283.
- [25] E . Lunghi, A . M asiero, I . Scim em i, L . Silvestrini hep-ph/9906286.
- [26] L . J . Hall, V . A . Kostolecki and S . Raby, Nucl. Phys. B 267 (1986) 415.
- [27] F . G abbiani and A . M asiero, Nucl. Phys. B 322 (1989) 235.
- [28] J . S . Hagelin, S . Kelley and T . Tanaka, Nucl. Phys. B 415 (1994) 293-331;  
E . G abrielli, A . M asiero and L . Silvestrini, Phys. Lett. B 374 (1996) 80;  
J . A . Bagger, K . T . M atchev and R . Zhang, Phys. Lett. B 412 (1997) 77-85;  
M . C iuchini et al, JHEP 10 (1998) 008;  
R . Contino and I . Scim em i, EurPhysJ.C 10 (1999) 347.
- [29] F . G abbiani, E . G abrielli, A . M asiero and L . Silvestrini, Nucl. Phys. B 477 (1996) 321.

- [30] J. A. Casas and S. Dimopoulos, *Phys. Lett. B* 387 (1996) 107.
- [31] A. Ali, P. Ball, L. T. Handoko, G. Hiller [hep-ph/9910221](#).
- [32] T. A. Older et al. (CDF Collaboration), *Phys. Rev. Lett.* 83 (1999) 3378.
- [33] R. Casalbuoni et al., *Phys. Rep.* 281 (1997) 145.
- [34] C. Greub, A. Ioannissian and D. Wyler, *Phys. Lett. B* 346 (1995) 149.
- [35] C. Caso et al. *Eur. Phys. J. C* (1998) 1.
- [36] K. Chetyrkin, M. Misiak and M. Munz, *Phys. Lett. B* 400 (1997) 206-219;  
Erratum *ibid.* B 425 (1998) 414.
- [37] N. Cabibbo and L. Maiani, *Phys. Lett. B* 79 (1978) 109;  
C. S. Kim and A. D. Martin, *Phys. Lett. B* 225 (1989) 186.
- [38] A. Ali, G. Hiller *Eur. Phys. J. C* 8 (1999) 619.
- [39] C.-S. Huang, L. Wei [hep-ph/9908246](#).

## List of Tables

|   |  |    |
|---|--|----|
| 1 | Central values of physical constants used in the phenomenological analysis .   | 17 |
| 2 | Central SM values for physical observables . . . . .   | 17 |
| 3 | Inclusive decays $B \rightarrow X_s \gamma$ . Maximum ( $M$ ) and minimum ( $m$ ) values of the BR and of the FB asymmetries (scenario A). $C_i^{SU(2)_L}$ are the total SU(2) <sub>L</sub> contributions to the Wilson coefficients at the $M_B$ scale. . . . . | 18 |
| 4 | Exclusive decays $B \rightarrow (K; \bar{K}) \gamma$ . Maximum ( $M$ ) and minimum ( $m$ ) values of the BR and of the FB asymmetries (scenario A). . . . .  | 18 |
| 5 | Inclusive decays $B \rightarrow X_s \gamma$ . Maximum ( $M$ ) and minimum ( $m$ ) values of the BR and of the FB asymmetries for the LL-only case (scenario B). . . .  | 19 |
| 6 | Exclusive decays $B \rightarrow (K; \bar{K}) \gamma$ . Maximum ( $M$ ) and minimum ( $m$ ) values of the BR and of the FB asymmetries for the LL-only case (scenario B). . .   | 19 |
| 7 | Inclusive decays $B \rightarrow X_s \gamma$ . Maximum ( $M$ ) and minimum ( $m$ ) values of the CP asymmetries (scenario A) in the regions of low $s$ ( $1 \leq s \leq 6 \text{ GeV}^2$ ) and high $s$ ( $14 \leq s \leq 23 \text{ GeV}^2$ ). . . . .            | 20 |
| 8 | Exclusive decays $B \rightarrow (K; \bar{K}) \gamma$ . Maximum ( $M$ ) and minimum ( $m$ ) values of the CP asymmetries (scenario A) in the regions of low $s$ ( $1 \leq s \leq 6 \text{ GeV}^2$ ) and high $s$ ( $14 \leq s \leq 23 \text{ GeV}^2$ ). . . . .   | 21 |

## List of Figures

|   |  |    |
|---|--|----|
| 1 | Inclusive decay $B \rightarrow X_s^+$ . Scatter plots of the Integrated FB asymmetries ( $A_{FB}$ and $A_{FB}^0$ ) versus the BR (in units of $10^{-6}$ ). Scenario A . . . . .            | 22 |
| 2 | Exclusive decay $B \rightarrow K^+$ . Scatter plots of the Integrated FB asymmetries ( $A_{FB}$ and $A_{FB}^0$ ) versus the integrated BR (in units of $10^{-6}$ ). Scenario A . . . . .   | 23 |
| 3 | Inclusive decay $B \rightarrow X_s^+$ . Scatter plots of the Integrated FB asymmetries ( $A_{FB}$ and $A_{FB}^0$ ) versus the integrated BR (in units of $10^{-6}$ ). Scenario B . . . . . | 24 |
| 4 | Exclusive decay $B \rightarrow K^+$ . Scatter plots of the Integrated FB asymmetries ( $A_{FB}$ and $A_{FB}^0$ ) versus the integrated BR (in units of $10^{-6}$ ). Scenario B . . . . .   | 25 |
| 5 | Branching Ratio for the inclusive decay $B \rightarrow X_s^{*+}$ plotted in units of $10^{-6}$ in the low $\{s$ and high $\{s$ regions. . . . .  | 26 |
| 6 | Inclusive decay $B \rightarrow X_s^{*+}$ . Scatter plots of $A_G^{BR}$ and $D_{A^0}$ versus the integrated BR (in units of $10^{-6}$ ) in the low $\{s$ region. Scenario A . . . . .       | 27 |
| 7 | Inclusive decay $B \rightarrow X_s^{*+}$ . Scatter plots of $A_G^{BR}$ and $D_{A^0}$ versus the integrated BR (in units of $10^{-6}$ ) in the high $\{s$ region. Scenario A . . . . .      | 28 |
| 8 | Inclusive decay $B \rightarrow X_s^+$ . Scatter plots of $A_G^{BR}$ versus the integrated BR (in units of $10^{-6}$ ) in the low $\{s$ region. Scenario B . . . . .                        | 29 |
| 9 | Inclusive decay $B \rightarrow X_s^+$ . Scatter plots of $A_G^{BR}$ versus the integrated BR (in units of $10^{-6}$ ) in the high $\{s$ region Scenario B . . . . .                        | 30 |

|                    |           |
|--------------------|-----------|
| $m_t$              | 173.8 GeV |
| $m_b$              | 4.8 GeV   |
| $m_c$              | 1.4 GeV   |
| $m_s$              | 125 MeV   |
| $M_B$              | 5.27 GeV  |
| $\sin^2(\theta_W)$ | 0.2334    |
| $\alpha_{em}(m_Z)$ | 1/128.9   |
| $\sin^2(\theta_W)$ | 0.2334    |

Table 1: Central values of physical constants used in the phenomenological analysis

|                    |                     |                      |                     |
|--------------------|---------------------|----------------------|---------------------|
| $BR(e; X_s)$       | $9.6 \cdot 10^{-6}$ | $BR(\mu; X_s)$       | $6.3 \cdot 10^{-6}$ |
| $A_{FB}(e; X_s)$   | 23.0%               | $A_{FB}(\mu; X_s)$   | 23.0%               |
| $A_{FB}^0(e; X_s)$ | 7.1%                | $A_{FB}^0(\mu; X_s)$ | 11%                 |
| $BR(e; K)$         | $5.7 \cdot 10^{-7}$ | $BR(\mu; K)$         | $5.7 \cdot 10^{-7}$ |
| $BR(e; K^*)$       | $2.3 \cdot 10^{-6}$ | $BR(\mu; K^*)$       | $1.9 \cdot 10^{-6}$ |
| $A_{FB}(e; K)$     | 17.0%               | $A_{FB}(\mu; K)$     | 17.0%               |
| $A_{FB}^0(e; K)$   | 19.8%               | $A_{FB}^0(\mu; K)$   | 24.4%               |

Table 2: Central SM values for physical observables

|                             |   |      | $C_7^{\text{SUSY}}$ | $C_9^{\text{SUSY}}$ | $C_{10}^{\text{SUSY}}$ |
|-----------------------------|---|------|---------------------|---------------------|------------------------|
| $10^6 \text{BR}(\epsilon)$  | M | 22:2 | 0:734 0:086I        | 1:17 0:182I         | 0:326+ 6:69I           |
|                             | m | 4:19 | 0:0429 0:0052I      | 1:72 0:863I         | 4:23 1:05I             |
| $10^6 \text{BR}(\ )$        | M | 17:4 | 0:734 0:086I        | 1:17 0:182I         | 0:326+ 6:69I           |
|                             | m | 1:72 | 0:0429 0:0052I      | 1:72 0:863I         | 4:23 1:05I             |
| $A_{\text{FB}}$             | M | 33:5 | 0:613+ 0:254I       | 1:37+ 0:881I        | 0:149 1:89I            |
|                             | m | 33:6 | 0:567 0:305I        | 1:98 0:722I         | 7:65 1:52I             |
| $A_{\text{FB}}^0(\epsilon)$ | M | 23:7 | 0:612+ 0:0328I      | 0:522 0:88I         | 2:9+ 0:911I            |
|                             | m | 19:  | 0:561+ 0:165I       | 2:05+ 0:664I        | 7:74+ 1:65I            |
| $A_{\text{FB}}^0(\ )$       | M | 27:3 | 0:678+ 0:154I       | 1:15+ 0:805I        | 0:514 1:46I            |
|                             | m | 27:  | 0:561+ 0:165I       | 2:05+ 0:664I        | 7:74+ 1:65I            |

Table 3: Inclusive decays  $B \rightarrow X_s^{*\prime} \ell$ . Maximum (M) and minimum (m) values of the BR and of the FB asymmetries (scenario A).  $C_i^{\text{SUSY}}$  are the total SUSY contributions to the Wilson coefficients at the  $M_B$  scale.

|                                  |   |        | $C_7^{\text{SUSY}}$ | $C_9^{\text{SUSY}}$ | $C_{10}^{\text{SUSY}}$ |
|----------------------------------|---|--------|---------------------|---------------------|------------------------|
| $10^6 \text{BR}(\epsilon; K)$    | M | 1:35   | 0:0265 0:0448I      | 0:665 0:101I        | 2:13 5:75I             |
|                                  | m | 0:0979 | 0:0359+ 0:0051I     | 1:72+ 0:872I        | 4:67 0:25I             |
| $10^6 \text{BR}(\ ; K)$          | M | 1:35   | 0:0265 0:0448I      | 0:665 0:101I        | 2:13 5:75I             |
|                                  | m | 0:0976 | 0:0359+ 0:0051I     | 1:72+ 0:872I        | 4:67 0:25I             |
| $10^6 \text{BR}(\epsilon; K^*)$  | M | 5:44   | 0:734 0:086I        | 1:17 0:182I         | 0:326+ 6:69I           |
|                                  | m | 0:742  | 0:0429 0:0052I      | 1:72 0:863I         | 4:23 1:05I             |
| $10^6 \text{BR}(\ ; K^*)$        | M | 4:84   | 0:657 0:079I        | 0:572 0:218I        | 3:51 1:59I             |
|                                  | m | 0:421  | 0:0493+ 0:112I      | 1:7 0:51I           | 5:59 0:501I            |
| $A_{\text{FB}}(K^*)$             | M | 25:3   | 0:678+ 0:154I       | 1:15+ 0:805I        | 0:514 1:46I            |
|                                  | m | 25:2   | 0:561+ 0:165I       | 2:05+ 0:664I        | 7:74+ 1:65I            |
| $A_{\text{FB}}^0(\epsilon)(K^*)$ | M | 33:3   | 0:612+ 0:0328I      | 0:522 0:88I         | 2:9+ 0:911I            |
|                                  | m | 29:5   | 0:561+ 0:165I       | 2:05+ 0:664I        | 7:74+ 1:65I            |
| $A_{\text{FB}}^0(\ )(K^*)$       | M | 36:    | 0:678+ 0:154I       | 1:15+ 0:805I        | 0:514 1:46I            |
|                                  | m | 35:6   | 0:561+ 0:165I       | 2:05+ 0:664I        | 7:74+ 1:65I            |

Table 4: Exclusive decays  $B \rightarrow (K; K^*)^{*\prime} \ell$ . Maximum (M) and minimum (m) values of the BR and of the FB asymmetries (scenario A).

|                             |   |      | $C_7^{\text{SUSY}}$ | $C_9^{\text{SUSY}}$ | $C_{10}^{\text{SUSY}}$ |
|-----------------------------|---|------|---------------------|---------------------|------------------------|
| $10^6 \text{BR}(\epsilon)$  | M | 19:3 | 0:647 + 0:265I      | 0:57 + 0:377I       | 5:8 0:386I             |
|                             | m | 8:98 | 0:0545 0:0218I      | 0:272 0:031I        | 4:93 + 0:0318I         |
| $10^6 \text{BR}(\mu)$       | M | 14:6 | 0:647 + 0:265I      | 0:57 + 0:377I       | 5:8 0:386I             |
|                             | m | 6:59 | 0:0181 0:00779I     | 0:376 0:0111I       | 4:83 + 0:0113I         |
| $A_{\text{FB}}$             | M | 32:4 | 0:645 0:235I        | 0:568 0:335I        | 5:79 + 0:342I          |
|                             | m | 18:9 | 0:0742 + 0:0316I    | 0:456 + 0:0449I     | 4:75 0:0459I           |
| $A_{\text{FB}}^0(\epsilon)$ | M | 22:8 | 0:605 0:0741I       | 0:511 0:105I        | 5:74 + 0:108I          |
|                             | m | 2:76 | 0:0741 0:0173I      | 0:455 0:0247I       | 4:75 + 0:0252I         |
| $A_{\text{FB}}^0(\mu)$      | M | 26:8 | 0:605 0:0741I       | 0:511 0:105I        | 5:74 + 0:108I          |
|                             | m | 4:88 | 0:0741 0:0173I      | 0:455 0:0247I       | 4:75 + 0:0252I         |

Table 5: Inclusive decays  $B \rightarrow X_s \ell^+ \ell^-$ . Maximum (M) and minimum (m) values of the BR and of the FB asymmetries for the LL (only case (scenario B)).

|                                 |   |       | $C_7^{\text{SUSY}}$ | $C_9^{\text{SUSY}}$ | $C_{10}^{\text{SUSY}}$ |
|---------------------------------|---|-------|---------------------|---------------------|------------------------|
| $10^6 \text{BR}(\epsilon; K)$   | M | 0:927 | 0:65 + 0:253I       | 0:574 + 0:36I       | 5:8 0:368I             |
|                                 | m | 0:48  | 0:0741 0:0173I      | 0:455 0:0247I       | 4:75 + 0:0252I         |
| $10^6 \text{BR}(\mu; K)$        | M | 0:925 | 0:65 + 0:253I       | 0:574 + 0:36I       | 5:8 0:368I             |
|                                 | m | 0:478 | 0:0741 0:0173I      | 0:455 0:0247I       | 4:75 + 0:0252I         |
| $10^6 \text{BR}(\epsilon; K^*)$ | M | 4:65  | 0:647 + 0:265I      | 0:57 + 0:377I       | 5:8 0:386I             |
|                                 | m | 2:2   | 0:0545 0:0218I      | 0:272 0:031I        | 4:93 + 0:0318I         |
| $10^6 \text{BR}(\mu; K^*)$      | M | 4:01  | 0:647 + 0:265I      | 0:57 + 0:377I       | 5:8 0:386I             |
|                                 | m | 1:76  | 0:0741 0:0173I      | 0:455 0:0247I       | 4:75 + 0:0252I         |
| $A_{\text{FB}}(K)$              | M | 24:3  | 0:605 0:0741I       | 0:511 0:105I        | 5:74 + 0:108I          |
|                                 | m | 13:9  | 0:0742 + 0:0316I    | 0:456 + 0:0449I     | 4:75 0:0459I           |
| $A_{\text{FB}}^0(\epsilon)(K)$  | M | 32:8  | 0:605 0:0741I       | 0:511 0:105I        | 5:74 + 0:108I          |
|                                 | m | 13:7  | 0:0742 + 0:0316I    | 0:456 + 0:0449I     | 4:75 0:0459I           |
| $A_{\text{FB}}^0(\mu)(K)$       | M | 35:5  | 0:605 0:0741I       | 0:511 0:105I        | 5:74 + 0:108I          |
|                                 | m | 18:6  | 0:0742 + 0:0316I    | 0:456 + 0:0449I     | 4:75 0:0459I           |

Table 6: Exclusive decays  $B \rightarrow (K; K^*) \ell^+ \ell^-$ . Maximum (M) and minimum (m) values of the BR and of the FB asymmetries for the LL (only case (scenario B)).

|                          |   |         | $C_7^{\text{SUSY}}$ | $C_9^{\text{SUSY}}$ | $C_{10}^{\text{SUSY}}$ |
|--------------------------|---|---------|---------------------|---------------------|------------------------|
| $A_G^{\text{BR}}$ (low)  | M | 1:31    | 0:248 + 0:347I      | 1:78 + 0:652I       | 5:32 0:206I            |
|                          | m | 1:54    | 0:122 0:271I        | 1:67 0:791I         | 4:63 + 0:0593I         |
| $A_G^{\text{BR}}$ (high) | M | 43:2    | 0:0359 + 0:0051I    | 1:72 + 0:872I       | 4:67 0:25I             |
|                          | m | 44:1    | 0:122 0:271I        | 1:67 0:791I         | 4:63 + 0:0593I         |
| $D_{A^0}$ (low)          | M | 0:00585 | 0:0439 0:188I       | 1:81 + 1:09I        | 5:14 4:I               |
|                          | m | 0:00619 | 0:027 + 0:144I      | 1:82 1:17I          | 5:22 + 3:33I           |
| $D_{A^0}$ (high)         | M | 0:392   | 0:0146 + 0:198I     | 1:47 0:439I         | 4:69 2:72I             |
|                          | m | 0:401   | 0:0514 + 0:00682I   | 1:69 + 0:49I        | 4:97 + 2:19I           |
| $A_L^{\text{BR}}$ (low)  | M | 0:292   | 0:248 + 0:347I      | 1:78 + 0:652I       | 5:32 0:206I            |
|                          | m | 0:344   | 0:122 0:271I        | 1:67 0:791I         | 4:63 + 0:0593I         |
| $A_L^{\text{BR}}$ (high) | M | 4:24    | 0:0359 + 0:0051I    | 1:72 + 0:872I       | 4:67 0:25I             |
|                          | m | 3:81    | 0:122 0:271I        | 1:67 0:791I         | 4:63 + 0:0593I         |
| $D_A$ (low)              | M | 0:00131 | 0:00781 + 0:0518I   | 1:52 + 0:0931I      | 5:21 3:01I             |
|                          | m | 0:00143 | 0:027 + 0:144I      | 1:82 1:17I          | 5:22 + 3:33I           |
| $D_A$ (high)             | M | 0:116   | 0:0514 + 0:0821I    | 1:64 0:785I         | 4:94 1:56I             |
|                          | m | 0:123   | 0:0199 + 0:0504I    | 1:76 0:777I         | 4:87 + 1:68I           |

Table 7: Inclusive decays  $B \rightarrow X_s^{*+} \gamma$ . Maximum (M) and minimum (m) values of the CP asymmetries (scenario A) in the regions of low  $s$  ( $1 \leq s \leq 6 \text{ GeV}^2$ ) and high  $s$  ( $14 \leq s \leq 23 \text{ GeV}^2$ ).

|                                   |   |         | $C_7^{\text{SUSY}}$ | $C_9^{\text{SUSY}}$ | $C_{10}^{\text{SUSY}}$ |
|-----------------------------------|---|---------|---------------------|---------------------|------------------------|
| $A_G^{\text{BR}}(K, \text{low})$  | M | 1:24    | 0:248 + 0:347I      | 1:78 + 0:652I       | 5:32 0:206I            |
|                                   | m | 1:45    | 0:122 0:271I        | 1:67 0:791I         | 4:63 + 0:0593I         |
| $A_G^{\text{BR}}(K, \text{high})$ | M | 40:3    | 0:0359 + 0:0051I    | 1:72 + 0:872I       | 4:67 0:25I             |
|                                   | m | 40:7    | 0:122 0:271I        | 1:67 0:791I         | 4:63 + 0:0593I         |
| $D_{A^0}(K, \text{low})$          | M | 0:00758 | 0:027 + 0:144I      | 1:82 1:17I          | 5:22 + 3:33I           |
|                                   | m | 0:0072  | 0:0439 0:188I       | 1:81 + 1:09I        | 5:14 4:I               |
| $D_{A^0}(K, \text{high})$         | M | 0:383   | 0:0514 + 0:00682I   | 1:69 + 0:49I        | 4:97 + 2:19I           |
|                                   | m | 0:374   | 0:0146 + 0:198I     | 1:47 0:439I         | 4:69 2:72I             |
| $A_L^{\text{BR}}(K, \text{low})$  | M | 0:225   | 0:167 + 0:362I      | 1:68 + 0:568I       | 5:53 + 0:539I          |
|                                   | m | 0:263   | 0:122 0:271I        | 1:67 0:791I         | 4:63 + 0:0593I         |
| $A_L^{\text{BR}}(K, \text{high})$ | M | 8:3     | 0:0359 + 0:0051I    | 1:72 + 0:872I       | 4:67 0:25I             |
|                                   | m | 7:91    | 0:122 0:271I        | 1:67 0:791I         | 4:63 + 0:0593I         |
| $D_A(K, \text{low})$              | M | 0:00136 | 0:027 + 0:144I      | 1:82 1:17I          | 5:22 + 3:33I           |
|                                   | m | 0:00126 | 0:00781 + 0:0518I   | 1:52 + 0:0931I      | 5:21 3:01I             |
| $D_A(K, \text{high})$             | M | 0:0369  | 0:0199 + 0:0504I    | 1:76 0:777I         | 4:87 + 1:68I           |
|                                   | m | 0:0356  | 0:0514 + 0:0821I    | 1:64 0:785I         | 4:94 1:56I             |
| $A_G^{\text{BR}}(K, \text{low})$  | M | 0:986   | 0:315 + 0:385I      | 1:55 + 0:885I       | 4:+ 0:441I             |
|                                   | m | 1:08    | 0:122 0:271I        | 1:67 0:791I         | 4:63 + 0:0593I         |
| $A_G^{\text{BR}}(K, \text{high})$ | M | 33:     | 0:0359 + 0:0051I    | 1:72 + 0:872I       | 4:67 0:25I             |
|                                   | m | 34:1    | 0:122 0:271I        | 1:67 0:791I         | 4:63 + 0:0593I         |
| $A_L^{\text{BR}}(K, \text{low})$  | M | 0:177   | 0:315 + 0:385I      | 1:55 + 0:885I       | 4:+ 0:441I             |
|                                   | m | 0:195   | 0:122 0:271I        | 1:67 0:791I         | 4:63 + 0:0593I         |
| $A_L^{\text{BR}}(K, \text{high})$ | M | 7:01    | 0:0359 + 0:0051I    | 1:72 + 0:872I       | 4:67 0:25I             |
|                                   | m | 6:95    | 0:122 0:271I        | 1:67 0:791I         | 4:63 + 0:0593I         |

Table 8: Exclusive decays  $B \rightarrow (K; K^*) \ell^+ \ell^-$ . Maximum (M) and minimum (m) values of the CP asymmetries (scenario A) in the regions of low  $s$  ( $1 \leq s \leq 6 \text{ GeV}^2$ ) and high  $s$  ( $14 \leq s \leq 23 \text{ GeV}^2$ ).




Figure 1: Inclusive decay  $B \rightarrow X_s^+$ . Scatter plots of the Integrated  $F\{B$  asymmetries ( $A_{FB}$  and  $A_{FB}^0$ ) versus the BR (in units of  $10^6$ ). Scenario A.




Figure 2: Exclusive decay  $B \rightarrow K^+$ . Scatter plots of the Integrated FB asymmetries ( $A_{FB}$  and  $A_{FB}^0$ ) versus the integrated BR (in units of  $10^6$ ). Scenario A.




Figure 3: Inclusive decay  $B \rightarrow X_s^+$ . Scatter plots of the Integrated FB asymmetries ( $A_{FB}$  and  $A_{FB}^0$ ) versus the integrated BR (in units of  $10^6$ ). Scenario B.




Figure 4: Exclusive decay  $B \rightarrow K^+$ . Scatter plots of the Integrated FB asymmetries ( $A_{FB}$  and  $A_{FB}^0$ ) versus the integrated BR (in units of  $10^6$ ). Scenario B.




Figure 5: B branching Ratio for the inclusive decay  $B \rightarrow X_s^{*+} \gamma$  plotted in units of  $10^{-6}$  in the low  $s$  and high  $s$  regions.




Figure 6: Inclusive decay  $B \rightarrow X_s^{*+}$ . Scatter plots of  $A_G^{BR}$  and  $D_{A^0}$  versus the integrated BR (in units of  $10^6$ ) in the low  $\{s$  region. Scenario A.




Figure 7: Inclusive decay  $B \rightarrow X_s^+ \gamma$ . Scatter plots of  $A_G^{BR}$  and  $D_{A^0}$  versus the integrated BR (in units of  $10^6$ ) in the high  $\sqrt{s}$  region. Scenario A.




Figure 8: Inclusive decay  $B \rightarrow X_s + \gamma$ . Scatter plots of  $A_G^{BR}$  versus the integrated BR (in units of  $10^6$ ) in the low  $\sqrt{s}$  region. Scenario B.




Figure 9: Inclusive decay  $B \rightarrow X_s^+$ . Scatter plots of  $A_G^{BR}$  versus the integrated BR (in units of  $10^6$ ) in the high  $\sqrt{s}$  region Scenario B.