Can we live in a self-tuning universe?

Sean M . Carroll and Laura M $ersin \hat{\mathcal{I}}$

¹Enrico Ferm i Institute and Department of Physics, University of Chicago 5640 S.Ellis Avenue, Chicago, IL 60637, USA carroll@theory.uchicago.edu

²Scuola Norm ale Superiore, Piazza dei Cavalieri 7, I-56126 Pisa, Italy mersini@cibs.sns.it

Abstract

The self-tuning brane scenario is an attempt to solve the cosm ological constant problem in the context of extra dimensions. Rather than making the vacuum energy small, this approach proceeds by removing the gravitational elect of vacuum energy on the expansion of the universe. Such behavior is only possible through changing the Friedmann equation of conventional cosm ology, and we discuss di culties in obtaining cosm ological evolution compatible with observation in this context. Speci c models considered include a bulk scalar eld coupling to the brane via a conform al transform ation of the brane metric, and via a rescaling of the brane volum element.

1 Introduction

The fact that the observed cosm ological constant is much smaller than the expected value [1, 2, 3] may provide a crucial clue in our attempts to understand the nature of spacetime. Most attempts to solve the problem can be characterized as making the vacuum energy much smaller than its natural value. A lternatively, however, we can in agine keeping a large vacuum energy, but changing the gravitational dynamics in such a way that the vacuum does not act as a (signi cant) source of spacetime curvature. Since it is only through its gravitational in uence that the vacuum energy can be measured, such an arrangement could reconcile the naive estimates vac $(10^{18} \text{ GeV})^4$ with the observationally favored result vac $(10^{3} \text{ eV})^4$ [4, 3].

The idea of brane-worlds and large extra dimensions [5] opens up a new set of ways to think about the cosm ological constant problem. In these scenarios, our observed fourdimensional theory of gravity is descended from a higher-dimensional embedding, and in principle the resulting dynamics can dier dram atically from a straightforward four-dimensional expectation.

An example of such an altered dynamics is provided by the idea of self-tuning branes [6,7]. Here, matter elds on a three-brane with a single extra dimension are coupled to a bulk scalar eld. W ith an appropriate choice of couplings, M inkow skian solutions on the brane can be found with any brane cosm ological constant. In this paper, we put aside fundamental issues of the feasibility of the self-tuning idea (e.g., the role of singularities) to concentrate on whether this kind of scenario can be made compatible with conventional cosm ology. (See [8, 9] for investigations of brane-world cosm ology, and [10, 11] for studies of the self-tuning scenario.)

On a self-tuning brane, there is vacuum energy (essentially the tension of the brane), but the spacetime geometry is nevertheless at. Somehow, then, the geometry is insensitive to the vacuum energy. It is necessary, however, that spacetime respond to at least some sorts of energy density; in particular, the success of B ig B ang Nucleosynthesis (BBN) [12] provides evidence in favor of the standard picture in the radiation-dom inated era. It is therefore necessary to recover at least some portion of conventional cosm ology, while rem oving the e ects of the cosm ological constant.

In this work we attempt to characterize the empirical challenges to a successful self-tuning cosm ology. In the next section we discuss in general how the self-tuning mechanism may be understood in terms of the energy and pressure on the brane, pointing out the distinctions

with conventional Friedm ann cosm ology. We then consider two specian models of self-tuning, and derive elective Friedm ann-like equations relating the Hubble parameter to the energy and pressure. In section 4 we compare these models with what we know about the universe, and discuss whether they may be brought into agreement with observation. Although such agreement seems unlikely, we are unable to rule it out entirely.

2 The secret of self-tuning cosm ology

In this section we consider how a theory of gravity m ay in principle be insensitive to vacuum energy while allowing other forms of energy-m om entum to in uence spacetime curvature. We consider a at Robertson-Walkermetric in 3 + 1 dimensions,

$$ds^2 = dt^2 + a^2 (t) dx^2$$
; (1)

where a(t) is the scale factor and dx^2 is the at Euclidean metric. An energy-momentum tensor consistent with a Robertson-Walkermetric will be spatially isotropic, taking the form

$$T = diag(; p; p; p);$$
 (2)

where is the energy density and p the pressure. The conventional Friedm ann equation of general relativity is then

$$H^{2} \qquad \frac{a}{a}^{2} = \frac{8 G}{3}$$
; (3)

where H is the Hubble parameter and G is the (four-dimensional) Newton's constant.

In a self-tuning model, it is possible to nd a M inkowski-spacetime solution (H = 0) regardless of the value of $_{vac}$ (when all other energy densities vanish). Clearly, for this to be possible (3) will have to be modiled, either in its explicit form or in the denition of . If, however, we consider theories which arise from varying a specified action with respect to the metric tensor to derive gravitational eld equations, the energy density and pressure (in this coordinate system) are dened by

$$= T_{0}^{0} = 2 \frac{1}{p \overline{q}} g^{00} \frac{\partial L_{m \text{ atter}}}{\partial q^{00}}; \quad p = T_{1}^{1} = 2 \frac{1}{p \overline{q}} g^{11} \frac{\partial L_{m \text{ atter}}}{\partial q^{11}}; \quad (4)$$

where g is the absolute value of the determ inant of the metric and L_{matter} is the matter Lagrange density. These de nitions do not discrim inate between dierent forms of energy density; there is now ay for the gravitational eld to tell the dierence between energy density from the vacuum and energy density from any other source. How, then, can we modify the theory to allow at solutions in the presence of vacuum energy, while remaining sensitive to the in uence of other sources on the expansion rate? G iven the metric (1) and energy-momentum tensor (2), we seek an equation written in terms of , p, a and its derivatives (which appear in the curvature tensor), and possibly explicit additional elds. A lthough vacuum energy should enter any such equation in the same way as other energy, the vacuum does have a distinguishing characteristic, namely its equation of state:

$$p_{vac} = vac : (5)$$

W ith the ingredients at our disposal, this relation suggests a form for a modi ed Friedmann equation:

$$H^{2} = f(;p)(+p) + other term s;$$
 (6)

where f(;p) is a well-behaved function at p =. Such a relation would allow for a M inkowski solution (H = 0) in the presence of arbitrary vacuum energy.

A lthough (6) di ers from the conventional Friedmann law, it is not necessarily incom – patible with observation. Consider for example a hypothetical relation

$$H^{2} = 2 G (+p) :$$
 (7)

During a matter-dom inated era, p = 0, and the Hubble parameter will dier by a factor p = 3=2 0.87 for a given value of the energy density. However, since the scaling of H with is as in the conventional theory, and p = 3=2 is close to unity, it would be hard to distinguish between the behavior predicted by (7) and the usual Friedmann equation during matter dom ination. During radiation dom ination, p = -3, and (7) precisely recovers the conventional expectation (by construction). This is in portant, as our most precise quantitative evidence in favor of conventional cosm ology comes from BBN, which occurs while the universe is radiation-dom inated.¹ Finally, this relation would solve the cosm ological constant problem, as H = 0 when p = -3.

Unfortunately, we do not have a theory that predicts (7). As we shall see below, the speci c self-tuning scenarios we consider lead to additional derivatives of the metric, as well as explicit dependence on the bulk scalar eld, and extra terms which are quadratic in the

¹The anisotropy spectrum in the cosm icm icrow ave background (CMB), although it does provide precision constraints on the expansion of the universe, does not do so in a model-independent way. Since a novel theory of gravity which predicted a di erent Friedmann equation could also predict di erent behavior for the evolution of large-scale density perturbations, it is impossible to compare directly a phenom enological relationship such as (7) to CMB observations. In any speci c full theory, CMB anisotropies are likely to provide an interesting test.

energy and pressure. Generally, it seems unlikely that a theory which did predict a linear dependence of H² on (+p) would both get the correct coe cient 2 G (or very close to it) and successfully recover N ewton's law $F = Gm_1m_2=r^2$ in the solar system. However, this unlikelihood does not seem so great that searching for such a theory would be a waste of time.

3 SpecicExamples

3.1 Setup

W e start with an action of the form [6, 7]

$$S = S_5[;g^{(5)}] + S_4[;g^{(4)};_i]:$$
(8)

Here, S_5 is the bulk action and S_4 that of the brane; is a scalar eld in the bulk with interactions on the brane, while the 's are matter elds con ned to the brane. The metric in the bulk is $g_{ab}^{(5)}$, and we choose coordinates such that the induced metric on the brane is

$$g^{(4)} = {}^{a} {}^{b}g_{ab} :$$
(9)

The indices a; b run over f0;1;2;3; yg and ; run over f0;1;2;3g.

The bulk action can be written

$$S_{5} = {}^{Z} d^{5}x L_{5} = {}^{Z} d^{5}x {}^{q} \overline{g^{(5)}} {}^{M} {}^{3}_{5} R (r)^{2} ; \qquad (10)$$

with a coe cient which we have unspecied for the moment, M $_5$ is the ve-dimensional P lanck mass, R is the ve-dimensional R icci scalar, and the brane action as

$$S_{4} = \overset{Z}{d^{5}} x L_{4} = \overset{Z}{d^{5}} x \overset{q}{\overline{g^{(4)}}} E_{4} \quad (\underline{y}) = \overset{Z}{d^{4}} x \overset{q}{\overline{g^{(4)}}} E_{4} :$$
(11)

We will have the actual form of f_4 (;g⁽⁴⁾; i) unspecied in this section, and consider dierent possibilities in sections 3.2 and 3.3. We will always consider geometries that have a Z_2 symmetry with the brane at the xed point, so that we need not include a boundary term in the brane action.

The 5-dimensional Einstein's equations are

$$G_{ab} = M_5^{3} T_{ab}$$
; (12)

where the energy-m om entum tensor is de ned by

$$T_{ab} = 2q \frac{1}{q^{(5)}} \frac{QL}{Qq^{ab}_{(5)}} :$$
(13)

W e choose the metric

$$ds_{(5)}^{2} = n^{2} (y;t) dt^{2} + a^{2} (y;t) dx^{2} + b^{2} (y;t) dy^{2} ; \qquad (14)$$

where $x = fx^1; x^2; x^3g$ are the spatial coordinates along the brane, and y is the transverse spatial dimension. The Einstein tensor is then

$$G_{00} = 3 \frac{a^2}{a^2} + \frac{ab}{ab} \frac{n^2}{b^2} \frac{a^0}{a} + \frac{a^{0^2}}{a^2} \frac{a^0b^{0^2}}{ab}; \qquad (15)$$

$$G_{0i} = 0 \tag{16}$$

$$G_{0y} = 3 \quad \frac{\underline{a}^{0}}{\underline{a}} + \frac{\underline{a}^{0}\underline{b}}{\underline{a}\underline{b}} + \frac{\underline{a}\underline{n}^{0}}{\underline{a}\underline{n}}; \qquad (17)$$

$$G_{ij} = \frac{a^{2}}{n^{2}} - 2\frac{a}{a} - \frac{b}{b} - \frac{a^{2}}{a^{2}} - 2\frac{ab}{ab} + 2\frac{an}{an} + \frac{bn}{bn}^{!} _{ij} + \frac{a^{2}}{b^{2}} - 2\frac{a^{0}}{a} + \frac{n^{0}}{n} + \frac{a^{0^{2}}}{a^{2}} - 2\frac{a^{0}b^{0}}{ab} + 2\frac{a^{0}n^{0}}{an} - \frac{b^{0}n^{0}}{bn}^{!} _{ij};$$
(18)

$$G_{iy} = 0 \tag{19}$$

$$G_{yy} = 3 \frac{b^2}{n^2} - \frac{a}{a} - \frac{a^2}{a^2} + \frac{an}{an} + \frac{a^{0^2}}{a^2} + \frac{a^0 n^{0^{"}}}{an}$$
 (20)

The energy-momentum tensor decomposes into contributions from the bulk and the brane, 0 = 1

$$T_{ab} = T_{ab}^{(5)} + T_{ab}^{(4)} = 2q \frac{1}{q^{(5)}} \frac{0}{2} \frac{0}{2} \frac{0}{2} \frac{1}{g^{(5)}} + \frac{0}{2} \frac{1}{2} \frac{0}{2} \frac{1}{2} \frac{1}$$

The components of the bulk $T_{ab}^{\ (5)}$ receive contributions only from the scalar $\ eld$, given by

$$T_{00}^{(5)} = n^2 \frac{-2}{n^2} + \frac{-2}{b^2}$$
 (22)

$$T_{0Y}^{(5)} = 2 - {}^{0}$$
(23)

$$T_{ij}^{(5)} = a^2 \frac{2}{n^2} \frac{b^2}{b^2} ij$$
 (24)

$$T_{yy}^{(5)} = b^2 \frac{a^2}{n^2} + \frac{a^2}{b^2}$$
 : (25)

The contribution to the energy-m on entum tensor from the brane will depend on the chosen brane action; in term s of f_4 de ned in (11), we have

$$T_{ab}^{(4)} = \stackrel{V}{\overset{U}{t}} \frac{\overline{g^{(4)}}}{g^{(5)}} \stackrel{0}{\theta} g^{(4)} \stackrel{1}{E}_{4} = 2 \frac{\frac{\theta E_{4}}{\theta g_{(4)}}}{\frac{\theta g_{(4)}}{\theta g_{(4)}}} \quad (y)_{ab} :$$
(26)

In addition to Einstein's equations, we need the equation of motion for $\$. With the action (10)-(11), the equation is

$$r_{a}\frac{@L_{5}}{@(r_{a})} = \frac{@L_{4}}{@}; \qquad (27)$$

since we will be considering brane actions that depend on but not its derivatives. The left-hand side is

$$r_{a}\frac{@L_{5}}{@(r_{a})} = 2^{p}\overline{g_{5}}2; \qquad (28)$$

yielding an equation of motion

$$2 = \frac{1}{2} \frac{1}{b} \frac{0}{6} \frac{1}{2} \frac{1}{b} \frac{0}{6} (y) ; \qquad (29)$$

where ${\rm \mathring{E}}_4$ is de ned by (11) and the D 'A lembertian on scalars is given by

$$2 = \frac{1}{n^2} + \frac{n}{n} + 3\frac{a}{a} + \frac{b}{b} - + \frac{1}{b^2} + \frac{n^0}{n} + 3\frac{a^0}{a} + \frac{b^0}{b} + \frac{a^0}{a} + \frac{a^0}{b} + \frac{$$

The equations of motion generally involve distributional sources localized on the brane. It is therefore convenient to separate the equations into distinct relations valid in the bulk and on the brane. The bulk equations are simply the full equations with distributional sources om itted; they include the bulk E instein equations:

$$3 \frac{a^{2}}{a^{2}} + \frac{ab}{ab} \frac{n^{2}}{b^{2}} \frac{a^{0}}{a} + \frac{a^{0^{2}}}{a^{2}} \frac{a^{0}b^{0}}{ab} = \frac{1}{M_{5}^{3}}n^{2} \frac{a^{2}}{n^{2}} + \frac{b^{2}}{b^{2}}$$
(31)

$$3 \quad \frac{\underline{a}^{0}}{a} + \frac{\underline{a}^{0}}{a\underline{b}} + \frac{\underline{a}\underline{n}^{0}}{a\underline{n}} = \frac{2}{M_{5}^{3}} - {}^{0}$$
(32)

$$\frac{a^{2}}{n^{2}} = 2\frac{a}{a} + \frac{b}{b} + \frac{a^{2}}{a^{2}} = 2\frac{ab}{ab} + 2\frac{an}{an} + \frac{bn}{bn} + \frac{a^{2}}{b^{2}} = 2\frac{a^{0}}{a} + \frac{n^{0}}{n} + \frac{a^{0^{2}}}{a^{2}} = 2\frac{a^{0}b^{0}}{ab} + 2\frac{a^{0}n^{0}}{ap} + \frac{b^{0}n^{0}}{bn} = -\frac{1}{M^{\frac{3}{2}}}a^{2} - \frac{a^{2}}{n^{2}} + \frac{b^{0}}{b^{2}}$$
(33)

and the bulk scalar eld equation

$$\frac{1}{n^{2}} + \frac{n}{n} + 3\frac{a}{a} + \frac{b}{b} - + \frac{1}{b^{2}} + \frac{n^{0}}{n} + 3\frac{a^{0}}{a} + \frac{b^{0}}{b} = 0:$$
(35)

The corresponding equations on the brane can be expressed without explicit -functions by integrating them in the vicinity of the brane to obtain jump conditions [8]. A sum ing Z₂ symmetry, the jump conditions relate second derivatives with respect to y to the coe cients of -function sources. That is, expressions of the form

$$f^{(0)}(y) = A (y) + B$$
 (36)

im ply, on the brane,

$$f_0^0 = \frac{1}{2} A ; (37)$$

where a subscript 0 indicates that a quantity is evaluated on the brane by taking the lim it as y ! 0^+ . For the scalar eld, (29) and (30) in ply

$${}_{0}^{0} = \frac{1}{4} b_{0} \frac{e \dot{E}_{4}}{e} :$$
(38)

For the metric, it is convenient to express the jump conditions for a_0^0 and n_0^0 separately for each model, which we do below.

To get a Friedm ann-like equation on the brane, we consider the yy component of E instein's equations (34), evaluated on the brane. (There is no -function in T_{yy} , so the bulk equation (34) holds true on the brane as well.) We can scale our time coordinate such that $n_0 = 1$, $\underline{n}_0 = 0$ (although n will generally vary o the brane). We then have

$$\frac{a_0}{a_0} + \frac{a_0^2}{a_0^2} = \frac{a_0^{02}}{a_0^2 b_0^2} + \frac{a_0^0 n_0^0}{a_0 b_0^2} - \frac{\frac{0^2}{2}}{3M_5^3} \frac{\frac{0^2}{b_0^2}}{b_0^2} - \frac{3M_5^3}{3M_5^3} - \frac{2}{0} :$$
(39)

The next step is to express the spatial derivatives on the right hand side of (39) in terms of energy and momentum on the brane, which requires a speci cation of the brane Lagrangian L_4 . In the original papers on self-tuning branes [6, 7], the matter Lagrangian was taken to be a cosm ological constant times an exponential of :

$$L_4 = \frac{q}{g^{(4)}}e^2$$
 (y); (40)

with a coupling constant. In what follows we consider two distinct generalizations of this form to actions with dynamical elds: conform alcoupling, in which matter elds on the brane

couple exclusively to a rescaled metric $g = e g^{(4)}$, or volum e element rescaling, in which the four-dimensional volum e element is taken to be $d^4x g^{(4)}e^2$ (and the four-dimensional metric is otherwise simply the induced metric from ve dimensions). Both approaches reduce to (40) when brane tension is the only contribution to the four-dimensional action, but give dimensional for other form s of matter.

3.2 Conform alcoupling

In this section we consider an action of the form

$$S = S_5[;g_{ab}^{(5)}] + S_4[_i;e g^{(4)}];$$
(41)

where the ve-dimensional action is as in (10), and the four-dimensional action is

$$S_{4} = {}^{Z} d^{5}x L_{4} = {}^{Z} d^{5}x {}^{q} - ({}_{i};g) (y) = {}^{Z} d^{4}x {}^{q} - ({}_{i};g) :$$
(42)

Here, $_{i}$ represents an unspecied set of matter elds, and the metric to which matter on the brane couples is related to the induced metric by a conformal transformation,

$$g = e g^{(4)};$$
 (43)

equivalent to $\dot{E}_4 = e^2 f(i;g)$ in the notation de ned by (11).

In this model, test particles on the brane move along geodesics of g; this is the only metric perceived by observers on the brane. It therefore is sensible to de ne all brane quantities in terms of this metric. The energy-momentum tensor as measured by observers living on the brane takes the form

$$\mathbf{\hat{r}} \quad (\mathbf{y}) = 2\mathbf{p} \frac{1}{\mathbf{g}} \frac{\mathbf{\hat{e}} \mathbf{L}_4}{\mathbf{\hat{e}} \mathbf{g}} :$$
(44)

The energy density and pressure as seen by brane observers will be, in our coordinate system,

$$\mathbf{e} = \mathbf{g}^{00} \mathbf{\hat{F}}_{00} \tag{45}$$

and

$$\mathbf{p} = \mathbf{g}^{11} \mathbf{\hat{T}}_{11} ;$$
 (46)

so we have

$$f_{00}^{e} = n^{2}e e$$
 (47)

$$\hat{\mathbf{T}}_{ij} = \mathbf{a}^2 \mathbf{e} \ \mathbf{p}_{ij} : \tag{48}$$

The energy-momentum tensor to which the ve-dimensional Einstein's equations couple is given by (21). Since the brane action depends on g only through g = e g, we can write

$$\frac{@L_4}{@g} = \frac{@g}{@g} \frac{@L_4}{@g} = \frac{1}{2} e^{-\frac{q}{g}} \frac{-}{g} (y) :$$
(49)

Using $p \overline{g} = na^{3}b$ and $p \overline{g} = e^{2} na^{3}$, from (21) we get

$$T_{ab}^{(4)} = e \frac{g}{g} \hat{T}_{ab} (y)$$
$$= \frac{e}{b} \hat{T}_{ab} (y); \qquad (50)$$

or, more explicitly,

$$T_{00}^{(4)} = \frac{n^2}{b}e^2 e^2 (y)$$
 (51)

$$T_{ij}^{(4)} = \frac{a^2}{b}e^2 \mathbf{p} (\mathbf{y})_{ij};$$
 (52)

with other components vanishing.

Our prim ary interest is in the cosm ological equation (39), where we are now in a position to evaluate the rst derivatives of the metric coe cients a and n on the brane. These are derived using the general relation (37) applied to E instein's equations $G_{ab} = M_5^3 T_{ab}$, with G_{ab} given by (15-20) and the relevant components of T_{ab} by (51-52). The jump conditions for the metric components then yield

$$\frac{a_0^0}{a_0} = \frac{1}{6M_5^3} b_0 e^2 \quad {}^{\circ}e ;$$
 (53)

$$\frac{n_0^0}{n_0} = \frac{1}{6M_5^3} b_0 e^2 \quad ^{\circ} (2e + 3e) :$$
 (54)

Finally we need the jump condition for , derived from the equation of motion (29). In the case of conform all coupling the brane source for can be expressed in terms of e and p by using the fact that L₄ depends on only through g to write

$$\frac{\partial L_4}{\partial t} = \frac{\partial g}{\partial t} \frac{\partial L_4}{\partial g} = (g) \frac{1}{2} \frac{1}$$

Putting it together gives

$$\frac{@E^{b}}{@} = \frac{s}{2} \frac{g}{g} (e \quad 3e) (v) :$$
(56)

The equation is thus

$$2 = \frac{e^2}{4 - b} (e^{-3}p) (v);$$
 (57)

where the D'A lem bertian is given by (30). The jump equation for is therefore

$${}_{0}^{0} = \frac{1}{8} b_{0} e^{2} e^{0} (e^{3} e) ;$$
 (58)

Now we plug in the brane equations to (39), yielding

$$\frac{a_0}{a_0} + \frac{a_0^2}{a_0^2} = \frac{1}{576M_5^6} e^{4} \circ 16 + 3 - M_5^3 e^2 + 48 \quad 18 - M_5^3 e^2 + 27 - M_5^3 e^2 \frac{4}{3M_5^3} - \frac{2}{3} e^2$$
(59)

For a pure cosm ological constant on the brane, we have e = p = . Then (59) becomes

$$\frac{a_0}{a_0} + \frac{a_0^2}{a_0^2} = \frac{1}{36M_5^6} e^4 \circ 2 \cdot 3 - M_5^2 \cdot 2 \cdot \frac{3}{3M_5^3} - \frac{2}{3M_5^3} - \frac{2}{3M_5^3} \cdot \frac{2}{3M_5^3}$$
(60)

We see that this can vanish for any value of $f_{-0} = 0$ and we choose

$$\frac{2}{-M} M_5^3 = \frac{2}{3}$$
 (61)

This is the condition for self-tuning, as derived by [6, 7]. (Of course, making this choice is a kind of ne-tuning, as we discuss brie y in the Appendix.)

In this case, our cosm obgical equation (59) for m ore general m atter sources becom es

$$\frac{a_0}{a_0} + \frac{a_0^2}{a_0^2} = \frac{1}{32M_5^6} e^4 \quad {}^{\circ} (e + p)^2 \quad \frac{1}{2} \quad {}^{2} - \frac{2}{0} :$$
(62)

If the brane tension is , we can decompose the energy density and pressure into tension plus dynamical energy-momentum (e.g. from matter and radiation on the brane): $e = + d_{yn}$, $p = + p_{dyn}$. From (62), however, the tension simply cancels out, and we are left with an identical equation relating for the dynamical density/pressure as we have for the total density/pressure. A sthis equation is quadratic rather than linear in the energy density, it will not yield anything close to conventional Friedm ann behavior, and can't be taken seriously as a description of the real world. (W e note that matter on the brane does not respond directly to the scale factor a, but to the conform ally-transform ed scale factor; how ever, for constant

the functional dependence on the energy density will still be quadratic.)

3.3 Volum e-elem ent coupling

In this section we consider a model in which enters only in an overall factor multiplying the Lagrange density, not in every appearance of the metric; in the notation of (11) we have

$$\mathbf{E}_{4}^{b} = \mathbf{e}^{2} \quad \mathbf{f}(\mathbf{g}^{(4)}; \mathbf{j}) :$$
 (63)

This is equivalent to modifying the four-dimensional spacetime volume element:

$$d^{4}x \stackrel{q}{\overline{g^{(4)}}} ! \quad d^{4}x \stackrel{q}{\overline{je}} \quad g^{(4)} = d^{4}x e^{2} \stackrel{q}{\overline{g^{(4)}}};$$
(64)

while otherwise coupling to the induced metric g⁽⁴⁾. Unlike the example of conformal coupling, this form of the action can be disrupted by quantum corrections; on the other hand, we shall see that the volum e-coupling ansatz leads to a som ew hat more acceptable cosm ological model, and is worth exploring for that reason.

Since now couples non-universally to matter, it is most sensible to think of g as the metric to which matter responds, and the coupling to e as an interaction. The energy-momentum tensor therefore takes the conventional form

$$\Gamma = 2q \frac{1}{q^{(4)}} g^{(4)} \frac{\partial L_4}{\partial g^{(4)}} = diag(;p;p;p):$$
(65)

This is related to $T_{ab}^{(4)}$, the brane contribution to the energy-m on entum tensor appearing in the ve-dimensional Einstein equations, by

77

$$T_{ab}^{(4)} = 2q \frac{1}{q^{(5)}} \frac{\varrho_{L_4}}{\varrho_{g^{(5)}ab}} = t \frac{\psi_{u}}{g^{(6)}} \frac{\overline{g^{(4)}}}{\overline{g^{(5)}}} T_{ab} :$$
(66)

The specic components of $T_{ab}^{(4)}$ are thus:

$$T_{00}^{(4)} = \frac{n^2}{b} \quad (y) \tag{67}$$

$$T_{ij}^{(4)} = \frac{a^2}{b} p (y)_{ij}$$
: (68)

These di er from (51)–(52) of the previous section by the replacement ($e^2 e;e^2 p$)! (;p). As a consequence, the jump conditions for the metric coe cients are

$$\frac{a_0^0}{a_0} = \frac{1}{6M_5^3} b_0 ; \qquad (69)$$

$$\frac{n_0^0}{n_0} = \frac{1}{6M_5^3} b_0 (2 + 3p) :$$
 (70)

To express the jump condition for in terms of and p, we need to calculate $\mathbb{C}_4=\mathbb{C}$. We can do this by considering our four-dimensional action to be that of a perfect uid, corresponding to

$$\dot{E}_4 = e^2 f(g^{(4)}; i) = p(;s) \frac{n}{2} g^{(4)} + i^2$$
 (71)

(see [11, 13] for a discussion). In (71), the dynamical degrees of freedom with respect to which we vary the action to obtain equations of motion include the entropy s, the enthalpy

, a Lagrange multiplier n, the three C lebsch potentials $\ , \ , \ ,$ and the therm asy $\ ;$ these last four scalars de ne the four-vector $\$ via

$$= 0 + 0 + 0s:$$
 (72)

Note that varying with respect to n gives the constraint $+^{2} = 0$; hence, on-shell the Lagrange density is simply equal to the pressure p. Therefore, from (29) the equation of motion for is

$$2 = \frac{1}{2} \frac{1}{b} \frac{0}{6} \frac{b_4}{6} \quad (y)$$
(73)

$$= -\frac{1}{b}p(y):$$
(74)

This corresponds to a jump condition describing the behavior of in the vicinity of the brane,

$${}_{0}^{0} = \frac{1}{2} b_{0} p :$$
 (75)

Now we plug in these jump conditions to (39), yielding

$$\frac{a_0}{a_0} + \frac{a_0^2}{a_0^2} = \frac{1}{36M_5^6} + 3p + 3p + 3m_5^2 + 3p^2 + 3p^2 + 3m_5^3 - \frac{2}{3} + 3m_5^3$$

W e see that the self-tuning condition is satis ed again only for

$$-\frac{2}{M} \frac{3}{5} = \frac{2}{3};$$
 (77)

in which case our cosm ological equation (76) for more general matter sources becomes

$$\frac{a_0}{a_0} + \frac{a_0^2}{a_0^2} = \frac{1}{36M_5^6} (+p) (+2p) \frac{1}{2} \frac{2}{-0}^2 :$$
(78)

The contribution of the brane tension to the Lagrangian enters in the combination $e^2 \circ . W$ e therefore decompose the energy density and pressure into tension plus dynam ical sources in the following manner:

$$= e^{2} + dyn;$$
 (79)

$$p = e^{2} + p_{dyn}$$
 : (80)

Then, unlike the cosm ological equation (62) obtained for conform al coupling, in the case of volum e-element coupling we obtain terms on the right-hand side which are linear in the dynam ical energy density and pressure:

$$\frac{a_0}{a_0} + \frac{a_0^2}{a_0^2} = \frac{1}{36M_5^6} e^2 \circ (a_{yn} + p_{dyn}) \circ (a_{yn}^2 + 3 a_{yn} p_{dyn} + 2p_{dyn}^2)^i \cdot \frac{1}{2} e^2 - \frac{2}{0} :$$
(81)

An equivalent equation was derived by M ennim and Battye [11].

A n equation of this type stands a chance of describing the real world. The term squadratic in energy/m on entum can presum ably be neglected at late times; if we assume that the $\frac{2}{0}$ term is negligible we are left with a relation which has some resemblance to the hoped-for equation (7). With this in mind, we turn now to comparison with observation.

4 D iscussion

Let us consider whether, under favorable circum stances, (81) could be consistent with what we know about the universe.

A time-dependent bulk scalar would generally lead to observable time-dependence in the four-dimensional Newton's constant, and therefore must be very small. A lthough the models under consideration do not include any mechanism for stabilizing , we will proceed optim istically and imagine that the bulk scalarm ay be approximated as independent of time,

$$(y;t) = (y) :$$
 (82)

W e in agine further that the y-dependence of the metric coe cients may be factored out, and the coe cient b^2 of dy^2 set to unity by an appropriate rescaling (which may always be done if b^2 is independent of t):

$$ds^{2} = ! (y) [dt^{2} + a_{0}^{2} (t) dx^{2}] + dy^{2} :$$
(83)

Referring back to the jump conditions (70) and (75), such ansatze do not seem unreasonable; in (70) and (75) the symbols and prefer to the entire energy and pressure, which will be dom inated by the brane tension term, which in turn remains constant. Thus, it is reasonable to approximate the bulk solutions for and ! by on their at-space values as derived in [6, 7],

$$(\mathbf{y}) = {}_{0} \frac{1}{2} \ln [! (\mathbf{y})]$$
(84)

and

$$! (y) = \frac{q}{1 \quad y = y_c};$$
(85)

where y_c represents the location of the boundary opposite our brane, corresponding to a singularity,

In order to compare the modi ed Friedmann relation (81) to observations, it is necessary to express the four-dimensional Planck mass $M_4 = 1 = \frac{p}{8} \frac{1}{6} \frac{1}{$

$$M_{4}^{2} = M_{5}^{3} \int_{0}^{Z_{y_{c}}} dy ! (y) = \frac{M_{5}^{6}}{2} e^{2}$$
 (87)

(In fact, we are again ignoring a subtlety: due to the presence of the scalar eld, general relativity is not exactly recovered in four dimensions, and the de nition of G has to be specified more carefully. It is possible that solar-system tests of gravity would rule out these simple models more de nitively than the cosm ological scenario considered here.)

For simplicity we drop the subscript \dyn" from and p for the remainder of this section. In terms of the Hubble parameter $H = \underline{a}=a$, we have $a=a + (\underline{a}=a)^2 = H_+ 2H^2$. Then setting $-_0 = 0$ and ignoring quadratic terms in ;p, our cosm obgical equation (81) becomes

$$H_{-} + 2H^{2} = \frac{2}{9} \frac{G}{9} (+p)$$
 : (88)

Unlike the ordinary Friedmann equation, which is a constraint relating the value of the Hubble parameter to the energy density, this is a dimensional equation for H, which will involve an additional integration constant in its solution. In fact we can integrate (88) explicitly to obtain

$$H^{2} = \frac{4}{9} a^{4} a^{3} (+p) da :$$
 (89)

Consider a universe dom inated by a combination of matter ($_M$ / a 3 , p_M = 0) and radiation ($_R$ / a 4 , p_R = $_R$ =3). The Hubble parameter then obeys

$$H^{2} = \frac{4}{9} \frac{G}{M} + \frac{16}{27} \frac{G}{R} \ln \frac{a}{a} ; \qquad (90)$$

where a is the integration constant alluded to above.

It is this relation (90) which is to be compared to the conventional Friedmann law $H^2 = (8 G = 3)$. In a matter-dom insted era, the functional dependence of H on a is conventional, leading to the fam iliar a / t^{2-3} behavior. The coe cient of , however, is di erent, suggesting a possible em pirical test of the model: in this theory, the Hubble parameter during matter dom ination will be smaller by a factor of 1=60:41 than the conventional expectation (at xed). Unfortunately, we do not have very precise em pirical information about the expansion rate during the matter-dominated era. Given that our current universe is apparently dom inated by a smooth component causing it to accelerate, we cannot directly constrain the coe cient of $_{\rm M}$ by contemporary observations. Since recombination occurred during the matter-dom inated era, it would be possible in principle to constrain this coe cient via observations of the CMB an isotropy power spectrum; it is rst necessary, however, to reliably calculate what that power spectrum should be, taking into account possible long-range deviations from general relativity due to the extra dimension. Since we have not undertaken this task, we are unable to say whether $H^2 = (4 G = 9)_{M}$ is consistent with the real universe.

In the radiation-dom inated era, quantitative constraints on the behavior of the Hubble parameter m ay be derived from Big-Bang Nucleosynthesis (BBN) [12]. Expanding the logarithm in (90), we have

$$H_{R}^{2} = \frac{16 G}{27} R \ln a + C a^{4}; \qquad (91)$$

where C is an integration constant. This constant is not merely a nuisance that can be set to zero, but rather relects the normalization of a; if we set a = 1 today, the rst term is always negative in the past, which is clearly unworkable. On the other hand, we can choose C such that the second term is dominant | and with the correct magnitude to be compatible with observation | during BBN. This seems a like an unlikely bit of ne-tuning, although it cannot be rigorously excluded. Therefore we see no way to de nitively state that the self-tuning cosm ologies we have considered are incompatible with observation, although it would require a mysterious coincidence to predict the correct light-elem ent abundances. Since these m odels were invented to solve the ne-tuning problem associated with the cosm ological constant, this must be considered as a strike against them.

Nevertheless, it is important to keep in m ind that we have only dealt with speci c toy m odels of self-tuning, which perhaps it is too optim istic to expect would lead to completely realistic cosm ologies. The general idea that the cosm ological constant problem m ay be solved not by m aking the vacuum energy sm all, but by m aking the metric insensitive to its value, is

an interesting one, and it seems worth the e ort to attempt to construct self-tuning models with more acceptable cosmological behavior.

A cknow ledgm ents

W e would like to thank Pierre Binetruy, Gregory G abadadze, Steve Giddings, M onica Guica, Sham it K achru, N em anja K aloper, Finn Larsen, Joe Lykken, Ricardo R attazzi, E va Silverstein, and D avid W ands for useful conversations. This work was supported in part by the U.S.Dept. of Energy, the A lfred P.Sloan Foundation, and the D avid and Lucile Packard Foundation.

Appendix: Two observations

In this Appendix we very brie y mention two issues of som ewhat related interest to this work: rst, the possibility of in ationary behavior in self-tuning cosm ologies, and second, the ability to tune away other equations of state.

An obvious question which arises in any model which would make the metric insensitive to vacuum energy is, how are we to explain the apparent nonzero value of the cosm ological constant today [3, 4], or implement an in ationary scenario in the early universe? O focurse if a realistic self-tuning theory is found, it is conceivable that the currently observed vacuum energy is simply a relation of an imperfectly tuned universe, once all aspects of the theory are taken into account. On the other hand, it is interesting to note that there is no di culty in obtaining \accelerating" solutions in the presence of slow ly-rolling scalar leds. Consider a minimally coupled four-dimensional scalar , with potential V (). The energy density and pressure are given by

$$= \frac{1}{2} - \frac{2}{2} + V(); \quad p = \frac{1}{2} - \frac{2}{2} + V(); \quad (92)$$

or

$$+ p = -2^{2}$$
: (93)

For a slow ly-rolling scalar with — const, (89) then in plies H const, just as in conventional theory with a nonzero vacuum energy. Thus there is no obstacle in principle to obtaining accelerated expansion either today or in the early universe. Of course the usual tuning problem s associated with getting the correct nonzero value of the apparent vacuum energy are as severe in such a hypothetical model as they are in conventional quintessence and in ation scenarios.

A nother interesting issue is the possibility of choosing parameters which work to tune away the e ects of a more general energy component with equation of state p = w, where w is not necessarily 1. We know of no compelling reason why this should happen, but the exercise illustrates the extent to which there really is some tuning going on in our choice of parameters.

Vacuum energy was tuned away by choosing our parameters and to be related by (77) (taking the case of volum e-element coupling for de niteness). Let us in agine that we instead take

$$-\frac{2}{M} \int_{5}^{3} = \frac{2}{3} \mathbf{x};$$
 (94)

where x is a parameter to be chosen. The cosm ological equation (76) becomes (setting -0 = 0 for simplicity),

$$\frac{a_0}{a_0} + \frac{a_0^2}{a_0^2} = \frac{1}{32M_5^6} + 3p + 2xp^2 :$$
(95)

For a universe dom instead by a component with p = w, the right hand side will autom atically vanish if we choose

$$x = \frac{1+3w}{2w^2} :$$
 (96)

The choice x = 1 tunes away vacuum energy (w = 1) as well as an exotic component with p = -2. With the exception of matter (w = 0), the expansion can be made insensitive to any speci c equation of state by an appropriate choice of x. Once again, we have no reason to suggest that such a possibility will occur (nor see any need for it); how ever, this phenom enon serves as an illustration that the speci c choice (77) represents a kind of ne-tuning in its own right.

References

- [1] S.W einberg, \The Cosm ological Constant Problem, "Rev.Mod.Phys. 61, 1 (1989).
- [2] E.W itten, \The cosm ological constant from the viewpoint of string theory," hepph/0002297.
- [3] S.M. Carroll, \The cosm ological constant," Living Reviews in Relativity, 2001-1 (2001) [astro-ph/0004075].

- [4] A.G.Riess et al. [Supernova Search Team Collaboration], Astron.J.116, 1009 (1998)
 [astro-ph/9805201]; S.Perlm utter et al. [Supernova Cosm ology Project Collaboration],
 Astrophys.J.517, 565 (1999) [astro-ph/9812133].
- [5] N. Arkani+Hamed, S. D in opoulos and G. Dvali, Phys. Lett. B 429, 263 (1998) [hep-ph/9803315]; I. Antoniadis, N. Arkani+Hamed, S. D in opoulos and G. Dvali, New dimensions at a millimeter to a Fermi and superstrings at a TeV, "Phys. Lett. B 436, 257 (1998) [hep-ph/9804398]; K. R. D ienes, E. Dudas and T. G herghetta, Phys. Lett. B 436, 55 (1998) [hep-ph/9803466]; L. Randall and R. Sundrum, \An alternative to compacti cation," Phys. Rev. Lett. 83, 4690 (1999) [hep-th/9906064].
- [6] N. Arkani-Hamed, S. Dimopoulos, N. Kaloper and R. Sundrum, \A small cosmological constant from a large extra dimension," Phys. Lett. B 480, 193 (2000) [hepth/0001197].
- [7] S.Kachru, M. Schulz and E. Silverstein, \Selftuning at dom ain walls in 5d gravity and string theory," Phys. Rev. D 62, 045021 (2000) [hep-th/0001206]; S.Kachru, M. Schulz and E. Silverstein, \Bounds on curved dom ain walls in 5d gravity," hep-th/0002121.
- [8] P.Binetruy, C.De ayet and D.Langlois, \N on-conventional cosm ology from a braneuniverse," Nucl. Phys. B 565, 269 (2000) [hep-th/9905012].
- [9] H.A. Cham blin and H.S.Reall, \D ynam ic dilatonic dom ain walls," Nucl. Phys.B 562, 133 (1999) [hep-th/9903225]; C.C saki, M.G raesser, C.K olda and J.Teming, \Cosmology of one extra dimension with localized gravity," Phys. Lett. B 462, 34 (1999) [hep-ph/9906513]; J.M. Cline, C.G rojean and G.Servant, \Cosmological expansion in the presence of extra dimensions," Phys.Rev.Lett.83, 4245 (1999) [hep-ph/9906523]; P.K anti, I.I.K ogan, K.A.O live and M.Pospelov, \Cosmological 3-brane solutions," Phys. Lett. B 468, 31 (1999) [hep-ph/9909481]; L.M ersini, \D ecaying cosmological constant of the in a ting branes in the R andall-Sundrum -O dam odel," hep-ph/9909494; P.B inetruy, C.D e ayet, U.Ellw anger and D.Langbis, \B rane cosmological evolution in a bulk with cosmological constant," Phys.Lett.B 477, 285 (2000) [hep-th/9910219]; P.K raus, \D ynam ics of anti-de Sitter dom ain walls," JHEP9912, 011 (1999) [hep-th/9910149]; T.Shirom izu, K.M aeda and M.Sasaki, \The Einstein equations on the 3-brane world," Phys.Rev.D 62, 024012 (2000) [gr-qc/9910076]; E.E.F lanagan, S.H.Tye and I.W asserm an, \Cosmological expansion in the R andall-Sundrum

brane world scenario," Phys. Rev. D 62, 044039 (2000) [hep-ph/9910498]; P. Kanti, I. I. Kogan, K. A. O live and M. Pospelov, \Single-brane cosm ological solutions with a stable com pact extra dimension," Phys. Rev. D 61, 106004 (2000) [hep-ph/9912266]; L. Mersini, \Radion potential and brane dynamics," hep-ph/0001017.

- [10] S. Forste, Z. Lalak, S. Lavignac and H. P. Nilles \A comment on self-tuning and vanishing cosm ological constant in the brane world" Phys. Lett. B481, 360 (2000) [hep-th/0002164]; C.Csaki, J.Erlich, C.Grojean and T.Hollowood, \General properties of the self-tuning domain wall approach to the cosm ological constant problem," Nucl. Phys. B 584, 359 (2000) [hep-th/0004133]; G. T. Horowitz, I. Low and A. Zee, \Self-tuning in an outgoing brane wave model," hep-th/0004206; B. Grinstein, D.R.Nolte and W.Skiba, \Adding matter to Poincare invariant branes," hepth/0005001; C. Zhu, \A self-tuning exact solution and the non-existence of horizons in in 5d gravity-scalar system s," JHEP0006,034 (2000) [hep-th/0005230]; V.Barger, T.Han, T.Li, J.D.Lykken and D.Marfatia, \Cosmology and hierarchy in stabilized warped brane m odels," Phys. Lett. B 488, 97 (2000) [hep-ph/0006275]; P. B inetruy, J.M. Cline and C.G ropan, D ynam ical instability of brane solutions with a self-tuning $\cos m o \log calconstant," hep-th/0007029; K . M aeda and D . W ands, \D ilaton-gravity on$ the brane," hep-th/0008188; L.E.M endes and A.M azum dar, \B rans-D icke brane cosmology," gr-qc/0009017; Z.Kakushadze, \'Self-tuning' and conform ality," M od. Phys. Lett. A 15, 1879 (2000) [hep-th/0009199]; C.K ennedy and E.M. Prodanov \Standard cosm ology on a self-tuning dom ain wall," hep-th/0010202; A.Kehagias and K.Tam vakis, \A self-tuning solution of the cosm ological constant problem ," hep-th/0011006; J.E.Kim, B.Kyae and H.M.Lee \A model for self-tuning the cosm ological constant," hep-th/0011118; T. Padm anabhan and S. Shankaranarayanan, \Vanishing of cosm ological constant in nonfactorizable geometry," Phys. Rev. D 63, 105021 (2001) hep-th/0011159]; J.E.K in , B.K yae and H.M. Lee \Self-Tuning Solution of the cosmological constant problem with antisymmetric tensor eld," hep-th/0101027; P.Brax and A.C.D avis, \Cosm ological evolution on self-tuned branes and the cosm ological constant," hep-th/0104023.
- [11] A.Mennim and R.A.Battye, \Cosmological expansion on a dilatonic brane-world," hep-th/0008192.
- [12] K.A.Olive, G. Steigman and T.P.Walker, \Primordial nucleosynthesis: Theory and observations," Phys. Rept. 333-334, 389 (2000) [astro-ph/9905320]; S.Burles,

K.M.Nollett, J.N.Truran and M.S.Turner, \Sharpening the predictions of big-bang nucleosynthesis," Phys.Rev.Lett.82, 4176 (1999) [astro-ph/9901157]; D.Tytler, JM. O'M eara, N.Suzuki and D.Lubin, astro-ph/0001318.

[13] D.S.Salopek and J.M. Stewart, Class. Quant. Grav. 9, 1943 (1992).