

A Brief Comment on “The Pyrotechnic Universe”³

Justin Khoury¹, Burt A. Ovrut², Paul J. Steinhardt¹ and Neil Turok³

¹ Joseph Henry Laboratories, Princeton University, Princeton, NJ 08544, USA

² Department of Physics, University of Pennsylvania, Philadelphia, PA 19104-6396, USA

³ DAMTP, CMS, Wilberforce Road, Cambridge, CB3 0WA, UK

We respond to the criticisms by Kallosh, Kofman and Linde concerning our proposal of the ekpyrotic universe scenario. We point out a number of errors in their considerations and argue that, at this stage, the ekpyrotic model is a possible alternative to inflationary cosmology as a description of the very early universe.

We have recently proposed an alternative to inflationary cosmology,^{1,2} entitled the “ekpyrotic universe,”³ designed to resolve the horizon, flatness, and monopole puzzles of standard hot big bang cosmology and to generate a nearly scale-invariant spectrum of adiabatic energy density perturbations needed to seed structure formation. The basic notion, motivated by string theory and M-theory,^{4,5} is that the hot big bang universe is produced by the collision in an extra-dimensional space-time between a three-dimensional brane in the bulk space with another brane or a bounding orbifold plane. The collision heats the universe to a high but finite temperature, from which point the hot big bang phase begins. Whereas the inflationary scenario relies on an extended period of exponential hyperexpansion prior to the hot big bang phase, the ekpyrotic model relies on extremely slow evolution over an exceedingly long time.

In a recent preprint,⁶ Kallosh, Linde, and Kofman (KKL) raised a number of criticisms of the ekpyrotic universe. In this comment we briefly explain why we respectfully disagree with each of their seminal points (italics, below). Where a full response requires a technical calculation, the details are given in the separate publications to which we refer.

We first consider the criticisms of the superstring and M-theoretic underpinnings. We then discuss the description of density perturbations in our scenario and finally, the criticisms of the initial conditions.

KKL: One of the central points of the ekpyrotic cosmological scenario is that we live on a negative tension brane. However, the tension of the visible brane in Hořava-Witten theory, as well as in relevant cases of non-standard embedding, is positive. Hence, there is a problem in the assignment of signs to the brane tensions.

This criticism is factually incorrect. The claim is based on experience with the standard embedding and other specific examples that have appeared in the literature previously, rather than on mathematical analysis. The fact is that there never was a cohomology requirement that the visible brane have positive tension to obtain realistic models. We have constructed numerous examples⁷ of heterotic M-theory models in which the visible brane has negative tension and is endowed with a much smaller

gauge group (*e.g.*, SU(5)) than that of the positive tension hidden brane (*e.g.*, E_7).

It is the standard embedding that motivated the minor variant suggested by Kallosh, *et al.*, dubbed by them the “pyrotechnic” universe. Ironically, this particular choice is disallowed since bulk branes are mathematically forbidden, and so there could no brane collision to ignite the hot big bang phase.

We should also emphasize that, contrary to KKL’s assertion, we never claimed that the assignment of negative tension to the visible brane is a “central point” of the ekpyrotic concept. We did not intend our proposal to be interpreted so narrowly. It is a feature of our specific example, and Ref. 7 demonstrates that this example is possible. For this example, we explained the role that negative tension plays in the energetics, fluctuations, and ultimate expansion of the universe. But, we also stated that the general principles could be adapted to numerous set-ups.³ Indeed, in Refs. 3 and 11, we derive the perturbation spectrum in a 4d static background in a limit which is totally insensitive to the assignment of brane tensions. The Kallosh *et al.* example, which is based on the same physical principles and mathematical equations, but entails flipping a few signs in the action, has no substantive differences to our original scenario (provided the embedding problem is fixed).

Kallosh *et al.* not only criticize the choice of sign for the tensions, but also the magnitudes:

KKL: In examples considered in the literature, the contribution to the cohomology constraint from the bulk brane is of the same order as the one from the boundaries, whereas the ratio is extremely small (4×10^{-5}) in the ekpyrotic model. More precisely, the claim is that the ratio of the bulk tension β to the boundary tension α in our example is incompatible for realistic models with what is required by the cohomology constraint.

This criticism again rests on examples of Hořava-Witten models in the literature rather than on mathematical analysis. In Ref. 7, it is shown that there is a wide range of freedom for β/α and that the value in our example is mathematically consistent with the cohomology constraint.

Furthermore, we did not intend our proposal to be in-

terpreted so narrowly. Let us not confuse an example with general principles. We never claimed that a small ratio is required. Indeed, viable and observationally consistent examples with ratios greater than $1/10$ are discussed in Ref. 3.

Other criticisms about the string-theoretic underpinnings include:

- *questioning the presence of the 4-form in the heterotic M-theory action:* The 4-form formulation of the action is equivalent to the action presented in Ref. (5). This is easily seen by eliminating the 4-form using its equation of motion. This formulation is particularly useful in heterotic vacua with bulk branes, such as in ekpyrotic cosmology.
- *questioning the origin of the bulk 3-brane:* The 3-brane is simply an M5 brane wrapped on a holomorphic curve of the Calabi-Yau threefold.
- *stabilization of moduli:* Granted, this is a deep, unsolved problem of string theory. We are presuming its solution does not interfere with the ekpyrotic scenario. The same presumption must be made in inflationary theory. Indeed, in this case, there are well-documented difficulties that arise with inflation if moduli are not stabilized prior to inflation (which may prevent inflation occurring⁸), difficulties which are not applicable to the ekpyrotic model.

In addition, Kallosh *et al.* criticize the sign of the potential, its parameters and the notion that the potential is zero after collision. Granted, assuming the potential to be zero after collision is fine-tuning. However, let us recognize it for what it is – the well-known cosmological constant problem. Precisely the same fine-tuning of the true vacuum energy is required in inflationary cosmology.

As for criticisms about tuning of parameters, this skepticism is based largely on a search of the current literature, all of which was written prior to the appearance of the ekpyrotic proposal, and on the one example in our paper. There is no serious physics argument, or mathematical analysis, or analytical discussion of what range of parameters is required or what emerges naturally from string theory. Indeed, our choice of parameters and potential appears to us to be well within the bounds one might expect from string theory. However, to make their point, Kallosh *et al.* reparameterize the action in variables that are not natural to string theory and then argue that the potential is finely-tuned. This seems a strange and unnecessary procedure.

More generally, any conclusions about parameters are premature since there has been no serious analysis of what range is required other than the point made in our paper Ref. 3 that there appears to be a great deal of flexibility. Perhaps some historical perspective is useful. A

similar search of the literature prior to first computations of density perturbations in 1982 would reveal no examples of slow-roll potentials with dimensionless couplings of order 10^{-14} . History shows that inflationary cosmology stimulated the search for these models, and now many examples are found in the literature. We would suggest that some patient, serious analysis is required before the ekpyrotic scenario can be judged against inflation in this respect.

KKL: The mechanism for the generation of density perturbations in this scenario is a particular limiting case of the mechanism of tachyonic preheating.

The basic notion of the ekpyrotic model is that the universe begins in a quasi-static (non-expanding) state, a concept that dates back to ancient philosophy. In pursuing this idea, a basic challenge is to generate fluctuations which are in accord with the impressive and precise measurements of the cosmic microwave background and large-scale structure which lend strong support for a nearly scale-invariant spectrum of linear, adiabatic density perturbations. Our paper shows that such perturbations are generated in a quasi-static multi-brane universe for certain simple potentials including negative exponentials and inverse power laws.

Tachyonic preheating,⁹ a concept introduced by Kofman, Linde and collaborators, concerns phenomena not directly related to the generation of large scale density perturbations, in a rapidly expanding universe just following inflation. It is legitimate within its own context. But the potentials needed for scale invariance do not appear anywhere in the tachyonic preheating paper, and we think that it obfuscates rather than clarifies the issue to conflate our fluctuation generation mechanism with tachyonic pre-heating. There is no more relation to tachyonic preheating than there is to fluctuation generation in inflation. In both cases, there are some common elements but also major differences.

KKL: Inflation removes all previously existing inhomogeneities and is robust, whereas the ekpyrotic model makes the homogeneity problem much worse.

The critique refers to the fact that inflation is a mechanism that can make an inhomogeneous universe more homogeneous, whereas homogeneity in the ekpyrotic model is part of the initial condition. Behind this critique lies a substantive point concerning the different assumptions of the inflationary and ekpyrotic scenarios. However, a more objective comparison is warranted.

Inflation assumes that the universe begins in a high energy state of no particular symmetry that is rapidly expanding from the start. For such general initial conditions, it is essential that there be a dynamical attractor mechanism that makes the universe more homogeneous as expansion proceeds. Superluminal expansion provides that mechanism. The ekpyrotic model, on the other hand, is built on the principle that the initial state is quasi-static with properties dictated by symmetry, par-

ticularly supersymmetry. So, by construction, the initial state is special, but special in a way that is well-motivated physically. In this case, while a dynamical attractor mechanism may be possible (see below), it is not essential. One can envisage the possibility that the initial conditions are simply the result of some selection rule that dictates a state of maximal supersymmetry and low energy.

Within the context of superstring theory and M-theory, the natural choice of initial state is the BPS (Bogolmon'yi-Prasad-Sommerfeld) state.⁵ The BPS property is already required from particle physics in order to have a low-energy, four-dimensional effective action with $\mathcal{N} = 1$ supersymmetry, necessary for a realistic phenomenology. For our purposes, the BPS state is ideal because, not only is it homogeneous, as one might suppose, but it is also flat. That is, the BPS condition links curvature and homogeneity. It requires the two boundary branes to be parallel.

We do not rule out the possibility of a dynamical attractor mechanism that drives the universe towards the BPS state from some more general initial condition. Such parallelism would be a natural consequence of all the branes emerging from one parent brane.¹² Alternatively, one can imagine beginning with two boundary branes only and no bulk-brane or interbrane potential. Perhaps curvature can be dissipated by radiating excitations tangential to the branes and having them travel off to infinity. The branes might settle into a BPS ground state until, through some rare process, a bulk brane is nucleated. Depending on the gauge structure on the branes, the bulk brane may be drawn towards the visible brane.

Beginning in an empty, quasi-static state solves some problems, but one should not underestimate the remaining challenges: how to generate a hot universe, and how to generate perturbations required for large-scale structure. The remarkable feature of the ekpyrotic picture is that brane collision naturally serves both roles.

Inflationary cosmology, based upon a powerful attractor mechanism, is very appealing philosophically, and we do not seek to discredit it by proposing an alternative. But, we would hesitate to characterize inflation as “robust.” If robustness means that a Universe such as the one we observe is *nearly inevitable* in the context of an inflationary model, one must disagree. The presence of large initial inhomogeneities and black holes can always prevent inflation from occurring. There is no rigorous or even postulated measure of the basin of attraction. Claims that inflation works for generic chaotic initial conditions are simply ill-defined.

There are other longstanding, unresolved issues. A successful inflationary scenario requires setting couplings of the inflaton to itself and to other fields at levels more than 12 orders of magnitude below the natural expectation.¹³ Furthermore, inflation is far from being a rigid theory. While flatness is certainly the simplest possi-

bility, at least one of the authors of⁶ has accepted that open universes¹⁴ are also possible.¹⁵ Rigorous methods for deciding the relative probability of patches with different physical properties (*e.g.*, open versus flat) are not known.¹⁶ Finally, inflation has not yet been embedded in a theory of quantum gravity in a consistent manner. Inflationary quantum fluctuations emanate from exponentially sub-Planckian scales, and the physics on such scales is very incompletely understood. All of these features suggest that it may be something of an exaggeration to describe inflation as “robust” at this stage.

We do not raise these points to argue that inflation is flawed or should be discarded. On the contrary, inflation is an outstandingly successful and appealing idea. However, if one is comparing the inflationary and ekpyrotic approaches to the homogeneity and flatness problems, it is sensible to maintain a fair and open-minded attitude, and to present the features and unsettled issues in both cases as objectively as possible.

We see no reason to rush to judgment. Inflationary theory is based on quantum field theory, a well-established theoretical framework. The model took several years to develop, and it has been carefully studied and vetted for twenty years. It is still in many respects incomplete. Our proposal is based on unproven ideas in string theory and entails many features which are new. A sober examination of the ekpyrotic approach will prove useful whichever way it turns out. If the model is found to be theoretically inconsistent, then our confidence in the inflationary solution will be enhanced. If the model proves to be a legitimate alternative, then there is a new possibility to consider and nature’s choice will be decided observationally by measurements of the gravitational wave background.³

-
- [1] A. H. Guth, *Phys. Rev. D* **23**, 347, (1981).
 - [2] A. D. Linde, *Phys. Lett.* **108B**, 389 (1982); A. Albrecht and P. J. Steinhardt, *Phys. Rev. Lett.* **48**, 1220 (1982).
 - [3] J. Khoury, B.A. Ovrut, P.J. Steinhardt and N. Turok, hep-th/0103239.
 - [4] P. Hořava and E. Witten, *Nucl. Phys.* **B460** (1996) 506; **B475** (1996) 94.
 - [5] A. Lukas, B.A. Ovrut and D. Waldram, *Nucl. Phys.* **B532** (1998) 43; *Phys. Rev. D* **57** (1998) 7529; A. Lukas, B.A. Ovrut, K.S. Stelle and D. Waldram, *Phys. Rev. D* **59** 086001 (1999).
 - [6] R. Kallosh, L. Kofman, and A. Linde, hep-th/0104073.
 - [7] R. Donagi, J. Khoury, B.A. Ovrut, P.J. Steinhardt and N. Turok, in preparation.
 - [8] R. Brustein and P. J. Steinhardt, *Phys. Lett. B* **302**, 196 (1993); T. Banks, M. Berkooz, G. Moore, S. H. Shenker,

- P. J. Steinhardt, *Phys.Rev. D***52**, 3548 (1995); G. Huey, B.A. Ovrut, P.J. Steinhardt and D. Waldram, *Phys.Lett. B***476**(2000) 379-386, (2000).
- [9] G. Felder, J. Garcia-Bellido, P.B. Greene, L. Kofman, A. Linde, and I. Tkachev, hep-ph/0012142.
- [10] J.M. Bardeen, P.J. Steinhardt and M.S. Turner, *Phys. Rev. D* **28** (1983) 679; A.H. Guth and S.-Y. Pi, *Phys. Rev. Lett.* **49** (1982) 1110; S. W. Hawking, *Phys. Lett. B***115**, 295 (1982). V. Mukhanov and G. Chibisov, *Pis'ma Zh. Eksp. Teor. Fiz.* **33**, 549, (1981); A. A. Starobinskii, *Phys. Lett. B***117**, 175 (1982).
- [11] J. Khoury, B.A. Ovrut, P.J. Steinhardt and N. Turok, in preparation.
- [12] J. Khoury, B.A. Ovrut, N. Seiberg, P.J. Steinhardt, N. Turok, in preparation.
- [13] P.J. Steinhardt and M.S. Turner, *Phys. Rev. D***29**, 2162-71 (1984).
- [14] M.A. Bucher, A.F. Goldhaber and N. Turok, *Phys. Rev. D***52**, 3314 (1995).
- [15] A. Linde, *Phys. Rev. D***59**, 023503 (1999); J. Garcia-Bellido and A. Linde, *Phys. Rev. D***55**, 7480 (1997).
- [16] A. Guth, astro-ph/0101507.