Free-eld realisation of boundary states and boundary correlation functions of m in im alm odels

Shinsuke Kawai

Theoretical Physics, D epartment of Physics, University of Oxford, 1 Keble Road, Oxford OX1 3NP, UK, and Helsinki Institute of Physics, P.O. Box 64, FIN-00014 University of Helsinki, Finland

(January 25, 2022)

We propose a general form alism to compute exact correlation functions for C ardy's boundary states. U sing the free-eld construction of boundary states and applying the C oulom b-gas technique, it is shown that charge-neutrality conditions pick up particular linear combinations of conform al blocks. As an example we study the critical Ising model with free and xed boundary conditions, and demonstrate that conventional results are reproduced. This form alism thus directly associates algebraically constructed boundary states with correlation functions which are in principle observable or num erically calculable.

PACS numbers: 11.10 Kk, 11.25 Hf, 68.35 Rh

I. IN TRODUCTION

Modular invariance of partition functions plays extremely important roles in two-dimensional conformal eld theory (CFT). The ADE classi cation of modular invariants by Cappelli, Itzykson and Zuber [1,2] (see also [3(5)] is obtained by considering CFTs on the torus. The classi ed m odular invariants correspond to particular sets of operators, which are supposed to model critical system s in certain universality classes. Sim ilar consideration also applies to CFTs de ned on a manifold with boundary. For a CFT on the cylinder, the constraints from modular invariance lead to a classi cation of boundary states. Thism ethod was invented by Cardy [6] in the eighties and developed by many in the nineties [7{12]. Such a classi cation of boundary states with consistent m odular properties has recently attracted m uch attention along with the developm ent of D-brane / string theory and various application of boundary CFT to statistical physics.

Boundary states with consistent modular properties, or consistent boundary states, are normally believed to represent boundary conditions which may be physically imposed on D-branes or borders of statistical systems. In order to understand the behaviour of observables in the presence of such a boundary, we need to nd correlation functions for consistent boundary states which are de ned through Cardy's classi cation. In principle this can be accomplished by operator product expansions (OPEs), that is, by nding boundary operators for a given boundary state, solving the constraints satis ed by coupling constants, and then obtaining the correlation functions by OPEs using the boundary operators. The correlation functions obtained in this way are perturbative, that is, in the form of series expansion. For practical use, we often need to know exact correlation functions and in such a case we normally solve dierential equations to nd conform al blocks and x their coe cients by physical considerations [13]. In this di erential equation m ethod, how ever, the relation to C ardy's classi cation of boundary states is not quite evident.

In this paper, we present an alternative formalism for nding boundary correlation functions directly using boundary states obtained by Cardy's method. It is well known in string theory that correlation functions are simply given by inserting vertex operators within am plitudes (with or without boundaries). This picture is generalised to non-bosonic (c \in 1) CFTs using the Coulom b-gas form alism on arbitrary Riem ann surfaces (without boundaries) [14{17]. The purpose of this paper is to present a m ethod to calculate boundary correlation functions based on the C oulom b-gaspicture, by using the free-eld representation of boundary states developed in [18]. Driven by a sim ilar motivation, Coulom b-gas system on the halfplane is discussed in [19], where the Ising m odel conform alblocks are reproduced using the contour integration technique of [20] and insertion of boundary operators. The key ingredient of the form alism proposed in the present paper is boundary states having boundary charges, which account both for the contour integral expression of conform alblocks and for their coe cients. The role of such boundary states has not been fully investigated so far in this context, and our form alism allows a systematic study on the relation between correlation functions and algebraically de ned boundary states. In the following we shall mainly consider C oulom b-gas systems on the unit disk, where Felder's charged bosonic Fock space (CBFS) construction [14] is readily used. A though our method itself is quite general, we shall focus

W e use this term for correlation functions of bulk operators in the presence of boundary.

on presenting the ideas in simplest cases and show that it reproduces known results obtained by the conventional approach.

We organise the rest of this paper as follows. In the next section we x our notations and review the free-eld construction of boundary states [18]. We describe in Sec. III our m ethod to compute correlation functions on the disk and on the half plane. In Sec. IV we illustrate the m ethod using the Ising m odel and show that it reproduces the results of [13]. Finally in Sec. V we sum m arise and conclude.

II.BOUNDARY STATES IN COULOM B-GAS FORM ALISM

Let us start, for the sake of self-containedness, by sum marising the C oulom b-gas construction of boundary states [18]. The idea is to de ne coherent states in the charged bosonic Fock spaces (CBFSs) and nd conditions for their di eom orphism invariance and m odular consistency.

A.Coulom b-gas and charged bosonic Fock space

In the C oulom b-gas form alism [21], the V irasoro m inim alm odels are realised by the action,

$$S = \frac{1}{8} d^{2}x^{p} \overline{g} (0 \ 0 + 2^{p} \overline{2}_{0} iR); \quad (1)$$

where the scalar eld (x) is assumed to decouple into two chiral sectors, (z;z) = '(z) + '(z). The energymom entum tensor is obtained by the variation of the action (1) as

$$T(z) = 2 T_{zz} = \frac{1}{2} : 0' 0' :+ i \frac{p}{2} 0 0^{2'} :$$
 (2)

The central charge of this system is

$$c = 1 \quad 24 \quad {}^2_0$$
: (3)

The vertex operators,

$$V (z) = :e^{i^{\frac{p}{2}} \cdot (z)} ;; \qquad (4)$$

are chiral elds of conform ald in ensions $h = {}^{2} 2_{0}$: In particular, the prim ary elds _{r;s} ($0 < r < p^{0}, 0 < s < p$) of a m in in alm odel are realised by the vertex operators V _{r;s} (z) with

$$r_{r;s} = \frac{1}{2} (1 \quad r)_{+} + \frac{1}{2} (1 \quad s)_{+} ;$$
 (5)

where $_{+} = \frac{p}{p=p^{0}}$, $= \frac{p}{p^{0}=p}$, and p and p^{0} (we assume $p > p^{0}$) are the two coprime integers characterising the m inim alm odel. The conform ald in ensions of the

operators are

$$h_{r;s} = \frac{1}{4} (r_{+} + s_{-})^{2} \qquad {}_{0}^{2};$$
 (6)

which agree with the K ac form ula.

The holomorphic chiral boson eld is expanded in modes as

$$'(z) = '_{0} \quad ia_{0} \ln z + i \qquad X \\ a_{n \in 0} \frac{a_{n}}{n} z^{n}; \qquad (7)$$

and likew ise for the antiholom orphic counterpart,

$$'(z) = '_{0}$$
 ia₀ ln z + i $\frac{a_{n}}{n} z^{n}$: (8)

A swe try to consider boundary CFT in the C oulom b-gas picture, a subtlety arises in the treatment of zero-m ode, since the zero-m ode of (z;z) does not naturally decouple into the holom orphic and antiholom orphic sectors. In our form alism, we shall sim ply split it into two identical and independent copies. In this sense, the boundary theory we shall consider is not exactly a non-chiral theory on a manifold with boundary but rather a chiral theory on its Schottky double^V. We thus have two copies of H eisenberg operators, satisfying the algebra

$$[a_{m};a_{n}] = m_{m+n;0}; ['_{0};a_{0}] = i; [a_{m};a_{n}] = m_{m+n;0}; ['_{0};a_{0}] = i;$$

$$(9)$$

with no interaction,

$$[a_{m};a_{n}] = 0; ['_{0};a_{0}] = ['_{0};a_{0}] = ['_{0};'_{0}] = 0: (10)$$

In term s of the H eisenberg operators, the V irasoro operators are w ritten as

$$L_{n \in 0} = \frac{1}{2} \sum_{k \ge Z}^{X} a_{n \ k} a_{k} \qquad p - \frac{1}{2} a_{n \ (n + 1)} a_{n}; \qquad (11)$$

$$L_{0} = \sum_{k=1}^{X} a_{k}a_{k} + \frac{1}{2}a_{0}^{2} \qquad p_{-2}a_{0}a_{0}; \qquad (12)$$

and likew ise for the antiholom orphic counterparts.

The Hilbert space of CFT is realised in CBFSs with BRST projection [14]. For a chiral theory the CBFS F; $_0$ is de ned as a space obtained by operating with the creation operators $a_{n<0}$ on the chiral highest weight state j; $_0$ i (see [14,18,23]). Since the boundary intertwines the two chiral sectors, we need to construct non-chiralFock spaces F; $_0$ in order to describe boundary states. W e denote the M obius invariant non-chiral vacua

^yThe Schottky double is a Riem ann surface obtained by doubling the manifold except for boundaries. See, e.g., [22].

with background charge $_0$ as^z h0;0; $_0$ j and j0;0; $_0$ i, and de ne highest weight vectors h; ; $_0$ j and j; ; $_0$ i as

$$j;;_{0}i = e^{i^{2} ' \circ e^{i^{2} ' \circ j}};0;_{0}i:$$
 (14)

The state h; ; $_0jhas$ holom orphic and antiholom orphic charges and , respectively. Likewise, j; ; $_0ihas$ holom orphic and antiholom orphic charges

and $% \left({{\mathbf{F}}_{\mathbf{n}}} \right)$. Using the H eisenberg algebra (9) it is easy to verify that these states satisfy

h;;
$$_{0}\dot{j}a_{0} = h;; _{0}\dot{j}\frac{p}{2}2;$$
 (15)

h;;
$$_{0}\dot{j}a_{0} = h;; _{0}\dot{j}\overline{2};$$
 (16)

$$a_{0}j;;_{0}i = 2j;;_{0}i;$$
 (17)

$$a_0j;; ; _{0}i = \frac{p_2}{2}j;; ; _{0}i:$$
 (18)

T here highest weight vectors are eigen states of the V i-rasoro zero-m odes:

$$L_0j;;_0i=(^2 2 _0)j;;_0i;$$
 (19)

$$L_0j;;;_0i = (\begin{array}{ccc} 2 & 2 & _0)j;;;_0i:$$
 (20)

The states j; ; _0i are annihilated by $a_{n>0}$, $a_{n>0}$, $L_{n>0}$, $L_{n>0}$, $L_{n>0}$, and h; ; _0 jare annihilated by $a_{n<0}$, $a_{n<0}$, $L_{n<0}$, $L_{n<0}$. The non-chiralCBFSF ; _0 is built on the highest weight vector j; ; _0 iby operating with $a_{n<0}$ and $a_{n<0}$. The dual space F ; _0 is de ned similarly, by acting with $a_{n>0}$ and $a_{n>0}$ on h; ; _0 j. The non-chiral CBFSs thus de ned are essentially the direct products of chiralCBFSs, F ;; _0 = F ; _0 F ; _0.

The inner products of highest weight vectors are subject to charge neutrality, i.e. they are non-vanishing only if the net charges in the two sectors are both zero. The norm alisation of the highest weight vectors must be in accordance with this condition and thus we have

In particular, the vacua are norm alised as

$$h0;0; _{0};0; _{0}i = :$$
 (22)

In unitary theories the constant is usually positive and we norm alise it to unity. A sthe C oulom b-gas system m ay

well include non-unitary theories, can be negative. In that case we choose = 1. Thus we have

$$_{\rm U}$$
 h;; $_{\rm 0}$ j;; $_{\rm 0}$ i_U = ; ; (23)

$$_{N}h$$
;; $_{0}j$;; $_{0}i_{N} = ;$; (24)

The subscripts U and N stand for unitary and nonunitary sectors, respectively. Those two sectors have no intersection.

B.Di eom orphism invariant boundary states

Boundary states appearing in CFT are di eom orphism invariant in the following sense [6,24]. Let us consider a CFT on the upper half plane Im 0, where is a com – plex coordinate, = x + iy, x, $y \ge R$. The boundary is y = 0, or = . Since the antiholom orphic coordinate dependence on the upper half plane m ay be m apped into the holom orphic dependence on the lower half plane [13], we often identify with . Now, once the boundary is xed, the conform al symmetry of the theory should be restricted so that the boundary is kept xed. In other words, as the conform altransform ation is generated by the energy-m om entum tensor, the energym om entum ow across the boundary m ust vanish,

$$T() T() = 0:$$
 (25)

This may be translated into a condition on boundary states by mapping a semiannular domain on the upper half -plane into a full-annulus on the z-plane by $z = exp(2 \ i = L)$, $= (T =) \ln$. The boundary of the plane is mapped to the two concentric circles $j_{2}j = 1$, $exp(2 \ T = L)$ bordering the annulus on the z-plane. Since the z-plane allow s radial quantisation, (25) is written as the condition on the boundary states β i (on $j_{2}j = 1$),

$$(L_k \ L_k) = 0$$
: (26)

This condition, often called the Ishibashi condition, must then be satis ed by any boundary state in CFT.

We may follow the standard construction of boundary states in open string theory [25{27] and nd boundary states on CBFS by starting from the coherent state ansatz,

$$hB_{;;0}; j = h;;0j \exp \frac{1}{k}a_{k}a_{k};$$

$$(27)$$

$$B_{;0}; i = Y \exp \frac{1}{k}a_{k}a_{k} j;;0i:$$

$$(28)$$

The subscript is either U or N, specifying the norm alisation of the vacuum . A swe have expressions of the V irasoro operators (11), (12) written in terms of the H eisenberg operators, one can see how the V irasoro m odes L_n

^zHere we give the same background charge $_{0}$ to both sectors. Even if we relax this condition and start by allocating di erent background charges $_{0}$ and $_{0}$ to holom orphic and antiholom orphic sectors, respectively, the condition (26) restricts either $_{0} = _{0}$. For $_{0} = _{0}$ we have = 1 and $_{2}$ $_{0} = 0$ instead of (29) and (30). This merely ips

the sign of all antiholom orphic charges and thus does not give any new results.

and L $_{n}$ operate on the coherent states β ; $_{0}$; i by an explicit computation. It is shown [18] that the condition (26) is satis ed if

and

+
$$2_0 = 0$$
: (30)

Similarly, we see that hB ;; $_{0}$; j(L_n L_n) = 0 as long as (29) and (30). In the following we shall only consider such manifestly dieomorphism invariant boundary states satisfying (29) and (30), and for simplicity we denote,

$$hB()j = hB_{;2}, j = 1j;$$
 (31)

$$\beta$$
 () $i = \beta$; 2 , ; ; ; 1 i: (32)

We note that the sum of the (holom orphic + antiholom orphic) boundary charges agrees with the topological background charge on the Schottky double. Due to the condition (30), an inner boundary (on the z-plane) contributes 2 $_{\rm 0}$ to the sum of the charges, and an outer boundary contributes 2 0. When we consider an annulus whose Schottky double is a torus, the sum of the boundary charges is zero $(2_0 \ 2_0 = 0)$, which coincides with the background charge of the torus expected from the Gauss-Bonnet theorem (the Euler num ber of a torus vanishes). For a disk, there is only an outer boundary which gives a charge 2_0 . This agrees with the background charge of a sphere, which is the Schottky double of the disk. Thus, the geometry of bulk can be assumed to be at everywhere, since the curvature of the manifold is concentrated on boundary.

${\tt C}$. Ishibashi states in free- eld representation

As the basis of boundary states in CFT is normally spanned by Ishibashi states, we need to construct Ishibashi states in term s of our Fock space representation in order to translate existing results of boundary CFT into C oulom b-gas language. At least for the diagonal minimalm odels, it is shown [18] that the Ishibashi states are expressible as linear combinations of β () i, as far as partition functions on the cylinder are concerned.

Ishibashi states are de ned for chiral representations of CFT and diagonalise the cylinder am plitudes (overlaps) to give characters,

$$lhij(q^{1=2})^{L_0 + L_0 - c=12} jji = _{ij - j} (q):$$
(33)

Here, we are considering a cylinder of length T and circum ference L, or equivalently, an annulus on the z-plane with 1 jzj exp(2 T=L). As the H am iltonian is written as H = (2 =L) (L₀ + L₀ c=12), the left hand side of

(33) is ling TH jji with $q = e^{4 T = L}$. The characters of the minimal models are given by Rocha-Caridi [28],

$$(r;s) (q) = Tr_{(r;s)} q^{L_0} c^{-24}$$

$$= \frac{pr p^0 s; pp^0 () pr + p^0 s; pp^0 ()}{()};$$
(34)

where ; () and () are the Jacobi theta function and the Dedekind eta function, de ned as ; () $P_{k2Z}^{2} q^{(2 \ k+)^{2}=4}$ and () $q^{=24} Q_{n-1}^{2} (1 \ q^{n})$, with $q = e^{2 \ i}$. Thus, the cylinder am plitudes (33) are power series in q, divided by (~).

As our boundary states β ()i are de ned in a Fock space representation, we may explicitly compute the cylinder am plitudes between such states. They are [18],

$$hB()je^{TH} B()i$$

$$= hB()j(q^{1-2})^{L_0+L_0} c^{-12}B()i$$

$$= \frac{q^{(0)^2}}{(c)}; ; ; (35)$$

The am plitudes between unitary and non-unitary sectors (e.g. $_{\rm U}$ hB ()je TH ()i $_{\rm N}$) vanish because these sectors do not intersect. Note that q^{(0)²} = () is the character ; $_{0}$ (g) of the chiral CBFSF; $_{0}$. In this sense, the state () i m ay be regarded as the Ishibashi state of F; $_{0}$. W e m ay now compare the expressions (33) and (35) to nd a possible free- eld representation of the Ishibashi states of m inim alm odels. D e ning

$$h(r;s) j = U ha_{r;s} j + N ha_{r;s} j; \qquad (36)$$

and

$$j(r;s)ii = ja_{r;s}i_{U} + ja_{r;s}i_{N};$$
 (37)

with

$${}_{U} ha_{r;s} j = \sum_{k \ge Z}^{X} {}_{U} hB ({}_{r;s} + k^{p} \overline{pp^{0}}) j;$$
(38)

$${}_{N} ha_{r; s} j = {}_{N} hB (r; s + k pp^{0}) j;$$
(39)

$$\dot{\mathbf{j}}_{r;s} \dot{\mathbf{i}}_{U} = \sum_{k \ge Z}^{X} \dot{\mathbf{j}}_{k} \left(\mathbf{r}_{;s} + \mathbf{k}^{p} \ \overline{pp^{0}} \right) \dot{\mathbf{i}}_{U}; \qquad (40)$$

where $r_{;s}$ are given by (5), one can show that the states j(r;s) ii diagonalise the overlaps and give m inim alm odel characters,

$$\hbar(\mathbf{r};\mathbf{s})\,\mathbf{j}(\mathbf{q}^{1=2})^{L_0+L_0}\,\,^{c=12}\,\mathbf{j}(\mathbf{r}^0;\mathbf{s}^0)\,\mathbf{\ddot{n}} = \,_{\mathbf{r}\mathbf{r}^0}\,_{\mathbf{s}\mathbf{s}^0}\,_{(\mathbf{r};\mathbf{s})}\,(\mathbf{q})\,\mathbf{:}$$
(42)

The appearance of the non-unitary sector even in unitary CFTs may seem odd, but this is necessary to describe cylinder diagram s where otherwise unphysical states would propagate. On the disk the non-unitary sector decouples and does not contribute to correlation functions (Sec. III). The states (36), (37) are a good candidate for the m inim alm odel Ishibashi states as far as the m odular properties are concerned. We, however, im m ediately notice that such Ishibashi' states are not unique since j(r;s)i and $j(p^0 r;p s)i$ give rise to the same character but are perpendicular to each other. In order to have a unique Ishibashi state for each prim ary eld (r;s) ($p^0 r;p s$) of m inim alm odels, we de ne the symmetrized states,

$$\begin{split} \ln_{r;s} j &= h_{p^{0} r;p} s j \\ &= \frac{1}{p \frac{1}{2}} (h(r;s)j + h(p^{0} r;p s)j); \quad (43) \\ j_{r;s} ii &= j_{p^{0} r;p} s ii \end{split}$$

$$= \frac{1}{p} (j(r;s) \ddot{u} + j(p^{0} r;p s) \ddot{u}); \qquad (44)$$

and shall regard them as our free-eld realisation of the Ishibashi states. Although this 'symmetrisation' was not considered in [18], such a prescription to ensure the equivalence of (r; s) (p^0 r; p s) is necessary.

D.Cardy's consistent boundary states

Physical boundary states in CFT are not only dieomorphism invariant, but must also satisfy an extra constraint called C ardy's consistency condition. W e consider a cylinder of length T and circum ference L as before and assume that boundary conditions ~ and ~ are im posed on the two boundaries. Then, depending on how we de ne the direction of time, the partition function on this cylinder may be written in two di erent ways. We may rst regard the cylinder as an open string propagating in the periodic direction of time, with boundary conditions \sim and \sim imposed at the two ends of the string. The partition function is then a sum of the chi- $\int_{1} n_{2}^{j} dq$, where j(q) is ral characters, Z_{2} (q) = the character for the chiral representation j and n_{j}^{j} is a non-negative integer representing the multiplicity of the representations. We have de ned q = $e^{\quad \ \ L=T}$. We m ay also see the cylinder as a closed string propagating from one boundary (with boundary condition ~) to the other (with \sim). Then the partition function is sim ply the cylinder am plitude between the two boundaries, $h \sim je^{TH} j_{p}$ i. Due to the equivalence of the two pictures, $_{j}n_{_{\sim}}^{j}$ $_{j}(q) = h \sim j(q^{1=2})^{L_{0} + L_{0}} c^{=12} j^{-1}$. This is we have the consistency condition which needs to be satis ed by the boundary states h~jand j~i.

If we have an appropriate basis of the boundary states, the right hand side of the consistency equation may be expanded using the basis states flag, fpig as

$$X = N_{-}^{j} (q) = X_{-}^{j} h_{-} j a i h a j (q^{1-2})^{L_{0} + L_{0}} c^{-12} j b i h b j^{-} i;$$

$$j = a_{ib}$$
(45)

Solving this equation, the consistent boundary states fj~ig are found as linear combinations of the basis states. It is convenient to use the Ishibashi states for such a basis. For diagonal minimal models, using the property (33) of the Ishibashi states and the modular transformation of the characters $_{i}(q) = _{j}S_{ij} _{j}(q)$ under ! ~ = 1 =, we have, by equating the coe cients of the characters,

ie characters, x

$$n_{\sim}^{i}S_{ij} = h^{jilhjj^{i}}$$
(46)

A sum ing the existence of a state \Im such that $n_{\Im^{-}}^{i} = n_{\Box^{-}}^{i}$, (46) was solved by Cardy [6] as,

$$j \sim i = \begin{cases} X \\ j \\ j \end{cases} j = \begin{cases} X \\ p \\ j \\ j \end{cases} = \begin{cases} X \\ p \\ s_{0j} \\ s_{0j} \end{cases} j = (47)$$

Now, as the minimal model Ishibashi states have been written in the free-eld representation (43) (44), Cardy's consistent boundary states (47) can also be expressed using our coherent boundary states by substituting (44) into (47).

For the convenience of later discussions, let us spell out these in the speci c example of the Ising model. The Ising model is the simplest non-trivial minimal model having the central charge c = 1=2 and is characterised by the two coprime integers p = 4 and $p^0 = 3$. There are three operators, the identity I, the energy and the spin , having the conformal dimensions 0, 1=2, and 1=16, respectively, and are identied in the K actable as I = $_{1;1} = _{2;3}$, $= _{2;1} = _{1;3}$ and $= _{1;2} = _{2;2}$. Using the modular transform ation matrices for the characters, C ardy's boundary states are written as [6]

$$\dot{\tau}_{1} = 2^{1=2} \exists \ddot{u} + 2^{1=2} \exists \ddot{u} = 2^{1=4} \exists \ddot{u};$$
 (49)

It is argued that the rst two states correspond to the xed (up and down) boundary conditions since they di er only by the sign of j ii which is associated with the spin operator. The last state (50) is then identi ed as the free boundary state. U sing (37) and (44), these states are written in our free- eld representation as

$$\begin{aligned} \mathbf{j}^{T}\mathbf{i} &= 2^{-1} \left(\mathbf{j}_{1;1} \mathbf{i}_{U} + \mathbf{j}_{1;1} \mathbf{i}_{N} + \mathbf{j}_{2;3} \mathbf{i}_{U} + \mathbf{j}_{2;3} \mathbf{i}_{N} \right. \\ &+ \mathbf{j}_{2;1} \mathbf{i}_{U} + \mathbf{j}_{2;1} \mathbf{i}_{N} + \mathbf{j}_{1;3} \mathbf{i}_{U} + \mathbf{j}_{1;3} \mathbf{i}_{U} + \mathbf{j}_{1;3} \mathbf{i}_{N} \right) \\ &+ 2^{-3=4} \left(\mathbf{j}_{1;2} \mathbf{i}_{U} + \mathbf{j}_{1;2} \mathbf{i}_{N} + \mathbf{j}_{2;2} \mathbf{i}_{U} + \mathbf{j}_{2;2} \mathbf{i}_{U} + \mathbf{j}_{2;2} \mathbf{i}_{N} \right); \end{aligned}$$

$$(51)$$

$$j \cdot i = 2^{1} (j_{a_{1};1} i_{U} + j_{a_{1};1} i_{N} + j_{a_{2};3} i_{U} + j_{a_{2};3} i_{N} + j_{a_{2;1}} i_{U} + j_{a_{2};1} i_{N} + j_{a_{1};3} i_{U} + j_{a_{1};3} i_{N}) 2^{3=4} (j_{a_{1;2}} i_{U} + j_{a_{1};2} i_{N} + j_{a_{2};2} i_{U} + j_{a_{2};2} i_{N});$$
(52)

$$j_{\nu}i = 2 \, {}^{1=2} (j_{a_{1};1}i_{U} + j_{a_{1};1}i_{N} + j_{a_{2};3}i_{U} + j_{a_{2};3}i_{N}
j_{a_{2};1}i_{U} j_{a_{2};1}i_{N} j_{a_{1};3}i_{U} j_{a_{1};3}i_{N});$$
(53)

where the states on the right hand sides are de ned by (40), (41). They are superpositions of countably many coherent states with di erent boundary charges.

III. BOUNDARY CORRELATION FUNCTIONS

Now let us discuss how to compute boundary correlation functions in our Coulom b-gas picture. After giving the general form alism, we shall focus on the one point function of $r_{;s}(z;z)$ and the two point function of 1:2 (z; z), and derive their explicit expressions on the unit disk. Once correlators on the disk are obtained, it is straightforward to map them on the halfplane. At the end of this section, we compare our free-eld approach and OPE com putation of boundary correlation functions.

A . Screened vertex operators and BRST states

In the CBFS language, correlation functions on the full plane are described as follows [14]. We de ne chiral screened vertex operators V^m_{r;s}ⁿ (z) as

$$V_{r;s}^{m,in}(z) = \begin{pmatrix} 1 & Y^{n} & Y^{n} \\ du_{1} & dv_{j}V_{r;s}(z)V_{+}(u_{1}) & + M_{m} \end{pmatrix}$$

$$i=1 \quad j=1 \\ V \quad (v_{1}) \quad (M_{n}); \quad (54)$$

where for conciseness we have denoted V $_{r:s}$ (z) as V $_{r;s}$ (z) and V (z) as V (z), and the integration contours are those of Felder's, all going through z and encircling the origin (Fig.1 of [14]). Such an operator is a primary eld of conform aldimension h_{r;s}. We also denote the chiral CBFS F $_{r;s}$; $_{0}$ as F $_{r;s}$. The operator $V_{r;s}^{m,in}$ (z) de nes a m ap from one Fock space to another,

$$V_{r;s}^{m;n}(z)$$
: $F_{r_0;s_0}$! F_{r_0+r} $2m$ 1; s_0+s $2n$ 1: (55)

The p-point correlator,

is then seen as a sequence of m appings,

F_{1:1} !

$$F_{r_p 2m_p;s_p 2n_p}!$$
; (57) and

and the nalstate $V_{r_1;s_1}^{m_1;n_1}$ (z₁) ^m_{rp};yⁿ_p (z_p)j); ₀imust belong to F1;1 in order to have a non-trivial innerproduct with h0; $_{0}j2 F_{1;1}$ (the dualm odule of $F_{1;1}$). The same p-point correlator m ay be expressed di erently, as

$$h_{p^{0} 1;p 1; 0} \mathcal{Y}_{r_{1};s_{1}}^{m_{1}^{0};m_{1}^{0}}(z_{1}) \qquad r_{p} \mathcal{X}_{p}^{m_{p}^{0}}(z_{p}) \mathcal{D}; 0 i; (58)$$

but this is in fact proportional to (56). A key object in this form alism is the BRST operator,

$$Q_{r} = \frac{e^{2 i \frac{2}{r} r} 1}{r(e^{2 i \frac{2}{r}} 1)} \int_{i=1}^{I} Y^{r} du_{i}V_{+}(u_{1}) + (u_{1}); \quad (59)$$

which maps Fr;s to F r;s. The BRST operator is nilpotent, $Q_r Q_{p^0} = 0$, and physical states are realised as the cohom ology space, K erQ r=Im Q p⁰ r. An im portant point which is evident in this picture is that all interm ediate states appearing in the correlator are BRST states since the vacuum j); 0 i is a BRST state and the screened vertex operators m ap BRST states to BRST states [14].

W e shall com bine the above m achinery and the boundary states of the previous section to compute correlation functions on the unit disk. The in-state of the correlators is the non-chiral vacuum (in the unitary sector) at the origin, $10i = 10;0; _0i_U$, which is a BRST state. For the out-state, we choose a boundary state $_{\rm U}$ hB ()j of (31) with a xed boundary charge . We only consider the unitary sector since the non-unitary sector does not give non-trivial inner products with the in-state vacuum . The correlators are then obtained by inserting non-chiral screened vertex operators,

$$V_{(r_{i};s_{i});(r_{i};s_{i})}^{(m_{i};n_{i});(m_{i};n_{i})}(z_{i};z_{i}) = V_{r_{i};s_{i}}^{m_{i};n_{i}}(z_{i})V_{r_{i};s_{i}}^{m_{i};n_{i}}(z_{i});$$
(60)

between $_{\rm U}$ hB ()j and jD;0; $_{\rm 0}$ i $_{\rm U}$. As we focus on diagonal theories, the antiholom orphic vertex operators have the same conform aldimensions as the holom orphic counterparts, i.e., either $(r_i; s_i) = (r_i; s_i)$ or $(r_i; s_i) =$ $(p^0 r_i; p s_i)$. Note that, in this construction, all the interm ediate states are manifestly BRST invariant because no spurious states arise. A ctual boundary p-point correlation functions for physical boundary conditions are obtained by sum ming the xed boundary-charge correlators,

$$U hB () j V_{r_{i};s_{i}}^{m_{i};n_{i}}(z_{i}) V_{r_{i};s_{i}}^{m_{i};n_{i}}(z_{i}) j 0;0; 0 i_{U};$$
(61)

over the boundary charges according to the linear com binations (such as (51) - (53)) given by Cardy's method. Note that (61) is non-vanishing only for certain con gurations of screening charges. The net holom orphic and antiholom orphic charges are respectively,

h

which must vanish independently. These charge neutrality conditions associate the allowed values of with the numbers of holom orphic and antiholom orphic screening operators.

In the computation of the expression (61), $V_{r;s}(z)V_{r;s}(z)$, $V_{r;s}(z)V_{p^0} r;p s(z)$, $V_{p^0} r;p s(z)V_{r;s}(z)$ and $V_{p^0} \ _{r \not\!\!\! r} \ _s (z) V_{p^0} \ _{r \not\!\!\! r} \ _s (z)$ all correspond to a nonchiral eld r;s (z;z) and one may use whichever com binations one wishes to use. This is ensured by the fact that (61) is essentially a chiral 2p-point function where the equivalence of (r;s) \$ $(p^0 r;p s)$ (after the truncation of unphysical states) is guaranteed [14,21]. In particular, we are allowed to use $V_{r;s}(z)$ as the holom orphic and $V_{p^0} r_{p^*} s$ (z) as the antiholom orphic (in irror in age') part of a single non-chiral operator (the analogy of a m irror and a boundary of CFT is based on the conform alW ard identity [13] which does not distinguish $V_{r,s}$ (z) from $V_{p^0} r_{p} s(z)$. Due to this equivalence, apparently di erent choices of vertex operators should all lead to a same conform alblock function. In practice, likewise to the case of the C oulom b-gas com putation without boundary, we shall choose such vertex operators that the num ber of screening operators is m in im ised. In the following subsections we shall evaluate the expression (61) for particular cases of one and two point functions.

B.Boundary one point functions

For evaluation of the boundary one point correlator,

$$_{\rm U}$$
 hB () $j_{\rm r;s}^{\rm m;n}$ (z) $V_{\rm r;s}^{\rm m;n}$ (z) $j_{\rm 0};0;_{\rm 0}i_{\rm U};$ (64)

It is convenient to choose $(r;s) = (p^0 r;p s)$ (the other choice (r;s) = (r;s) should lead to the same result but involves complicated integral expressions). A coording to (55), the holom orphic part of the CBFS is mapped as

$$F_{1:1} ! F_r _{2m:s} _{2n};$$
 (65)

and the antiholom orphic part is mapped as

$$F_{1;1} ! F_{p^0} r_{2m,p} s_{2n}$$
: (66)

Since $_{U}$ hB ()j 2 F; $_{0}$ F₂₀; $_{0}$, the correlator is non-vanishing only when F; $_{0}$ = F_{r 2m}; $_{2n}$ and F₂₀; $_{0}$ = F_{p⁰ r 2m; $_{p}$ s 2n, that is,}

$$= \frac{1}{2} (1 r + 2m) + \frac{1}{2} (1 s + 2n) ; \quad (67)$$

and

2
$$_{0} = \frac{1}{2} (1 p^{0} + r + 2m) + \frac{1}{2} (1 p + s + 2n) : (68)$$

Sum m ing (67), (68) and using $_{+} = p = p^{0}$, $= p^{0} = p^{0}$, we have $(m + m)_{+} + (n + n)_{-} = 0$, or


$$m + m = 0; n + n = 0;$$
 (69)

in plying no screening charges. Then from (67) we have = r_{13} , and the correlator (64) is evaluated as

$$_{\rm U}$$
 hB ($_{\rm r;s}$) $\mathcal{Y}_{\rm r;s}^{0;0}$ (z) $\mathcal{V}_{\rm r;s}^{0;0}$ (z) $\mathcal{D};0;$ $_{\rm I}$ = (1 zz) ^{2h}; (70)

where $h = r_{;s} (r_{;s} 2_0)$. As we shall see later, this is proportional to the two point correlator on the full plane.

(a) Conform alblock I_I.






FIG.1. The two conform alblocks I_I and I_{II} in the boundary two point function of $_{1,2}$ on the disk. The sphere represents the double of the disk, and the upper and lower hem is spheres stand for the holom orphic and antiholom orphic sectors, which are glued on the boundary (the equator). The lower hem isphere coordinates z_i are obtained from z_i by the doubling $(z_i; z_i)$! $(z_i; z_i)$. In the case of I_I where the screening operator Q lie in the holom orphic sector, the integration contour can be deformed into the Pochhammer type around z_1 and z_2 . The integral is then proportional to the one from z_2 to z_1 (a). Similarly, the screening contour of I_{II} is in the antiholom orphic sector and the integral is proportional to the one from z_2 to z_1 (b).

C.Boundary two point functions of 1;2

As a less trivial case, we consider the two point correlator of the primary eld $_{1;2}$. For the convenience of calculation we de ne one of the operators as $_{1;2}(z_1;z_1) = V_{1;2}(z_1)V_{p^0 \ 1;p \ 2}(z_1)$ and the other as $_{1;2}(z_2;z_2) = V_{1;2}(z_2)V_{1;2}(z_2)$. The expression (61) then becomes

$$V_{1;2}^{m_{2},n_{2}} (z_{2}) V_{1;2}^{m_{2},n_{1}} (z_{1}) V_{p^{0}}^{m_{1},n_{1}} (z_{1}) (z_{p^{0}}^{m_{2},n_{2}} (z_{1}) V_{1;2}^{m_{2},n_{2}} (z_{2}) D_{p^{0}} (z_{1}) (z_{1}$$

In the holomorphic and antiholomorphic sectors, the CBFSs are mapped as

$$F_{1;1} ! F_{1 \ 2m_2;2 \ 2n_2} ! F_{1 \ 2m_1 \ 2m_2;3 \ 2n_1 \ 2n_2};$$
 (72)

$$F_{1;1} ! F_{1 \ 2m_{2};2} \ 2n_{2} ! F_{p^{0} \ 2m_{1}} \ 2m_{2} \ 1;p \ 2n_{1} \ 2n_{2} \ 1;$$
(73)

and hence, in order that the correlator be non-vanishing we must have

$$= (m_1 + m_2) + (n_1 + n_2 1) ; \qquad (74)$$

$$2_{0} = (m_{1} + m_{2} + 1)_{+} + (n_{1} + n_{2} + 1) :$$
(75)

Adding the above two expressions we have (m + m) + (n + n - 1) = 0, where $m = m_1 + m_2$ and $n = n_1 + n_2$ are the numbers of positive (+) and negative (-) screening operators in the holom orphic sector, and similarly $m = m_1 + m_2$ and $n = n_1 + n_2$ for the antiholom orphic counterparts. This charge neutrality condition in plies

$$m + m = 0; n + n = 1:$$
 (76)

Then we have two possibilities:

(I)
$$m = m = n = 0; n = 1;$$
 (77)

$$(II) m = m = n = 0; n = 1:$$
(78)

From (74), we have the boundary charge = $_{1;1} = 0$ for (I) and = $_{1;3} =$ for (II).

For the charge con guration (I), we have one screening operator

т

$$Q = dvV (v);$$
 (79)

in the holom orphic sector and thus the correlator (71) takes the form , $% \left(\mathcal{T}^{(1)}_{1}\right) =0$

$$I_{I} = {}_{U} hB ({}_{1;1})j dvV_{1;2} (z_{1})V_{p^{0}} {}_{1;p} {}_{2} (z_{1})V (v)$$

$$V_{1;2}(z_2)V_{1;2}(z_2)J;0; _0 i_U;$$
 (80)

which, by an explicit calculation, reduces to

$$dv (1 \quad z_{1} z_{1})^{a} (1 \quad z_{1} z_{2})^{b} (1 \quad v z_{1})^{c} (1 \quad v z_{2})^{d}$$

$$(1 \quad z_{2} z_{1})^{a} (1 \quad z_{2} z_{2})^{b} (z_{1} \quad v)^{d}$$

$$(z_{1} \quad z_{2})^{b} (v \quad z_{2})^{d} (z_{1} \quad z_{2})^{a}; \qquad (81)$$

Ι

with a = 2 $_{1,2}$ (2 $_0$ $_{1,2}$), b = 2 $_{1,2}^2$, c = 2 (2 $_0$ $_{1,2}$), and d = 2 $_{1,2}$. As this expression is analytic, we may deform the integration contour as long as it is closed and non-contractable. In this case the screening operator must lie entirely on the holom orphic part and Felder's contour can be deformed into the Pochhammer type, going around z_1 and z_2 . The correlator is then proportional to the integration from z_2 to z_1 , and is written as

$$I_{I} = N_{I} (1 \quad z_{1} z_{1})^{a} (1 \quad z_{1} z_{2})^{b} (1 \quad z_{2} z_{1})^{a}$$

$$Z_{z_{1}} (1 \quad z_{2} z_{2})^{b} (z_{1} \quad z_{2})^{b} (z_{1} \quad z_{2})^{a}$$

$$dv (1 \quad v z_{1})^{c} (1 \quad v z_{2})^{d} (z_{1} \quad v)^{d} (v \quad z_{2})^{d}:$$

$$z_{2}$$
(82)

Here, N_I is a constant arising from the deformation of the contour. Note that, at this point, the expression is similar (in fact, proportional) to the integral representation of a chiral four point conformal block [21] (without boundary). One may then proceed in the standard manner, namely, by xing the projective SL (2;C) gauge, performing the integration, and then recovering the co-ordinate dependence. We thus have,

$$I_{I} = N_{I} (1 \quad z_{1} z_{1})^{a} (1 \quad z_{2} z_{2})^{a} [(1 \quad 1)^{a}]$$
$$\frac{(1 \quad 2)^{2}}{(2 \quad 2^{2})^{a}} F (2a; 1 \quad 2; 2 \quad 2^{2};); \qquad (83)$$

where $F = {}_2F_1$ is the hypergeometric function of the Gaussian type, and is de ned as

$$= \frac{(z_1 \quad z_2)(z_2 \quad z_1)}{(1 \quad z_1 z_1)(1 \quad z_2 z_2)}:$$
(84)

The calculation for (II) goes similarly. A swe have one screening operator

$$Q = dvV (v); \qquad (85)$$

in the antiholom orphic sector, the correlator is written as I

$$I_{II} = {}_{U} h B ({}_{1;3}) j dv V_{1;2} (z_1) V_{p^0} {}_{1;p} {}_{2} (z_1) V (v)$$

$$V_{1;2} (z_2) V_{1;2} (z_2) \mathfrak{D}; 0; _{0} i_{U}$$

$$= N_{II} (z_1 \quad z_2)^{b} (z_1 \quad z_2)^{a} (1 \quad z_1 z_1)^{a}$$

$$(1 \quad z_1 z_2)^{b} (1 \quad z_2 z_1)^{a} (1 \quad z_2 z_2)^{b}$$

$$Z_{z_1}$$

$$dv (z_1 \quad v)^{c} (v \quad z_2)^{d} (1 \quad z_1 v)^{d} (1 \quad z_2 v)^{d};$$

$$z_2$$
(86)

We have again deformed Felder's integration contour (this time in the antiholomorphic sector) into the Pochhammer contour around z_1 and z_2 . The resulting integral is proportional to the one from z_2 to z_1 , and N_{II} is a constant. Performing the integration we have

$$I_{II} = N_{II} (1 \quad z_1 z_1)^a (1 \quad z_2 z_2)^a [(1)^a (1)^{a} (1)^{$$

The two correlators $I_{\rm I}$ and $I_{\rm II}$ with xed boundary charges are represented schematically (in the Schottky double picture) in Fig.1. They correspond to the two conform alblocks of the chiral four point function.

D.Correlation functions on the half plane

Boundary correlation functions obtained on the unit disk arem apped on the halfplane by the global conform al transform ation,

$$w = iy_0 \frac{z}{z+1}; \quad w = iy_0 \frac{z}{z+1};$$
 (88)

which takes the unit circle $j_z j = 1$ on the z-plane to the in nite line Im w = 0 on the w-plane, and the origin z = 0to the point $w = iy_0$, $y_0 \ge R$. Under this transform ation the holom orphic coordinate dependence on the unit disk is mapped on to the upper half w-plane, and the antiholom orphic dependence is mapped on to the lower half w-plane. U sing the transform ation (88), p-point correlation functions on the half plane are written using those on the z-plane, i.e. on the disk, as

$$h_{1}(w_{1};w_{1}) = (w_{p};w_{p})i_{UHP}$$

$$= \frac{Y^{p}}{\sum_{i=1}^{dw_{i}} \frac{dw_{i}}{dz_{i}}} = \frac{M^{p}}{\sum_{i=1}^{h_{i}} \frac{dw_{i}}{dz_{i}}} = \frac{Y^{p}}{\sum_{i=1}^{2} \frac{2Y_{0}}{(z_{i}+1)(z_{i}+1)}} = \frac{Y^{p}}{h_{1}(z_{1};z_{1})} = \frac{2y_{0}}{h_{1}(z_{1};z_{1})} = \frac{2y_{0}}{h_{1}(z_{1};z_{$$

where we have assumed $h_i = h_i$. The parameter of (84) is mapped under this transform ation as

$$=\frac{(z_1 \quad z_2)(z_2 \quad z_1)}{(1 \quad z_1z_1)(1 \quad z_2z_2)}=\frac{(w_1 \quad w_2)(w_1 \quad w_2)}{(w_1 \quad w_1)(w_2 \quad w_2)}; \quad (90)$$

which is an anharm onic ratio of the four points w_1 , w_2 , w_1 and w_2 .

Now, the boundary one point function of $_{\rm r;s}$ on the upper half plane is easily found by using (70) as

$$h_{r;s} (w; w) i_{B(r;s)} = \frac{2y_{0} (1 zz)}{(z+1) (z+1)}$$

$$= [i(w w)]^{2h}$$

$$= (2y)^{2h}; \qquad (91)$$

where $h = h_{r;s} = r_{r;s} (r_{r;s} 2_0)$ is the conform ald in ension of the operator $r_{r;s}$, and w = x + iy, w = w = x iy, x;y 2 R. Note that the result is y_0 -independent. The two point function of 1;2 on the disk is mapped on to the half plane similarly. For the conform alblock (I) we have

$$h_{1;2} (w_{1}; w_{1})_{1;2} (w_{2}; w_{2}) i_{B (1;1)} = \frac{4y_{0}^{2}}{(z_{1} + 1) (z_{1} + 1) (z_{2} + 1) (z_{2} + 1)} I_{I}$$

$$= N_{I} \frac{(w_{1} + w_{1}) (w_{2} + w_{2})}{(w_{1} + w_{2}) (w_{1} + w_{2}) (w_{1} + w_{2}) (w_{1} + w_{2})} I_{I}$$

$$= \frac{(1 + 2)^{2}}{(2 + 2)^{2}} F (4h; 1 + 2; 2 + 2) (w_{1} + w_{2}) (w_{$$

and for (II) we have

$$h_{1;2} (w_{1}; w_{1})_{1;2} (w_{2}; w_{2}) i_{B_{(1;3)}}$$

$$= \frac{4y_{0}^{2}}{(z_{1} + 1)(z_{1} + 1)(z_{2} + 1)(z_{2} + 1)} I_{II}$$

$$= N_{II} \frac{(w_{1} w_{1})(w_{2} w_{2})}{(w_{1} w_{2})(w_{1} w_{2})(w_{1} w_{2})(w_{1} w_{2})} {(1 - 2)(3 - 2)(w_{1} w_{2})(w_{1} w_{2})}$$

$$= \frac{(1 - 2)(3 - 2)}{(2 - 2)} ((1 - 2)^{2h + 2})$$

$$= F(1 - 2)(3 - 2)(2 - 2)$$

$$(93)$$

where $h = h_{1,2} = 1,2 (1,2,2)$. Physical correlation functions are linear sum s of these conform alb locks where the coecients are given by Cardy's states.

E.Boundary states and conform alblocks

Before illustrating in specic examples, we mention how the above description of boundary correlation functions ts into the conventional discussion of [7,8], and see the validity and limitation of the C oulom b-gas approach.

The charge neutrality conditions (67) (68) for a onepoint function pick up the coe cient of the corresponding Ishibashi state from a Cardy state, since = r_{rs} is the only boundary charge which gives a non-vanishing term. This agrees with our understanding that the coe cients of Cardy's state are essentially the one-point coupling constants of bulk (closed string vertex) operators to the boundary (brane) [7,8]. Once one-point coupling constants are known, it is in principle possible to compute boundary multipoint functions since they reduce to one point functions after repeated use of bulk OPEs (Fig 2a). In particular, we may start such a procedure from the farthest point from the boundary (in the radial ordering sense), approaching the boundary by perform ing OPE with the farthest remaining point one by one. Due to naturality of CFT, such OPEs are translated into the fusions of operators,

.

$$\begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix};$$

$$\begin{bmatrix} 1 \\ 3 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix};$$

$$\begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix};$$

$$\begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix};$$

$$\begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix};$$

$$\begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix};$$

$$\begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix};$$

$$\begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix};$$

$$\begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix};$$

$$\begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix};$$

$$\begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix};$$

which de ne a conformal block with no subchains (Fig 2b). As the fusion of primary operators is equivalent to the map (55) between CBFSs restricted to BRST subspaces [14], the conform alblock of Fig 2b is represented by our xed boundary-charge correlator (61) with corresponding to $[j_{p-1}]$. The Ishibashi state lhj_{p-1} jacts as a liter (or a half mirror) transmitting only the V irasoro representation $[j_{p-1}]$. This property of Ishibashi states is captured by boundary charges and charge neutrality.

Due to the absence of internal channels, two point functions on the disk (or half plane) are com pletely determ ined by a boundary state, apart from the norm alisation of conform alblocks. In the case of p-point functions with 3, however, states in the internal channels $j_{i < p}$ 1 α cannot in general be determ ined uniquely even if the state in the boundary channel j_{p-1} is xed (an example is the spin three point function of the Ising model with $\dot{p} = -$, where j_1 can be I or). Corresponding to this, the contours of (61) with p 3 may be deformed in several di erent ways to give independent convergent functions which are expected to reproduce conform alblocks with di erent internal states. The relative coe cients of such conform alblocks in a boundary correlation function cannot be determined by the boundary state (as these conform al blocks belong to the same Ishibashi state) but should be constrained by inform ation of the bulk.




FIG.2. OPE and conform all blocks of boundary p-point functions. Repeating OPEs in the bulk, the boundary correlation function reduces to one point functions (a). This can also be seen as conform all blocks with internal channels $j_{i and boundary channel <math>j_{p \ 1}$ (b).

IV.ISING MODEL

We shall illustrate the method presented in the previous section in the example of the critical Ising model. Before starting actual calculations we notice from the discussions of the previous section that the non-unitary sector and the boundary states with charges outside the K ac table do not contribute to correlation functions. Neglecting such unnecessary terms in (51) - (53), for the bra-boundary states of the Ising model we have,

$$\begin{array}{l} h\Gamma j & 2 \\ & {}^{1} \left({}_{0} hB \left({}_{1;1} \right) j + {}_{U} hB \left({}_{2;3} \right) j \right. \\ & + {}_{U} hB \left({}_{2;1} \right) j + {}_{U} hB \left({}_{1;3} \right) j \right. \\ & + 2 \\ & {}^{3=4} \left({}_{U} hB \left({}_{1:2} \right) j + {}_{U} hB \left({}_{2:2} \right) j \right);$$
 (95)

$$h \sim j \quad 2^{-1} (_{U} hB (_{1;1})j + _{U} hB (_{2;3})j + _{U} hB (_{2;1})j + _{U} hB (_{1;3})j \\ 2^{-3=4} (_{U} hB (_{1;2})j + _{U} hB (_{2;2})j); \quad (96)$$

h~j 2
$$^{1=2}$$
 ($_{U}$ hB ($_{1;1}$) j+ $_{U}$ hB ($_{2;3}$) j
 $_{U}$ hB ($_{2;1}$) j $_{U}$ hB ($_{1;3}$) j: (97)

Since these are linear combinations of xed boundarycharge states $_{\rm U}$ hB ($_{\rm r;s}$); the correlation functions on the disk with physical boundary conditions I, ~ and ~ are given by linear combinations of xed boundary-charge correlators (61). U sing the global conform altransform ation explained in the last section, we shall obtain correlation functions on the half plane and com pare them with existing results.

A.One point functions of spin and energy operators

Let us rst consider the spin one point function. We may choose either = $_{1;2}$ or = $_{2;2}$. Let, for definiteness, (r;s) = (1;2) in the holom orphic part and in order to use the result of the last section, (r;s) = (2;2) in the antiholom orphic part. From (70) we immediately notice that only the state $_{\rm U}$ hB ($_{1;2}$) j contributes to the one point function, all other states giving vanishing correlators. For the boundary condition Γ , the one point function on the disk is

$$hI'j (z;z) Di$$

$$= hI'jV_{1;2} (z)V_{2;2} (z) D;0; _0 i_U$$

$$= 2^{-3=4} B (_{-1;2}) JV_{1;2} (z)V_{2;2} (z) D;0; _0 i_U$$

$$= 2^{-3=4} (1 - zz)^{-1=8} :$$
(98)

P roperly norm alised one point function is then,

$$\frac{hfj(z;z)ji}{hfji} = 2^{1=4} (1 zz)^{1=8} :$$
 (99)

This is mapped on to the half plane by using the conformal transformation (91), as

h (w; w)
$$i_r = h$$
 (y) $i_r = 2^{1-4}$ (2y) $^{1-8}$; (100)

where y is the distance from the boundary. Likewise, boundary spin correlation functions for the conditions ~ and ~ are obtained simply by picking up the coe cients of $_{U}$ hB ($_{1,2}$) jin (96) and (97), and are norm alised using h~ $\mathfrak{Pi} = 1=2$ and h~ $\mathfrak{Pi} = 1=2$. On the half plane, they are

h (w; w) i = h (y) i =
$$2^{1-4}$$
 (2y) $^{1-8}$; (101)

and

h (w; w)
$$i_{-} = h$$
 (y) $i_{-} = 0$: (102)

Hence, I and ~ are indeed the xed (up and down) and ~ is the free boundary condition, as is stated in [6,7]. In our C oulom b-gas form alism the relation between one point functions and the coe cients of Ishibashi states is explained by the neutrality of charges.

Putting, say, (r;s) = (2;1) and (r;s) = (1;3), the energy one point function is obtained similarly. On the half plane we have,

h
$$(y)_{i} = h (y)_{i} = (2y)^{-1};$$
 (103)

h (y)i =
$$(2y)^{-1}$$
: (104)

B. Spin two point function

Next, let us consider the spin two point function. Since = $_{1;2}$, we can use the result of Subsec C of the preceding section. There are only two values of boundary charges, $_{1;1}$ and $_{1;3}$, which give non-trivial contributions to the correlator. The two corresponding states $_{U}$ hB ($_{1;1}$) jand $_{U}$ hB ($_{1;3}$) jgive rise to the two conform all blocks I_I and I_{II}, respectively, and the correlation function is a linear combination of these conform all blocks with coe cients given by Cardy's states (95) - (97). Then,

$$\frac{hTj(z_1;z_1)(z_2;z_2)Di}{hTDi} = \frac{hrj(z_1;z_1)(z_2;z_2)Di}{hrDi}$$
$$= I_T + I_{TT};$$
(105)

$$\frac{h \sim j (z_1; z_1) (z_2; z_2) j j j}{h \sim j j} = I_I I_{II};$$
(106)

where the actual form s of I_I and I_{II} are given by (83) and (87), with a = 1=8, b = 3=8 and ² = 3=4. In this case the hypergeom etric functions reduce to,

$$_{2}F_{1}\left(\begin{array}{c}\frac{1}{4};\frac{1}{4};\frac{1}{2};\end{array}\right) = \frac{p}{\frac{1+p}{1+\frac{1}{2}}};$$
 (107)

$$_{2}F_{1}(\frac{3}{4};\frac{1}{4};\frac{3}{2};) = \frac{P(2(1-P(1-1)))}{P(1-1)};$$
 (108)

Using (107), (108) and the conformal transformation (88), we nd the two point functions on the half plane,

h (w₁;w₁) (w₂;w₂)i_r
= h (w₁;w₁) (w₂;w₂)i.
=
$$\frac{N}{P_{\overline{2}}} \frac{(w_1 w_1)(w_2 w_2)}{(w_1 w_2)(w_1 w_2)(w_1 w_2)(w_1 w_2)(w_1 w_2)}$$
N_I $\frac{q}{p} \frac{1}{1} + 1 + N_{II} \frac{q}{p} \frac{p}{1} \frac{1}{1} ; (109)$

where N[•] = $(1=4)^2$ = (1=2). Studying the behaviours away from the boundary (we accept the convention that two point functions of bulk operators are normalised as h_i(w₁)_j(w₂)i = _{ij}(w₁ w₂)^{2h_i}) and comparing the leading terms with the OPE coe cients of [8], the normalisation of the conformal blocks are xed as N_I = N_{II} = $(1=2)=(1=4)^2$.

This result was obtained long time ago [13], by solving a di erential equation to nd the two conformal blocks I_I , I_{II} , and then xing the coe cients by considering asymptotic behaviours of the correlation function. The (relative) coe cients of the conformal blocks are now attributed to the coe cients of C ardy's states in our C culom b-gas approach, although we have used the asymptotic behaviours to x the normalisation of each conform alblock.

C.Energy two point function

Finally we derive the energy two point function in our form alism. As $= _{2;1}$, the calculation is parallel to the case of the spin two point function. From the charge neutrality condition we nd that non-vanishing correlators arise from the states $_{\rm U}$ hB ($_{1;1}$) j and $_{\rm U}$ hB ($_{3;1}$) j. How-ever, as none of the boundary states (95) - (97) contain $_{\rm U}$ hB ($_{3;1}$) j only $_{\rm U}$ hB ($_{1;1}$) j gives non-trivial contribution to the correlation function. Hence, the energy two point function does not depend on boundary conditions. A fler a simple calculation we nd, on the half plane,

h
$$(w_1; w_1)$$
 $(w_2; w_2) i_{T; \sim; \sim}$
= N $\frac{(w_1 \ w_1)(w_2 \ w_2)}{(w_1 \ w_2)(w_1 \ w_2)(w_1 \ w_2)(w_1 \ w_2)(w_1 \ w_2)}$
 $\frac{(1=3)^2}{(2=3)}$ F (2; 1=3; 2=3;): (111)

The norm alisation constant N is determined as N = $(2=3)=(1=3)^2$, by considering the o-boundary behaviour. The hypergeom etric function turns out to be an

algebraic function F (2; 1=3; 2=3;) = 1 + ², and using the coordinates $w_i = x_i + iy_i$, $w_i = w_i = x_i$ iy_i (x_i, y_i 2 R), the correlation function (111) is written as

h
$$(x_1; y_1)$$
 $(x_2; y_2)i_{T_{i}\sim i^{\sim}}$

$$= \frac{4y_1y_2}{[(x_1 \quad x_2)^2 + (y_1 \quad y_2)^2][(x_1 \quad x_2)^2 + (y_1 + y_2)^2]}$$

$$+ \frac{1}{4y_1y_2}:$$
(112)

This agrees with the result in [13].

V.SUMMARY

In this paper we have described a novelm ethod to calculate correlation functions of two dimensional CFT on the disk and on the halfplane. We have used the free-eld construction of boundary states developed in [18], and derived boundary correlation functions for the boundary states classified by C ardy's method. The key feature of our form alism is the neutrality of bulk and boundary charges, which associates the coe cients in C ardy's boundary states directly with the linear com binations of conform alblocks. Thus we could unify the two important pieces of boundary CFT, namely, boundary correlation functions [13] and consistent boundary states of C ardy [6], by using the C oulom b-gas picture. We have checked the form alism in the Ising model, and shown that our m ethod reproduces the known results.

C ardy's classi cation of boundary states has been generalised by Lewellen [8] and Pradisi, Sagnotti, Stanev [9] beyond the diagonal models, and the Coulomb-gas technique is also known to be applicable to more general CFTs, such as W ZNW models [29] and CFTs with W – algebra [30{32]. We therefore expect that the method discussed in this paperm ay be applied to such CFTs relatively easily. In particular, from a string theory point of view, application to W ZNW theories seem s to be quite fruitful since it would provide an alternative method to nd correlation functions with D-branes on a group manifold. We shall discuss such issues in separate publications [33].

ACKNOW LEDGMENTS

I thank Dr. John W heater for useful discussions and constant encouragement. I am also grateful to Prof. John Cardy, Dr. M atthias G aberdiel and Dr. Ian K ogan for helpful discussions.

- [L] A. Cappelli, C. Itzykson and J.B. Zuber, Nucl. Phys. B 280 (1987) 445.
- [2] A. Cappelli, C. Itzykson and J. B. Zuber, Commun. M ath. Phys. 113 (1987) 1.
- [3] T.Gannon, Rev.Math.Phys.12 (2000) 739.
- [4] T.Gannon, Annales Poincare Phys. Theor. 65 (1996) 15.
- [5] T.Gannon, Nucl.Phys.B 491 (1997) 659.
- [6] J.L.Cardy, Nucl. Phys. B 324 (1989) 581.
- [7] J.L.C ardy and D.C.Lewellen, Phys.Lett.B 259 (1991) 274.
- [8] D.C.Lewellen, Nucl. Phys. B 372 (1992) 654.
- [9] G. Pradisi, A. Sagnotti and Y. S. Stanev, Phys. Lett. B 381 (1996) 97.
- [10] I.Runkel, Nucl. Phys. B 549 (1999) 563; ibid. 579 (2000) 561.
- [11] R.E.Behrend, P.A.Pearce, V.B.Petkova and J.B.Zuber, Phys. Lett. B 444 (1998) 163; Nucl. Phys. B 570 (2000) 525 Nucl. Phys. B 579 (2000) 707].
- [12] J. Fuchs and C. Schweigert, Nucl. Phys. B 558 (1999) 419; ibid. 568 (2000) 543.
- [13] J.L.Cardy, Nucl. Phys. B 240 (1984) 514.
- [14] G.Felder, Nucl. Phys. B 317 (1989) 215 Erratum -ibid. B 324 (1989) 548].
- [15] G.Felder and R.Silvotti, Phys.Lett. B 231 (1989) 411; Commun.M ath.Phys.144 (1992) 17.
- [16] P. Di Vecchia, F. Pezzella, M. Frau, K. Homfeck, A.Lerda and A.Sciuto, Nucl. Phys. B 322 (1989) 317.
- [17] M. Frau, A. Lerda, J. G. M cCarthy and S. Sciuto, Phys. Lett. B 228 (1989) 205; Nucl. Phys. B 338 (1990) 415.
- [18] S.Kawai, Nucl. Phys. B 630 (2002) 203.
- [19] J.Schulze, Nucl. Phys. B 489 (1997) 580.
- [20] S.D.Mathur, Nucl. Phys. B 369 (1992) 433.
- [21] V. S. Dotsenko and V. A. Fateev, Nucl. Phys. B 240 (1984) 312; ibid. 251 (1985) 691.
- [22] C.Schweigert, J.Fuchs and J.W alcher, \Conformal eld theory, boundary conditions and applications to string theory," [hep-th/0011109].
- [23] P.D iFrancesco, P.M athieu and D. Senechal, \Conform al eld theory," Springer, New York (1997).
- [24] N. Ishibashi, Mod. Phys. Lett. A 4 (1989) 251.
- [25] C.G.Callan, C.Lovelace, C.R.Nappiand S.A.Yost, Nucl. Phys. B 293 (1987) 83.
- [26] J. Polchinski and Y. Cai, Nucl. Phys. B 296 (1988) 91.
- [27] N.Ishibashiand T.Onogi, Nucl. Phys. B 318 (1989) 239.
- [28] A.Rocha-Caridi, \Vacuum vector representations of the Virasoro algebra," in Vertex operators in mathematics and physics, J. Lepowsky, S. Mandelstam and I. M. Singer (eds.), Springer, Berlin (1984), p. 451.
- [29] A. Gerasim ov, A. Morozov, M. Olshanetsky, A. Marshakov and S. L. Shatashvili, Int. J. Mod. Phys. A 5 (1990) 2495.
- [30] P.Bouwknegt and K.Schoutens, Phys. Rept. 223 (1993) 183.
- [31] V.A.Fateev and S.L.Lukyanov, Int.J.M od.Phys.A 3 (1988) 507; JETP 67 (1988) 447.
- [32] S.M izoguchi, Int. J.M od. Phys. A 6 (1991) 133.
- [33] A.Caldeira, S.K awai and J.F.W heater, in preparation.