

A Matrix Model for a Class of Fractional Quantum Hall States

A. Jellal^{1y}, E. H. Saidi², H. B. Geyer³ and R. A. Romer⁴¹ Institut für Physik Technische Universität, D-09107, Chemnitz, Germany² Lab/UFR, High Energy Physics, Physics Department, Mohammed VI University, Av. Ibn Battouta, P.O. Box. 1014, Rabat, Morocco³ Institute for Theoretical Physics, University of Stellenbosch, Private Bag X1, Matieland 7602, South Africa⁴ Department of Physics, University of Warwick, Coventry CV4 7AL, UK

(Received July 18, 2002)

KEYWORDS: Non-commutative Chern-Simons, Matrix Model Theory, Fractional Quantum Hall (FQH) Fluids

1. Introduction

Recently, Susskind¹⁾ showed that an Abelian non-commutative Chern-Simons theory at level k is actually equivalent to Laughlin theory.²⁾

$$S = \frac{k}{4} \int d^3y \, A \star \partial A + \frac{2}{3} A \star A \star A \quad (1)$$

where the star-product is the usual Moyal product with parameter θ . Therefore, he obtained the filling factor

$$\nu = \frac{1}{k} \quad (2)$$

He also pointed out that the above theory can be formulated in terms of a matrix model involving classical Hermitian matrix variables $A_0; X^i, i = 1; 2$. The Lagrangian for the matrix theory is

$$L = B \text{Tr}_{ij} (X^{-i} + iA_0; X^j) X^j + 2 A_0 \quad (3)$$

B is the magnetic field. The equation of motion for the coordinate A_0 (Gauss law constraint) is

$$[X^1; X^2] = i \quad (4)$$

which can only be solved if the matrices are in finite dimensional. This corresponds to an finite number of electrons on an finite plane.

For a finite system, Polychronakos³⁾ has introduced an additional set of bosonic degrees of freedom $m, m = 1; 2; \dots; M$, such that $\nu = (1; \dots; M)$,

$$L = \sum_y (i - A_0) : \quad (5)$$

Considering $L + L$, Polychronakos³⁾ found a quantum correction to Susskind's filling factor such that

$$\nu = \frac{1}{k+1} \quad (6)$$

In this case, the Gauss law constraint becomes

$$[X^1; X^2] = i \left(1 - \frac{1}{k+1} \right) \quad (7)$$

Later Helleman and Van Raamsdonk⁴⁾ built the cor-

responding wavefunctions for $L + L$,

$$\psi_k = \frac{1}{n!} \epsilon_{i_1 \dots i_n} (y_{i_1})_{i_1} (y_{i_2})_{i_2} \dots (y_{i_n})_{i_n} \quad (8)$$

where $\epsilon_{i_1 \dots i_n}$ is the fully antisymmetric tensor. These are similar to Laughlin's wavefunction.²⁾ Subsequently, three of us generalised⁵⁾ the above results to any filling factor given by

$$\nu_{k_1 k_2} = \frac{1}{k_1} + \frac{1}{k_2}; \quad k_2 > k_1 \quad (9)$$

In what follows, we propose a matrix model to describe such FQH states that are not of Laughlin type.

2. $\nu_{k_1 k_2}$ fractional quantum Hall states

Although the $\nu = \frac{2}{5}$ FQH state is not of the Laughlin type, it shares some basic features of Laughlin fluids. The point is that from the standard definition of the filling factor $\nu = \frac{N}{N}$, the state $\nu = \frac{2}{5}$ can naively be thought of as corresponding to $\nu = \frac{N}{N}$ where the number N of flux quanta is given by a fractional amount of the electron number; that is

$$N = (3 \frac{1}{2})N : \quad (10)$$

In fact this way of viewing things reflects the original idea of a hierarchical construction of FQH states for general filling factor $\frac{p}{q}$. In Haldane's hierarchy,⁶⁾ the elements of the series

$$\nu_{p_1 p_2} = \frac{p_2}{p_1 p_2 - 1} \quad (11)$$

correspond to taking N as given by a specific rational factor of the electron number, i.e.,

$$N = (p_1 \frac{1}{p_2})N : \quad (12)$$

Upon setting

$$k_1 = p_1; \quad k_2 = k_1 (k_1 p_2 - 1) \quad rk_1 \quad (13)$$

we have $\nu_{p_1 p_2} = \nu_{k_1 k_2}$. For $\nu = \frac{2}{5}$, e.g.,

$$\nu = \frac{2}{5} = \frac{1}{3} + \frac{1}{15} \quad (14)$$

3. Matrix model analysis

To describe FQH fluids at $\nu_{k_1 k_2}$, we consider the following action for a system of $N = N_1 + N_2$ particles⁵⁾

$$S = \int dt \sum_{i=1}^N \left[\frac{1}{2} \dot{\mathbf{r}}_i^2 + \frac{1}{4} \text{Tr} \left(\mathbf{Z}_i^\dagger \mathbf{D} \mathbf{Z}_i \right) + \frac{1}{2} \mathbf{Z}_i^\dagger \mathbf{Z}_i + 2 \sum_{a=1}^{N_1} \mathbf{A}_{0,i} \cdot \mathbf{r}_i + \frac{1}{2} \sum_{a=1}^{N_1} \mathbf{r}_i \cdot \mathbf{A}_{0,i} + \frac{1}{2} \sum_{a=1}^{N_1} \mathbf{r}_i \cdot \mathbf{A}_{0,i} + \frac{1}{2} \sum_{a=1}^{N_1} \mathbf{r}_i \cdot \mathbf{A}_{0,i} \right] \quad (15)$$

where $\mathbf{1} = \mathbf{r}_1, \dots, \mathbf{r}_{N_1}, \mathbf{1} = \mathbf{a}; \mathbf{b} = \mathbf{r}_{N_1+1}, \dots, \mathbf{r}_N, \mathbf{Z}_1 = \mathbf{X}_1 + i\mathbf{X}_1^2$ and $\mathbf{A}_{0,i}$ the gauge for the i th particle. The $J^{(1)}$ and $J_{aa}^{(2)}$ currents (Gauss law constraints) read as

$$\begin{aligned} J^{(1)} &= [\mathbf{Z}_1; \mathbf{Z}_1] + \frac{1}{2k_1} \sum_{a=1}^{N_1} \mathbf{r}_a \cdot \mathbf{A}_{0,a} J_0^{(1)} \\ J_{aa}^{(2)} &= [\mathbf{Z}_2; \mathbf{Z}_2]_{aa} + \frac{1}{2k_2} \sum_{a=1}^{N_2} \mathbf{r}_a \cdot \mathbf{A}_{0,a} J_0^{(2)} \end{aligned} \quad (16)$$

where the two $U(1)$ charge operators $J_0^{(1)}$ and $J_0^{(2)}$ are

$$J_0^{(1)} = J_0^{(2)} = J_0 = \sum_{a=1}^{N_1} \mathbf{r}_a \cdot \mathbf{A}_{0,a} : \quad (17)$$

The wavefunctions ψ describing the $(N_1 + N_2)$ system of electrons on the non-commutative plane \mathbb{R}^2 with filling factor $\nu_{k_1 k_2}$ should obey the constraint⁵⁾

$$J_0 \psi = \frac{N}{k_1 k_2} \psi : \quad (18)$$

Once we know the fundamental state $\psi_{k_1 k_2}^{(0)}$, excitations are immediately determined by applying the usual rules. Upon recalling the coordinate operators as

$$\mathbf{Z}_1 = \frac{1}{2} \mathbf{r}^+; \quad \mathbf{Z}_{2aa} = \frac{1}{2} \mathbf{s}_{aa}^+ \quad (19)$$

the total Hamiltonian H may be treated as the sum of a free part given by

$$H_0 = \frac{1}{2} (2N_1 + 2N_2 + N_1^2 + N_2^2) \quad (20)$$

where $N_1 = \sum_{a=1}^{N_1} \mathbf{r}_a^\dagger \mathbf{r}_a$ and $N_2 = \sum_{a=1}^{N_2} \mathbf{s}_{ab}^\dagger \mathbf{s}_{ba}$ are the operator numbers counting the N_1 and N_2 particles respectively, and an interacting part

$$H_{\text{int}} = \sum_a \mathbf{r}_a^\dagger \mathbf{s}_{ab} + \text{h.c.} \quad (21)$$

describing couplings between the two sectors.⁵⁾ The creation and annihilation operators $\mathbf{r}, \mathbf{s}_{aa}$, and \mathbf{s}_a satisfy the Heisenberg algebra

$$\begin{aligned} [\mathbf{r}_a, \mathbf{r}_b^\dagger] &= \delta_{ab}; \quad [\mathbf{s}_a, \mathbf{s}_b^\dagger] = \delta_{ab} \\ [\mathbf{r}_a, \mathbf{s}_b] &= 0; \quad [\mathbf{s}_a, \mathbf{r}_b^\dagger] = 0 \end{aligned} \quad (22)$$

all others are given by commuting relations.

4. Wavefunctions

A way to build the spectrum of the Hamiltonian H_0 is given by help of the special condensate operators

$$\mathbf{A}_a^{+(n,m)} = \mathbf{s}_a^{+n} \mathbf{r}_a^{+m} : \quad (23)$$

The wavefunctions for the vacuum $|0\rangle$ of H_0 read as

$$\psi = \prod_{j=1}^N \mathbf{O}^{(j)} |0\rangle \quad (24)$$

where the $\mathbf{O}^{(j)}$'s are building blocks and given by

$$\mathbf{O}^{(j)} = \mathbf{A}_a^{+(jN_2+1)} \cdots \mathbf{A}_a^{+(j+1)N_2} \mathbf{A}_a^{+(1,j)} \cdots \mathbf{A}_a^{+(j+1)N_2} : \quad (25)$$

The corresponding energy spectrum $E_c(k_1 k_2)$ is

$$E_c = k_1 \frac{(N_2-1)(N_2-2)}{2} + \frac{(N_1-1)(N_1-2)}{2} N_2 + \frac{N_1+N_2}{2} : \quad (26)$$

Note that for large value of N_1 and N_2 ($N_1 = rN_2$), $E_c(k_1 k_2)$ behaves quadratically in N_2 ,

$$E_c(k_1 k_2) \sim \frac{k_2}{2} N_2^2 : \quad (27)$$

This energy relation is less than the total energy $E_d(k_1)$ of the decoupled configuration $(j_1; \mathbf{v}_{k_1}) \rightarrow (j_2; \mathbf{v}_{k_2})$:

$$E_d(k_1) = E_{\frac{1}{k_1}} + E_{\frac{1}{k_2}} = \frac{k_2(r+1)}{2} N_2^2 : \quad (28)$$

Therefore, we have the following relation

$$E_d = (r+1) E_c : \quad (29)$$

For the example of the FQH state at $\nu = \frac{2}{5}$, the energy of the decoupled representation reads as

$$E = \frac{1}{3} + E_{\frac{1}{15}} = 45 N_2^2 \quad (30)$$

while that of the interacting one is

$$E_c = \frac{2}{5} = \frac{15}{2} N_2^2 \quad (31)$$

leading to

$$E = \frac{1}{3} + E_{\frac{1}{15}} = 6 E_c = \frac{2}{5} : \quad (32)$$

5. Conclusion

We have developed a matrix model for FQH states at filling factor $\nu_{k_1 k_2}$ going beyond the Laughlin theory. To illustrate our idea, we have considered an FQH system of a finite number $N = (N_1 + N_2)$ of electrons with filling factor $\nu_{k_1 k_2} = \frac{p_2}{p_1 p_2 - 1}$; p_1 is an odd integer and p_2 is an even integer. The $\frac{p_2}{p_1 p_2 - 1}$ series corresponds just to the level two of the Haldane hierarchy; it recovers the Laughlin series $\frac{1}{p_1}$ by going to the limit p_2 large and contains several observable FQH states such

$$as = \frac{2}{3}; \frac{2}{5}; \quad .$$

1) L. Susskind: hep-th/0101029.

2) R. B. Laughlin: Phys. Rev. Lett. 50 1395 (1983).

3) A. P. Polychronakos: JHEP 0104 (2001) 011.

4) S. Hellerman and M. Van Raamsdonk: JHEP 0110 (2001) 039.

5) A. Jella, E. H. Saidi and H. B. Geyer: hep-th/0204248.

6) F. D. M. Haldane: Phys. Rev. Lett. 51 (1983) 605.