

Higher Toda Mechanics and Spectral Curves

Liu Zhao, Wangyun Liu^y

Institute of Modern Physics,
Northwest University, Xi'an 710069, China

December 16, 2021

Abstract

For each one of the Lie algebras \mathfrak{gl}_n and \mathfrak{sl}_n , we constructed a family of integrable generalizations of the Toda chains characterized by two integers m_+ and m_- . The Lax matrices and the equations of motion are given explicitly, and the integrals of motion can be calculated in terms of the trace of powers of the Lax matrix L . For the case of $m_+ = m_-$, we find a symmetric reduction for each generalized Toda chain we found, and the solution to the initial value problems of the reduced systems is outlined. We also studied the spectral curves of the periodic $(m_+; m_-)$ -Toda chains, which turns out to be very different for different pairs of m_+ and m_- . Finally we also obtained the nonabelian generalizations of the $(m_+; m_-)$ -Toda chains in explicit form.

1 Introduction

The very first research interests in classical many body systems can be traced back to the dates from Galileo to Newton, when the classical mechanics for the celestial bodies in the solar system was the major physical system of concern. However, it turns out that it is extremely difficult, or even impossible in most cases, to get exact solutions to the equations of motion for a given many body mechanical system. Perhaps the major breakthrough in this direction was made since the late 1960's and the early 1970's, when a number of exactly solvable many body systems in one dimension were discovered and solved by means of the inverse scattering method, among which the classical Toda chains [1, 2], the Calogero-Moser systems [3, 4]

e-mail: lzhaonw@nwnu.edu.cn

^ye-mail: wyl@phy.nwnu.edu.cn

and the Ruijsenaars-Schneider models [5, 6] are the most famous and important examples.

The recent surge of interests in these classical mechanical systems is mostly due to the work of Donagi and Witten [7] and the followers [8], which revealed a remarkable and unexpected relationship between such solvable mechanical systems and certain $N = 2$ supersymmetric gauge theories. To be more specific, the exact integrability of the above many body systems are characterized by their Lax matrices, whose spectral curves in turn are identified with the so-called Seiberg-Witten curves [9] of the $N = 2$ SUSY gauge theories, while the complete information about the moduli space parameters and the prepotentials of the $N = 2$ SUSY gauge theories are encoded in the equation defining the Seiberg-Witten curves.

More recently, Dijkgraaf and Vafa made another important progress [10] in finding nonperturbative information about the moduli structure of SUSY gauge theories. They found a remarkable correspondence between the nonperturbative chiral effective superpotentials and the effective potentials of certain matrix integrals in the planar diagram limit. Their results fully confirmed the earlier results of Dorey [11], who identified the same chiral superpotentials with the values of the Hamiltonians of certain integrable classical many body systems in the stationary point configuration. Although this last step toward the nonperturbative treatment of $N = 1$ SUSY gauge theories is mathematically less complete compared to the corresponding Seiberg-Witten theory for $N = 2$ cases, it has caught quite massive attention because the $N = 1$ SUSY theories are considered to be more realistic than the $N = 2$ ones.

So far, the classical many body systems known to play certain role in the studies in both the $N = 2$ and $N = 1$ gauge theories is restricted within the above mentioned three families, i.e. the classical Toda chains, the Calogero-Moser systems and the Ruijsenaars-Schneider models. It seems an interesting question to ask whether there exist other pairs of SUSY gauge theory and classical many body systems and whether the known classes of SUSY gauge theories may be related to other many body systems besides the ones just listed and vice versa. In this paper we shall try to construct some novel exactly solvable many body systems which fall in the family of Toda mechanics, but with off-diagonal extensions (the meaning of off-diagonal extension will be clear in the main context), and study their spectral curves. We hope some day such systems will be proved useful in describing certain SUSY gauge theory in a nonperturbative way, following either the Seiberg-Witten or the Dijkgraaf-Vafa routes. But now we shall mostly concentrate on the integrable aspect of these extended Toda mechanical systems.

This paper is organized as follows. In Section 2 we give the Lax matrix construction of the extended Toda mechanics, referred to as $(m_+; m_-)$ -Toda chains. The construction applies for finite chains, infinite chains as well as periodic chains. The Liouville integrability of the infinite $(m_+; m_-)$ -Toda chains is discussed via the

construction for integrals of motion and the definition for the fundamental Poisson structure. For the case of $m_+ = m_-$, we show that there is a symmetric reduction to the $(m_+; m_-)$ -Toda chains and the solution to their initial value problems are described. Section 3 is devoted to a brief introduction for the spectral curves corresponding to the extended periodic chains. It is shown that there are two different ways for describing the spectral curves, each has a correspondence in the cases of standard Toda chains. In Section 4, we generalize our construction for the extended Toda chains to the nonabelian cases and explicit equations of motion for generic nonabelian $(m_+; m_-)$ -Toda chains are obtained. The paper is then concluded in Section 5, where we present some discussions and speculations for further studies.

2 Construction of the system

In this section we shall mainly work on the construction of the extended Toda mechanics. We shall follow the route of Lax integrability during the construction, give necessary explanation for the construction and briefly discuss the Liouville integrability of the systems. The systems we shall be considering can be divided into three classes: the infinite chains, the finite chains and the periodic chains. It is the periodic chains which have nontrivial spectral curves. We leave the detailed study on the spectral curve to the next section.

2.1 Infinite chain

Let us first introduce some auxiliary notations. We shall need an infinite dimensional vector space with basis vectors $f_{ji}; i \in \mathbb{Z}$ and their duals, $f_{ji}^*; i \in \mathbb{Z}$. The inner product between the basis vectors and their duals is specified by

$$f_{ji}^* f_{ji} = \delta_{ij}.$$

Let us remind that such vectors are introduced purely for notational conveniences, they have nothing to do with the Fock space basis vectors for the Heisenberg algebras.

With the above notations in mind, we are now ready to introduce the Lax matrices for the infinite chain of extended Toda mechanics. By Lax matrices we mean the pair $(L; M)$ of matrices appearing in the Lax equation

$$\dot{L} = [M; L]; \tag{1}$$

where, as usual, a dot over the mechanical variables means taking a time derivative. For the case of infinite chains we shall introduce, the matrix L is a formal infinite

matrix with a strip form, i.e. all nonzero entries of L are at positions close to the main diagonal. More concretely, L may be written as follows,

$$L = L^{(0)} + L^{(>)} + L^{(<)}; \quad (2)$$

$$L^{(0)} = \sum_{i \in \mathbb{Z}} q_i j_i h_i j; \quad (3)$$

$$L^{(>)} = \sum_{i \in \mathbb{Z}} \sum_{k=1}^{\infty} a_i^{(k)} j_{i+k} h_i + k j; \quad (4)$$

$$L^{(<)} = \sum_{i \in \mathbb{Z}} \sum_{k=1}^{\infty} c_i^{(k)} j_i + k i h_j; \quad (5)$$

Each L matrix is characterized with a pair of fixed integers (m_+, m_-) , with $m_+ + m_- + 1$ being the width of the nonzero strip in the L matrix. We shall see that for $m_+ = m_- = 1$, the matrix L would yield the standard infinite Toda chain. Therefore we may call the system with generic values of m_+ an (m_+, m_-) -th extended Toda chain. In what follows, we shall always assume that $m_+ = m_-$, without loss of generality.

Now let us proceed to specify the M matrix which is needed to obtain the equation of motion for a Lax integrable system via the Lax equation (1). The M matrix for the present case is simply given as

$$M = L^{(>)} - L^{(<)}; \quad (6)$$

Therefore, the right hand side of the Lax equation may be written as

$$[M; L] = [L^{(>)} - L^{(<)}; L^{(0)}] + 2[L^{(>)}; L^{(<)}];$$

Straightforward calculations yield the following results,

$$[L^{(>)}; L^{(0)}] = \sum_{i \in \mathbb{Z}} \sum_{k=1}^{\infty} (a_i^{(k)} q_{i+k} - q_i a_i^{(k)}) j_i h_i + k j;$$

$$[L^{(<)}; L^{(0)}] = \sum_{i \in \mathbb{Z}} \sum_{k=1}^{\infty} (q_{i+k} c_i^{(k)} - c_i^{(k)} q_i) j_i + k i h_j;$$

$$[L^{(>)}; L^{(<)}] = \sum_{i \in \mathbb{Z}} \sum_{k=1}^{\infty} \sum_{k^0=1}^{\infty} a_i^{(k)} c_{i-k}^{(k^0)} - c_{i-k}^{(k^0)} a_i^{(k)} j_i h_j$$

$$+ \sum_{i \in \mathbb{Z}} \sum_{k=1}^{\infty} \sum_{k^0=1}^{\infty} a_i^{(k+k^0)} c_{i+k}^{(k^0)} - c_{i-k^0}^{(k^0)} a_{i-k^0}^{(k+k^0)} A j_i h_i + k j$$

$$+ \sum_{i \in \mathbb{Z}} \sum_{k=1}^{\infty} \sum_{k^0=1}^{\infty} a_{i+k}^{(k^0)} c_i^{(k+k^0)} - c_{i-k^0}^{(k+k^0)} a_{i-k^0}^{(k^0)} j_i + k i h_j;$$

Therefore, the equations of motion obtained from the Lax equation read

$$q_i = 2 \sum_{k=1}^{\infty} a_i^{(k)} c_i^{(k)} - c_{i-k}^{(k)} a_{i-k}^{(k)}; \quad (7)$$

$$a_i^{(k)} = a_i^{(k)} q_{i+k} - q_i a_i^{(k)} + 2 \sum_{k^0=1}^{m \text{ in } (m \times k \text{ in } m)} a_i^{(k+k^0)} c_{i+k}^{(k^0)} - c_{i-k^0}^{(k^0)} a_{i-k^0}^{(k+k^0)}; \quad (1 \leq k \leq m+1) \quad (8)$$

$$c_i^{(k)} = q_{i+k} c_i^{(k)} - c_i^{(k)} q_i + 2 \sum_{k^0=1}^{m \times k} a_{i+k}^{(k^0)} c_i^{(k+k^0)} - c_{i-k^0}^{(k^0)} a_{i-k^0}^{(k+k^0)}; \quad (1 \leq k \leq m+1) \quad (9)$$

$$a_i^{(m+1)} = a_i^{(m+1)} q_{i+m+1} - q_i a_i^{(m+1)}; \quad (10)$$

$$c_i^{(m)} = q_{i+m} c_i^{(m)} - c_i^{(m)} q_i; \quad (11)$$

Notice that while writing the above equations we have been very careful in keeping the order of the variables in their essential positions, with the hope of generalizing to the case of nonabelian variables in mind. However, at present, let us proceed with the view point that all variables $q_i, a_i^{(k)}$ and $c_i^{(k)}$ are usual abelian variables. Then it is easy to see that the last pair of the equations of motion, eqs.(10), (11) can be integrated explicitly, yielding

$$a_i^{(m+1)} = \exp(q_{i+m+1} - q_i); \quad c_i^{(m)} = \exp(q_{i+m} - q_i); \quad (12)$$

Similarly, we can rewrite the variables $a_i^{(k)}$ ($1 \leq k \leq m+1$) and $c_i^{(k)}$ ($1 \leq k \leq m$) in the form

$$a_i^{(k)} = \exp(q_{i+k} - q_i) a_i^{(k)}; \quad c_i^{(k)} = \exp(q_{i+k} - q_i) c_i^{(k)}; \quad (13)$$

so that the equations (7-9) are turned into the following form in terms of the new variables $a_i^{(k)}$,

$$q_i = 2 \sum_{k=1}^{\infty} a_i^{(k)} a_i^{(k)} \exp[2(q_{i+k} - q_i)] - c_{i-k}^{(k)} a_{i-k}^{(k)} \exp[2(q_i - q_{i-k})]; \quad (14)$$

$$a_i^{(k)} = 2 \sum_{k^0=1}^{m \text{ in } (m \times k \text{ in } m)} a_i^{(k+k^0)} c_{i+k}^{(k^0)} \exp[2(q_{i+k+k^0} - q_{i+k})] + c_{i-k^0}^{(k^0)} a_{i-k^0}^{(k+k^0)} \exp[2(q_i - q_{i-k^0})]; \quad (1 \leq k \leq m+1) \quad (15)$$

$$a_i^{(k)} = 2 \sum_{k^0=1}^{m \times k} a_{i+k}^{(k^0)} c_i^{(k+k^0)} \exp[2(q_{i+k+k^0} - q_{i+k})] + c_{i-k^0}^{(k^0)} a_{i-k^0}^{(k+k^0)} \exp[2(q_i - q_{i-k^0})]; \quad (1 \leq k \leq m+1); \quad (16)$$

In the last equations, we have supplemented the substitution rule (13) with $q_i^{(m_-)} = 1$ according to (12). The coupled system of equations (14)–(16) constitute the set of equations of motion for our $(m_+; m_-)$ -th extended Toda chain.

Among all possible pairs of integers $(m_+; m_-)$, the special choices $m_+ = m_- = m$ have some special features. In these cases, the equations of motion for the $(m; m)$ -th extended Toda chain is completely symmetric under $q_i^{+(k)} \leftrightarrow q_i^{(k)}$ and thus admit a symmetric reduction $q_i^{+(k)} = q_i^{(k)} = q_i^{(k)}$. The reduced system of equations of motion reads

$$q_i = 2 \prod_{k=1}^{m-1} q_i^{(k)^2} \exp[2(q_{i+k} - q_i)] \prod_{k=1}^{m-1} q_{i-k}^{(k)^2} \exp[2(q_i - q_{i-k})] \\ + 2 \exp[2(q_{i+m} - q_i)] \exp[2(q_i - q_{i-m})] g; \quad (17)$$

$$-1^{(k)} = 2 \prod_{k^0=1}^{m-k} q_i^{(k+k^0)} q_{i+k}^{(k^0)} \exp[2(q_{i+k+k^0} - q_{i+k})] \prod_{k^0=1}^{m-k} q_{i-k^0}^{(k+k^0)} q_{i-k^0}^{(k^0)} \exp[2(q_i - q_{i-k^0})] ; \\ (1 - k - m - 1): \quad (18)$$

In particular, the special case of $m = 1$ gives rise to the standard infinite Toda chain.

To give the readers some more intuitive idea about the structure for the $(m_+; m_-)$ -th extended Toda chain, let us write explicitly several more cases with concrete values for m_+ and m_- , i.e. the cases for $m_+; m_- = 3$. These include, beside the simplest case of $m_+ = m_- = 1$ (which is automatically a symmetric reduced case, i.e. the standard Toda chain), the following cases:

$m_+ = 2; m_- = 1$. The corresponding equations of motion read

$$q_i = 2 \prod_{i=1}^n q_i^{+(1)} \exp[2(q_{i+1} - q_i)] \prod_{i=1}^n q_{i-1}^{+(1)} \exp[2(q_i - q_{i-1})] ; \\ -1^{+(1)} = 2 \exp[2(q_{i+2} - q_{i+1})] \exp[2(q_i - q_{i-1})] g;$$

$m_+ = 2; m_- = 2$. The corresponding equations of motion read

$$q_i = 2 \prod_{i=1}^n q_i^{+(1)} q_i^{(1)} \exp[2(q_{i+1} - q_i)] \prod_{i=1}^n q_{i-1}^{(1)} q_{i-1}^{+(1)} \exp[2(q_i - q_{i-1})] \\ + 2 \exp[2(q_{i+2} - q_i)] \exp[2(q_i - q_{i-2})] g; \\ -1^{+(1)} = 2 \prod_{i=1}^n q_{i+1}^{(1)} \exp[2(q_{i+2} - q_{i+1})] \prod_{i=1}^n q_{i-1}^{(1)} \exp[2(q_i - q_{i-1})] ; \\ -1^{(1)} = 2 \prod_{i=1}^n q_{i+1}^{+(1)} \exp[2(q_{i+2} - q_{i+1})] \prod_{i=1}^n q_{i-1}^{+(1)} \exp[2(q_i - q_{i-1})] ;$$

This case admits a symmetric reduction, which reads

$$\begin{aligned} q_i &= 2 \sum_{i=1}^{(1)} \exp [2(q_{i+1} - q_i)] + \sum_{i=1}^{(1)} \exp [2(q_i - q_{i-1})] \\ &\quad + 2 \exp [2(q_{i+2} - q_i)] - \exp [2(q_i - q_{i-2})] g; \\ -\frac{1}{i} &= 2 \sum_{i+1}^{(1)} \exp [2(q_{i+2} - q_{i+1})] - \sum_{i=1}^{(1)} \exp [2(q_i - q_{i-1})] ; \end{aligned}$$

$m_+ = 3; m_- = 1$. We have

$$\begin{aligned} q_i &= 2 \sum_{i=1}^{n+1} \exp [2(q_{i+1} - q_i)] + \sum_{i=1}^{(1)} \exp [2(q_i - q_{i-1})] ; \\ -\frac{1}{i} &= 2 \sum_{i=1}^{n+1} \exp [2(q_{i+2} - q_{i+1})] + \sum_{i=1}^{(2)} \exp [2(q_i - q_{i-1})] ; \\ -\frac{1}{i} &= 2 \exp [2(q_{i+3} - q_{i+2})] - \exp [2(q_i - q_{i-1})] g; \end{aligned}$$

$m_+ = 3; m_- = 2$. The corresponding equations of motion read

$$\begin{aligned} q_i &= 2 \sum_{i=1}^{n+1} \exp [2(q_{i+1} - q_i)] + \sum_{i=1}^{(1)} \exp [2(q_i - q_{i-1})] \\ &\quad + 2 \sum_{i=1}^{n+1} \exp [2(q_{i+2} - q_i)] + \sum_{i=2}^{(2)} \exp [2(q_i - q_{i-2})] ; \\ -\frac{1}{i} &= 2 \sum_{i=1}^{n+1} \exp [2(q_{i+2} - q_{i+1})] + \sum_{i=1}^{(1)} \exp [2(q_i - q_{i-1})] \\ &\quad + 2 \exp [2(q_{i+3} - q_{i+1})] - \exp [2(q_i - q_{i-2})] g; \\ -\frac{1}{i} &= 2 \sum_{i+2}^{(1)} \exp [2(q_{i+3} - q_{i+2})] - \sum_{i=1}^{(1)} \exp [2(q_i - q_{i-1})] ; \\ -\frac{1}{i} &= 2 \sum_{i+1}^{n+1} \exp [2(q_{i+2} - q_{i+1})] + \sum_{i=1}^{(1)} \exp [2(q_i - q_{i-1})] ; \end{aligned}$$

$m_+ = 3; m_- = 3$. We have

$$\begin{aligned} q_i &= 2 \sum_{i=1}^{n+1} \exp [2(q_{i+1} - q_i)] + \sum_{i=1}^{(1)} \exp [2(q_i - q_{i-1})] \\ &\quad + 2 \sum_{i=1}^{n+1} \exp [2(q_{i+2} - q_i)] + \sum_{i=2}^{(2)} \exp [2(q_i - q_{i-2})] \\ &\quad + 2 \exp [2(q_{i+3} - q_i)] - \exp [2(q_i - q_{i-3})] g; \\ -\frac{1}{i} &= 2 \sum_{i=1}^{n+1} \exp [2(q_{i+2} - q_{i+1})] + \sum_{i=1}^{(1)} \exp [2(q_i - q_{i-1})] \\ &\quad + 2 \sum_{i+1}^{(2)} \exp [2(q_{i+3} - q_{i+1})] - \sum_{i=2}^{(2)} \exp [2(q_i - q_{i-2})] ; \\ -\frac{1}{i} &= 2 \sum_{i+2}^{(1)} \exp [2(q_{i+3} - q_{i+2})] - \sum_{i=1}^{(1)} \exp [2(q_i - q_{i-1})] ; \\ -\frac{1}{i} &= 2 \sum_{i=1}^{n+1} \exp [2(q_{i+2} - q_{i+1})] + \sum_{i=1}^{(1)} \exp [2(q_i - q_{i-1})] \\ &\quad + 2 \sum_{i+1}^{(2)} \exp [2(q_{i+3} - q_{i+1})] + \sum_{i=2}^{(2)} \exp [2(q_i - q_{i-2})] ; \\ -\frac{1}{i} &= 2 \sum_{i+2}^{(1)} \exp [2(q_{i+3} - q_{i+2})] - \sum_{i=1}^{(1)} \exp [2(q_i - q_{i-1})] ; \end{aligned}$$

This system of equations also admits a symmetric reduction, for which the equations of motion read

$$\begin{aligned}
\dot{q}_i &= 2 \sum_{i=1}^{(1)} \exp [2 (q_{i+1} - q_i)] - \sum_{i=1}^{(1)} \exp [2 (q_i - q_{i-1})] \\
&+ 2 \sum_{i=1}^{(2)} \exp [2 (q_{i+2} - q_i)] - \sum_{i=2}^{(2)} \exp [2 (q_i - q_{i-2})] \\
&+ 2 f \exp [2 (q_{i+3} - q_i)] - \exp [2 (q_i - q_{i-3})] g; \\
\dot{\tau}_i^{(1)} &= 2 \sum_{i=1}^{(2)} \sum_{i+1}^{(1)} \exp [2 (q_{i+2} - q_{i+1})] - \sum_{i=1}^{(2)} \sum_{i-1}^{(1)} \exp [2 (q_i - q_{i-1})] \\
&+ 2 \sum_{i+1}^{(2)} \exp [2 (q_{i+3} - q_{i+1})] - \sum_{i-2}^{(2)} \exp [2 (q_i - q_{i-2})]; \\
\dot{\tau}_i^{(2)} &= 2 \sum_{i+2}^{(1)} \exp [2 (q_{i+3} - q_{i+2})] - \sum_{i-1}^{(1)} \exp [2 (q_i - q_{i-1})] :
\end{aligned}$$

All these systems of equations are integrable in the Lax sense by construction. However, to prove their Liouville integrability, we need to truncate the number of degrees of freedom into a finite integer. There are two possible ways to make the truncation, i.e. finite chain reduction and the periodic reduction. We shall go into details of both kinds of reductions in the following.

2.2 Finite chain

As just mentioned, the fact that equations (14)–(16) contain an infinite number of variables makes it difficult to consider their Liouville integrability. Therefore we now turn to a finite chain variant.

The Lax matrix L for the finite chain case is given by $(m = n - 1)$

$$\begin{aligned}
L &= L^{(0)} + L^{(>)} + L^{(<)}; \\
L^{(0)} &= \sum_{i=1}^n q_i j_i h_i j; \\
L^{(>)} &= \sum_{k=1}^n \sum_{i=1}^k \exp (q_{i+k} - q_i) \sum_{i=1}^{(k)} j_i h_{i+k} j; \\
L^{(<)} &= \sum_{k=1}^n \sum_{i=1}^k \exp (q_{i+k} - q_i) \sum_{i=1}^{(k)} j_{i+k} h_i j;
\end{aligned} \tag{19}$$

In other words, L is obtained from L by setting $j_i = 0$ for $i < 1$ and $i > n$. The corresponding M matrix is again given by the difference between $L^{(>)}$ and $L^{(<)}$,

$$M = L^{(>)} - L^{(<)};$$

The equations of motion now follow from a finite version of the Lax equation (1),

$$L_- = [M, L]; \quad (20)$$

and it turns out that they again take the form of equations (14)–(16), but now the number of variables is restricted to be finite, i.e.

$$\begin{aligned} q_i &= 0 & \text{if } i < 1 \text{ or } i > n; \\ q_i^{(k)} &= 0 & \text{if } i < 1 \text{ or } i + k > n; \end{aligned} \quad (21)$$

In what follows, we shall refer to the finite $(m_+; m_-)$ -chain characterized by the chain length n as the $(m_+; m_-)$ -th extended Toda n -chain, or simply $(m_+; m_-)$ -Toda n -chain. It should be remarked that the finite (nonsymmetric) $(2; 1)$ - and $(2; 2)$ -chains respectively can be regarded as dimensional reductions of the so-called heterotic Toda field [12] and the bosonic superconformal Toda field [13] theories studied by one of the authors with others about ten years ago. On the other hand, the particular cases of symmetrically reduced $(m; m)$ -Toda n -chain have already been discussed by Deift and collaborators [14], in particular, the symmetric $(2; 2)$ -chain was studied by them in detail.

The $(m_+; m_-)$ -Toda n -chain is simpler than the $(m_+; m_-)$ -th extended infinite Toda chain (or the $(m_+; m_-)$ -Toda 1-chain) in that its Liouville integrability can be established explicitly, and its initial value problem can also be solved systematically. To show this we remind that the operators $j_{ij}h_{jm}$ may be identified with the generator e_{ij} of the Lie algebra gl_n . Therefore, we may outline the Liouville integrability of the $(m_+; m_-)$ -Toda n -chain using the knowledge about the Lie algebra gl_n . The following proposition is a straightforward generalization of the corresponding proposition for the standard Toda chain (in the present terminology, these are just the $(1; 1)$ -Toda n -chain), which is known to the integrable systems community for over twenty years:

Proposition 1 Let L be any matrix in gl_n which has a strip form. Let M be also a matrix in gl_n whose upper triangular part coincides with that of L , while its lower triangular part differs from that of L by a total sign, i.e.

$$M = L^{(>)} - L^{(<)};$$

Then, $L; M$ satisfy the Lax equation (20) and the quantities $H_k(L) := \frac{1}{k} \text{tr}(L^k)$; $k \in \mathbb{Z}$ are integrals of motion for the matrix dynamics defined by the Lax equation.

For instance, the second integral of motion for the generic $(m_+; m_-)$ -Toda n -chain is easy to be calculated, yielding

$$\begin{aligned} H_2(L) &= \frac{1}{2} \text{tr} (L^{(0)})^2 + \text{tr} L^{(>)} L^{(<)} \\ &= \frac{1}{2} \sum_{i=1}^n q_i^2 + \sum_{i=1}^n \sum_{k=1}^{\infty} \exp [2 (q_{i+k} - q_i)] q_i^{+ (k)} q_i^{(-k)}; \end{aligned} \quad (22)$$

The higher integrals of motion can also be constructed using the explicit expression for L , but here we shall not list the explicit result since they are quite complicated.

Moreover, by slightly generalizing the Hamiltonian description for the standard Toda chain, we can see that the L matrix satisfy the well known fundamental Poisson relations for integrable systems, i.e.

$$fL; Lg = [r; L - 1 + 1 - Lg; \quad (23)$$

where r is the r matrix for standard Toda chains, now written as

$$r^+ = \sum_{i,j=1}^{X^n} j i h i j - j j i h j j + \sum_{\substack{i,j=1 \\ i < j}}^{X^n} j i h j j - j j i h i j$$

$$r^- = \sum_{i,j=1}^{X^n} j i h i j - j j i h j j + \sum_{\substack{i,j=1 \\ i > j}}^{X^n} j i h j j - j j i h i j$$

This is achieved as follows. First, let us introduce the linear operator R which acts on elements of gl_n in the following way,

$$R(X) = \text{tr}_2(r^+ + r^-; 1 \otimes X); \quad 8X \in 2 \otimes gl_n;$$

where tr_2 means taking the trace over the second tensor component of the tensor product space $gl_n \otimes gl_n$. Then for any linear function f over gl_n , we can see that the bracket [15]

$$ff(X); f(Y)g = f([X; Y]_R) \quad (24)$$

with

$$[X; Y]_R = [R(X); Y] + [X; R(Y)]$$

is a Poisson bracket. In particular, if f is chosen to be the Lax matrix L , then the Poisson bracket (24) is exactly (23). By the way, we can see that $R(L) = L^{(p)}$ $L^{(q)} = M$. This establishes the Hamiltonian description for the finite $(m_+; m_-)$ -Toda n-chain.

For the symmetrically reduced n-chain, it is easy to prove that the following proposition holds, giving the unique solution to the initial value problem :

Proposition 2 The unique solution to the Lax system (20) with symmetric matrix L and initial data $L(0) = L_0$ is given by

$$L(t) = A(t)L_0A(t)^{-1};$$

where $A(t)$ is the orthogonal component in the unique matrix decomposition

$$\exp(tL_0) = A(t)N(t);$$

where $N(t)$ is a lower triangular matrix.

For generic $(m_+; m_-)$ -Toda n-chain without symmetric reduction, the solution to the initial value problem is quite complicated in practice. However, the underlying idea is the same: one can make a decomposition for the matrix $\exp(tL_0)$ such that $\exp(tL_0) = A(t)N(t)$ and $A(t)$ must satisfy

$$A(t)A^{-1}(t)^{(>)} = A(t)L_0A^{-1}(t)^{(>)}; \quad A(t)A^{-1}(t)^{(<)} = A(t)L_0A^{-1}(t)^{(<)}; \quad (25)$$

Then $L(t) = A(t)L_0A^{-1}(t)$ will solve the initial value problem with initial data $L(0) = L_0$. The explicit solution for the matrix $A(t)$ can be obtained by first writing $A(t)$ in the form of Gauss decomposition and then solve (25) order by order in terms of the heights of the roots of \mathfrak{g}_n .

2.3 Periodic chain

The Lax matrix L of the periodic $(m_+; m_-)$ -Toda n-chain is given as follows,

$$\begin{aligned} L &= L^{(0)} + L^{(>)} + L^{(<)}; \\ L^{(0)} &= \sum_{i=0}^{X-1} q_i j_i i h_i j; \\ L^{(>)} &= \sum_{k=1}^X \sum_{i=0}^{n-X^k-1} \exp(q_{i+k} - q_i)^{+ (k)} j_i i h_i + k j \\ &\quad + w^{-1} \sum_{k=1}^X \sum_{i=n-k}^{X-1} \exp(q_{i+k} - q_i)^{+ (k)} j_i i h_i - (n-k) j; \\ L^{(<)} &= \sum_{k=1}^X \sum_{i=0}^{n-X^k-1} \exp(q_{i+k} - q_i)^{(k)} j_i + k i h_i j \\ &\quad + w \sum_{k=1}^X \sum_{i=n-k}^{X-1} \exp(q_{i+k} - q_i)^{(k)} j_i - (n-k) i h_i j; \end{aligned} \quad (26)$$

where w is a complex spectral parameter, and the M matrix is the difference between $L^{(>)}$ and $L^{(<)}$ given in (26),

$$M = L^{(>)} - L^{(<)};$$

The equations of motion following from the Lax equation still take the form of (14)–(16), but now the sum of the variables q_i and $i^{(k)}$ is to be understood modulo n , i.e.

$$q_{i+n} = q_i; \quad i+n^{(k)} = i^{(k)}; \quad i = 0, 1, \dots, n-1; \quad r \in \mathbb{Z}; \quad (27)$$

Of course, just like the cases of finite chain and finite n-chain, the periodic n-chain also admits a symmetric reduction.

The Lax matrix (26) may be considered to be defined over the loop algebra \mathfrak{gl}_n . Its key difference from the case of finite chains is the appearance of the complex spectral parameter w , which can give rise to a complex curve called spectral curve. The Liouville integrability and the Hamiltonian description for the periodic chains can be outlined in a similar fashion like the case of finite chains, the only difference being that we need to replace all the objects from the finite dimensional Lie algebra \mathfrak{gl}_n by objects from the loop algebra \mathfrak{gl}_n .

Among all periodic $(m_+; m_-)$ chains, the special cases of $(m_+; m_-) = (1; 1)$, $(2; 1)$, $(2; 2)$ are the simplest ones. The periodic $(1; 1)$ -chain is just the standard Toda mechanics which we shall not pay any more words. That the $(2; 1)$ - and $(2; 2)$ -chains are much simpler than the generic cases is not only because that they contain less mechanical variables but also due to the fact that the dynamics of these two special chains is completely determined by the second integral of motion. More concretely we may take the second integral of motion (notice that the lower and upper bounds for the summations over i are different from the finite chain cases)

$$H_2^{(2;1)}(L) = \frac{1}{2} \sum_{i=0}^{N-1} q_i^2 + \sum_{i=0}^{N-1} \exp [2(q_{i+1} - q_i)] q_i^{(1)};$$

$$H_2^{(2;2)}(L) = \frac{1}{2} \sum_{i=0}^{N-1} q_i^2 + \sum_{i=0}^{N-1} \exp [2(q_{i+1} - q_i)] q_i^{(1)} q_i^{(1)} + \exp [2(q_{i+2} - q_i)] q_i^{(1)} q_i^{(1)}$$

as the Hamiltonian, and the equations of motion follow as Hamiltonian flows for the variables $q_i; q_i^{(1)}$ (recall that $q_i^{(m)} = 1$) provided the nonzero Poisson brackets for these variables are specified as

$$\{q_i; q_j\} = \delta_{ij};$$

$$\{q_i^{(1)}; q_j^{(1)}\} = 2(\delta_{i,j-1} - \delta_{i,j+1});$$

$$\{q_i^{(1)}; q_j^{(1)}\} = 2(\delta_{i,j-1} - \delta_{i,j+1});$$

The Hamiltonian structure together with the Lax representation constitute a description of the $(2; 1)$ - and $(2; 2)$ -chains as integrable Hamiltonian systems.

3 Spectral problem of the periodic chain: an example

As stated earlier in the introduction, a key object of interests in the study of many body mechanical systems from the modern view point is the spectral curves of the Lax matrix L , which we now study for the case of periodic $(m_+; m_-)$ -Toda chain.

First of all, the Lax equation can be viewed as the compatibility condition for

the pair of linear equations

$$\frac{d}{dt} M = 0; \quad (28)$$

$$L = \lambda I; \quad (29)$$

in which the second equation, (29), defines the eigenvalue problem of L .

On the other hand, since the mechanical variables appeared in L defined in (26) is periodic in the sites with periodicity n , we can define the monodromy operator T which shifts the sites of these mechanical operators by n and consequently acts on q_i and $p_i^{(k)}$ as identity operator,

$$T q_i = q_{i+n} = q_i; \quad T p_i^{(k)} = p_{i+n}^{(k)} = p_i^{(k)};$$

However, the form of the L matrix implies that T acts on the vector states j_i and h_j respectively in a nontrivial way,

$$T j_i = j_{i+n} = w j_i; \quad (30)$$

$$T h_j = h_{j+n} = w^{-1} h_j; \quad (31)$$

and hence the action of T on $\prod_{i=0}^{n-1} j_i$ is also nontrivial,

$$T \prod_{i=0}^{n-1} j_i = w \prod_{i=0}^{n-1} j_i;$$

The actions of L and T on $\prod_{i=0}^{n-1} j_i$ commutes, $[L; T] = 0$, which implies the existence of certain relationship between L and T ,

$$P(L; T) = 0;$$

Such a relationship can be formulated exactly by the spectral curve

$$P(\lambda; w) = 0;$$

with the map P specified by the characteristic equation of the L matrix,

$$P(\lambda; w) = \det(L(\lambda, w) - \lambda I); \quad (32)$$

The explicit form of the spectral map $P(\lambda; w)$ for general periodic $(m_+; m_-)$ -Toda n-chain is very complicated, and we shall not go into the detailed description for the generic cases. Instead, we shall illustrate the structure of the spectral curve only for the simplest cases of periodic $(2; 1)$ -chains.

For $(m_+; m_-) = (2; 1)$, we can rewrite (32) in explicit matrix form,

$$P^{(2;1)}(\lambda; w) = \det \begin{pmatrix} q_0 & a_0^{(1)} & a_0^{(2)} & & & & & & \\ & c_0^{(1)} & q_1 & a_1^{(1)} & a_1^{(2)} & & & & \\ & & c_1^{(1)} & q_2 & \ddots & \ddots & & & \\ & & & \ddots & \ddots & \ddots & & & \\ & & & & \ddots & \ddots & & & \\ & & & & & \ddots & & & \\ & & & & & & a_n^{(2)} & & \\ & & & & & & a_n^{(1)} & & \\ w^{-1} a_n^{(2)} & & & & & & & & \\ w^{-1} a_n^{(1)} & & & & & & c_n^{(1)} & q_{n+1} & \end{pmatrix}; \quad (33)$$

with entries $a_i^{(k)}, c_i^{(k)}$ given by equation (13), where s_i stands for the variables $q_i, -s_i^{(k)}$ are to be understood modulo n . A direct evaluation of (33) yields

$$P^{(2;1)}(\lambda; w) = (\lambda - 1)^{n-1} w + \frac{Q_{n-3}(\lambda)}{w} + \frac{1}{w^2} + P_n(\lambda) = 0; \quad (34)$$

where $Q_{n-3}(\lambda)$ and $P_n(\lambda)$ respectively are some polynomials in λ of order $n-3$ and n , with coefficients being polynomials in $q_i; \exp(q_i)$ and $s_i^{+(1)}$ which are symmetric under exchange of s_i . It should be remarked that the form of the spectral curve (34) for (2;1)-Toda chains is very different from that of the (1;1)-chains. Therefore, it is interesting to learn what kind of SUSY gauge theories might correspond, taking (34) as its Seiberg-Witten curve. We leave this problem for future study.

Another way of looking at the spectral problems for the $(m_+; m_-)$ -Toda chains can be specified as follows. First, let us define an analogous eigenvalue problem to eq. (29) before making the periodic reduction. This amounts to introduce formally an eigenvector $\vec{p} = \sum_{i \in \mathbb{Z}} p_i \vec{j}_i$ for the L matrix (2), i.e.

$$L \vec{p} = \lambda \vec{p}; \quad (35)$$

Using the explicit form of L we can rewrite (35) as

$$\begin{aligned} \sum_{i \in \mathbb{Z}} p_i \vec{j}_i &= \sum_{i \in \mathbb{Z}} q_i p_i \vec{j}_i \\ &+ \sum_{k=1}^{\infty} \sum_{i \in \mathbb{Z}} \exp(q_{i+k} - q_i) s_i^{+(k)} p_{i+k} \vec{j}_i \\ &+ \sum_{k=1}^{\infty} \sum_{i \in \mathbb{Z}} \exp(q_i - q_{i-k}) s_{i-k}^{(k)} p_{i-k} \vec{j}_i; \end{aligned} \quad (36)$$

Since the states \vec{j}_i are all linearly independent, the last equation is in fact a collection of equations which provide a recursive relationship between the vector components p_i , i.e.

$$\begin{aligned} p_{i+m_+} &= \left(\prod_{k=1}^{m_+} q_i \right) \sum_{k=1}^{\infty} \exp(q_{i+k} - q_{i+m_+}) s_i^{+(k)} p_{i+k} \\ &+ \sum_{k=1}^{\infty} \exp(2q_i - q_{i-k} - q_{i+m_+}) s_{i-k}^{(k)} p_{i-k}; \quad i \in \mathbb{Z}; \end{aligned} \quad (37)$$

Therefore, for generic values of m_+ and m_- , such a set of equations can be recast into matrix form with coefficient matrix of dimension $m_+ + m_-$. More concretely, writing

$$\vec{p} = \begin{pmatrix} p_{i+m_+-1} & p_{i+m_+-2} & \dots & p_{i-m_++1} & p_{i-m_+} \end{pmatrix}^T$$

and defining the recursive matrix L_i in terms of the matrix equation

$$L_{i+1} = L_i L_i;$$

we may read out the matrix L_i from (37),

$$L_i = \begin{pmatrix} 0 & & & & & 1 \\ & B & & & & \\ & 1 & 0 & & & \\ & 0 & 1 & 0 & & \\ & \vdots & \ddots & \ddots & \ddots & \\ & 0 & & 0 & 1 & 0 \end{pmatrix} \begin{matrix} C_0 \\ C_1 \\ \vdots \\ C_{i-1} \\ A \end{matrix};$$

where the C_j 's are just the coefficients of ψ_j 's appeared in (37). The form of the above matrices L_i is very similar to the Lax matrix of the KdV hierarchy in the Drinfeld-Sokolov gauge [16], which reminds us that it may be gauge transformed into a form with the only nontrivial elements (i.e. the ones besides the 1's on the subdiagonal) being on the main diagonal following the standard procedure of Miura transformation. However, at present, we shall not need to make such a transformation.

Now acting on, say, ψ_0 , from the left by L_0 and then by L_1, L_2, \dots until L_{n-1} , we have

$$\psi_n = L_{n-1} L_{n-2} \dots L_0 \psi_0;$$

It is now a good place to recall the periodic reduction condition (30) which implies the relationship

$$\psi_n = W \psi_0;$$

Writing

$$T(\lambda) = L_{n-1} L_{n-2} \dots L_0(L) \quad (38)$$

where explicit dependence on the parameter λ is emphasized, we have

$$T(\lambda) \psi_0 = W \psi_0;$$

We thus have an alternative spectral problem

$$\det(T(\lambda) - W) = 0; \quad (39)$$

This last equation gives another expression for the spectral curves associated with the periodic (m_+, m_-) -Toda n-chain.

As an illustrating example, we shall give the explicit form of the matrices L_i for the case of $(m_+, m_-) = (2; 1)$. In this case, we have

$$L_i^{(2;1)} = \begin{pmatrix} 0 & & 1 \\ & B & \\ & 1 & 0 \\ & 0 & 1 & 0 \\ & \vdots & \ddots & \ddots & \ddots \\ & 0 & & 0 & 1 & 0 \end{pmatrix} \begin{matrix} C_0 \\ C_1 \\ \vdots \\ C_{i-1} \\ A \end{matrix};$$

and consequently

$$L_i^{(2;1)} = \begin{pmatrix} 0 & \exp(q_{i+1} - q_{i+2}) & a_i^{(1)} & 0 \\ \exp(q_i - q_{i+2}) & 1 & 0 & 0 \\ \exp(2q_i - q_{i+1} - q_{i+2}) & 0 & 0 & 1 \end{pmatrix} A :$$

It is then a simple practice to get $T^{(2;1)}(\lambda)$ using (38), and then get the spectral curve via (39). That the two spectral curves (32) and (39) are identical is a well known fact for the case of standard Toda chains (i.e. the (1;1)-chains).

4 Nonabelian generalizations

The extended Toda chain we studied so far can also be generalized along another line, i.e. the nonabelian generalizations. By nonabelian generalization we mean that the fundamental mechanical variables q_i , $a_i^{(k)}$ and $c_i^{(k)}$ can be replaced by nonabelian objects, e.g. q_i may be replaced by $\mathfrak{N}_i \times \mathfrak{N}_i$ matrices Q_i , while $a_i^{(k)}$ and $c_i^{(k)}$ are to be replaced by some generic rectangular matrices $A_i^{(k)}$ of size $\mathfrak{N}_i \times \mathfrak{N}_{i+k}$ and $C_i^{(k)}$ of size $\mathfrak{N}_{i+k} \times \mathfrak{N}_i$. Such nonabelian versions of the extended Toda chain is expected to be useful in the study of moduli structure of $N = 2$ and $N = 1$ SUSY gauge theories with non-semisimple or partially broken gauge symmetries.

Now let us give explicitly the Lax matrices and the equations of motion for the nonabelian extended Toda chain. For the case of infinite chain, the L matrix is given as

$$\begin{aligned} L &= L^{(0)} + L^{(>)} + L^{(<)}; \\ L^{(0)} &= \sum_{i \in \mathbb{Z}} Q_i j_i h_i j; \\ L^{(>)} &= \sum_{i \in \mathbb{Z}} \sum_{k=1}^{\infty} A_i^{(k)} j_i h_i + k j; \\ L^{(<)} &= \sum_{i \in \mathbb{Z}} \sum_{k=1}^{\infty} C_i^{(k)} j_{i+k} h_i j; \end{aligned}$$

The M matrix is still the difference between $L^{(>)}$ and $L^{(<)}$,

$$M = L^{(>)} - L^{(<)} :$$

The equations of motion follow directly from the Lax equation $L_- = [M, L]$ and they

read

$$Q_i = 2 \sum_{k=1}^n A_i^{(k)} C_i^{(k)} - C_{i+k}^{(k)} A_{i+k}^{(k)}; \quad (40)$$

$$A_{-1}^{(k)} = A_i^{(k)} Q_{i+k} - Q_{-1} A_i^{(k)} + 2 \sum_{m=1}^{m \wedge (m+k)} A_{i+k}^{(k+k^0)} C_{i+k}^{(k^0)} - C_{i+k^0}^{(k^0)} A_{i+k^0}^{(k+k^0)}; \quad (1 \leq k \leq m+1) \quad (41)$$

$$G_{-1}^{(k)} = Q_{i+k} C_i^{(k)} - C_i^{(k)} Q_{-1} + 2 \sum_{k^0=1}^{m \wedge k} A_{i+k}^{(k^0)} C_i^{(k+k^0)} - C_{i+k^0}^{(k^0)} A_{i+k^0}^{(k+k^0)}; \quad (1 \leq k \leq m+1) \quad (42)$$

$$A_{-1}^{(m+)} = A_i^{(m+)} Q_{i+m+} - Q_{-1} A_i^{(m+)}; \quad (43)$$

$$G_{-1}^{(m+)} = Q_{i+m} C_i^{(m+)} - C_i^{(m+)} Q_{-1}; \quad (44)$$

The Lax integrability for the system of equations (40)–(44) is guaranteed by construction. However, for generic matrix variables $Q_i; A_i^{(k)}$ and $C_i^{(k)}$, the form of these equations cannot be simplified any further. In particular, the appearance of Q_{-1} on the right hand side of eqs. (41)–(44) is inevitable.

Although the construction given above corresponds to infinite nonabelian chains only, it is extremely easy to get the equations of motion for the finite and periodic chains out of (40)–(44) by imposing constraints over the mechanical variables $Q_i; A_i^{(k)}$ and $C_i^{(k)}$ following the similar fashion as made in eqs. (21) and (27). Concretely, the equations of motion for the finite nonabelian $(m+; m)$ -Toda n-chain are given by (40)–(44) together with the constraint conditions

$$Q_i = 0 \quad \text{if } i < 1 \text{ or } i > n; \\ A_i^{(k)} = C_i^{(k)} = 0 \quad \text{if } i < 1 \text{ or } i+k > n;$$

The equations of motion for the periodic $(m+; m)$ -Toda n-chain are also given by (40)–(44) together with the following constraint conditions,

$$Q_{i+m} = Q_i; \quad A_{i+m}^{(k)} = A_i^{(k)}; \quad C_{i+m}^{(k)} = C_i^{(k)}; \quad r \in \mathbb{Z}:$$

For the size of the matrices to be consistent, we must also require

$$A_{i+m} = A_i; \quad r \in \mathbb{Z}:$$

In order that the nonabelian generalizations take a similar form as (14)–(16), we now consider to replace the matrices Q_{-1} by certain symmetric right invariant vector field over $GL(N_i)$, namely

$$Q_{-1} = B_{-1} B_i^{-1}; \quad B_{-1} B_i^{-1}{}^T = B_{-1} B_i^{-1}; \quad B_i \in GL(N_i):$$

Such a replacement of variables is a kind of reduction to the original system of equations, and hence the resulting system

$$\begin{aligned}
\frac{d}{dt} B_i B_i^{-1} &= 2 \sum_{k=1}^{\infty} A_i^{(k)} C_i^{(k)} - C_i^{(k)} A_i^{(k)}; \\
\frac{d}{dt} A_i^{(k)T} &= B_{i+k} B_{i+k}^{-1} A_i^{(k)T} - A_i^{(k)T} B_i B_i^{-1} \\
&\quad + 2 \sum_{k^0=1}^{m \text{ in } (m-k^0)} C_{i+k}^{(k^0)T} A_i^{(k+k^0)T} - A_{i+k^0}^{(k+k^0)T} C_{i+k^0}^{(k^0)T}; \\
(1 \leq k \leq m-1) \\
\frac{d}{dt} C_i^{(k)} &= B_{i+k} B_{i+k}^{-1} C_i^{(k)} - C_i^{(k)} B_i B_i^{-1} \\
&\quad + 2 \sum_{k^0=1}^{m-k} A_{i+k}^{(k^0)} C_i^{(k+k^0)} - C_{i+k^0}^{(k+k^0)} A_{i+k^0}^{(k^0)}; \quad (1 \leq k \leq m-1) \\
\frac{d}{dt} A_i^{(m)T} &= B_{i+m} B_{i+m}^{-1} A_i^{(m)T} - A_i^{(m)T} B_i B_i^{-1}; \\
\frac{d}{dt} C_i^{(m)} &= B_{i+m} B_{i+m}^{-1} C_i^{(m)} - C_i^{(m)} B_i B_i^{-1}
\end{aligned}$$

is not strictly equivalent to the original one. In the last equations, we used the symbol X^T to denote the transpose of the matrix X . After the above reduction, we may further let

$$A_i^{(k)T} = B_{i+k}^{-1} A_i^{(k)T} B_i; \quad C_i^{(k)} = B_{i+k}^{-1} C_i^{(k)} B_i;$$

so that in terms of the new variables B_i , $A_i^{(k)}$ and $C_i^{(k)}$, the reduced system of

equations reads

$$\frac{d}{dt} B_i B_i^{-1} = 2 \sum_{k=1}^{\infty} B_i^{-1 T} {}^{(k)}_i (B_{i+k})^T B_{i+k} {}^{(k)}_i B_i^{-1} \\ B_i {}^{(k)}_i B_i^{-1} B_i^{-1 T} {}^{(k)}_i (B_i)^T; \quad (45)$$

$$\frac{d}{dt} {}^{(k)}_i = 2 (B_i)^T \sum_{k^0=1}^{\infty} B_i^{-1 T} {}^{(k+k^0)}_i (B_{i+k+k^0})^T B_{i+k+k^0} {}^{(k^0)}_{i+k} B_{i+k}^{-1} \\ B_i {}^{(k^0)}_{i+k^0} B_i^{-1} B_i^{-1 T} {}^{(k+k^0)}_{i+k^0} (B_{i+k})^T (B_{i+k})^{-1 T}; \quad (46)$$

$$\frac{d}{dt} {}^{(k)}_i = 2 (B_{i+k})^{-1} \sum_{k^0=1}^{\infty} B_{i+k}^{-1 T} {}^{(k^0)}_{i+k} (B_{i+k+k^0})^T B_{i+k+k^0} {}^{(k+k^0)}_i B_i^{-1} \\ B_{i+k} {}^{(k+k^0)}_{i+k^0} B_i^{-1} B_i^{-1 T} {}^{(k^0)}_{i+k^0} (B_i)^T B_i; \quad (47)$$

$$\frac{d}{dt} {}^{(m_+)}_i = 0; \quad \frac{d}{dt} {}^{(m_-)}_i = 0; \quad (48)$$

We may also consider the nonabelian analogues of the symmetric reductions when $m_+ = m_- = m$. The condition for the symmetric reduction reads

$${}^{(k)}_i = {}^{(k)}_i{}^T {}^{(k)}_i;$$

and the reduced equations read

$$\frac{d}{dt} B_i B_i^{-1} = 2 \sum_{k=1}^{\infty} B_i^{-1 T} {}^{(k)}_i (B_{i+k})^T B_{i+k} {}^{(k)}_i{}^T B_i^{-1} \\ B_i {}^{(k)}_i{}^T B_i^{-1} B_i^{-1 T} {}^{(k)}_i (B_i)^T; \quad (49)$$

$$\frac{d}{dt} {}^{(k)}_i = 2 (B_i)^T \sum_{k^0=1}^{\infty} B_i^{-1 T} {}^{(k+k^0)}_i (B_{i+k+k^0})^T B_{i+k+k^0} {}^{(k^0)}_{i+k}{}^T B_{i+k}^{-1} \\ B_i {}^{(k^0)}_{i+k^0}{}^T B_i^{-1} B_i^{-1 T} {}^{(k+k^0)}_{i+k^0} (B_{i+k})^T (B_{i+k})^{-1 T}; \quad (50)$$

$$\frac{d}{dt} {}^{(m)}_i = 0; \quad (51)$$

It is also interesting to study the spectral curves and stationary point configurations for the nonabelian generalizations. However, the corresponding calculations

would be much more complicated than the abelian cases, and we shall not go into any details along this direction.

5 Conclusion and discussion

In this paper, we studied the integrability of generalized Toda chains based on the Lie algebras \mathfrak{gl}_n (including the case $n = 1$) and \mathfrak{gl}_n in full generality. For each one of these Lie algebras we find a family of integrable generalizations of Toda chains characterized by two integers m_+ and m_- , with the Lax matrices and equations of motion given explicitly. For the case of $m_+ = m_-$ there is a symmetric reduction for the equations of motion, and the solution to the initial value problem is outlined. For each pair of m_+ and m_- , we also constructed a family of nonabelian generalizations of the Toda chains.

Another problem that we considered in this paper, but not in full generality, is the spectral curves of the abelian periodic $(m_+; m_-)$ -Toda chains. We outlined two different ways for the description of the spectral problems, and presented the form of the spectral curve in the special case of $(2; 1)$ -chains.

We should remark that for $n = 1$, the (abelian) generalized Toda chains we get may be considered as dimensional reductions of the well-known Toda lattice hierarchy [17], which is a collection of infinite many integrable field theoretic models in $(1+1)$ -dimensions. However, the equations of motion for the Toda lattice hierarchy were never written in such an explicit and unified way as we did in this paper.

Although the general construction we made in this paper is based on the Lie algebras \mathfrak{gl}_n and \mathfrak{gl}_n only, the results is ready to be applied to the simple Lie algebras \mathfrak{sl}_n and \mathfrak{sl}_n by simply imposing a traceless condition on the L matrices. Further generalizations to the cases of \mathfrak{so}_n , \mathfrak{sp}_n and their loop algebras is also possible, which we leave for later study. Another important problem which we also leave for future study is the application of spectral curves for the periodic $(m_+; m_-)$ -Toda chains in $n = 2$ SUSY gauge theories. The stationary point configurations for the periodic chains is also of great interests from the $N = 1$ gauge theoretic point of view.

Acknowledgement

LZ is grateful to Ian MacIntosh for introducing to him the technique for the construction of initial value problem for the standard periodic Toda chain and bring his attention to [14]. This work is supported in part by the NSFC.

References

- [1] M .Toda, \Vibration of a chain with nonlinear interaction", J.Phys.Soc.Japan, Vol22, (1967) 431-436; \Wave Propagation in Anharmonic Lattices", J.Phys. Soc. Japan, Vol23, (1967) 501-506.
- [2] H .F laschka, \The Toda Lattice II. Existence of integrals" Phys.Rev B 9 (1974) 1924-1925; \On the Toda Lattice II. Inverse Scattering Solution", Prog.Theor. Phys.51 (1974) 703-716.
- [3] F .C alogero, \Solution of the one-dimensional N-body problem with quadratic and/or inversely quadratic pair potentials", J.M ath.Phys. 12, 419-436 (1971); Erratum , ibidem 37 (1996) 3646 ; \Exactly solvable one-dimensional many-body problem s", Lett.Nuovo C im ento 13, 411-416 (1975).
- [4] J. M oser, \Three integrable Hamiltonian systems connected with isospectral deformations", Adv.M ath. 16 (1975) 197-220 ; \Various aspects of integrable Hamiltonian system s", in: Dynam ical System s, Prog. in M ath.8, Birkhauser, Basel, 1980, pp. 233-289.
- [5] S.N .M .Ruijsenaars and H .Schneider, \A new class of integrable system s and its relation to solitons", Ann.Phys. (NY) 170 (1986) 370-405.
- [6] S. N .M .Ruijsenaars, \Complete integrability of relativistic Calogero–Moser system s and elliptic function identities", Comm un. M ath. Phys. 110 (1987) 191-213.
- [7] R D onagi and E W itten, \Supersym metric Yang–M ills System s And Integrable System s", Nucl.Phys. B 460 (1996) 299-334, hep-th/9510101.
- [8] See, e.g. A .M arshakov, \Seiberg–W itten Theory and Integrable System s", World Scienti c Publishing, 1999 and references therein.
- [9] N .Seiberg and E .W itten, \Monopole Condensation, And Con nement In N= 2 Supersym metric Yang–M ills Theory", Nucl. Phys B 426 (1994) 19-52, hep-th/9407087.
- [10] R .Dijkgraaf and C .Vafa, \Matrix M odels, Topological Strings, and Supersym – metric Gauge Theories", Nucl Phys. B 644 (2002) 3-20, hep-th/0206255; \On Geometry and Matrix M odels", Nucl Phys. B 644 (2002) 21-39, hep-th/0207106; \A Perturbative W indow into Non-Perturbative Physics", hep-th/0208048.
- [11] N D orey, \An Elliptic Superpotential for Softly Broken N= 4 Supersym metric Yang–M ills Theory", JHEP 9907 (1999) 021, hep-th/9906011.
- [12] L .Chao and B-Y .Hou, \Heterotic Toda elds", Nucl. Phys. B 436 (1995) 638-658.

- [13] B.-Y. Hou and L. Chao, "From integrability to conformality: Bosonic superconformal Toda theories", *Int. J. Mod. Phys. A* 8 (6) (1993) 1105-1123.
- [14] P. Delft, L.C. Li, T. Nanda and C. Tomei, "The Toda Flow on a Generic Orbit is Integrable", *Commun. on Pure and Appl. Math.*, Vol. XXXIX (1986) 183-232.
- [15] O. Babelon, C.-M. Viallet, "Integrable models, Yang-Baxter equations and quantum groups", Part I, *Ref. SISSA* 54 EP (1989)
- [16] V.G. Drinfeld, Sokolov, *J. Sov. Math. Phys.* 30 (1984) 1975.
- [17] K. Ueno, K. Takasaki, "Toda lattice hierarchy", *Advanced Studies in Pure and Applied Mathematics* 4 (1984) 1-95.