

# A “general boundary” formulation for quantum mechanics and quantum gravity

Robert Oeckl

*Centre de Physique Théorique, CNRS Luminy, 13288 Marseille, France*

(Dated: August 13, 2003)

I propose to formalize quantum theories as topological quantum field theories in a generalized sense, associating state spaces with boundaries of arbitrary (and possibly finite) regions of space-time. I further propose to obtain such “general boundary” quantum theories through a generalized path integral quantization. I show how both, non-relativistic quantum mechanics and quantum field theory can be given a “general boundary” formulation. Surprisingly, even in the non-relativistic case, features normally associated with quantum field theory emerge from consistency conditions. This includes states with arbitrary particle number and pair creation. I also note how three dimensional quantum gravity is an example for a realization of both proposals and suggest to apply them to four dimensional quantum gravity.

One obstacle in the quest for a quantum theory of gravity appears to be the fact that the foundations of quantum mechanics (QM) are inherently non-covariant. On the other hand, quantum field theory (QFT) can be formulated in a covariant way. The price for this is a globality (manifest in the path integral expression for  $n$ -point functions) that fixes space-time to be Minkowskian. Well known difficulties result from this already for the extension of QFT to curved space-times.

I propose an approach to formulating quantum theories that is at the same time local and inherently compatible with special or general covariance. The main idea is firstly, to associate state spaces with boundaries of general regions of space-time. Secondly, amplitudes are determined by a complex function for each region and associated state space. Crucially, (and contrary to standard QM) connected boundaries of compact regions are the main focus of attention. In this sense the formulation is “holographic”, i.e. the information about the interior of a region is encoded through the states on the boundary. These structures are required to be coherent in the sense of topological quantum field theory (TQFT) [1]. This does not mean that the underlying structure is necessarily topological. For QM and QFT the relevant background structure is the metric. Only for quantum gravity would the theory be topological (more precisely differentiable) in the usual sense. Note that this does not imply a lack of local degrees of freedom (see [2]).

Since the association of states with possibly time-like hypersurfaces is a quite radical step for QM it is crucial to understand their physical meaning. This is particularly true for particle states in QFT. Thus it makes sense even with the goal of quantum gravity in mind first to reformulate non-relativistic quantum mechanics (NRQM) and QFT in the “general boundary” sense. This is the main focus of this letter.

To achieve this goal I make a second proposal in the form of a quantization scheme. It turns out that the path integral approach [3] is particularly suitable and my scheme is a rather straightforward generalization of

it. It is designed so as to produce “general boundary” theories of the type described above. It is “holographic” not only in the sense mentioned above but also in the sense that the underlying classical configuration space on the boundary should be chosen such that it uniquely encodes a solution of the equations of motions in the interior.

I show that both NRQM of particles in a potential as well as scalar perturbative QFT are obtainable as holographic quantizations of the relevant classical theories. This then allows the generalization to the “general boundary” formulation. Surprisingly, in the case of NRQM features of quantum field theory emerge, such as the necessity for states of any particle number and the suggestion of pair creation. Finally, I remark on the fact that quantum gravity in three dimensions is a TQFT, i.e. has a “general boundary” description. What is more, it can be obtained precisely as a holographic quantization and thus fits perfectly into the scheme.

Another main motivation for this work comes from an analysis of the measurement problem in quantum gravity. As I have argued in [2], this suggests precisely a “general boundary” formulation for quantum gravity. Interpretational implications of the “general boundary” formulation are also investigated there.

## “GENERAL BOUNDARY” FORMULATION

Let me explain what the “general boundary” formulation, i.e. a (suitably adapted notion of) TQFT is. Let  $M$  be a region of space-time (i.e. a four-dimensional manifold) and  $S$  its boundary hypersurface. (For the moment I do not specify what background structure this entails.) (T1) Associated with each such boundary  $S$  is a vector space  $H_S$  of *states*. (T2) If  $S$  decom-

poses into disconnected components<sup>1</sup>  $S = S_1 \sqcup \dots \sqcup S_n$  then the state space decomposes into a tensor product  $H_S = H_{S_1} \otimes \dots \otimes H_{S_n}$ . (T3) For a given boundary  $S$ , changing its orientation, i.e. the side on which it bounds a region  $M$ , corresponds to replacing  $H_S$  with the dual space  $H_S^*$ . (T4) Associated with  $M$  is a complex function  $\mathcal{M} : H_S \rightarrow \mathbb{C}$  which associates an *amplitude* to a state. One may also dualize boundaries. This means that one may convert  $\mathcal{M} : H_{S_1} \otimes \dots \otimes H_{S_n} \rightarrow \mathbb{C}$  to a function  $\mathcal{M} : H_{S_1}^* \otimes \dots \otimes H_{S_k}^* \otimes H_{S_{k+1}} \otimes \dots \otimes H_{S_n}$ , replacing spaces with dual spaces. Mathematically both versions of  $\mathcal{M}$  are equivalent, giving one determines the other (hence the same notation). A crucial property is the *composition rule*. (T5) Let  $M_1$  and  $M_2$  be two regions of space-time that share a common boundary  $S$ . Let  $M_1$  also have a boundary  $S_1$  and  $M_2$  a boundary  $S_2$ . Consider  $\mathcal{M}_1 : H_{S_1} \rightarrow H_S$  and  $\mathcal{M}_2 : H_S \rightarrow H_{S_2}$ . (The state spaces are chosen with respect to suitable orientations of the boundaries.) Then gluing gives  $M = M_1 \sqcup M_2$  with boundaries  $S_1$  and  $S_2$ . The composition rule demands the equality  $\mathcal{M} = \mathcal{M}_2 \circ \mathcal{M}_1$ .

For quantum gravity the background structure coming with space-time regions and their boundaries is just a differentiable structure. One obtains essentially a proper TQFT. For QM and QFT the background structure is a fixed metric (usually that inherited from Minkowski space).

## HOLOGRAPHIC QUANTIZATION

Consider a classical field theory with a given set of fields  $(\phi)$  and an action  $S[\phi]$  so that the equations of motion can be derived from a minimization of  $S$ . (For simplicity I use the notation of a single scalar field.) Now let  $K_S$  be the space of field configurations on a hypersurface  $S$  bounding a region  $M$ . The guiding principle is here, that the amount of boundary data encoded in  $K_S$  should be such that it essentially uniquely determines a classical solution inside  $M$  in a generic situation (e.g.  $M$  a 4-ball). (Q1) The space of states  $H_S$  associated with  $S$  is the space of complex valued functions  $\mathbb{C}(K_S)$  on  $K_S$ . This means adopting a *state functional* picture. (Q2) The amplitude  $\mathcal{M}$  for a state  $\phi \in H_S$  is given by the expression

$$\mathcal{M}(\phi) = \int_{K_S} D\phi \int_{\mathcal{B}=0} D\phi_0 e^{\frac{i}{\hbar} S[\phi]};$$

The first integral is over field configurations  $\phi_0$  on  $S$ . The second integral is over all (not necessary classical) field

configurations  $\phi$  inside  $M$  that match the boundary data  $\phi_0$  on  $S$ .

Note that (Q1) gives a prescription for (T1) and ensures (T2) since for  $S = S_1 \sqcup S_2$  a disjoint union,  $K_S = K_{S_1} \sqcup K_{S_2}$  and hence  $\mathbb{C}(K_S) = \mathbb{C}(K_{S_1}) \otimes \mathbb{C}(K_{S_2})$ . (T4) is determined by (Q2). The dualization of boundaries corresponds simply to leaving the evaluation with a state on those boundaries open. Let  $M$  have boundaries  $S_1$  and  $S_2$  and consider states  $\phi_1 \in H_{S_1}$  and  $\phi_2 \in H_{S_2}$ . Then  $\mathcal{M}(\phi_1)$  is an element of  $H_{S_2}^*$ , i.e. a linear map  $H_{S_2} \rightarrow \mathbb{C}$  by mapping  $\phi_2$  to

$$\int_{K_{S_2}} D\phi_2 \int_{K_{S_1}} D\phi_1 \mathcal{M}(\phi_1, \phi_2) = \int_{K_{S_2}} D\phi_2 e^{\frac{i}{\hbar} S[\phi]};$$

This also explains (T3). The composition property (T5) is also rather obvious: Consider an integral over all field configurations in two regions with fields fixed on a common boundary and integrate also over the boundary values. Then this is the same as doing the unrestricted integral over field configurations in the union of the regions.

This heuristic quantization procedure based on the path integral thus leads to “general boundary” type quantum theories. The TQFT-like axioms (T1) - (T5) are automatically satisfied. I will refer to it as “holographic quantization” for the reasons given above. The quantization prescription is meant to be neither precise nor complete. In particular, one would usually “divide out” symmetries either from the configuration space  $K_S$  or from the functions  $\mathbb{C}(K_S)$  on it to arrive at the “physical” state space  $H_S$ . In quantum mechanics an example is the symmetrization or antisymmetrization for identical particles (see below). In quantum gravity an important step would be to divide out diffeomorphism symmetry (see also below).

## QUANTUM MECHANICS

Let me show how the standard formalism of quantum mechanics fits into the “general boundary” scheme. Space-time is now Euclidean or Minkowski space. I denote a point in space-time by coordinates  $(\mathbf{x}; t)$ . The background structure is the metric which is inherited by regions and their boundaries. The regions  $R$  we admit are time intervals  $[t_1; t_2]$  extended over all of space. The boundaries  $S$  are thus pairs of time-slices  $S = S_1 \sqcup S_2$  with  $S_1$  at  $t_1$  and  $S_2$  at  $t_2$ . According to (T2)  $H_S = H_{S_1} \otimes H_{S_2}$  and because of (T3)  $H_{S_2} = H_{S_1}^*$ . Indeed, let  $H_{S_1} = H$  be the Hilbert space of quantum mechanics and define  $U_R : H \rightarrow H$  to be the time-evolution operator  $e^{-\frac{i}{\hbar} H(t_2 - t_1)}$ . Then a state in  $H_S$  corresponds to a pair of states  $\phi \in H$  at time  $t_1$  and  $\psi \in H$  at time  $t_2$  (or a linear combination of such pairs). The transition amplitude between  $\phi$  and  $\psi$  is given by  $U_R$  via

$$\langle \psi | U_R(t_2 - t_1) | \phi \rangle = U_R(\phi, \psi); \quad (1)$$

<sup>1</sup> One might also want to admit the decomposition of connected boundaries, but this will not be discussed here.

The composition property (T5) encodes the composition of time evolutions. The inverse operation corresponds to the insertion of a complete set of states  $\int |j\rangle \langle j|$  at a given time.

So far we have only reformulated quantum mechanics. To show that a generalization to a “general boundary” formulation is possible it is necessary to consider a specific theory. The idea is to show that this theory arises as a holographic quantization. Then, the generalization should be straightforward, being determined by (Q1) and (Q2).

Let me specialize to the NRQM of scalar particles. Start with just one free particle. The action is  $S[\dot{x}] = \int_{t_1}^{t_2} dt \frac{1}{2} m \dot{x}^2(t)$  for a path  $\dot{x}(t)$  and a classical solution of the equations of motions is a straight line in space-time. It intersects each time-slice exactly once. Thus for a region  $R$  determined by a time interval  $[t_1; t_2]$  as above, denote the intersections with the boundaries  $S_1; S_2$  by  $\dot{x}_1; \dot{x}_2$ . The configuration space on the boundary  $K_S$  associated with  $S = S_1 \cup S_2$  which determines a classical solution uniquely is thus the space of pairs  $(\dot{x}_1; \dot{x}_2)$ , i.e.  $\mathbb{R}^3 \times \mathbb{R}^3$ . According to (Q1) we should set  $H_S = C(K_S) = C(\mathbb{R}^3 \times \mathbb{R}^3)$ . For the disconnected components we get that  $H_{S_1}$  and  $H_{S_2}$  can be identified with  $C(\mathbb{R}^3)$ . Indeed, this is the correct state space for a particle at time  $t$ , namely the space of fixed-time wave functions  $\psi_t(\dot{x})$ . An element  $(\dot{x}_1; \dot{x}_2)$  of  $H_S$  is generally a linear combination of products  $\psi_{t_1}(\dot{x}_1) \psi_{t_2}(\dot{x}_2)$ . (Q2) tells us that (2) is given by

$$Z = \int_{\mathbb{R}^3 \times \mathbb{R}^3} d\dot{x}_1 d\dot{x}_2 \int_{\substack{\dot{x}(t_1) = \dot{x}_1 \\ \dot{x}(t_2) = \dot{x}_2}} \mathcal{D}\dot{x} e^{\frac{i}{\hbar} S[\dot{x}]} \quad (2)$$

Here the second integral is over all paths  $\dot{x}(t)$  in the interval  $[t_1; t_2]$  with the given boundary values. This is indeed the correct expression for the transition amplitude (1) in the path integral formulation of NRQM. Note that we can easily generalize (2) to include a potential in the action.

The extension to several particles is rather obvious. For example, for two particles,  $K_S$  would be the space of quadruples  $(\dot{x}_1; \dot{y}_1; \dot{x}_2; \dot{y}_2)$ , while  $K_{S_1}$  would be given by pairs  $(\dot{x}_1; \dot{y}_1)$  etc.  $H_{S_1}$  would be given by  $C(K_{S_1}) = C(\mathbb{R}^3 \times \mathbb{R}^3)$ , i.e. fixed-time wave functions  $\psi(\dot{x}; \dot{y})$  of two particles. (2) is generalized in the obvious way with the path integral now over one path for each particle. This gives the correct results for *distinguishable* particles. For *identical* (and bosonic) particles we have to take for  $H_{S_1}$  the subspace of symmetric functions in  $(\dot{x}; \dot{y})$ . A different way to look at this is to replace the space  $K_{S_1}$  of ordered pairs by the space of unordered pairs. Of course this is not something coming out of the quantization prescription sketched above, but compatible with it. Note also that the path integral in (2) does not have to be explicitly symmetrized as it is always evaluated with symmetrized functions.




FIG. 1: A contribution to the path integral over  $R = A \cup B$  between boundaries  $S_1$  and  $S_2$ . The inner region  $B$  is crossed twice, corresponding to a two-particle state on its boundary  $S$ .

The next step is to see how using the rules (Q1) and (Q2) the NRQM of particles extends to a “general boundary” formulation. Consider a 4-ball shaped region  $B$  in space-time with boundary  $S$ . As with the spatial slices, a classical particle trajectory intersects  $S$  exactly twice. Thus, the configuration space is essentially  $K_S = S \times S$ . However, the entry time of the particle into  $B$  is necessarily earlier than the exit time. Thus  $K_S$  is really the subspace of  $S \times S$  where one point (say the first one) has a smaller time coordinate. By (Q1) then  $H_S$  is the space of functions  $(z_{in}; z_{out})$  on  $S \times S$  with this restriction. Here  $z$  denotes a parameterization of the hypersurface  $S$ . By (Q2) then we have a function  $\psi_B$  that associates amplitudes with such a generalized wave function. The physical interpretation is that of the amplitude of a particle being sent into the region  $B$  at  $z_{in}$  (which includes a time coordinate  $t_{in}$ ) and being observed emerging from  $B$  at time and place  $z_{out}$ .

So far everything seems straightforward. However, consider the following situation. Choose a time interval  $[t_1; t_2]$  containing  $B$  and the region of space-time  $R$  defined by it. Call its boundaries  $S_1; S_2$ . Cut out  $B$  from  $R$  and call the remainder  $A$ . Then, as  $R = A \cup B$  the composition rule (T5) requires that  $\psi_R$  equals the composition of  $\psi_A$  and  $\psi_B$ . Say we consider a one-particle state on  $S_1 \cup S_2$ . Then  $\psi_R$  contains an integral over paths from  $S_1$  at  $t_1$  to  $S_2$  at  $t_2$ , see Figure 1. Such a path may cross the inner region  $B$  an arbitrary number of times. However, we have taken above as the state space  $H_S$  associated with the boundary  $S$  between  $A$  and  $B$  the one for one particle. This only accounts for the paths in the integral that cross  $B$  exactly once. The composition rule seems to be violated. What went wrong?

Looking back at the definition of the one-particle state space  $H_S$  for  $S$  we see that already there is a problem. Namely, the path integral in the expression for  $\psi_B$  only constrains paths at their starting point and end point. There is no a priori restriction for them to lie entirely




FIG. 2: Allowing to “forget” connectivities introduces pair creation and annihilation. The connections (a;b) and (c;d) on the left are changed to (a;d) and (c;b) on the right.

inside B. However, we only want to allow to integrate inside B. Thus, how do we deal with paths that would leave B in between? The answer is rather obvious now. This corresponds to states with several particles on B. We need to let  $H_S$  be a direct sum of state spaces *for any number of particles*, i.e.  $H_S = H_S^0 \oplus H_S^1 \oplus H_S^2 \oplus \dots$  (the superscript indicates particle number). Now one

may restrict the path integral to paths inside B. The occurrence of all paths in a composition (such as with A) is ensured by the summation over all numbers of particles on S. This way we do obtain a consistent formalism, as guaranteed by the composition rule.

The surprising result is that one needs states with all possible numbers of particles even in NRQM. Perhaps this is not too surprising after all. The parallel to relativistic quantum mechanics is rather apparent. Namely, we can think of a boost there as tilting a fixed-time boundary (such as  $S_1$ ). The induced departure from fixed-time boundaries in the original frame has essentially the effect I have just described.

Note another subtlety. In the configuration spaces for the multi-particle states, one has to keep track which classical endpoint is connected to which other one by paths in the integral. If we remove this restriction we introduce pair creation and annihilation of particles. Consider Figure 2. On the left the connectivity inside B of the intersection points on the boundary S is described by the pairs (a;b) and (c;d). On the right this is exchanged to (a;d) and (c;b). The emerging picture might be interpreted as admitting an extra virtual particle that is created and subsequently annihilated.

Apart from the fact that we know this to happen in QFT there is good physical reason to admit this situation also here. Namely, being an observer on S, I see a particle coming into B at a and c and a particle emerging at b and d. If the particles are identical I have no way to say if it was *the same* particle that I saw twice or two different particles. In summary, it turns out that a “general boundary” formulation even of NRQM recovers essential features of *relativistic* quantum mechanics.

From the point of view of the holographic quantization prescription the “accident” of the failure of the naive (one-particle only) quantization is also explicable. The starting point of that prescription is a field theory. NRQM may be regarded as a field theory, but with the unusual

property that the value of the field  $\phi(\mathbf{x}(t))$  is at the same time a coordinate in space. This resulted then in the problems with the boundaries of the path integrals that had to be fixed.

## QUANTUM FIELD THEORY

Let us move to QFT. Space-time is Minkowski space and I denote coordinates by  $x = (\mathbf{x}; t)$ . I start by considering regions R determined by time intervals  $[t_1; t_2]$ . The encoding of Hilbert spaces and time evolutions in terms of the TQFT language is as described generically for quantum mechanics above. The first goal will be to show that the holographic quantization scheme reduces to the usual Feynman path integral quantization.

Consider a theory with a massive scalar field  $\phi(x)$ , for the moment free. The classical action is thus  $S[\phi] = \int_R dx \mathcal{L}(\phi, \partial\phi) = \int_R dx \left( \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \frac{1}{2} m^2 \phi^2 \right)$ . Considering two time-slices  $S_1; S_2$  at  $t_1; t_2$ , a classical solution to the field equations in the region R in between is essentially uniquely determined by the values of the field on  $S_1$  and  $S_2$ . Thus, we declare this to be the configuration space  $K_S$  associated with the boundary  $S = S_1 \cup S_2$  of R. It decomposes into  $K_S = K_{S_1} \cup K_{S_2}$  where each is the space of field configurations on the respective boundary. According to (Q1) the state space  $H_S$  is the space of functions  $C(K_S)$  on  $K_S$ . It decomposes into  $C(K_S) = C(K_{S_1}) \oplus C(K_{S_2})$ . Consider  $\phi_{x_1, \dots, x_n} \in H_{S_1}$  and  $\phi_{y_1, \dots, y_n} \in H_{S_2}$  given by  $\phi_{x_1, \dots, x_n}(1) = \phi_1(x_1) \dots \phi_1(x_n)$  and  $\phi_{y_1, \dots, y_n}(2) = \phi_2(y_1) \dots \phi_2(y_n)$ . Then, by (Q2) the amplitude of the corresponding state in  $H_S$  is given by

$$Z_{K_{S_1} \cup K_{S_2}} = \int_{K_{S_1}} \int_{K_{S_2}} D\phi_1 D\phi_2 \phi_1(x_1) \dots \phi_1(x_n) \phi_2(y_1) \dots \phi_2(y_n) e^{-S[\phi]} \quad (3)$$

We recognize this as essentially the transition amplitude between the “in” state  $|\phi(x_1; t_1) \dots \phi(x_n; t_1)\rangle$  at  $t_1$  and the “out” state  $|\phi(y_1; t_2) \dots \phi(y_n; t_2)\rangle$  at  $t_2$ . We may switch on even a perturbative interaction in the time interval  $[t_1; t_2]$  by modifying the action in the path integral accordingly.

It turns out that one recovers the complete quantum field theory correctly. To see this in terms of particle states one has to repeat the reduction procedure of Lehman, Symanzik and Zimmermann [4], adapted to the situation where initial and final states are at fixed times  $t_1; t_2$  and not in the infinite past or future. This will be detailed elsewhere. An essential role plays the concept of the vacuum as will be explained in [5].

Having once established the formalism for the special time-slice boundaries the generalization to arbitrary boundaries is straightforward following (Q1) and (Q2).

Here, no “accident” as for NRQM can happen as the fields have nothing to do with coordinates in space-time. States on time-slices can be “pulled-back” to any kind of boundary using the composition rule (T5), as ensured by the form of (Q2). The resulting description is not only local but also natural in terms of typical experimental setups.

Consider for example a scattering experiment in high energy physics. A typical detector has roughly the form of a sphere with the scattering happening inside (e.g. a collision of incoming beams). The entries for particles and the individual detection devices are arranged on the surface. At some time  $t_1$  the beam is switched on and at  $t_2$  it is switched off. The space-time region  $M$  relevant for the experiment is the region inside the sphere times the time interval  $[t_1; t_2]$ . The particle inflow and detection happens on the boundary  $S$  of  $M$ . What seems unusual is that the parts of  $S$  that are really interesting and carry the particle states are its *timelike* components. On the spacelike components at  $t_1$  and  $t_2$  there are no particles (we imagine the switching to happen smoothly). Concerning the interaction term in the Lagrangian it is now natural to turn it on only inside  $M$ . Indeed, the particles detected on the boundary  $S$  should (as usual) be thought of as free.

For calculational reasons it will usually be still advantageous to use particle states that are asymptotic. Indeed, the difference should have negligible effect on the resulting amplitudes as will be discussed elsewhere [5]. However, there no longer seems to be a *fundamental* reason to do this. This becomes rather important for the construction of a non-perturbative theory of quantum gravity. There, “asymptotic states” in terms of a Minkowski space are not expected to be a useful fundamental concept. The advantage of a local description is thus crucial.

## QUANTUM GRAVITY

Both, the “general boundary” formulation of quantum theories as well as the holographic quantization prescription are mainly designed for a quantum theory of gravity. Then, the background structure for the TQFT-type axioms is just that of differentiable manifolds and their boundaries. Going down to three dimensions it is well known that pure quantum gravity is a TQFT [6], i.e. satisfies (T1) - (T5). What is more, this TQFT is ob-

tainable by following the quantization prescription given by (Q1) and (Q2). Using connection variables the configuration space  $K_S$  associated with a boundary  $S$  of a region  $M$  is basically the space of flat spin connections on  $S$ . The path integral (Q2) is rigorously defined through a discretization of  $M$  as a spin foam model.

The role of diffeomorphism invariance is the following here: If we think of the spin connection as specified by a connection 1-form  $A(x)$  then  $K_S$  is the space of equivalence classes of such 1-forms under general gauge transformations. These general gauge transformations are now both, the  $SU(2)$  gauge transformations and diffeomorphisms.

I propose to approach also four-dimensional quantum gravity using the quantization prescription (Q1) and (Q2). Indeed, the path integral approach to quantum gravity is well established [7, 8]. The crucial new ingredient is the admission of arbitrary (in particular connected) boundaries and their interpretation [2]. A promising context for a non-perturbative realization of this appear to be spin foam models [9] in connection with a renormalization procedure [10].

I thank Carlo Rovelli for stimulating discussions and Thomas Schücker for a careful reading of the manuscript. This work was supported through a Marie Curie Fellowship of the European Union.

---

Electronic address: oeckl@cpt.univ-mrs.fr

- [1] M. Atiyah, Inst. Hautes Études Sci. Publ. Math. pp. 175–186 (1989).
- [2] R. Oeckl, *Schrödinger’s cat and the clock: Lessons for quantum gravity*, to appear in Class. Quantum Grav., Preprint gr-qc/0306007.
- [3] R. P. Feynman, Rev. Mod. Phys. **20**, 367 (1948).
- [4] H. Lehmann, K. Symanzik, and W. Zimmermann, Nuovo Cimento **1**, 205 (1955).
- [5] F. Conrady, L. Doplicher, R. Oeckl, C. Rovelli, and M. Testa, *Minkowski vacuum in background independent quantum gravity*, Preprint gr-qc/0307118.
- [6] H. Ooguri, Nucl. Phys. **B 382**, 276 (1992).
- [7] C. W. Misner, Rev. Mod. Phys. **29**, 497 (1957).
- [8] S. W. Hawking, in *General Relativity: An Einstein Centenary*, edited by S. W. Hawking and W. Israel (Cambridge University Press, Cambridge, 1979).
- [9] A. Perez, Class. Quantum Grav. **20**, R 43 (2003).
- [10] R. Oeckl, Nucl. Phys. **B 657**, 107 (2003).