

Superstring Orientifolds with Torsion: O5 Orientifolds of Torus Fibrations and their Massless Spectra

Michael B. Schulz

California Institute of Technology 452-48

Pasadena, CA 91125 USA

Using a "Superstrings with Torsion" type description, we study a class of IIB orientifolds in which space-filling O5 planes and D5 branes wrap the T^2 fiber in a warped modification of the product of 4D Minkowski space and a T^2 fibration. For the case that the base is T^4 , we provide examples that preserve 4D $N = 1, 2,$ and 3 supersymmetry, both with internal RR flux, and with a combination of internal RR and NS flux. In these examples, the internal geometries admit integrable complex structure; however, the almost complex structure selected by the supersymmetry conditions is nonintegrable in the case that there is NS flux. We indicate explicitly the massless spectrum of gauge fields and moduli in each example. In a previous investigation, this class of orientifolds was studied using T-duality. Here, we extend the previous analysis, first by providing an intrinsic description that does not rely on duality, and then by elaborating on details of the T-duality map, which we use to check our results.

Contents

1. Introduction	1
2. O5O orientifolds of T^2 Fibrations	9
3. Equations of Motion and Integrability Constraints	14
4. Supersymmetry Conditions	27
5. Relation to T-Dual O3O orientifolds	41
6. $N = 2$ Examples	48
7. $N = 3$ Example: The Warped Iwasawa Manifold	62
8. $N = 1$ Example	71
9. Preview: $N = 2$ Calabi-Yau Duals without Flux	78
10. Conclusions and Outlook	81
Appendices	84

1. Introduction

String theory admits an enormous set of seemingly consistent Minkowski vacua, yet the number of classes of compact geometries that have appeared in these vacua is small. With a few noteworthy exceptions [1,2,3], all supersymmetric compactifications of string theory to 4D Minkowski space studied until the past two years have involved only five types of special holonomy manifolds. The canonical building blocks include Calabi-Yau threefolds, K3 surfaces, and tori. Beyond these, the list also includes G_2 manifolds for M theory compactifications and Calabi-Yau fourfolds for F theory. It is natural to ask to what extent there exist sensible 4D Minkowski vacua based on other compactification geometries. The goal of this paper is to make contact between certain exotic orientifold vacua deduced via string dualities [4] and other work based on a "Superstrings with Torsion" type description [2].

In the absence of flux, when the geometry is a direct product of $R^{3,1}$ and a compact manifold, 4D $N = 1$ supersymmetry of the low-energy supergravity action demands that the compact manifold be a product of the special holonomy manifolds listed above [5]. In the case of type IIB string theory, a similar statement can also be made for a more general class of warped compactifications with internal flux and chiral spinor constraints. Here, the size of spacetime is allowed to vary over the internal manifold, and the data defining the compactification is enlarged to include a discrete choice of branes and internal NS and RR flux. For the class that has received the most attention, chiral (Becker-type) constraints are also imposed on the spinors generating the supersymmetries; these are the constraints associated with D3 branes and D7 branes [6].

Such D 3/D 7 warped compactifications are attractive for a number of reasons. They provide a stringy embedding [7,8,9] of the Randall-Sundrum approach to the hierarchy problem [10], as well as a mechanism for perturbative stabilization of moduli [3,9,11,12,13,14,15]. (See Refs. [16,17,18] for a gauged supergravity perspective; also, see Ref. [19], for a discussion of local models that includes a worldsheet perspective.) In addition, they are amenable to a statistical analysis as an ensemble of vacua [20], and with a few new ingredients, have led to a construction of metastable de Sitter vacua of string theory [21].¹

However, for compactification to 4D Minkowski space they require the same special holonomy manifolds as do conventional compactifications, up to an overall conformal rescaling by the warp factor [6]. Schematically, the reason is as follows. For supersymmetric IIB vacua, the gravitino variation is

$$\delta \psi_M = (D_M + \alpha \text{ux}) \psi + (\beta \text{ux}) \psi = 0;$$

where ψ is a 10D Weyl spinor. In the case of Becker-type constraints, the α and β terms separately vanish. The β term decomposes into the usual covariant constancy condition plus a relation between the RR flux F_5 and the warp factor. The α term gives conditions on the complex three-form $G_{(3)}$.

For other spinor constraints, or for vacua other than type IIB warped compactifications, the fluxes can mix with covariant derivatives. One obtains generalized covariant constancy conditions based on torsionful connections, whose solutions, if any, involve internal manifolds not appearing in the list above. These exotic compactification manifolds can be non-Kähler and even non-complex. However, while it is easy to write down the equations of motions and supersymmetry conditions, it is not so easy to find solutions. Given certain assumptions, one can prove no-go theorems on their existence [26,9,27].²

¹ An explicit model of this type with all Kähler moduli stabilized was recently exhibited in Ref. [22]. See also Ref. [23] for a nonexistence proof for one Kähler modulus, and Ref. [24] for a related discussion in the $T^6 = Z_2$ orientifold. For work on inflation and de Sitter space in models with D 3 branes and D 7 branes, see Ref. [25].

² The no-go theorems state that in the absence of negative-tension sources, the leading α' -order supergravity action has no solutions with internal flux and compact internal manifold. These theorems are evaded in string theory by the existence of negative-tension orientifold planes, by α' -R \wedge R D 7 brane worldvolume curvature couplings (which give D 7 branes negative D 3 brane charge and tension), and α' -R \wedge R corrections to the heterotic Bianchi identities.

1.1. Older Progress

Most of the progress toward understanding torsionful supersymmetric vacua builds on work performed by Strominger nearly two decades ago, in the context of the heterotic string with NS flux [2]. In the heterotic theories, the gravitino and dilatino variations are

$$\begin{aligned} \delta\psi_M &= \gamma_M \psi + \frac{1}{4} H_{(3)M} \lambda; \\ \delta\lambda &= \not{D}\lambda + \frac{1}{2} H_{(3)} \lambda; \end{aligned} \tag{1.1}$$

where ψ is a 10D Majorana-Weyl spinor, and $\lambda = (u + u^c) \frac{\gamma^{\underline{p}}}{2}$ in terms of a 4D Weyl spinor u and 6D Weyl spinor λ . (Strominger did not specialize to compactifications to 4D, but we will do so here). For supersymmetric vacua, the gravitino variation directly implies that ψ and λ are both covariantly constant with respect to the same connection of torsion $T_{bc}^a = H_{(3)}^a{}_{bc}$. If we define an almost complex structure (ACS)

$$J_a{}^b = i \gamma_a{}^b; \tag{1.2}$$

then Eqs. (1.1) together imply that the ACS is integrable. Lowering one index on $J_a{}^b$ gives the fundamental form 3J , which is related to the flux via

$$\not{D} e^2 J = {}_6H_{(3)}; \tag{1.3a}$$

$$d e^2 {}_6J = 0; \tag{1.3b}$$

Eqs. (1.3) are not quite the relations that appeared in Strominger's work [2], but are equivalent to them [28,27].

Strominger went on to obtain a complete set of geometrical conditions for supersymmetry, as well as a set of equations expressing the dilaton and flux in terms of the geometry. In addition, he provided a worldsheet description, and for compactifications to 6D was able to give exact solutions. For compactifications to 4D on a compact manifold other than a Calabi-Yau, he argued that the solutions could not extend to large volume, since his equations reduced to Calabi-Yau conditions in this limit. He did not provide any solutions, but did mention the Iwasawa manifold as an example of a complex non-Kähler manifold satisfying the topological condition $h^{3,0} = 1$, that could conceivably be used as the basis for a consistent torsionful supersymmetric solution.⁴

³ In the special case that J is closed, the internal manifold is Kähler and J is the Kähler form.

⁴ Heterotic compactifications on the Iwasawa manifold have been re-explored using the tools discussed in the first part of Sec. 1.2 [29]. In Sec. 7, we discuss orientifolds based on a warped Iwasawa manifold.

This is the way things stood for thirteen years until the first non-Kähler heterotic compactifications to 4D were constructed using string dualities [3]. The starting point was a IIB warped compactification, obtained as the orientifold limit of a warped F-theory compactification on $K3 \times K3$ with internal flux [30,31]. After two T-dualities,⁵ this theory becomes type I on a non-Kähler manifold with RR three-form flux. The heterotic theory is then obtained via S-duality. The resulting geometry takes the form of a warped T^2 fibration over $K3$, with the noteworthy property that the fiber volume is fixed in terms of the fiber complex structure modulus. So, lengths cannot simply be scaled by an overall factor to obtain a large volume solution, in agreement with Strominger's earlier observation forbidding an overall volume modulus.

1.2. More Recent Progress

Two important organizing principles that have emerged more recently are the notions of G-structures and generalized calibrations. These tools were first applied to studying supergravity solutions with background flux in Ref. [28]. Since then, the subject has flourished [32,29,33,27,34,35,36,37,38,39,40,41,42,43,44]. In the context of the heterotic string with 4D $N = 1$ supersymmetry, the appropriate group G is $SU(3)$, and the idea is as follows [32,29,27]. The existence of the covariantly constant spinor discussed above is equivalent to the statement that the compactification manifold X_6 has $SU(3)$ holonomy with respect to the torsionful connection. As a consequence of the existence of a privileged 6D spinor, the usual $SU(3)$ structures J and ω can be canonically defined in terms of this spinor, just as for a Calabi-Yau. However, these objects are no longer closed. A useful mathematical characterization of the precise sense in which X_6 differs from a Calabi-Yau comes from the fact that the $SU(3)$ structures induce a natural $SU(3)$ decomposition of the torsion into five torsion classes [45]:

$$T = \begin{matrix} W_1 \\ (1+1) \end{matrix} + \begin{matrix} W_2 \\ (8+8) \end{matrix} + \begin{matrix} W_3 \\ (6+6) \end{matrix} + \begin{matrix} W_4 \\ (3+3) \end{matrix} + \begin{matrix} W_5 \\ (3+3) \end{matrix} : \quad (1.4)$$

In terms of this decomposition,

$$\begin{aligned} dJ &= \frac{3}{4}i(W_1 - W_1) + W_3 + J \wedge W_4; \\ d\omega &= W_1 J \wedge J + J \wedge W_2 + \omega \wedge W_5; \end{aligned} \quad (1.5)$$

⁵ The geometry of the IIB orientifold is a warped version of $R^{3;1} \times K3 \times T^2 = (1)^{F_1} I_2$, where I_2 inverts the T^2 . The T-dualities are performed in the T^2 directions.

The supersymmetry conditions can then be translated into constraints on the W_1 . Conversely, once these constraints are known, we can instead start with a manifold satisfying the constraints, and then $H_{(3)}$ and e follow from Eqs. (1.3).

The connection to generalized calibrations arises when, following Ref. [28,27], we interpret Eq. (1.3a) as a consequence of the fact that (i) NS5 branes are a source of $H_{(3)}$ flux, and (ii) J is a generalized calibration [46] for the two-cycles on which we can wrap a supersymmetric NS5 brane probe. In some sense, we can think of the manifold X_6 as a fully backreacted geometry involving one or more wrapped NS5 branes. The appeal of this formalism is that it readily generalizes. Instead of considering the heterotic string on a 6D manifold with a torsionful connection of $SU(3)$ holonomy, we can consider type I, II, or the heterotic theories on a manifold of some other dimension, with torsionful connection of holonomy group G . The $SU(3)$ structures J and e just get replaced by the appropriate G -structures. The generalization of Eq. (1.3a) is an analogous relation between some NS or RR flux and the G -structure that calibrates the cycles on which we can wrap a corresponding NS or D brane probe.

Beyond G -structures, another important advance has been the reformulation of the heterotic moduli constraints in a way analogous to the usual formulation of the constraints for IIB Becker-type warped compactifications. For the latter, one defines a complex flux $G_{(3)} = F_{(3)} - \text{dil}H_{(3)}$. The supersymmetry conditions constrain the moduli so that $G_{(3)}$ is primitive⁶ and of type (2,1). The (2,1) condition follows from a superpotential $W = R G_{(3)} \wedge$. Finally, the scalar potential comes from Kaluza-Klein reduction of the kinetic term for $G_{(3)}$ in the 10D supergravity action. A similar structure has been exhibited in the heterotic theories, except that the scalar potential now comes from a combination of the $H_{(3)}$, Einstein-Hilbert, and dilaton kinetic terms in the 10D supergravity action [47]. The complex flux $H_{(3)} = e^2 \text{dil}^{-2} J$ is required to be (2,1) and primitive. It has been argued that the (2,1) condition follows from an analogous superpotential [48,47,49], although this is more subtle than in the IIB case.

The duality chains, which proved so useful in the first non-Kahler heterotic compactifications to four dimensions, have also been revisited. In the original construction [3], all analysis was performed in the dual IIB/F-theory description. Much effort has been devoted to providing an intrinsically heterotic description of these vacua and their generalizations, as well as to the mathematical properties of the compact manifolds involved [48,50,49].

⁶ Primitivity means that $J \wedge G_{(3)} = 0$.

From the results of Ref. [49]: the previously known class of admissible compact manifolds obtained via duality has been generalized to include manifolds with nonzero Euler characteristic, and a better understanding now exists of phenomenologically relevant details concerning the vector bundles on these manifolds and the numbers of fermion generations; moreover, through a different duality, there is now a relation to brane-box constructions. A particular duality symmetry that has proven extremely useful in the past is mirror symmetry, and steps have been taken toward generalizing Calabi-Yau mirror symmetry to a mirror symmetry of non-Kähler manifolds with nonvanishing NS flux [4,32,38]. Even for geometric transitions, the cornerstones in our modern understanding of topological string theory [51], there now exists a sequence of supergravity descriptions for a complete duality cycle connecting the various IIB, IIA, and M theory descriptions before and after the transition [52]. This duality cycle relates D-branes wrapped on cycles of non-Kähler manifolds to fluxes in other non-Kähler manifolds. For other interesting results involving dualities and torsion, see Ref. [53].

1.3. Work Reported Here

A natural way in which duality symmetries have been applied is in the construction of a new class of non-Kähler orientifold vacua [4]. These vacua will be our focus here. They are relatively simple to describe: the geometry is a warped torus fibration over a torus base, with O planes and D-branes wrapping the fiber and filling spacetime. In addition, there is internal flux. These vacua were constructed using a duality argument similar to the one used to obtain non-Kähler heterotic compactifications. However, the orientifold vacua that we discuss here do not have obvious geometrical heterotic duals. The starting point is a T^6 orientifold of type IIB with internal flux, where the orientifold inverts all of the torus directions. So, the initial theory contains $O3$ planes and possibly D3-branes, but no $O7$ planes or D7-branes. The non-Kähler orientifolds are obtained via T-duality. For supersymmetric flux, the maximum number of T-dualities that can be performed is either two or three,⁷ depending on the choice of flux. Therefore, we cannot relate these string vacua to type I with $O9$ planes, or subsequently to the heterotic theory by S-duality.

⁷ After this number of T-dualities, the metric has no further isometries that can be used to perform additional standard T-dualities. There does exist a generalized sense in which we can perform additional T-dualities. However, the result is a nongeometrical string compactification [54,4], involving duality twists that mix the metric and NSB-field, so that neither of these quantities is globally well-defined by itself. In contrast, standard T-dualities only result in geometric fibrations (Scherk-Schwarz [55] twists of the metric) and NS flux (Scherk-Schwarz twists of the NSB-field).

The main motivation for the present investigation is that the orientifolds just described have been analyzed in a way that relies almost exclusively on the original pre-duality theory, with little or no intrinsic description in the final non-Kähler orientifold. On the other hand, the geometry of these orientifolds is very similar to that of certain noncompact manifolds of $SU(2)$ or $SU(3)$ structure that have been described elsewhere, and used in consistent supergravity solutions with a "Superstrings with Torsion" type description [27]. The primary goal below is to provide such a description for torsionful orientifolds of type IIB string theory in which $O5$ planes are wrapped on the fiber of a T^2 fibration. A secondary goal is to study moduli stabilization in these orientifolds.

An outline of the paper is as follows. We begin in Sec. 2 by defining the class of $O5$ orientifolds that we consider. Sec. 2.1 contains a description of the 10D supergravity background, including the 10D metric, the orientifold operation, and fluxes. In Secs. 2.2 and 2.3 we discuss the fields and energy scales of the 4D effective field theory.

Sec. 3 is devoted to analyzing the model to the extent possible without any assumptions regarding supersymmetry. Here, and also in Sec. 4, we relax the ansatz that the base of the internal manifold is T^4 , and assume only that it is some compact manifold B . However, order to proceed systematically, we find it necessary to impose one restriction on the NS flux. This restriction is discussed at the end of Sec. 3.1. The main results of Sec. 3 are the Gauss's law constraint in Sec. 3.2, a pseudo-BPS constraint on the fluxes in Sec. 3.3, expressions for the dilaton and warp factor in Secs. 3.4 and 3.5, and the requirement that the base be Ricci flat. The pseudo-BPS constraint follows from an equality between an exact four-form on the base and sum of positive semidefinite squares of fluxes. Integration of this relation imposes a set of Hodge duality conditions on the fluxes. There is a naive paradox concerning the Gauss's law constraint. Due to the fibration, one can have $d\mathbb{F}_{(3)}$ supported on local sources, the sum of whose charges does not vanish. This paradox is resolved by correctly pushing forward the $\mathbb{F}_{(3)}$ Bianchi identity to the base. We conclude the section with a precise definition of the moduli, and a decomposition of the fluxes into a quantized discrete part and moduli dependent deformations.

Sec. 4 contains a discussion of the supersymmetry conditions. After a brief discussion in Sec. 4.1 of the constraints on the spinors from the orientifold projection, in Sec. 4.2 we analyze the case in which only the RR three-form flux is nonzero. Here, the discussion closely parallels that given by Strominger for the heterotic string with NS flux. The vacua that we describe are related by S-duality to dual vacua with NS-flux, NS5 branes, and $O5$ planes, that are very similar to those discussed by Gauntlett, Martelli, and Waldram

[27] in the case of 6D internal manifold. The S-duality map is discussed in Sec. 4.3. We then consider the case of more general flux in Sec. 4.4, and derive the corresponding generalization of the supersymmetry conditions. These conditions are stated in terms of the flux, the right-SU(3) structures J and \tilde{c} constructed from the right-moving part of the Killing spinors only (or alternatively, left-SU(3) structures), and the volume form Vol_6 on the T^2 fiber. Note that while we employ SU(3) structures in analyzing the supersymmetry conditions, we do not find it useful here to work in terms of SU(3) (or SU(2)) torsion classes in order to satisfy these conditions. Therefore, we will not compute torsion classes in this paper, except for a meeting instance in Sec. 6.3 to verify the nonintegrability of the almost complex structure. See, however, the discussion in Sec. 10.

In Sec. 5, we relate O5 orientifolds with T^4 base to T-dual O3 orientifolds with internal T^6 . Then, in Secs. 6–8 we study a number of specific classes of O5 backgrounds preserving various amounts of supersymmetry, and their O3 duals. Sec. 6.3 is devoted to a class with nonvanishing $F_{(1)}$ and $H_{(3)}$ flux, and nonintegrable complex structure, but the examples in Secs. 6–8 otherwise contain only $\mathbb{F}_{(3)}$ flux and integrable complex structure. Our discussion of the O3 duals extends the known results for the $T^6=\mathbb{Z}_2$ orientifold.

Sec. 9 contains a preview of work to appear elsewhere [56]. In the $N = 2$ case, our examples are dual to M theory on a Calabi-Yau threefold Y_6 times a circle. That is, they can be viewed as F theory on $Y_6 \times T^2$, where the F theory fiber is taken to be the product of an S^1 in the T^2 and an S^1 in Y_6 (with Y_6 viewed as a fibration). Here, we provide a few preliminary observations concerning this duality.

Finally, in Sec. 10, we conclude and discuss possibilities for future work.

The appendices contain auxiliary details not found in the body of the paper. App. A contains a summary of our conventions, mostly with regard to Dirac matrices. In App. B, we describe the twisted coordinate identifications that generalize the T^6 identifications $x^a = x^a + 1$ to a T^2 fibration over T^4 . In App. C, we derive the first equation of Sec. 3.3, from which the pseudo-BPS constraints follow. App. D is a reference for the IIB supersymmetry variations of fermions in various forms (string frame, Einstein frame, in terms of L, R and in terms of $\psi = L + iR$). In App. E, we discuss subtleties in applying the superpotential of Gukov, Vafa, and Witten [57] to the $T^6=\mathbb{Z}_2$ O3 orientifold in which the fluxes (partially) break $N = 4$ extended supersymmetry. The T-duality map of RR moduli, as defined in Secs. 3.9 and 5.1, is worked out in App. F. App. G contains a discussion of moduli space metrics for the $T^6=\mathbb{Z}_2$ orientifold; here, we first review the results of Frey and Polchinski for the case of $N = 3$ supersymmetry, and then consider the $N = 2;1$

cases relevant to Secs. 6 and 8. Lastly, in App. H we prove that a particular class of fluxes discussed in Sec. 8.2 reduces to a unique choice of flux modulo the discrete identifications of the axion-dilaton and T^6 complex structure.

2.05 Orientifolds of T^2 Fibrations

2.1. Supergravity Background

As a starting point, recall that the supergravity background for N coincident D5 branes in flat noncompact spacetime is [58]

$$ds_{\text{string}}^2 = Z^{-1/2} (dx^0 dx^0 + Z^{1/2} g_{mn} dx^m dx^n); \quad (2.1a)$$

$$e^{-2\phi} = g_s Z^{-1/2}; \quad (2.1b)$$

$$F_{(7)} = g_s^{-1} d(Z^{-1}) \wedge dx^0 \wedge \dots \wedge dx^5; \quad (2.1c)$$

where $\mu, \nu = 0, \dots, 5$ and $m, n = 6, \dots, 9$. The subscript "string" indicates that Eq. (2.1a) gives the string frame metric. The warp factor is

$$Z = 1 + N \frac{g_s}{2r^2}; \quad (2.2)$$

This function is harmonic away from $r = 0$ and is a solution to Poisson's equation

$$\nabla_m \nabla_m Z = N (2\pi)^2 g_s^{-4} \delta^{(4)}(x) \quad (2.3)$$

in the non-warped metric g_{mn} on the R^4 transverse to the D5 branes.

We would like to study the analogous supergravity backgrounds for orientifolds of type IIB string theory in which 4D spacetime-filling O5 planes and D5 branes wrap the T^2 fiber over a T^4 base.⁸ The new metric ansatz is

$$ds_{\text{string}}^2 = Z^{-1/2} (dx^0 dx^0 + ds_{T^2_{\text{fib}}}^2) + Z^{1/2} ds_{T^4_{\text{base}}}^2; \quad (2.4a)$$

where $ds_{T^2_{\text{fib}}}^2$ is the 4D Minkowski metric, and the fiber and base metrics are

$$ds_{T^2_{\text{fib}}}^2 = g^{(T^2_{\text{fib}})}_{\mu\nu} dx^\mu dx^\nu; \quad \mu, \nu = 0, 1; \quad \mu, \nu = 4, 5; \quad (2.4b)$$

$$ds_{T^4_{\text{base}}}^2 = g^{(T^4_{\text{base}})}_{mn} dx^m dx^n; \quad m, n = 5, \dots, 9; \quad (2.4c)$$

⁸ Although we focus on the case of T^4 base, most of the analysis in Secs. 3 and 4 is performed for arbitrary compact base B .

Here, we assume that $g^{(T_{\text{fib}}^2)}$ and $g_m^{(T_{\text{base}}^4)}$ are metrics, and that the warp factor depends only on the base coordinates. The fiber and base coordinates are identified via a twisted generalization of $x^m = x^m + 1$, $x^n = x^n + 1$ that is discussed in App. B. The quantity $A = A_m f x^n g dx^m$ is the fiber connection. The one-forms α^m are globally defined and are the generalizations of dx^m consistent with the twisting of the fibration. Nontrivial twisting is measured by nonvanishing cohomology classes

$$[F] \in 2H^2(T_{\text{base}}^4; 2Z); \quad (2.5)$$

where

$$F = dA; \quad (2.6)$$

These are the Chern classes of the x^m circle fibrations (or equivalently $U(1)$ fibrations) over T_{base}^4 .

For later convenience, we define a 6D metric

$$ds_6^2 = ds_{T_{\text{fib}}^2}^2 + Z ds_{T_{\text{base}}^4}^2; \quad (2.7)$$

so that

$$ds_{\text{string}}^2 = Z^{-1/2} (dx^m dx^m + ds_6^2); \quad (2.8)$$

We denote by X_6 the 6D space whose metric is (2.7).

In addition to this geometry, the model includes 16 O5 planes and $2M$ D5 branes. The Z_2 orientifold operation that defines the O5 planes is I_4 , where ω is worldsheet parity and I_4 inverts the base T^4 :

$$I_4: x^m \rightarrow -x^m; \quad (2.9)$$

The orientifold planes wrap the fiber and are located at the $2^4 Z_2$ fixed points on the base where $x^m = 0; 1=2$.⁹ The $2M$ D5 branes also wrap the fiber, and are located at M arbitrary points on the base together with their $M Z_2$ images. The $2Z$ rather than Z valued cohomology in Eq. (2.5) ensures that $[F] \in 2H^2(T^4=I_4; Z)$, which is needed to define the orientifold. (See Sec. 3.7).

⁹ The orientifold planes are assumed to be standard O5 planes in the terminology of [59], as opposed to the exotic O5⁺ or $\tilde{O}5$ planes that lead to shifted flux quantization conditions. Similarly, we assume that there is no localized F curvature at the I_4 fixed points.

Finally, the orientifold projection preserves the following even fluxes together with their 10D Hodge duals:¹⁰

$$\text{Even fluxes: } \mathbb{F}_{(1)m}^{\mathbb{E}}; \mathbb{F}_{(3)mn}^{\mathbb{E}}; \mathbb{F}_{(5)mnp}^{\mathbb{E}}; H_{(3)m}; H_{(3)mnp}: \quad (2.10)$$

By even, we mean that these components must be even functions of the transverse coordinates x^m . The orientifold projection also preserves the complementary set of components, provided these components are odd functions of the transverse coordinates. With the exception of the special case that $2M = 32$ and each O5 plane is coincident with exactly one D5 brane and its image (so that charges and tensions cancel locally), there do not exist string vacua with all odd fluxes set to zero. In order to satisfy the low energy equations of motion, we need to include at least the following odd flux:

$$\text{Odd flux: } \mathbb{F}_{(3)mnp}^{\mathbb{O}}: \quad (2.11)$$

We will limit our investigation to those backgrounds in which the other odd fluxes can be consistently set to zero, and in which all fields have functional dependence on the base coordinates only. (In other words, we consider the low energy effective field theory at energy scales below $1/R_b$; see Sec. 2.3 for further discussion.) In addition, we assume vanishing field strengths of the D5 worldvolume gauge fields.

The fluxes $\mathbb{F}_{(p)}$ are the gauge-invariant fluxes that appear in the kinetic terms of the bulk supergravity action. In the conventions in which the T-duality action on RR potentials is simplest,¹¹ the relation between the RR potentials $C_{(p)}$ and gauge-invariant fluxes $\mathbb{F}_{(p)}$ is

$$\mathbb{F}_{(p)} = \begin{cases} F_{(p)} & C_{(p-3)} \wedge H_{(3)} & p = 3, \\ F_{(p)} & & p = 1; 2, \end{cases} \quad (2.12)$$

where

$$F_{(p)} = dC_{(p-1)}: \quad (2.13)$$

In Sec. 3, we will see that the odd flux (2.11) is completely determined by the equations of motion. On the other hand, the even fluxes (2.10) contain both moduli-dependent and moduli-independent components, with the latter constrained only by Dirac quantization.

¹⁰ Here, all components of fluxes and potentials are given in the basis $dx^0; \dots; dx^m$. These components should not be confused with those in the basis $dx^m; dx^{\tilde{m}}; dx^{\tilde{m}}$.

¹¹ There is different convention, $\mathbb{F}_{(5)} = F_{(5)} + \frac{1}{2}B_{(2)} \wedge F_{(3)} - \frac{1}{2}C_{(2)} \wedge H_{(3)}$, in which the potentials $B_{(2)}$ and $C_{(2)}$ transform simply under $SL(2; Z)$ duality.

Therefore, the even fluxes, like F and M , contain discrete data that needs to be specified in order to fully define the model.

For trivial bration and in the absence of even flux, this orientifold is dual to type I on T^6 via T-duality in the four T_{base}^4 directions. In this case, the orientifold preserves 4D $N = 4$ supersymmetry. Also, since the D5 branes and O5 planes fill 4D spacetime, there is a Gauss's law constraint (RR tadpole cancellation condition in worldsheet language) that requires $2M = 32$ D5 branes. This constraint and its generalization to the case of nontrivial bration and flux are discussed in Sec. 3.2 below.

2.2. Low Energy Bosonic Field Content

For the class of 10D supergravity backgrounds just described, there exist deformations that continuously connect different consistent supergravity solutions. Let us restrict to deformations with no functional dependence on x . For the case of trivial bration and in the absence of even flux, one class of such deformations consists of the zero modes on X_6 of the even bosonic fields

$$\begin{aligned} \text{RR sector: } & C_{(2)} ; C_{(2)} \text{ scalar}; C_{(2)mn}; C_{(4)mnp}; \\ \text{NSNS sector: } & g ; g_{mn}; B_{(2)m}; \end{aligned} \quad (2.14a)$$

together with the zero modes on the T^2 fiber of the D5 worldvolume fields

$$\text{D5 worldvolume: } A_I ; \varphi_I^m ; I = 1, \dots, M : \quad (2.14b)$$

These are the deformations which, when promoted to 4D fields, become the moduli of the 4D $N = 4$ low energy effective field theory.

The quantity that plays the role of the axion-dilaton is (cf. Sec. 4.2)

$$\varphi_{\text{dil}} = \frac{1}{(2\pi\alpha')^2} \left(C_{(2)45} + \frac{i}{g_s} V_b \right) ; \quad (2.15)$$

where the volume of the T^2 fiber is

$$V_b = g^{(T_{\text{fib}}^2)} \varphi^2 ; \quad (2.16)$$

This expression for φ_{dil} is T-dual to the more familiar expression given App. D that is common to O3 orientifolds, O7 orientifolds, and pure IIB string theory.

A second class of such deformations consists of the zero modes on X_6 of the even fields

$$\begin{aligned}
\text{RR sector: } & C_{(2)} \ ; \ C_{(4) \ m} \ ; \\
\text{NSNS sector: } & V \ ; \ B_{(2)m} \ ; \\
\text{D5 worldvolume: } & A_I \ ; \ I = 1, \dots, M \ .
\end{aligned}
\tag{2.17}$$

These are the deformations which, when promoted to 4D fields, become the massless gauge bosons of the 4D $N = 4$ low energy effective field theory. Here, V is the deformation that corresponds to the Kaluza-Klein gauge boson for the translation isometry in the x^5 -direction. It arises by replacing A with $A + V dx^5$ in the 10D metric (2.4b).

Finally, a third class of such deformations shifts the 4D metric away from $g_{\mu\nu}$ and parametrizes the space of constant metrics on $R^{3,1}$. When promoted to 4D fields, these deformations become the 4D graviton.

Note that there is an important distinction between the allowed x^5 -independent deformations of the supergravity background and the fields of the 4D effective action. While it is easy to identify the former, the massless fields in the dimensional reduction to 4D are more complicated than those obtained by simply endowing these deformations with x^5 -dependence. The correct low energy 4D fields must also include warp factor dependence to prevent mixing with higher Kaluza-Klein modes [60,15,61].

In the case of nontrivial bration and flux, the supersymmetry and massless field content is reduced. The $U(1)^M$ $N = 4$ vector multiplets on the D5 branes remain massless, but many of the RR and NS fields in (2.14) and (2.17) are lifted. The lifting occurs through a supersymmetric Higgs mechanism. For the bosons, the equations of motion impose certain metric and axion-dilaton dependent Hodge duality relations on the fluxes that lift a subset of the NS moduli and the zero mode of the RR axion $C_{(2)45}$. The vevs of some of these moduli then break a subset of the gauge symmetries, and their axionic partners (zero modes of some of the $C_{(2)m_n}$ and $C_{(2)m_{np}}$) are eaten by massive vectors. The fermions also take part in the superHiggs mechanism, and the various possibilities for the resulting $N < 4$ low energy spectrum are determined by the masses of the $(4 - N)$ massive gravitini, as described in Ref. [17]. (See also Ref. [16]).

2.3. Energy Scales

There are two points of view that we can adopt regarding the moduli stabilization and symmetry breaking just described. If we assume that the fiber and base have roughly the

same length scale R , then these effects occur at the Kaluza-Klein scale $l=R$. However, if we assume a hierarchy $R_b \ll R_{\text{base}}$, then these effects occur at the scale $m = R_b = R_{\text{base}}^2$ [32], which is much smaller than $m_{\text{base}}^{\text{KK}} = 1/R_{\text{base}}$ and $m_b^{\text{KK}} = 1/R_b$:

$$m : m_{\text{base}}^{\text{KK}} : m_b^{\text{KK}} = 1 : R_{\text{base}} = R_b : R_{\text{base}} = R_b^2 : \quad (2:18)$$

In the former case, it only makes sense to talk about a low energy $N < 4$ theory with all massive fields integrated out. In the latter, there is a consistent low energy 4D supergravity theory within which $N = 4$ is softly broken to $N < 4$ by a supersymmetric Higgs mechanism. We will adopt the latter point of view below. In either case, we need $R_{\text{base}} \gg R_b$ in order to be able to neglect higher string modes.

3. Equations of Motion and Integrability Constraints

Before imposing the supersymmetry conditions, let us first determine the constraints that follow from equations of motion alone.

3.1. Equations of Motion

The trace-reversed Einstein equations are

$$R_{MN} = \mathbb{T}_{MN} ; \quad (3:1)$$

where $\mathbb{T}_{MN} = T_{MN} - \frac{1}{8}g_{MN}T$.

The components of the Ricci tensor in the metric (2:4) are

$$\begin{aligned} R_{\mu\nu} &= r_{\mu\nu} \log Z + \frac{1}{4}Z^{-1}r_{\mu\nu}^2 \log Z ; \\ R_{\mu B} &= r_{\mu B} \log Z + \frac{1}{4}Z^{-1}r_{\mu B}^2 \log Z + \frac{1}{4}F_{\mu\nu}^2 ; \\ R_{mn} &= r_{mn} \log Z - \frac{1}{4}Z^{-1}r_{mn}^2 \log Z - \frac{1}{2}F_{mn}^2 \\ &\quad - \frac{1}{2}(r_{Bm} \log Z)(r_{Bn} \log Z) + R_{Bmn} ; \end{aligned} \quad (3:2)$$

Here B denotes the 4D base of the torus fibration. Most of the results of Secs. 3 and 4 are independent of the choice $B = T^4$, so we work with arbitrary B . The operator r_B is the Levi-Civita covariant derivative on the base, and the Laplacian r_B^2 is contracted using the base metric

$$ds_B^2 = g_{Bmn} dx^m dx^n \quad (3:3)$$

that generalizes Eq. (2:4c).

In string frame, the trace-reversed stress tensor is $\hat{T}_{MN} = T_{MN}^{NS} + T_{MN}^{RR} + T_{MN}^{local}$, where

$$\begin{aligned} T_{MN}^{NS} &= 2r_M r_N + \frac{1}{2} \frac{1}{2!} H_{(3)MN}^2 ; \\ T_{MN}^{RR} &= \frac{1}{2} e^2 \sum_{p=1;3} \frac{1}{(p-1)!} F_{(p)MN}^2 - \frac{1}{2} g_{MN} \frac{1}{p!} F_{(p)}^2 + \frac{1}{4} e^2 \frac{1}{4!} F_{(5)MN}^2 - \frac{1}{2} g_{MN} \frac{1}{5!} F_{(5)}^2 ; \\ T_{MN}^{local} &= (2)^2 \frac{01}{2} e \sum_i (h_{MN}(x_i) - \frac{1}{2} g_{MN}) Q_i \frac{4(x-x_i)}{Z P g_B} ; \end{aligned} \quad (3:4)$$

Here, $h_{MN}(x_i)$ is the restriction of the 10D metric (2:4) to the point $x = x_i$ in the base,

$$ds_{h(x_i)}^2 = Z^{1=2} dx^6 dx^7 + g^{(T_{fib}^2)} dx^m dx^m : \quad (3:5)$$

The sum on i runs over D5 brane and O5 plane sources of charge Q_i in units of D5 brane charge, localized at points $x = x_i$ on the base. For $B = T^4$, this includes $2M$ D5 branes of charge $Q_i = +1$ and 16 O5 planes of charge $Q_i = -2$ located at the Z_2 fixed points. We work on the covering space of the orientifold, so that there are M independent D5 branes at $x_I, I = 1; \dots; M$, and M image D5 branes located at $x_{\bar{I}}$. The O5 plane charge on the covering space is twice the O5 charge of the "downstairs" picture.

The dilaton equation of motion is

$$r^2 \Delta(r^2) = e^2 F_{(1)}^2 + \frac{1}{2} \frac{1}{3!} F_{(3)}^2 - \frac{1}{2} \frac{1}{3!} H_{(3)}^2 + (2)^2 \frac{01}{2} e \sum_i Q_i \frac{4(x-x_i)}{Z P g_B} : \quad (3:6)$$

Finally, the Bianchi identity for $F_{(3)}$ is

$$dF_{(3)} = F_{(1)} \wedge H_{(3)} - (2)^2 \text{Vol}_B \sum_i Q_i \frac{4(x-x_i)}{P g_B} ; \quad (3:7)$$

where Vol_B is the volume form on the base,

$$\text{Vol}_B = (g_B)^{1=2} dx^6 \wedge dx^7 \wedge dx^8 \wedge dx^9 : \quad (3:8)$$

There are additional equations of motion and Bianchi identities for the fluxes. These are discussed in Sec. 3.8 below.

In order to make the equations of motion tractable, we will now impose one further restriction on supergravity backgrounds that we consider. We demand that $H_{(3)mn} = 0$.

This condition is equivalent to restricting to the subset of vacua of this orientifold that are related to D3/O3 orientifold vacua via T-duality in the fiber directions.¹² We have not succeeded in constructing supergravity solutions with $H_{(3)} \neq 0$. One of the complications associated with this case is that the pseudo-BPS constraints of Sec. 3.3 apparently no longer hold. The flux $H_{(3)}$ contributes negatively to an otherwise positive semidefinite sum of squares. (See App. C for further discussion of this point). More significant qualitative differences implied by $H_{(3)} \neq 0$ are (i) $B_{(2)}$ is nonzero, so that there is noncommutativity on the D5 branes, and/or (ii) there does not exist a consistent truncation of the supergravity action to the lowest Kaluza-Klein modes. See Ref. [62] for a discussion of the gauge algebra for certain supergravity backgrounds with $H_{(3)} \neq 0$ and positive 4D vacuum energy.

3.2. Gauss's Law Constraint

Since the O5 planes and D5 branes fill all of the noncompact Minkowski directions, we expect the Bianchi identity (3.7) to imply an integrability condition that roughly demands that the total D5 charge vanish.

In order to derive this constraint, let us first introduce a notation that will be useful throughout the paper. For any p -form $\omega_{(p)}$ on X_6 , we write

$$\omega_{(p)} = \omega_{(p)}^0 + \omega_{(p)}^1 + \omega_{(p)}^2; \quad (3:9)$$

where the component $\omega_{(p)}^i$ is of rank i on the T^2 fiber. That is,

$$\omega_{(p)}^0 = \frac{1}{p!} \omega_{(p)m_1 \dots m_p} dx^{m_1} \wedge \dots \wedge dx^{m_p}; \quad (3:10a)$$

$$\omega_{(p)}^1 = \omega_{(p)}^1 \wedge \omega_{(p)}^1; \quad \omega_{(p)}^2 = \frac{1}{2} \omega_{(p)}^2 \wedge \omega_{(p)}^2; \quad (3:10b)$$

where

$$\omega_{(p)}^1 = \frac{1}{(p-1)!} \omega_{(p)m_2 \dots m_p} dx^{m_2} \wedge \dots \wedge dx^{m_p} \quad (3:11)$$

$$\omega_{(p)}^2 = \frac{1}{(p-2)!} \omega_{(p)m_3 \dots m_p} dx^{m_3} \wedge \dots \wedge dx^{m_p};$$

From our restrictions on the flux (including the additional assumption $H_{(3)} = 0$ made at the end of the last section) we have

$$F_{(1)} = F_{(1)}^0; \quad H_{(3)} = H_{(3)}^0; \quad \text{and} \quad \mathbb{F}_{(3)} = \mathbb{F}_{(3)}^0 + \mathbb{F}_{(3)}^1; \quad (3:12)$$

¹² The result of T-dualizing $H_{(3)}$ in the directions of two of its indices is a nongeometrical compactification [54,4].

Since the first two decompositions are trivial, we drop the superscripts on $F_{(1)}$ and $H_{(3)}$ everywhere except in App. C, where the assumption $H_{(3) m} = 0$ is temporarily relaxed. We also drop the tilde on $F_{(3)}$ since $F_{(3)}^1 = F_{(3)}^1$ for $H_{(3)}$ even. We will not need to assume anything about $F_{(5)}$ in order to show in the next section that it must vanish. We note only that $\int_{(5)}^0 = 0$ identically for any 5-form on X_6 . Therefore,

$$F_{(5)} = F_{(5)}^1 + F_{(5)}^2 : \quad (3.13)$$

The utility of these definitions is that forms (3.10a) and (3.11) can be interpreted as (pullbacks of) forms on the base. Noting that $dF_{(3)}^1 = F_{(1)} \wedge F_{(3)} + dF_{(3)}$, the Bianchi identity (3.7) becomes

$$dF_{(3)}^0 = F_{(1)} \wedge F_{(3)} + F_{(1)} \wedge H_{(3)} - (2)^0 \text{Vol}_B \sum_i^X Q_i \frac{(x - x_i)}{g_B}; \quad (3.14a)$$

together with

$$dF_{(3)} = 0 : \quad (3.14b)$$

All forms in (3.14a) are now (pullbacks of) forms on the base. So, the entire equation can be integrated over the base to give the Gauss's law constraint

$$0 = N_{ux} + \sum_i^X Q_i; \quad (3.15)$$

where

$$N_{ux} = \frac{1}{(2)^0} \int_B F_{(1)} \wedge F_{(3)} + F_{(1)} \wedge H_{(3)} : \quad (3.16)$$

For the case that the base B is T^4 ,

$$\sum_i^X Q_i = 2M - 32 : \quad (3.17)$$

Note that this implies that N_{ux} must be an even integer. This is guaranteed by the Dirac quantization conditions, as will be shown in Sec. 3.9.

The results of this subsection resolve a naive paradox concerning the Bianchi identity. In all but one of the examples that we consider in Secs. 6-8, the only nonzero ux is the RR three-form ux and $2M < 32$. Naively, we can integrate both sides of Eq. (3.7) over a transverse four-cycle linking all of the sources of D5 charge, to give zero on the LHS and $(2M - 32)(2)^0 < 0$ on the RHS. The resolution is that no such linking cycle exists.

The desired linking cycle would be a global section embedding the base in the fibration. However, unless the fibration is trivial ($F = 0$), there does not exist such global section.¹³ There only exist four-chains C that locally approximate the base in the sense that they project to the base, $C = B$, but globally necessarily have boundary. Consequently, $\int_C d\mathbb{F}_{(3)} = \int_B \mathbb{F}_{(3)} \neq 0$. On the other hand, it is true that $\int_C d\mathbb{F}_{(3)}^0 = 0$, since this integral is the pullback of $\int_B d!_{(3)}$, where $\mathbb{F}_{(3)}^0 = !_{(3)}$. Note that the nonexistence argument just given only applies to cycles linking all sources of D5 charge. There do exist (homologically trivial) four-cycles linking only individual D5 branes or O5 planes. Therefore, the charge of these objects is still well-defined.

3.3. Pseudo-BPS Constraints

As a consequence of the dilaton equation (3.6), the Bianchi identity (3.14a), and the Einstein equation for $g_{\text{R}} + g_{\text{R}}$, it is shown in App. C that

$$\begin{aligned}
d(Z^2 g_s^2 \mathbb{F}_{(3)}^0 + Z^2 \int_B d g_s^1 Z^{-1}) &= \\
\frac{1}{2} Z^2 g_s^1 (F_{(1)} + \int_B g_s^1 H_{(3)} \wedge \int_B F_{(1)} + \int_B g_s^1 H_{(3)} & \\
+ 2Z^{-1} \mathbb{F}_{(3)}^0 + Z^2 \int_B d g_s^1 Z^{-1} \wedge \int_B \mathbb{F}_{(3)}^0 + Z^2 \int_B d g_s^1 Z^{-1} & \\
+ g^{(T_{\text{fib}}^2)} F_{(3)} - g^{(T_{\text{fib}}^2)} \int_B g_s^1 F_{(3)} \wedge \int_B F_{(3)} - g^{(T_{\text{fib}}^2)} \int_B g_s^1 F_{(3)} & \\
+ g^{(T_{\text{fib}}^2)} 2Z^{-1} \mathbb{F}_{(5)}^1 \wedge \int_B \mathbb{F}_{(5)}^1 + g^{(T_{\text{fib}}^2)} \mathbb{F}_{(5)}^2 \wedge \int_B \mathbb{F}_{(5)}^2 & :
\end{aligned} \tag{3.18}$$

Here, g_s is defined by the equation

$$e = g_s Z^{-1/2} \tag{3.19}$$

(cf. Eq. (2.1b)) and is not necessarily constant at this point, although that will be shown soon. Since Eq. (3.18) involves only (pullbacks of) forms on the base B , we can integrate both sides over B . The integral of the LHS vanishes, while the RHS is positive semidefinite

¹³ This statement relies on the particular form of the T^2 fibration (2.4), in which there are no fiber degenerations associated with the shrinking of a $(p; q)$ S_1 in the fiber. It is true that there exist, for example, elliptically fibered Calabi-Yau manifolds with global section, but this relies on the existence of such degenerations.

and vanishes only if all terms vanish individually. Therefore, we obtain the Hodge duality relations

$$F_{(1)} = \star_B g_s^{-1} H_{(3)} ; \quad (3.20a)$$

$$F_{(3)}^0 = \star^2 \star_B d g_s^{-1} Z^{-1} ; \quad (3.20b)$$

$$F_{(3)} = g^{(T_{fib}^2)} \star_B g_s^{-1} F ; \quad (3.20c)$$

$$F_{(5)} = 0 ; \quad (3.20d)$$

We can alternatively write the first three relations as

$$F_{(3)}^0 = d e \star \text{Vol}_{h(x)} ; \quad (3.21a)$$

$$F_{(1)} = e \star \text{Vol}_{h(x)} \wedge H_{(3)} ; \quad (3.21b)$$

where \star is 10D Hodge star operator in the metric (2.4), and $\text{Vol}_{h(x)}$ is the volume form in the metric (3.5) with the point x on the base allowed to vary:

$$\text{Vol}_{h(x)} = \star^{3=2} dx^{0123} \wedge \text{Vol}_b ; \quad (3.22)$$

Here,

$$dx^{0123} = dx^0 \wedge dx^1 \wedge dx^2 \wedge dx^3 ; \quad (3.23)$$

and

$$\text{Vol}_b = V_b^{-4}(x) \wedge^{-5}(x) ; \quad (3.24)$$

with V_b given by Eq. (2.16). We will refer to the constraints (3.20) and (3.21) as pseudo-BPS conditions, since (i) they arise from the equations of motion alone, with no supersymmetry conditions imposed, and (ii) they give a proper subset of the supersymmetry conditions. The existence of such pseudo-BPS conditions is a consequence of the metric ansatz with 4D Minkowski space in the noncompact directions, together with a property of the low energy supergravity theory known as no-scale structure: consistency of the ansatz requires vanishing 4D vacuum energy, and the no-scale structure implies that the scalar potential is positive semidefinite.¹⁴

¹⁴ The canonical example of no-scale structure is the $N = 1$ no-scale structure of a theory whose superpotential is independent of Kahler moduli. Then, modulo D -terms, the scalar potential is $V = \sum_i |W_i|^2 - 3|W|^2 = \sum_i^0 |W_i|^2$, where i runs over all moduli and i^0 runs over all moduli other than the Kahler moduli. The models of Ref. [9] possess exactly this type of no-scale structure. In the 03 orientifolds of Refs. [60,11], the no-scale structure is an $N > 1$ analog of this. For the 05 theory discussed here, the only difference is that the 03 Kahler moduli are replaced by a more complicated subset of the 05 moduli. See Refs. [63,18,9].

Though the pseudo-BPS conditions were derived from the closed string sector equations of motion, they have a clear interpretation in the open string sector. The interpretation of Eq. (3.21a) is that Vol_b is a generalized calibration for fiber-class cycles wrapped by the D5 branes. The mass of a D5 brane that wraps a two-cycle in the homology class of the T^2 fiber is minimized when the two-cycle is the vertical fiber over a point in the base, as opposed to another cycle in the same homology class with horizontal components. For certain choices of flux, such as those in the examples that we present in Secs. 6.1 and 8.1, the S-dual of this calibration condition has already appeared elsewhere [28,27]. The S-duality is discussed briefly in Sec. 4.3. In the earlier treatment, the generalized calibration Vol_b was given a further interpretation as deriving from an $SU(2)$ structure on the 6D internal manifold.¹⁵

Eq. (3.21b) is the condition that the Myers D5 worldvolume coupling [64]

$$S_3 = \int_{\text{vol}_v} e^{-2\phi} \text{Vol}_{wv} H_{(3)mn} F_{(9)012345mnp} dx^{0123} \wedge dx^4 \wedge dx^5 \text{Tr}(F^m{}_n F^p) \quad (3.25)$$

vanish. Here Vol_{wv} is the volume form on the D5 worldvolume. When this condition is not satisfied, the D5 branes are polarized by the external $F_{(1)}$ and $H_{(3)}$ fluxes and blow up into D7 branes.

3.4. Dilaton

The dilaton equation of motion (3.6) can be combined with the g_{RR} Einstein equation to give

$$\begin{aligned} r_B^2 Z^{-1} e^{-2\phi} &= Z^{-1/2} e^{-2\phi} \left(g_{RR} + 2r \dot{r} + 2r^2 \ddot{r} - 2(\dot{r})^2 \right) \\ &= Z^{-1/2} \left(F_{(1)}^2 + \frac{1}{2} \frac{1}{5!} F_{(5)}^2 - e^{-2\phi} \frac{1}{3!} H_{(3)}^2 \right) : \end{aligned} \quad (3.26)$$

(See App. C, Eqs. (C.1) and (C.5)). The RHS vanishes by Eqs. (3.20), so g_s as defined in Eq. (3.19) is indeed constant.

¹⁵ The $SU(2)$ structure on X_6 is in this case defined by a triple of two-forms $J, \text{Re } \Omega_{(2)}$ and $\text{Im } \Omega_{(2)}$ on the 4D base, together with a pair of Killing one-forms K_1 and K_2 , such that $K_1 \wedge K_2 = \text{Vol}_b$.

3.5. Warp Factor

The warp factor Z is a solution to Poisson's equation with localized sources at the D5 branes and O5 planes, and constant D5 charge density from the flux and bration. From the result that g_s is constant,

$$dF_{(3)}^0 = \frac{1}{g_s} d_B dZ = \frac{1}{g_s} r_B^2 Z \text{Vol}_B : \quad (3.27)$$

It will be shown in Sec 3.7 and 3.8 that F , $F_{(3)}$, $H_{(3)}$, and $F_{(1)}$ are (pullbacks of) harmonic forms on B as a consequence of the geometrical conditions on F and the Bianchi identities/equations of motion for the fluxes. Therefore, the integrand in Eq. (3.16) is a harmonic representatives of $H^4(B; \mathbb{R})$, that is, a constant multiple of Vol_B . This allows us to replace Eq. (3.16) with the stronger relation

$$F \wedge F_{(3)} + F_{(1)} \wedge H_{(3)} = (2)^2 N_{\text{ux}} V_B^{-1} \text{Vol}_B : \quad (3.28)$$

Here, $V_B = \int_B \text{Vol}_B$ is the volume of the base. Note that from Eqs. (3.20a;c), we then have $N_{\text{ux}} \geq 0$. Therefore, the number of D5 branes is bounded above by (the absolute value of) the total charge from O5 planes, due to the constraint (3.15).

Combining Eqs. (3.27) and (3.28), the Bianchi identity (3.14a) becomes

$$\begin{aligned} \frac{1}{g_s} r_B^2 Z &= (2)^2 N_{\text{ux}} \frac{1}{V_B} + \sum_i^X Q_i \frac{\int_B d^4(x; x_i)}{\int_B \sqrt{g_B}} \\ &= (2)^2 \sum_i^X Q_i \frac{\int_B d^4(x; x_i)}{\int_B \sqrt{g_B}} \frac{1}{V_B} : \end{aligned} \quad (3.29)$$

The solution is

$$Z = 1 + (2)^2 \sum_i^X \frac{Q_i}{g_s} G_B(x; x_i); \quad (3.30)$$

where G_B is the Green's function for Poisson's equation on the base B ,

$$r_B^2 G_B(x; x^0) = \frac{\int_B d^4(x; x^0)}{\int_B \sqrt{g_B}} - \frac{1}{V_B} : \quad (3.31)$$

As part of our definition of the Green's function, we fix the constant component of G_B by requiring that $\int_B d^4 x \sqrt{g_B} G_B(x; x^0) = 0$. The leading integration constant of unity in Eq. (3.30) is conventional and ensures that the warp factor completely drops out of the metric (2.4) in the limit of zero gravitational coupling $g_s \rightarrow 0$, at fixed V_B . This fixes the rescaling ambiguity

$$g_s \rightarrow g_s; \quad Z \rightarrow Z^2 \quad \text{at fixed } e \quad (3.32)$$

in Eq. (3.19).

3.6. Base Geometry

The only other constraint that follows from the Einstein, dilaton, and $\mathbb{F}_{(3)}$ equations, and is not automatically satisfied as a consequence of Eqs. (3.20), (3.26), (3.14b), and (3.30) is

$$R_{Bmn} = 0; \tag{3.33}$$

from the $m n$ Einstein equations. For compact base B , this comes close to implying that B is T^4 or $K3$. Where it falls short is that Ricci-flatness implies only that $c_1(B)$ vanishes in $H^2(B; \mathbb{R})$. There can still be a torsion component in $H^2(B; \mathbb{Z})$, as is the case for an Enriques surface. Such manifolds are ruled out only after we impose supersymmetry. The supersymmetry conditions $\hat{e}^m = r_{Bm} \hat{e}^m = 0$ of Sec. 4.4 give $c_1(B) = 0$ as an integrability condition.

3.7. Geometrical Bianchi Identity and Quantization of the Fibration Curvature

For the fibration (2.4) to be globally well-defined, the fibration curvature must satisfy

$$dF = 0; \tag{3.34}$$

This is equivalent to the triple-overlap condition on the transition functions that relate coordinate patches. Under a fiber coordinate redefinition, $x \rightarrow x + f x^m g$, we have $F \rightarrow F + d(f g)$ (cf. App. B). Therefore, the exact part of F contains coordinate-gauge information, and the topology of the fibration is characterized by the cohomology class of F .

We require that $[F]$ be $2\mathbb{Z}$ valued (cf. Eq. (2.5)) for the following reason. First, ignore the orientifold operation. The subgroup of $H^2(X_6; \mathbb{Z})$ that comes directly from $H^2(B; \mathbb{Z})$ is the quotient group $H^2(B; \mathbb{Z}) = [F]_g$. This quotient is well-defined only if $[F] \in 2H^2(B; \mathbb{Z})$. Then, in order to define the orientifold, we further require that F describe a fibration over $B = \mathbb{Z}_2$. This is equivalent to the condition

$$[F] \in 2H^2(B; 2\mathbb{Z}); \tag{3.35}$$

which guarantees integer periods of $[F]$ over the cycles in $B = \mathbb{Z}_2$ that descend from half-cycles in B . It is possible that the $2\mathbb{Z}$ quantization condition can be replaced by a $2\mathbb{Z} + 1$ quantization condition if one includes localized fibration curvature at some or all of the \mathbb{Z}_2 fixed points. However, we do not consider such localized curvature here.

3.9. Discrete Data versus Moduli, and Additional Moduli Constraints

Beyond the choice of base manifold B and orientifold operation, the complete set of discrete data needed to define the model is the number $2M$ of D5 branes and their images, together with the quantized parts of the even fluxes and the bration curvature (3.35).

For later discussion of moduli stabilization, it is necessary to decompose the fluxes into a part that is moduli-dependent and a quantized part that only involves the discrete choice. For simplicity of exposition, we restrict to the case that $B = T^4$. (The results of this section are analogous in the case that $B = K3$, except that the absence of a first cohomology class for $K3$ implies that $F_{(1)} = 0$, and then from Eq. (3.20a), $H_{(3)} = 0$.) In this subsection, we also drop all functional dependence of the moduli on x^μ . That is, we analyze deformations of the supergravity background, and do not yet promote these deformations to 4D fields. We restrict to deformations that correspond to moduli and not to gauge bosons. In the next subsection, when we discuss the kinetic terms and gauge couplings, the x^μ -dependence and gauge bosons will be reintroduced.

Let us write all of the internal fluxes and potentials as the sum of a background value and a deformation. In the NS sector we write

$$\begin{aligned} B_{(2)} &= B_{(2)}^{bg} + b_{(2)}; \\ H_{(3)} &= H_{(3)}^{bg} + h_{(3)}; \end{aligned} \tag{3.41}$$

Since we have assumed that $H_{(3)} = H_{(3)}^0$ (in the notation of Sec. 3.2), we take $B_{(2)}^{bg}$ to have purely base components, $B_{(2)}^{bg} = B_{(2)}^{bg0}$. The deformation permitted by the orientifold projection is a shift $b_{(2)\ m}$ in the zero-mode, or constant component,¹⁶ of the even potential $B_{(2)\ m}$. Therefore,

$$\begin{aligned} b_{(2)} &= \hat{b}_{(2)}; \\ h_{(3)} &= db_{(2)} = F \wedge \hat{b}_{(2)}; \end{aligned} \tag{3.42}$$

In the RR sector, we similarly write

$$\begin{aligned} C_{(p)} &= C_{(p)}^{bg} + c_{(p)}; \\ F_{(p+1)} &= F_{(p+1)}^{bg} + f_{(p+1)}; \quad \mathbb{F}_{(p+1)} = \mathbb{F}_{(p+1)}^{bg} + \mathbb{f}_{(p+1)}; \end{aligned} \tag{3.43}$$

¹⁶ To justify this usage of the term zero-mode, note that the p -forms with constant coefficients are annihilated by a Laplacian operator on X_6 formed from a torsionful connection that forgets about the bration and warping (cf. Sec. 4.4).

In this case, the moduli are not quite the $c_{(p)}$. The latter are in general multivalued quantities on X_6 due to the action of the $C_{(p-2)}$ transition functions on $C_{(p)}$ when $H_{(3)} \neq 0$. This subtlety is discussed in Ref. [60]. Instead, the moduli are the combinations

$$e_{(2)} = c_{(2)} \quad h_{(2)} C_{(0)}^{bg}; \quad (3:44a)$$

$$e_{(4)} = c_{(4)} \quad h_{(2)} \wedge C_{(2)}^{bg1} + e_{(2)} \quad \frac{1}{2} b_{(2)} \wedge b_{(2)} \wedge C_{(0)}^{bg}; \quad (3:44b)$$

$$e_{(6)} = c_{(6)} \quad h_{(2)} \wedge e_{(4)} \quad \frac{1}{2} b_{(2)} \wedge b_{(2)} \wedge e_{(2)}; \quad (3:44c)$$

as can be verified by arguments analogous to those in Ref. [60]. Here, Eqs. (3:44a;b;c) are relations between quantities with internal 6D indices only.

The nonvanishing components of the background are

$$C_{(0)}^{bg}; \quad C_{(2)mn}^{bg}; \quad C_{(2)m}^{bg}; \quad C_{(4)mnpq}^{bg}; \quad C_{(4)n pq}^{bg}; \quad (3:45)$$

or a subset thereof, and the moduli are the zero-modes

$$e_{(2)mn}; \quad e_{(2)}; \quad e_{(4)mnp}; \quad e_{(6)mnpq}; \quad (3:46)$$

The last modulus is $e_{(6)456789}$. This is the scalar that is dual to the corresponding deformation $e_{(2)}$ of $C_{(2)}$, and will be more convenient to work with than $e_{(6)}$ when we discuss gauge couplings in Sec. 3.10 and the T-duality map in Sec. 5.2.

The flux deformations that follow from Eqs. (3:44a;b;c) are

$$f_{(3)} = \mathbb{F}_{(3)} = de_{(2)} + b_{(2)} \wedge F_{(1)}; \quad (3:47a)$$

$$\mathbb{F}_{(5)} = e_{(2)} \wedge H_{(3)}^{bg} + b_{(2)} \wedge F_{(3)}^{bg1} + de_{(2)} + \frac{1}{2} b_{(2)} \wedge b_{(2)} \wedge F_{(1)}; \quad (3:47b)$$

expressable in terms of the moduli and the gauge invariant fluxes. The second equality in Eq. (3:47a) implies that the only nonvanishing component of $\mathbb{F}_{(3)}$ is $\mathbb{F}_{(3)}$, with one fiber index; this, combined with the earlier observation that $\mathbb{F}_{(3)} = F_{(3)}$, gives the first equality. In Eq. (3:47b), $F_{(1)} = F_{(1)}^{bg}$, so we have dropped the superscript.

One consequence of Eqs. (3.42) and (3:47a) is that

$$F \wedge f_{(3)} + F_{(1)} \wedge h_{(3)} = 0; \quad (3:48)$$

from which Eq. (3.16) becomes

$$N_{ux} = \frac{1}{(2-)^2} \int_B^Z F \wedge F_{(3)}^{bg} + F_{(1)} \wedge H_{(3)}^{bg}; \quad (3:49)$$

independent of the moduli.

Dirac quantization constrains the closed moduli-independent fluxes $F_{(1)}$, $F_{(3)}^{bg}$, $F_{(5)}^{bg}$, and $H_{(3)}^{bg}$ to be representatives of $2\mathbb{Z}$ -valued cohomology:

$$F_{(1)} \in 2H^1(B; 2\mathbb{Z}); \quad F_{(3)}^{bg} \in 2(2)^2 H^2(B; 2\mathbb{Z}); \quad (3.50a)$$

$$H_{(3)}^{bg} \in 2(2)^2 H^2(B; 2\mathbb{Z}); \quad F_{(5)}^{bg} \in 2(2)^4 H^3(B; 2\mathbb{Z}): \quad (3.50b)$$

As in Eq. (3.35), the $2\mathbb{Z}$ quantization of periods on the covering space B ensures \mathbb{Z} quantization of periods over half cycles in B that descend to proper boundaryless cycles in $B = \mathbb{Z}_2$. We can now return to the issue of the even integer quantization of N_{ux} mentioned in Sec. 3.2. From Eq. (3.49), we see that N_{ux} depends only on the discrete choice (3.35) and (3.50). As a result, N_{ux} satisfies $N_{ux} \in 2\mathbb{Z}$, which is indeed an even integer, and is moduli independent, so no moduli constraints arise from the Gauss's law constraint.

While Dirac quantization alone permits a quantized five-form flux $F_{(5)}^{bg}$, it can be shown that $F_{(5)}^{bg} = 0$ in order to satisfy the equations of motion. We omit the direct proof, but Sec. 5.2 contains a proof via T-duality. From the constraint (3.20d), we then have

$$F_{(5)} = 0: \quad (3.51)$$

Without imposing any supersymmetry conditions, the complete set of moduli constraints that follow from the equations of motion consists of Eqs. (3.20a;c) and (3.51). When $F_{(1)} = H_{(3)}^{bg} = 0$, the constraints simplify. Eq. (3.20a) then implies that

$$0 = h_{(3)} = F_{(2)} \wedge b_{(2)}; \quad (3.52a)$$

and the constraint (3.51) becomes

$$b_{(2)} \wedge F_{(3)}^{bg} = 0: \quad (3.52b)$$

In this case, the complete set of moduli constraints is (3.20c) and Eqs. (3.52a;b).

3.10. Couplings to Gauge Bosons

In addition to the deformations of the supergravity background just discussed, which when promoted to 4D fields become moduli, there are other deformations of the supergravity background which when promoted to 4D fields become gauge bosons. In the closed

string sector, these deformations are the zero modes on X_6 of V , $B_{(2)m}$, $C_{(2)}$, and $C_{(4) m} + 6b_{(2)[m j}C_{(2)j]}$, and will be denoted by the lowercase symbols

$$v; b_{(2)m}; c_{(2)}; \text{ and } e_{(4) m}; \quad (3.53)$$

respectively. When the couplings to these gauge bosons are included and all deformations are promoted to 4D fields, the kinetic term for $e_{(2)m n}$ is the square of

$$\partial e_{(2)m n} - F_{(3) m n}^{bg} v + 2F_{(1)[m} b_{(2)n]} + F_{m n} c_{(2)}; \quad (3.54a)$$

that for $e_{(4) m n p}$ is the square of

$$\partial e_{(4) m n p} + 3F_{[m n j} e_{(4) j p]} + H_{(3) m n p}^{bg} c_{(2)} + 3F_{(3) [m n} b_{(2)p]}; \quad (3.54b)$$

and that for $e_{(6) m n p q}$ (the scalar that is dual to $e_{(2)}$) is the square of

$$\partial e_{(6) m n p q} - 3H_{(3) [m n p j}^{bg} e_{(4) j q]}; \quad (3.54c)$$

The axion $e_{(2)}$ of dil does not couple to gauge bosons.

Beyond the closed string sector gauge bosons, there are also gauge bosons that arise from the lowest Kaluza-Klein mode of the D5 worldvolume gauge fields A_I on the wrapped T^2 fiber.

4. Supersymmetry Conditions

The 10D type IIB dilatino and gravitino variations are given in App. D. From these fermion variations, we will now determine the conditions on the 6D geometry and internal flux for unbroken 4D $\mathcal{N} = 1$ supersymmetry.

4.1. Decomposition of 10D Supersymmetry Parameters

In 10D, the IIB supersymmetry transformations are parametrized by two Majorana-Weyl spinors $\epsilon_{L,R}$ (real and negative chirality in our conventions), which combine to form a single Weyl spinor $\epsilon = \epsilon_L + i\epsilon_R$. For compactification to 4D, it is desirable to decompose into 4D and 6D spinors.¹⁷ An arbitrary 10D negative chirality Weyl spinor can be written

$$\epsilon = u_1 + u_2; \quad (4.1)$$

¹⁷ In this discussion, we follow Gherghel and Polchinski [6].

where λ_1 and λ_2 are negative chirality 6D Weyl spinors, and u is a positive chirality 4D Weyl spinor.¹⁸ For each pair $(\lambda_1; \lambda_2)$ such that (4.1) gives vanishing dilatino and gravitino variation for all u , we obtain one 4D $N = 1$ supersymmetry generated by u .

Orientifold planes impose additional constraints on the spinors. Two broad classes of constraints are

$$\begin{aligned} \lambda_1 = 0 \quad \text{or} \quad \lambda_2 = 0 & \quad \text{Becker-type or chiral constraints,} \\ \lambda_2 = e^i \lambda_1; \quad \lambda_1 = \text{const} & \quad \text{Andy-type or Majorana-Weyl constraints.} \end{aligned} \tag{4.2}$$

Becker-type spinors are of definite 6D and 4D chirality, while Andy-type spinors are real up to an overall constant phase. Roughly speaking, Becker-type constraints are associated with $O(3)$ or $O(7)$ planes, and Andy-type constraints are associated with $O(5)$ or $O(9)$ planes or the heterotic theories. More precisely, $O(3)$ and $O(9)$ constraints require that the spinors be of pure Becker- or Andy-type; $O(7)$ and $O(5)$ constraints require that the spinors be a particular linear combination of two Becker- or two Andy-type spinors, such that for certain choices of u we can decompose the supersymmetry algebra into $N = 1$ subalgebras, each generated by a spinor for which one of the two terms in the linear combination vanishes. We will demonstrate this for the $O(5)$ case which is the focus our investigation, and for the $O(3)$ case which is needed in Sec. 5.

Independent of the details of the orientifold projection, λ_R is a real Majorana-Weyl spinor, and can therefore be written as

$$\lambda_R = \frac{1}{\sqrt{2}} u + u \quad ; \tag{4.3}$$

for some u and γ , where the leading factor of $1/\sqrt{2}$ is for compatibility with the normalization conventions

$$\gamma_{L;R}^Y \lambda_{L;R} = u^Y u = \gamma^Y = 1: \tag{4.4}$$

For the $O(5)$ planes of interest, the 10D supersymmetries preserved by the orientifold projection are generated by $\lambda_{L;R}$ such that

$$\lambda_L = \gamma_B \lambda_R: \tag{4.5}$$

Here, γ_B is the chirality operator formed from product of Dirac matrices in the directions transverse to the orientifold planes, normalized so that $\gamma_B^2 = 1$. Similarly, we define γ_b

¹⁸ Note also that for λ and u to be standard anticommuting spinors, λ_1 and λ_2 must be commuting spinors [30].

to be the chirality operator in the two compact directions wrapped by the $O(5)$ plane. That is, γ_b is the chirality operator on the T^2 fiber, and γ_B is the chirality operator on the base B . Using these chirality operators, we can decompose the 6D spinor into components of definite fiber and base chirality,

$$\psi = \psi_+ + \psi_-; \quad \text{where } \gamma_b \psi_{\pm} = \pm \psi_{\pm}; \quad \gamma_B \psi_{\pm} = \pm \psi_{\pm}; \quad (4.6)$$

Here, we have gone from uppercase 10D to lowercase 6D internal Dirac matrices using the relations $\gamma_b = 1 - \gamma_b$ and $\gamma_B = 1 - \gamma_B$. (See App. A for a precise statement of our conventions for Dirac matrices, chirality operators, and their decompositions under $SO(1,4) \times SO(6) \rightarrow SO(1,4) \times SO(2) \times SO(4)$).

Applying the decomposition (4.6) to Eq. (4.5), we obtain

$$\psi_L = \frac{1}{\sqrt{2}} (\psi_+ - \psi_-); \quad (4.7)$$

where

$$\psi_{\pm} = \psi_{\pm B} = \psi_{\pm+} + \psi_{\pm-}; \quad (4.8)$$

The 10D Weyl spinor $\psi = \psi_L + i\psi_R$ becomes

$$\psi = e^{i\pi/4} (\psi_{+} + \psi_{-}) + e^{-i\pi/4} (\psi_{+} - \psi_{-}); \quad (4.9)$$

which is the desired linear combination of two Andry-type spinors. In the case that $F_{(1)} = H_{(3)} = 0$, we will show in Sec. 4.2 that the supersymmetry conditions on ψ_{\pm} and decouple from one another, so that the space of 6D Killing spinors decomposes into a subspace on which $\psi_{\pm} = 0$ and a subspace on which $\psi_{\pm} \neq 0$. On either subspace, ψ is of pure Andry-type.

For spacetime filling $O(3)$ planes, the 10D supersymmetries are generated by $\psi_{L,R}$ such that

$$\psi_L = i^{(6)} \psi_R; \quad (4.10)$$

Here, $i^{(6)}$ is the chirality operator formed from the product of Dirac matrices in the six internal directions transverse to the orientifold planes. Using Eq. (4.3), this becomes Eq. (4.7) with

$$\psi_L = i^{(6)} \psi_R = i \psi_R; \quad (4.11)$$

So, the 10D Weyl spinor is

$$\psi = \frac{1}{\sqrt{2}} (\psi_L + i\psi_R); \quad (4.12)$$

of Becker-type, as claimed.

Although it is not directly relevant to this investigation, it is interesting to note that there exist supergravity backgrounds that are neither Andry-type or Becker type, but rather interpolate between the two [65,44]. These solutions are D3-like in some regions and D5/NS5-like in others. Similarly, there exist dielectric flow solutions, involving D3 branes that become polarized into D5/NS5 branes under radial flow in IIB, or M2 branes that become polarized into M5 branes under radial flow in M theory [66,67]. In Ref. [67], the technique of "Algebraic Killing spinors" was developed and repeatedly applied as an efficient method for obtaining solutions to the supergravity equations of motion. In the M theory case, these supergravity solutions fit nicely into classification of Ref. [34], which provides a general framework in the language of G-structures for describing arbitrary compactifications of M theory to $R^{2,1}$ and AdS_3 [34]. This framework also accommodates the M theory dual of Refs. [65,44]. A more restrictive treatment with applications, in particular, to the M theory lift of IIB pp-wave backgrounds has also been given [68]. Finally, recent work has shown that the correct description of the most general $R^{3,1}$ or AdS_5 compactifications of type IIB or M theory, should be given in terms of $SU(2)$ structures [41,36,42]. This work has brought us closer to understanding, for example, what the complete supergravity solution for the Polchinski-Strassler background might be, a subject that is currently under investigation [69].

4.2. Supersymmetry Conditions for RR Three-Form Flux Only

In the case that there is only $F_{(3)}$ RR flux and nontrivial bration, the supersymmetry conditions are very similar to those for the heterotic string with NS flux. By substituting the expression (4.9) for \hat{g}_s into the Eqs. (D.7) for the IIB fermion variations, and then demanding that the result vanish, we obtain

$$\begin{aligned} \frac{1}{2} \hat{g}_s^{-1} \hat{D} \hat{g}_s - \frac{1}{4} e^{\hat{F}_{(3)}} \hat{u} &= 0; \\ \hat{r}_M - \frac{1}{4} \hat{g}_s^{-1} \hat{D}_M \hat{g}_s - \frac{1}{4} e^{\hat{F}_{(3)M}} \hat{u} &= 0: \end{aligned} \quad (4.13)$$

After performing a Weyl rescaling

$$ds_{\text{string}}^2 = e^{(\hat{\phi})} ds^2; \quad \hat{g}_s = e^{(\hat{\phi})=4 \wedge}; \quad \hat{r}_M = e^{(\hat{\phi})=2 \wedge M}; \quad (4.14)$$

where $e^{\hat{\phi}} = g_s$, these equations become

$$\frac{1}{2} \hat{D} \hat{g}_s - \frac{1}{4} g_s \hat{F}_{(3)} \hat{u} \wedge = 0; \quad (4.15a)$$

$$\hat{r}_M - \frac{1}{4} g_s \hat{F}_{(3)M} \hat{u} \wedge = 0: \quad (4.15b)$$

Here, the hatted metric is

$$ds^2 = Z^{-1/2} e^{\hat{g}_{\mu\nu}} dx^\mu dx^\nu + ds_6^2 \quad (4.15c)$$

Eqs. (4.15) are a doubled version of the equations that formed the starting point for Strominger's "Superstrings with Torsion" analysis [2]. Since the constraints on $\hat{g}_{\mu\nu}$ and $\hat{H}_{(3)}$ are decoupled from one another, we are free to take $\hat{H}_{(3)} = \hat{H}_{(3)}^+$ or $\hat{H}_{(3)}^-$, and set the other spinor to zero. Then the starting points are identical. The results that follow from the same analysis are just Strominger's results, with $H_{(3)} = \pm g_6 F_{(3)}$, $\hat{H}_{(3)} = \pm \hat{H}_{(3)}$, and $\hat{H}_{(3)} = \pm \hat{H}_{(3)}$ compared to Sec. 1.1.¹⁹

$$Z^{-1/2} e^{\hat{g}_{\mu\nu}} = 1; \quad (4.16a)$$

$$e^2 d e^2 J = \pm g_6 F_{(3)}; \quad (4.16b)$$

$$d e^2 \pm g_6 J = 0; \quad (4.16c)$$

$$d e^2 = 0; \quad (4.16d)$$

where the SU(3) structures are²⁰

$$J_a{}^b = i \wedge^Y_a{}^b; \quad (4.17a)$$

$$\hat{H}_{abc} = \wedge^Y_{abc}; \quad (4.17b)$$

Just as for the heterotic string, $e^2 J$ is a holomorphic (3,0) form and the Nijenhuis tensor vanishes, so that the complex structure (4.17a) is integrable. Eq. (4.16a) reproduces the relation (3.19) between the dilaton and warp factor. Eqs. (4.17a;b) imply the relations [2,32]

$$J_a{}^b J_b{}^c = \delta_a{}^c; \quad (4.18a)$$

$$\frac{1}{3!} J \wedge J \wedge J = \frac{i}{8} \wedge^3 = \text{Vol}_6; \quad (4.18b)$$

where Vol_6 is the volume form associated with the metric (2.7),

$$\text{Vol}_6 = Z^2 g^{(\mathbb{T}_{\text{fib}}^2)} g_B^{-1/2} dx^4 \wedge dx^5 \wedge dx^6 \wedge dx^7 \wedge dx^8 \wedge dx^9; \quad (4.19)$$

¹⁹ The heterotic analogs of Eqs. (4.16a;d) and (4.17b) were suppressed from our short summary in Sec. 1.1, but follow from Strominger's analysis [2,28,27].

²⁰ The fact that $\hat{H}_{(3)}$ depends antiholomorphically on $\hat{g}_{\mu\nu}$ is a consequence of the negative ⁽⁶⁾ chirality of $\hat{g}_{\mu\nu}$.

In the next two paragraphs, we derive a number of results concerning the fiber and base decomposition of J and $\hat{\omega}$ that will be useful for the examples in Sec. 6-8. These paragraphs can be skipped over, if desired, without much loss to the understanding of this section.

Since $\hat{\omega}_+$ and $\hat{\omega}_-$ have definite fiber and base chirality, these spinors factorize into the product of a 2D Weyl spinor on the fiber and a 4D Weyl spinor on the base,

$$\hat{\omega} = \tilde{e}_b \tilde{e}^B; \text{ where } e_b \tilde{e}^b = \delta_b^b; e_B \tilde{e}^B = \delta_B^B; \quad (4.20)$$

(The tilded Dirac matrices are defined in App. A). Consequently, the Kähler form and (3,0) form decompose as

$$J = J^{(T_{\text{fib}}^2)} + Z J^B; \quad \tilde{\omega} = 2g_{11}^{(T_{\text{fib}}^2)} z^1 \wedge \tilde{z}^1 + Z^B \tilde{\omega}_B; \quad (4.21)$$

Here, the forms J^B and $\tilde{\omega}_B$ are the Kähler form and holomorphic (2,0) form on the base, defined via

$$J_m^B n^m = i^{BY} e_m^B n^m; \quad \tilde{\omega}_B m n = i^{BY} e_m^B n^m; \quad (4.22)$$

Also, in complex coordinates, the fiber metric (2.4b) is

$$ds_{T_{\text{fib}}^2}^2 = g_{11}^{(T_{\text{fib}}^2)} z^1 \tilde{z}^1 + g_{11}^{(T_{\text{fib}}^2)} z^1 \tilde{z}^1; \quad g_{11}^{(T_{\text{fib}}^2)} = g_{11}^{(T_{\text{fib}}^2)} = \frac{V_b}{2 \text{Im } \tau_1}; \quad (4.23)$$

where V_b is given by Eq. (2.16), and the fiber (1,0) form is

$$z^1 = dz^1 + A z^1; \text{ where } z^1 = x^4 + \tau_1 x^5; \quad A z^1 = A^4 + \tau_1 A^5; \quad (4.24)$$

in terms of the fiber complex structure modulus τ_1 .

The fiber complex structure is related to the spinors by

$$J^{(T_{\text{fib}}^2)} = i^{bY} e_b^Y; \quad (4.25)$$

which implies that

$$J^{(T_{\text{fib}}^2)} = \text{Vol}_b; \quad (4.26a)$$

where Vol_b was defined in Eq. (3.24), and

$$\text{Vol}_b = \text{sign}(\text{Im } \tau_1); \quad (4.26b)$$

Then, from Eqs. (4.18b) and (4.21),

$$\frac{1}{2} J^B \wedge J^B = \text{Vol}_b : \quad (4.26c)$$

So, the geometrical origin of the doubling in Eqs. (4.16) and (4.17) is that positive orientation of X_6 (4.18b) can correspond to either positive-positive or negative-negative orientation of the fiber and base (4.26a;c).

As a consequence of Eqs. (4.16a) and (4.17b), Eq. (4.16d) is automatically satisfied. The remaining conditions (4.16b;c) are equivalent to [29,48]

$$F_{(3)} = \frac{i}{g_s} e^2 d e^2 J \quad (2,1) \text{ and primitive.} \quad (4.27)$$

Using Eqs. (4.21) and (4.26a), and the decomposition (3.12), this becomes

$$F_{(3)}^0 = \text{Vol}_b g_s^{-1} dZ ; \quad (4.28a)$$

$$G_{(3)} \quad (2,1) \text{ and primitive,} \quad (4.28b)$$

where

$$G_{(3)} = F_{(3)}^1 \quad \frac{i}{g_s} d\text{Vol}_b : \quad (4.28c)$$

From the results of Sec. 3.9 applied to the case that $H_{(3)} = 0$, we can also write $G_{(3)}$ as

$$G_{(3)} = F_{(3)}^{bg1} \quad (2) \int^0 d\text{il} d^4 \wedge^5 ; \quad (4.28d)$$

where

$$d\text{il} = \frac{1}{(2)^2 \cdot 0} \quad e_{(2)45} + \frac{i}{g_s} V_b ; \quad (4.29)$$

which is the suitable generalization of Eq. (2.15) to the case of nonvanishing background flux.

Eqs. (4.28a;b;d) express the supersymmetry conditions in the form that we will find most convenient to apply in Secs. 6{8. Given a discrete choice of $F_{(3)}^{bg}$ and $F_{(3)}$, we can easily deduce the constraints on moduli via these equations. The number N of supersymmetries preserved is

$$N = N_+ + N_- ; \quad (4.30)$$

where N_- is the number of independent complex structures (i.e., the number of $\hat{\cdot}$) such that these conditions are satisfied. Eqs. (4.28) also imply the equations of motion (3.20).²¹ The maximum amount of supersymmetry is $N = 4, N_+ = N_- = 2$, since there are two 4D spinors of each chirality and one 2D spinor of each chirality, from which we must construct the negative chirality 6D spinors $\hat{\cdot}$. For nonvanishing flux , the supersymmetry is strictly less than this. That is, either $N_+ < 2$ and $N_- = 2$, or $N_+ = 2$ and $N_- < 2$.

4.3. S-dual Orientifolds with NS Flux Only

The class of supersymmetric vacua just discussed is S-dual to a class of vacua with NS flux $H_{(3)}$ only. Such vacua have received considerable attention, so it is desirable to state the precise connection between Sec. 4.2 and known results.

The result of applying S-duality to the class of orientifolds analyzed here is a class of dual orientifolds in which O5 planes are replaced by ON5 planes and D5 branes are replaced by NS5 planes [70]. The orientifold operation \mathbb{I}_4 in the original theory becomes $(-1)^{F_L} \mathbb{I}_4$ in the dual theory. In our S-duality conventions, the dual NS flux is related to the original RR three-form flux by

$$H_{(3)}^0 = F_{(3)} \quad (4.31)$$

The S-dual metric is

$$\begin{aligned} ds_{\text{string}}^2 &= e^{-(\phi - \phi_0)} ds_{\text{string}}^2 \\ &= dx^2 + ds_6^2; \end{aligned} \quad (4.32)$$

where we have used Eq. (3.19) (or equivalently, Eq. (4.16a)) in the second equality. We recognize this as the hatted metric (4.15c) that appeared in the previous section.

The S-dual dilaton is given by

$$e^{-(\phi - \phi_0)} = e^{-\phi_0} = z^{1/2}; \quad g_s^0 = g_s^{-1} \quad (4.33)$$

So, the 6D metric can be written

$$ds_6^2 = g^{(T_{\text{fib}}^2)} (dx^2 + A^2) + e^{2(\phi - \phi_0)} g_{B m n} dx^m dx^n; \quad (4.34)$$

²¹ The equations of motion (3.20b;c) require that G_3 be imaginary-selfdual (ISD): $\star_6 G_{(3)} = iG_{(3)}$. This is weaker than (4.28b). The space of ISD three-forms includes not only primitive (2,1) forms, but also (0,3) forms and non-primitive (1,2) forms $J \wedge !$, where $!$ is a (0,1) form.

which is a form that has appeared previously in the literature. Nearly identical vacua were described by Gauntlett, Martelli, and Waldram, in their classification of static supersymmetric backgrounds with NS flux only [27]. The one minor difference is that the local ON5 and NS5 sources were excluded from their discussion, and as a result, the base geometry was required to be noncompact to avoid contradicting the Gauss's law constraint (3.15).

In the conventions of Eq. (4.31), the 10D IIB supersymmetry parameters after the S-duality are

$$\begin{aligned} N_L^0 &= \frac{1}{2} u_{-+} + u_{++} ; \\ N_R^0 &= \frac{1}{2} u_{-+} + u_{+-} + u_{++} ; \end{aligned} \quad (4.35)$$

up to a possible overall sign in N_L^0 and/or N_R^0 that will not concern us here. Thus, N_+ and N_- , which before the S-duality counted the number of 6D Killing spinors of + and - chirality, map to the number of right and left Killing spinors, respectively, in the S-dual theory:

$$N_R^0 = N_+ ; \quad N_L^0 = N_- ; \quad (4.36)$$

In Ref. [27], the metric (4.34) was shown to describe supergravity backgrounds (with SU(2) structure in 6D) such that $N_R^0 > 0$ and $N_L^0 = 0$. On the other hand, for $N_R^0 = N_L^0 = 1$, the metric was shown to possess an almost product structure and take the form

$$ds_6^2 = g_{ab}^{(4)} dx^a dx^b + e^{2(\phi - \phi_0)} g_{cd}^{(2)} dx^c dx^d ; \quad a,b = 4;5;6;7 ; \quad c,d = 8;9 ; \quad (4.37)$$

Here, $g_{ab}^{(4)}$ and ϕ can depend on all six coordinates, but $g_{cd}^{(2)}$ can only depend on x^8 and x^9 .

In contrast, for the orientifold backgrounds studied here, we have argued that the S-dual metric is of the form (4.34), without reference to N_- . For the two results to agree, it must be true that for $N_+ ; N_- = 1$, the 6D metric of the O5 theory takes a form compatible with both (4.34) and (4.37). In Sec. 6.2, we present an example with $N_+ = N_- = 1$, in which this is indeed the case. In this example, the base metric g_{Bmn} reduces to the product metric on T_{f67g}^2 times T_{f89g}^2 , and the fibration is such that we can take

$$A^4 = 2nx^8 dx^6 ; \quad A^5 = 2nx^8 dx^7 ; \quad (4.38)$$

Also, as is true throughout our investigation, $g^{(T_{fib}^2)}$ is constant and the dilaton depends only on the base coordinates. Therefore, the metric (4.34) can be cast in the form (4.37) by writing

$$\begin{aligned} g_{ab}^{(4)} dx^a dx^b &= g^{(T_{fib}^2)} (dx^4 + A^4)(dx^5 + A^5) + e^{2(\phi - \phi_0)} ds_{T_{f67g}^2}^2 ; \\ g_{cd}^{(2)} dx^c dx^d &= ds_{T_{f89g}^2}^2 ; \end{aligned} \quad (4.39)$$

Similar remarks apply to the example in Sec. 7.1, which preserves $N_+ = 1$, $N_- = 2$ supersymmetry.

4.4. Supersymmetry Conditions for More General Flux

The generalization of Eqs. (4:15a;b) to the case of arbitrary flux (2.10) and (2.11) is

$$\frac{1}{2}\hat{\mathcal{G}} + \frac{1}{4}g_s\hat{\mathcal{F}}_{(3)} \hat{u} + \frac{1}{4}e^{(\cdot)}\hat{H}_{(3)} + \frac{1}{2}g_s e^{(\cdot)}\hat{\mathcal{F}}_{(1)} \hat{u} = 0; \quad (4:40a)$$

$$\hat{r}_M + \frac{1}{4}g_s\hat{\mathcal{F}}_{(3)M} \hat{u} + \frac{1}{8}e^{(\cdot)}\hat{H}_{(3)}^{\hat{M}} + 2\hat{M}\hat{H}_{(3)} + \frac{1}{8}g_s e^{(\cdot)}\hat{\mathcal{F}}_{(1)}^{\hat{M}} + 2\hat{M}\hat{\mathcal{F}}_{(1)} + \frac{1}{16}g_s e^{(\cdot)}\hat{\mathcal{F}}_{(5)}^{\hat{M}} \hat{u} = 0; \quad (4:40b)$$

In analyzing the equations of motion in Sec. 3, we made the simplifying assumption that $H_{(3)} = 0$ and found that $\mathcal{F}_{(5)} = 0$ as a result. It can be shown that the same conclusion also follows if instead of using the equations of motion we impose the supersymmetry conditions plus Bianchi identities.²² However, in this section, we will simply take as a starting point that both $H_{(3)}$ and $\mathcal{F}_{(5)}$ vanish.

Since the analysis below is somewhat involved, let us first summarize the results. We will find that the supersymmetry conditions become a remnant of the Hodge duality conditions (3:20a;b;c), together with the condition that $\hat{u} = \hat{u}_+ + \hat{u}_-$ is constant on the fiber and base,

$$\partial \hat{u} = 0; \quad r_{Bm} \hat{u} = 0; \quad (4:41)$$

From the internal 6D point of view, Eq. (4.41) says that \hat{u} is covariantly constant with respect to a torsionful connection that simply forgets about the warping (Z) and fibration (A) in the metric (2.7). The remnant of Eqs. (3:20a;b;c) is

$$G_{(1)} \text{ of type } (0,1), \quad (4:42a)$$

$$\mathcal{G}_{(3)} \text{ (2,1) and primitive,} \quad (4:42b)$$

where

$$G_{(1)} = F_{(1)} - \frac{i}{2}(ZJ \wedge J)Y - \frac{1}{g_s} \text{Vol}_b \wedge H_{(3)}; \quad (4:42c)$$

$$\mathcal{G}_{(3)} = G_{(3)} - \frac{i}{2}Z^{-1}J \wedge G_{(1)}; \quad (4:42d)$$

²² When there is maximal unbroken supersymmetry, the supersymmetry conditions together with the Bianchi identities imply the equations of motion. For reduced supersymmetry, this is not necessarily the case, nevertheless, it does seem to be true for the backgrounds presented in [9,27,11,4], as well as those presented here. See Ref. [33] for a recent discussion.

and $G_{(3)}$ is given by Eqs. (4.28c).²³ Here, the symbol \cdot denotes contraction,

$$A_{(p)} Y B_{(q)} \cdot_{a_1 \dots a_q} \cdot_{p_1 \dots p_p} = \frac{1}{p!} A_{(p)}^{b_1 \dots b_p} B_{(q) b_1 \dots b_p a_1 \dots a_q} \cdot \quad (4.43)$$

To obtain these results, first note that Eqs. (4.40) can be further decomposed based on fiber and base chirality. Using the assumption that $H_{(3)} \cdot_m$ and $F_{(5)}$ vanish, the dilatino equation (4.40a) becomes

$$\frac{1}{2} \cdot^m \mathcal{L}_m \cdot \frac{1}{24} g_s F_{(3) m n p} \cdot^{m n p} \wedge = 0; \quad (4.44a)$$

$$\frac{1}{8} g_s F_{(3) m n} \cdot^{m n} \wedge + \frac{1}{2} g_s e^{(\cdot \cdot)} F_{(1) m} \cdot^m + \frac{1}{24} e^{(\cdot \cdot)} H_{(3) m n p} \cdot^{m n p} \wedge = 0; \quad (4.44b)$$

Eq. (4.40b) can be similarly decomposed. In the $M = 4$ equation, we can take the 4D spacetime spinor u to be constant on $R^{3,1}$; that is, $\mathcal{L}_u = 0$. (If the theory has 4D local supersymmetry, then it also has 4D global Poincaré supersymmetry). Then, this equation becomes

$$r_{B m} \cdot \left(+ \frac{1}{2} \log Z \right) \cdot^m \wedge = 0; \quad (4.45a)$$

$$\frac{1}{8} e^{(\cdot \cdot)} H_{(3) m n p} \cdot^{m n p} - g e^{(\cdot \cdot)} F_{(1) m} \cdot^m \wedge = 0; \quad (4.45b)$$

Here, the leading factors can be eliminated by contracting with $\frac{1}{4}$. Eq. (4.45a) is due to the spin connection $(\hat{\omega})_m$ in the metric (4.15c). It reproduces the condition that g_s as defined in Eq. (3.19) is constant, and it is equivalent to Eq. (4.16a), which states that the prefactor $Z^{-1/2} e^{(\cdot \cdot)}$ in the hatted metric (4.15c) is unity.

For $M = 6$, Eq. (4.40b) decomposes into $\mathcal{L} \cdot \wedge = 0$ and

$$\frac{1}{8} g^{(\tau_{\text{fib}}^2)} F_{m n} \cdot g F_{(3) m n} \cdot^{m n} \wedge = 0 \quad (4.46)$$

as a consequence of Eqs. (4.16a) and (4.45b).

²³ In case the reader is bothered by the explicit appearance of the warp factor Z in Eqs. (4.42c;d), we note that J and γ also contain implicit Z dependence, so that Z drops out of the moduli constraints implied by conditions (4.42a;b) (cf. Sec. 6.3). We can also remove explicit reference to Z from Eqs. (4.42c;d), if desired, by redefining J and γ in terms of the rescaled metric $Z^{-1/2} ds_6^2$, which is the restriction of the 10D string frame metric (2.4) to the internal space.

Finally, using Eqs. (4.16a), (4.44a), and (4.45b), the $M = n$ equation (4.40b) decomposes into $r_{Bn} \wedge = 0$ and

$$\frac{1}{4} g^{(T_{fib}^2)} F_{mn} - g_s F_{(3)nm} \wedge - \frac{1}{8} e^{(\quad)} H_{(3)np} \wedge - \frac{1}{4} g_s e^{(\quad)} F_{(1)n} \wedge = 0: \quad (4.47)$$

In Sec. 3, we found that the equations of motion and Bianchi identities implied a set of Hodge duality constraints (3.20). We can now reproduce two of these constraints from the supersymmetry conditions as follows. For any Hodge dual pair of odd degree forms on B ,

$$!_{(3)} = B !_{(1)}; \quad !_{(1)} = B !_{(3)}; \quad (4.48)$$

we have

$$\frac{1}{6} !_{(3)mp} \wedge = Z !_{(1)r} \wedge; \quad (4.49a)$$

$$\frac{1}{2} !_{(3)mnp} \wedge = Z !_{(1)r} \wedge; \quad (4.49b)$$

From Eq. (4.16a) and the first of these identities, Eqs. (4.44a) and (4.45b) become

$$dZ + g_s B F_{(3)m}^0 \wedge = 0; \quad (4.50a)$$

$$g_s F_{(1)} + B H_{(3)m} \wedge = 0: \quad (4.50b)$$

Therefore,

$$dZ = B g_s F_{(3)}^0; \quad (4.51a)$$

$$g_s F_{(1)} = B H_{(3)}; \quad (4.51b)$$

which are the first two constraints (3.20a;b).

To reproduce the third constraint and obtain the remaining primitivity condition, we will need to introduce $SU(3)_{L,R}$ structures. The three spinor conditions that remain are Eqs. (4.44b), (4.46), and (4.47). Using Eqs. (4.16a), (4.49a), and (4.51b), the first of these spinor conditions is implied by the other two. So, Eqs. (4.46) and (4.47) are the complete set of equations that remain. Since the latter couples $+$ and $-$, we cannot in general expect to find solutions with $+$ or $-$ set to zero. Therefore, we will write these constraints in terms of the spinor

$$\wedge = e^{(\quad)=4} = \wedge_+ + \wedge_- \quad (4.52)$$

(related to J_R by Eq. (4.3)), and the $SU(3)_R$ structures constructed from this spinor,

$$J_a^b = i \gamma^Y_a{}^b; \quad (4.53a)$$

$$J_{abc} = \gamma^Y_{abc}; \quad (4.53b)$$

Alternatively, we could express the constraints in terms of

$$\hat{J}_L = e^{(\theta)} \hat{J}_R = \hat{J}_+ + \hat{J}_- \quad (4.54)$$

(cf. Eqs (4.7) and (4.8)), and the corresponding $SU(3)_L$ structures

$$J_{L a}^b = i \gamma_L^Y_a{}^b; \quad (4.55a)$$

$$J_{L abc} = \gamma_L^Y_{abc}; \quad (4.55b)$$

However, since \hat{J}_L is related to \hat{J}_R by Eq. (4.8), the two $SU(3)$ structures are also related, so that there is no new information gained in doing this.²⁴ Note that in the case that $\theta = 0$ or $\theta = \pi$ vanishes, we have \hat{J}_R and \hat{J}_L equal up to a sign, and the $SU(3)_R$ and $SU(3)_L$ structures are identical. This was the case in Sec. 4.2.

Using Eqs. (4.16a), and the identities (4.49), the two remaining equations become²⁵

$$6F_{(3)} - \frac{1}{g_s} d\text{Vol}_b{}_{mn}{}^m{}^n = 0; \quad (4.56a)$$

$$F_{(3)} + \frac{i}{g_s} d\text{Vol}_b{}_{nm}{}^m + Z^{1=2} F_{(1)n} - \frac{1}{g_s} Z^{1=2} \epsilon_6(\text{Vol}_b \wedge H_3)_{m}{}^m{}^n = 0; \quad (4.56b)$$

With the sign conventions

$$J_{jk} = (ig_6)_{jk}; \quad (4.57)$$

(equivalent to Eq. (A.3b)), the nonzero components of the ACS are

$$J_j{}^k = i \delta_j{}^k; \quad J_{|}{}^k = i \delta_{|}{}^k; \quad (4.58)$$

Together with Eq. (4.53a), this implies that \hat{J}_L is annihilated by J_j and J^k , while $J_{|}$ and J^k act as creation operators. The spinors $\hat{\chi}, \hat{\chi}'$ form a basis for the space of 6D spinors

²⁴ One can easily show that, for example, $J_{Lmn} = J_{m n}$, $J_L = J$, and $J_{Lm} = J_m$.

²⁵ For $\theta = \pi$, the sign of the second term in Eq. (4.56b) is reversed.

of negative chirality, and $\hat{\cdot}$, $\hat{\cdot}^i$ form a basis for positive chirality 6D spinors. Using Eq. (4.53b) as well, it is then possible to prove the identities²⁶

$$\hat{bc} \wedge = \frac{1}{2} \hat{bc}_d \hat{d} \wedge \hat{iJ}^{bc} \hat{\cdot}; \quad (4.59a)$$

$$\hat{a} \hat{bc} = \hat{abc} \wedge + \frac{1}{4} \hat{d} \hat{bc} \hat{da}_f \hat{iJ}^{bc} \hat{a}_f \hat{\cdot}; \quad (4.59b)$$

The latter implies that for an arbitrary three-form $\hat{\cdot}_{(3)}$,

$$\frac{1}{3!} \hat{\cdot}_{(3)abc} \hat{abc} \wedge = \hat{Y} \hat{\cdot}_{(3)} \wedge \hat{iJY} \hat{\cdot}_{(3)} \hat{\cdot}^i; \quad (4.60)$$

where the contraction operator \hat{y} was defined in Eq. (4.43).

Now, define $G_{(3)}$ as in Eq. (4.28d), and define the complex one-form $\hat{\mu}_X$

$$G_{(1)a} = F_{(1)a} - \frac{i}{g_s} Z_{(6)} (\text{Vol}_b \wedge H_{(3)})_b J^b_a; \quad (4.61)$$

which is equivalent to Eq. (4.42c). Using the Hodge duality constraint (3.20b), the remaining spinor conditions (4.56a;b) contracted with $\hat{\cdot}$ and $\hat{\cdot}^n$, respectively, become

$$\hat{Y} G_{(3)} = 0; \quad \hat{JY} G_{(3)}^{(1;0)} = 0; \quad G_{(1)}^{(1;0)} = 0; \quad (4.62a)$$

$$\hat{Y} G_{(3)} = 0; \quad \hat{iJY} G_{(3)} + Z_{(1=2)} G_{(1)}^{(1;0)} = 0; \quad (4.62b)$$

where we have made use of the identity (4.59b). Eq. (4.42a) implies the Hodge duality condition (3.20b). The constraints on $G_{(3)}$ are equivalent to demanding that $\mathcal{E}_{(3)}$ be primitive and of type $(2;1) + (1;2)$, where $\mathcal{E}_{(3)}$ is given by Eq. (4.42d).

Using this result, the original noncontracted constraints (4.56) eliminate the $(1,2)$ primitive piece via identity (4.59a). Thus, our final condition on $G_{(3)}$ for supersymmetry becomes Eq. (4.42b). This implies, among other things, that $G_{(3)}$ is imaginary-selfdual (ISD), which in turn reproduces the remaining Hodge duality condition (3.20c).

To summarize, the complete set of conditions for solutions to the equations of motion with $N = 1$ supersymmetry consists of the constancy conditions (4.41), the constraints (3.20a) and (4.16a) relating the dilaton and $F_{(3)}^0$ to the warp factor, vanishing one-form $\hat{\mu}_X$ (3.51), and finally the flux constraints (4.42a;b).

²⁶ Similar identities have appeared in numerous places. See, for example ref. [71].

5. Relation to T-dual O3/O5 orientifolds

The O5 orientifolds just discussed are T-dual to O3 orientifolds on $T^2 \times K3$ for $B = K3$ and on T^6 for $B = T^4$. For the rest of the paper, we will restrict to the case that $B = T^4$.

5.1. Review of O3/O5 orientifolds with Internal T^6

In this subsection and the next, symbols with (without) a prime denote quantities associated with the O3 (O5) orientifold. The orientifold projection for O3 planes has already been discussed in connection with the decomposition of the 10D supersymmetry parameters $\epsilon_{L,R}^0$ in Sec. 4.1. The even internal axes preserved by the orientifold projection are $F_{(3)}^0$ and $H_{(3)}^0$. Except in the case that O3 and D3 charges cancel locally, the equations of motion also require odd $F_{(5)}^0$ flux. For compact internal manifold X_6^0 , the most general $N^0 = 1$ O3 orientifold background compatible with 4D Poincare invariance is [6]

$$ds_{\text{string}}^2 = Z^{0 \cdot 1=2} dx dx + Z^{0 \cdot 1=2} ds_6^2; \quad (5.1a)$$

$$e^0 = g_s^0 = \text{const}; \quad (5.1b)$$

$$\frac{1}{g_s^0} r_{X_6^0}^2 Z^0 = (2^{-1})^4 \omega^X \sum_i Q_i^0 \frac{\delta^6(x-x_i)}{g_6^0} \frac{1}{V_6^0}; \quad (5.1c)$$

$$F_{(5)}^0 = (1 + \epsilon_6) d g_s^{0 \cdot 1} Z^{0 \cdot 1} dx^0 \wedge dx^1 \wedge dx^2 \wedge dx^3; \quad (5.1d)$$

where X_6^0 is a Calabi-Yau threefold, $K3 = T^2$, or T^6 , and V_6^0 is the volume of X_6^0 .

The Gauss's law constraint is

$$N_{\text{ux}}^0 + \sum_i^X Q_i^0 = 0; \quad \text{where } N_{\text{ux}}^0 = \frac{1}{(2^{-1})^4 \omega^X} \int_{X_6^0} F_{(3)}^0 \wedge H_{(3)}^0; \quad (5.1e)$$

Here, $f(Q_i^0; x_i)$ is the set of charges and positions of local O3 and D3 sources, with Q_i^0 equal to 1 for a D3 brane. We assume that all O3 planes are standard O3 planes in the terminology of Ref. [59], and work on the covering space of the orientifold, so that there are M D3 branes and $M/2$ image branes. The orientifold operation is $(-1)^{F_1} I_6$, where I_6 inverts X_6^0 . For $X_6^0 = T^6$, the O3 planes are located at the 2^6 fixed points on the base where each x^a is equal to 0 or $1=2$, and each O3 plane has charge $Q_i^0 = 1=2$.

In the absence of three-form flux, the theory preserves 4D $N^0 = 4$ supersymmetry. The 4D moduli are the zero-modes on X_6^0 of

$$\text{dil}^0 = C_{(0)}^0 + i g_s^0; \quad g_{6ab}^0; \quad C_{(2)ab}^0; \quad \text{and} \quad C_{(4)abcd}^0; \quad (5.2)$$

together with the D3 worldvolume scalars $\phi_I, I = 1; \dots; M$. The massless 4D gauge bosons are the zero-modes of

$$B_{(2)a}^0 \quad \text{and} \quad C_{(2)a}^0; \quad (5:3)$$

together with the D3 worldvolume gauge bosons A_I .

In the presence of three-form flux, the supersymmetry and massless field content is reduced [9]. For the case that $X_6^0 = T^6$, this model has been analyzed in great detail [11,60]. The possible choices of NS and RR three-form flux are $H_{(3)}^0; F_{(3)}^0 \in (2,2)^0 H^3(T^6; 2Z)$. That is,

$$\begin{aligned} H_{(3)}^0 &= (2,2)^0 m_{[abc]} dx^a \wedge dx^b \wedge dx^c; \quad m_{[abc]} \in 2Z; \\ F_{(3)}^0 &= (2,2)^0 n_{[abc]} dx^a \wedge dx^b \wedge dx^c; \quad n_{[abc]} \in 2Z; \quad a;b;c = 1; \dots; 6; \end{aligned} \quad (5:4)$$

Without imposing any supersymmetry conditions, the equations of motion alone imply the ISD condition

$$*_6 G_{(3)}^0 = i G_{(3)}^0; \quad (5:5)$$

where

$$G_{(3)}^0 = F_{(3)}^0 - \text{dil}^0 H_{(3)}^0; \quad (5:6)$$

Since $H_{(3)}^0$ and $F_{(3)}^0$ are discrete, this is a constraint on dil^0 and metric moduli.

The condition for $N^0 = 1$ 4D (Poincare) supersymmetry further refines this to

$$G_{(3)}^0 \in (2,1) \text{ and primitive.} \quad (5:7)$$

The complex structure on T^6 can be parametrized by a complex 3×3 period matrix Q_{ij}^0 ,

$$z^i = x^i + Q_{ij}^0 y^j; \quad (5:8)$$

where $z^i = z^i + 1 = z^i + Q_{ij}^0$, for $i;j = 1;2;3$. The holomorphic three-form is

$$*_6 J^0 / dz^1 \wedge dz^2 \wedge dz^3; \quad (5:9)$$

which depends on the complex structure moduli up to cubic order in the period matrix.

The Kahler form is

$$J^0 = i g_{i\bar{j}}^0 dz^i \wedge d\bar{z}^{\bar{j}}; \quad (5:10)$$

The (2,1) supersymmetry condition is easily implemented by varying the superpotential of Gukov, Vafa, and Witten [57]

$$W_{\text{GVW}} = \int_{X_6^0} G_{(3)}^0 \wedge \text{dil}^0; \quad (5.11)$$

with respect to dil^0 and all complex structure moduli α_j^0 , and in addition imposing the condition $W_{\text{GVW}} = 0$. (The latter follows from $D W_{\text{GVW}} = 0$, where D is the complexified overall volume modulus [9,11].) For generic supersymmetric vacua, this fixes dil^0 and all of the complex structure moduli, but for nongeneric flux, some of these moduli are left unfixd. The primitivity condition then becomes a linear constraint on the $g_{i\bar{j}}^0$, with coefficients determined by the flux and axion-dilaton. One subtlety in this procedure for $X_6^0 = T^6$ is that not all of the (18 real) α_j^0 and (9 real) $g_{i\bar{j}}^0$ correspond to the (21 real) physical metric moduli. See App. E for further discussion. Due to the subtleties discussed in App. E concerning W_{GVW} in theories with extended supersymmetry, we have chosen in this paper to avoid any reference to a superpotential for the O5 orientifold, and instead to state the supersymmetry conditions directly in terms of conditions on the fields.

In the O3 orientifold, the decomposition of RR potentials into a background part $C_{(p)}^{\text{bg}}$ and deformation $c_{(p)}^0$,

$$C_{(p)}^0 = C_{(p)}^{\text{bg}} + c_{(p)}^0; \quad (5.12)$$

is cleaner than it was for the O5 orientifold. First, set the gauge fields to zero and focus on deformations corresponding to moduli. Then,

$$C_{(0)}^0 = c_{(0)}^0; \quad C_{(2)}^0 = C_{(2)}^{\text{bg}}; \quad (5.13a)$$

and it is only in $C_{(4)}^0$ that both contributions appear:

$$C_{(4)}^0 = C_{(4)}^{\text{bg}} + c_{(4)}^0; \quad (5.13b)$$

Here, $F_{(3)}^0$, $H_{(3)}^0$, and $F_{(5)}^0$, as given by Eqs. (5.4) and (5.1d), satisfy

$$\begin{aligned} F_{(3)}^0 &= dC_{(2)}^{\text{bg}} \\ F_{(5)}^0 &= dC_{(4)}^{\text{bg}} + C_{(2)}^{\text{bg}} \wedge H_{(3)}^0; \end{aligned} \quad (5.14)$$

In addition to the deformations just discussed, there are other deformations of the supergravity background which when promoted to 4D fields become gauge bosons. In the closed string sector, these deformations are the zero modes on X_6^0 of $B_{(2)d}^0$ and $C_{(2)d}^0$,

and will be denoted by the lowercase symbols $b_{(2)d}^0$ and $c_{(2)d}^0$, respectively. When the couplings to gauge bosons are included and all deformations are promoted to 4D fields, the kinetic term for $c_{(4)abcd}^0$ is the square of [60]

$$\partial \left[c_{(4)abcf}^0 + 4F_{(3)[abcj]}^0 b_{(2)jf}^0 \right] - 4H_{(3)[abcj]}^0 c_{(2)jf}^0 : \quad (5.15)$$

If we define complex gauge bosons

$$d_a^0 = c_{(2)a}^0 - \text{dil} \, b_{(2)a}^0 ; \quad \text{and} \quad \tilde{d}_a^0 = c_{(2)a}^0 + \text{dil} \, b_{(2)a}^0 ; \quad (5.16)$$

then Eq. (5.15) can also be written as

$$\partial \left[c_{(4)abcf}^0 - \frac{1}{\text{Im} \, \text{dil}^0} \text{Im} \, 4G_{(3)[abcj]}^0 d_{jf}^0 \right] : \quad (5.17)$$

5.2. T-Duality Map

In this subsection, as in the previous one, symbols with (without) a prime denote quantities associated with the $O(3,1)/O(5)$ orientifold. Let $X_6^0 = T^6$ and $B = T^4$. Then, the metric on X_6^0 can be written as a trivial T^2 fibration over B :

$$ds_6^2 = ds_B^2 + g^{(T_{\text{fib}}^2)^0} da^0 ; \quad \omega^0 = dx^0 + a^0 ; \quad (5.18)$$

where trivial means that $F^0 = da^0 = 0$. The use of a lowercase a^0 indicates that the a_{mn}^0 are moduli rather than a quantized background. Until the very end of this subsection, we set all gauge fields to zero.

For the $O(5)$ orientifold, we write NSB-field as $B_{(2)} = B_{(2)}^{bg} + b_{(2)}$, where, from Sec. 3.9,

$$\begin{aligned} B_{(2)}^{bg} &= B_{(2)}^{bg0} = \frac{1}{2} B_{(2)mn} dx^m \wedge dx^n ; \\ b_{(2)} &= b_{(2)}^1 = \omega^0 \wedge b_{(2)} ; \quad \text{with} \quad b_{(2)} = b_{(2)m} dx^m ; \end{aligned} \quad (5.19)$$

The NSB-field for the $O(3)$ orientifold is given by $B_{(2)}^0 = B_{(2)}^{bg}$, where $B_{(2)}^{bg} = B_{(2)}^{bg0} + B_{(2)}^{bg1}$, and

$$\begin{aligned} B_{(2)}^{bg0} &= \frac{1}{2} B_{(2)mn}^{bg} dx^m \wedge dx^n ; \\ B_{(2)}^{bg1} &= \omega^0 \wedge B_{(2)}^{bg} ; \quad \text{with} \quad B_{(2)m}^{bg} = B_{(2)m}^{bg} dx^m ; \end{aligned} \quad (5.20)$$

T-duality is an exact duality between string vacua that relates conformal field theories order by order in string perturbation theory. At the level of the conformal field theories, it has the simple interpretation as the sign reversal $X_R(z) \rightarrow X_R(\bar{z})$ of the right-moving worldsheet scalars in the duality directions. However, in the low energy supergravity

description, it involves an intermediate operation known as smearing. In the $O3$ orientifold, the locations of the $O3$ planes and $D3$ branes spontaneously break the translation isometries in each of the internal T^6 directions. The breaking is spontaneous in the sense that, on the Z_2 covering space, the entire $O3/D3$ system can be translated by an arbitrary amount along any of the T^6 circles, to produce an inequivalent vacuum with identical relative $O3/D3$ positions and identical physics. The space of vacua respects the isometry, but a particular vacuum does not. Smearing is the operation of averaging a supergravity background over all vacua related by such translations in order to restore an isometry. To perform the T-duality between the $O3$ and $O5$ supergravity backgrounds, we first smear in the 4 and 5 directions and then perform the supergravity analog of the conformal field theory T-duality in these directions. The information about $D3$ brane positions lost in the smearing is recovered in the Wilson lines of the $U(1)$ worldvolume gauge fields in the 4 and 5 directions.

The effect of the smearing is that the warp factors Z and Z^0 are related via

$$Z = Z_{\text{smear}}^0 = \frac{1}{V_b^0} \int_{T_{\text{fib}}^2} \text{Vol}_b^0 Z^0; \quad (5.21)$$

where

$$\text{Vol}_b^0 = V_b^0 \omega^4 \wedge \omega^5; \quad \text{and} \quad V_b^0 = g^{(T_{\text{fib}}^2)} \omega^{1=2} = (2\pi\alpha')^4 \omega^4 \omega^5 = V_b; \quad (5.22)$$

which is easily seen to relate solutions of the 6D Poisson equation (5.1c) to solutions of the 4D Poisson equation (3.29).

The T-duality action on the dilaton is

$$e^{2\phi} = Z^{-1} V_b = e^{\phi_0}; \quad (5.23)$$

or equivalently

$$g_s (2\pi\alpha')^2 = V_b = g_s^0; \quad (5.24)$$

The T-duality action on the NSB-field and metric interchanges the geometrical S^1 -bracket of connection a^0 with the formal S^1 -bracket of connection $B_{(2)}^{\text{bg}}$.²⁷ The relations are [38]

$$b_{(2)} = a^0 (2\pi\alpha')^2; \quad (5.25a)$$

$$A = B_{(2)}^{\text{bg}} = (2\pi\alpha')^2 a^0; \quad (5.25b)$$

$$B_{(2)}^{\text{bg}} = B_{(2)}^{\text{bg}0} + a^0 \wedge B_{(2)}^{\text{bg}}; \quad (5.25c)$$

²⁷ This point has been emphasized recently, first in Ref. [72] (and subsequent work [73]) and then in Ref. [38].

together with

$$g^{(T^2_{\text{fib}})} = g^{(T^2_{\text{fib}})0} \quad (2)^2 \quad 0^2; \quad (5.25d)$$

$$g_{B m n} = g_{B m n}^0; \quad (5.25e)$$

Note that the T-duality map does not leave the purely base component $B_{(2)}^0$ invariant.²⁸ Nevertheless, the correction term in Eq. (5.25c) has a straightforward interpretation. Let underscored indices $\underline{m}; \underline{n}$ denote components in the $dx^{\underline{m}}; dx^{\underline{n}}; dx^m$ basis as opposed to the bration-adapted bases involving \underline{m} or \underline{n} . Then Eq. (5.25c) is equivalent to

$$B_{(2)m n}^{bg} = B_{(2)\underline{m} \underline{n}}^{0bg}; \quad (5.26)$$

This illustrates an important rule. Many (but not all) of the T-duality relations connecting 03 and 05 orientifolds take the simplest form when expressed in terms of the $dx^{\underline{m}}; dx^{\underline{n}}; dx^m$ basis for the 05 orientifold and the $dx^{\underline{m}}; dx^{\underline{n}}; dx^m$ basis for the 03 orientifold. This is perhaps to be expected, since in the 03 orientifold there is nothing special about the $\underline{m} = 4; 5$ directions| any other pair of directions could have been used to define the fiber of a T^2 bration. The three-form fluxes of the 03 orientifold are of the moduli-independent form (5.4). Thus, $H_{(3)\underline{m} \underline{n}}^0$ and $H_{(3)m n r}^0$ are quantized and moduli-independent, whereas $H_{(3)m n r}^0$ involves a combination of quantized fluxes and the metric moduli $a^0_{m n}$. From the T-duality map (5.25), the relation between quantized NS sector fluxes is

$$F_{m n} = H_{(3)\underline{m} \underline{n}}^0 = (2)^2 \quad 0^0 \quad \text{and} \quad H_{(3)m n r}^{bg} = H_{(3)\underline{m} \underline{n} r}^0; \quad (5.27)$$

The relation between NS sector moduli has already been given explicitly in Eqs. (5.25a;d;e).

The rule described in the previous paragraph is particularly applicable to the T-duality action on RR fluxes and moduli. The T-duality relations between RR fluxes are [74,64,75]

$$\begin{aligned} F_{(n)}^0 &= \frac{1}{2} \quad F_{(n+2)}^{02} = (2)^2 \quad 0^0; \\ F_{(n)}^1 &= \quad F_{(n)}^{01}; \\ F_{(n)}^2 &= \quad F_{(n-2)}^{00} (2)^2 \quad 0^0; \end{aligned} \quad (5.28)$$

Here $\epsilon^{45} = \epsilon^{54} = 1$, and we have assumed that the ordering of the T-dualities in going from the 03 theory to the 05 theory is that we first T-dualize in the 4-direction and then in the 5-direction.

²⁸ I am grateful to A. Tomasiello for correspondence regarding this point.

In Eq. (5.28), both sides involve a combination of moduli-dependent and quantized contributions. It is desirable to disentangle the two types of contributions. When the RHS is expanded in terms of a^0 , $c_{(0)}^0$, and underscored flux components, the relations become

$$F_{(1)m} = F_{(3)}^0 \underset{(0)}{C}_{(0)} H_{(3) \underline{45m}}^0 = (2^-)^2 \cdot 0; \quad (5.29a)$$

$$F_{(3) \underline{m n}} = F_{(3)}^0 \underset{(0)}{C}_{(0)} H_{(3) \underline{m n}}^0 - 2a_{\underline{m j}}^0 F_{(3)}^0 \underset{(0)}{C}_{(0)} H_{(3) \underline{45 j n}}^0; \quad (5.29b)$$

$$\begin{aligned} \mathbb{F}_{(5) \underline{45m n r}} = & F_{(3)}^0 \underset{(0)}{C}_{(0)} H_{(3) \underline{m n r}}^0 - 3a_{\underline{m j}}^0 F_{(3)}^0 \underset{(0)}{C}_{(0)} H_{(3) \underline{j n r}}^0 \\ & + 6a_{\underline{m j}}^0 a_{\underline{j n}}^0 F_{(3)}^0 \underset{(0)}{C}_{(0)} H_{(3) \underline{45 j r}}^0 - (2^-)^2 \cdot 0; \end{aligned} \quad (5.29c)$$

In addition, the map of odd fluxes is

$$\mathbb{F}_{(3) \underline{m n r}} = \mathbb{F}_{(5) \underline{sm eared} \underline{45m n r}}^0 = (2^-)^2 \cdot 0; \quad (5.29d)$$

Here, the vector flux in the smeared supergravity background is

$$\mathbb{F}_{(5) \underline{sm eared}}^0 = \frac{1}{g_s^0} \text{Vol}_b \cdot \mathbb{D}z_{\underline{sm eared}}^0; \quad (5.30)$$

By Eqs. (5.21) and (5.24), Eq. (5.29d) agrees with Eq. (3.20b). In the remaining relations (5.29a;b;c), we can drop the components $H_{(3) \underline{45m}}^0$, which necessarily vanish for the T-duality map to exist (cf. Footnotes 7 and 12). Upon setting these terms to zero and using the map (5.25a), Eqs. (5.29c;d) can be identified, term for term, with Eqs. (3.43) and (3.47a;b). The result is that the T-duality map between quantized fluxes is

$$F_{(1)m} = F_{(3) \underline{45m}}^0 = (2^-)^2 \cdot 0; \quad (5.31a)$$

$$F_{(3) \underline{4m n}}^{\text{bg}} = F_{(3) \underline{5m n}}^0 \quad \text{and} \quad F_{(3) \underline{5m n}}^{\text{bg}} = F_{(3) \underline{4m n}}^0; \quad (5.31b)$$

$$F_{(5) \underline{45m n r}}^{\text{bg}} = F_{(3) \underline{m n r}}^0 - (2^-)^2 \cdot 0; \quad (5.31c)$$

and the axionic partner of the dilaton maps as

$$e_{(2) \underline{45}} = (2^-)^2 \cdot c_{(0)}^0; \quad (5.31d)$$

From $H_{(3) \underline{45m}}^0 = 0$ together with Eq. (5.5), the flux components $F_{(3) \underline{m n r}}^0$ vanish. Therefore,

$$F_{(5) \underline{45m n r}}^{\text{bg}} = 0; \quad (5.32)$$

which is a result that we stated without proof in Sec. 3.9. Eqs. (5.31d) and (5.24) together give

$$d_{\text{dil}} = d_{\text{dil}}^0; \quad (5.33)$$

where d_{dil} is defined in Eq. (4.29) and d_{dil}^0 in Eq. (5.2). See App. F for a discussion of the T-duality map between RR potentials.

The T-duality map between supersymmetry parameters is [75]

$$\begin{aligned} \epsilon_R &= \epsilon_R^0; \\ \epsilon_L &= g^{(T_{\text{fib}}^2)} \epsilon_L^0 = \epsilon_L^0; \end{aligned} \quad (5.34)$$

in agreement with the results of Sec. 4.1. Here, as explained in Ref. [75], given a vielbein e^A_M such that $g_{MN}^0 = \eta_{AB} e^A_M e^B_N$ in the O3 theory, one obtains two different vielbeine $(e_{L;R})^A_M$ in the dual O5 theory, and correspondingly two different Dirac matrix representations $(\gamma_{L;R})_M = (\gamma_{L;R})^A_M \gamma_A$, for fixed choice of representation of γ_A satisfying the algebra $\{\gamma_A, \gamma_B\} = 2\eta_{AB}$. The two vielbeine are related by a local Lorentz transformation. In writing Eq. (5.34), an arbitrary choice has been made to define

$$(\gamma_R)_M = (\gamma_L)_M = (\gamma_R)^A_M \gamma_A; \quad (5.35)$$

and to re-express $(e_L)^A_M$ in terms of $(e_R)^A_M$ wherever it appears. The net effect of this is to supplement T-duality with a local Lorentz transformation in the left moving worldsheet sector, which leads to nontrivial T-duality map of ϵ_L .

Finally, when the deformations of the supergravity background corresponding to 4D gauge bosons are reintroduced, the T-duality map between these deformations is

$$\begin{aligned} v &= v_{(2)}^0 = (2)^2 v^0; \quad b_{(2)m} = b_{(2)m}^0; \\ c_{(2)} &= e c_{(2)}^0; \quad e_{(4)45m} = c_{(2)m}^0 (2)^2 v^0; \end{aligned} \quad (5.36)$$

6. $N = 2$ Examples

In this section, we present three $N = 2$ examples in succession. The first two examples contain $F_{(3)}$ flux only, while the third example also contains $F_{(1)}$ and $H_{(3)}$ flux. The three examples are T-dual to the same O3 background, given in Sec. 6.4, via three different choices of the cycles on which to T-dualize.

6.1. Example 1: $N_+ = 2, N_- = 0$

Consider the choice ofibration curvature and background flux

$$F^4 = 2n dx^6 \wedge dx^8 + dx^7 \wedge dx^9; \quad F^5 = 0; \quad (6.1a)$$

$$F_{(3)}^{bg1} = (2m)^2 dx^4 \wedge dx^6 \wedge dx^8 + dx^7 \wedge dx^9; \quad (6.1b)$$

with $F_{(1)} = H_{(3)} = 0$. This choice gives a contribution to the Gauss's law constraint,

$$N_{ux} = 8mn; \quad (6.2)$$

so that Eq. (3.15) becomes

$$2M + 8mn = 32; \quad (6.3)$$

As discussed in Sec. 3.2, N_{ux} is nonnegative. So, m and n have the same sign, and the number of D5 branes is $2M = 0, 8, 16, 24$, or 32 , depending on the choice of integers m and n . We assume that $m, n \neq 0$, so that $2M$ is strictly less than 32 .

The complex three-form flux is

$$\frac{1}{(2m)^2} G_{(3)} = 2m dx^4 + (n=m) dx^5 \wedge dx^6 \wedge dx^8 + dx^7 \wedge dx^9; \quad (6.4)$$

If we define an almost complex structure via (cf. Eq. (4.24))

$$\begin{aligned} z^1 &= dx^4 + dx^5; \\ dz^i &= dx^{2i+2} + dx^j dx^{2j+3}; \quad i, j = 2, 3; \end{aligned} \quad (6.5)$$

then, using the decomposition (4.21), the type (2,1) condition on the flux is

$$G_1 = (n=m) dx^5 \quad \text{and} \quad \det_2 G_2 = 1; \quad (6.6)$$

Note that the first condition implies that only for large number complex structure $\text{Im } G_1 = 1$ can we simultaneously have $g_s = 1$ and $V_b = 0$. Eq. (6.6) is equivalent to demanding that the two factors appearing in $G_{(3)}$ be of definite Hodge type:

$$\begin{aligned} dx^4 & \quad (n=m) dx^5 \quad \text{of type } (1,0), \\ dx^6 \wedge dx^8 + dx^7 \wedge dx^9 & \quad \text{of type } (1,1). \end{aligned} \quad (6.7)$$

Thus, from Eq. (6.1a), the complexibration curvature $F^{z^1} = F^4 + dx^1 F^5$ is also of type (1,1). This provides a check of our formal result that the complex structure is integrable

in the case of RR three-form flux only. The vanishing of the (0,2) component of $F^z{}^1$ is all that is required for theibration to be holomorphic; that condition is satisfied.

Finally, the primitivity condition is

$$J^B \wedge dx^6 \wedge dx^8 + dx^7 \wedge dx^9 = 0; \quad (6:8)$$

Not all of the i_j and g_{Bij} left unixed by these constraints correspond to independent physical moduli. (This is the T^4 version of the ambiguity discussed in App. E.) We can ix the redundancy at the cost of breaking manifest $SU(2)$ covariance in the parametrization of the metric and complex structure on the base T^4 . We write the T^4 metric as a at T_{f67g}^2 bration over T_{f89g}^2 ,

$$ds_B^2 = \frac{V_2}{j\text{Im } 2j} e^6 + 2e^7{}^2 + \frac{V_3}{j\text{Im } 3j} dx^8 + 3dx^9{}^2; \quad (6:9)$$

where

$$\begin{aligned} e^6 &= \mathbf{e}^6_8 dx^8 + \mathbf{e}^6_9 dx^9; \\ e^7 &= \mathbf{e}^7_8 dx^8 + \mathbf{e}^7_9 dx^9; \end{aligned} \quad (6:10)$$

with $F^m = da^m = 0, m = 6;7$. We take the (1,0)-forms on X_6 to be

$$z^1 = 4 + 1 5; \quad e^{z^2} = e^6 + 2e^7; \quad \text{and} \quad dz^3 = dx^8 + 3dx^9; \quad (6:11)$$

instead of those given in Eq. (6.5). Then,

$$\begin{aligned} B &= \frac{V_2 V_3}{j\text{Im } 2j j\text{Im } 3j} {}^{1=2} e^6 + 2e^7 \wedge dx^8 + 3dx^9; \\ J^B &= (\text{sign } 2)V_2 e^6 \wedge e^7 + (\text{sign } 3)V_3 dx^8 \wedge dx^9; \end{aligned} \quad (6:12)$$

The (2,1) condition becomes

$$1 = (n=m)_{dil}; \quad (6:13a)$$

$$2 3 = 1; \quad (6:13b)$$

and the primitivity condition becomes

$$\mathbf{e}^6_9 = \mathbf{e}^7_8; \quad (6:14)$$

The constraints on NS B-eld moduli follow from Eqs. (3:52a;b). These equations become

$$0 = h_{(3)} = b_{(2)4m} F^4 \wedge dx^m; \quad (6:15a)$$

and

$$0 = b_{(2)5m} dx^m \wedge F_{(3)4}^{bg}; \quad (6:15b)$$

respectively. The first equation eliminates $b_{(2)4m}$, and the second $b_{(2)5m}$. So, there are no NSB-eld moduli.

The massless RR moduli are those that do not couple to the gauge elds. (Those that do are eaten via the supersymmetric Higgs mechanism). From the kinetic terms (3:54), they are

$$e_{(2)89}; e_{(2)78}; e_{(2)69}; e_{(2)67}; e_{(2)68} \quad e_{(2)79}; \text{ and } e_{(2)} \quad e_{(6)456789}; \quad (6:16)$$

Similarly, the massless gauge bosons are those that do not couple to RR scalars. From (3:54), they are the three linear combinations of v^4 , v^5 , $c_{(2)4}$, and $c_{(2)5}$ orthogonal to

$$(2)^2 \quad (m=n)v^4 + c_{(2)4}; \quad (6:17)$$

where orthogonality is defined by the metric appearing in the gauge kinetic terms. (See Sec. 6.4 for further discussion in the context of the dual $O3$ orientifold).

Finally, there are $6M$ massless scalars A_I , I^m and M massless gauge bosons A_I from the D5 branes.

In summary, the massless bosonic elds are one graviton, $3+M$ vectors, and $16+6M$ moduli. The moduli consist of

$$V_b, V_2, V_3, 2 \text{ indep } \quad , 3 \text{ indep } e^m_n; \\ 6 e_{(2)} \text{ scalars, and } 6M \text{ D5 scalars.}$$

These elds combine to form one 4D $N = 2$ gravity multiplet, $2+M$ vector multiplets, and $3+M$ hypermultiplets.

The amount of unbroken supersymmetry is

$$N_+ = 2; \quad N_- = 0; \quad (6:18)$$

in the notation of Sec. 4.2. To verify this, first note that since $V_b; g_s > 0$, we have $\text{Im}_{\text{dil}} > 0$. Then, from the moduli constraints, $\text{Im}_1 > 0$. So, $J^b = + \text{Vol}_b$, and we conclude that $= +$ and $N_- = 0$. Next, observe that there is exactly one antiholomorphic involution of X_6 compatible with the constraints on complex structure moduli:

$$1; \quad i_j \quad ! \quad 1; \quad i_j \quad (6:19)$$

(or equivalently, $1; 2; 3 \quad ! \quad 1; 2; 3$ in the complex structure (6.11)). Therefore, there are two independent 6D Killing spinors. The spinors are of the form

$$1+ = \quad +^b \quad +^B; \quad 2+ = \quad +^b \quad +^B; \quad (6:20)$$

for some e_+^b and e_+^B , and are related by complex conjugation of the base.

6.2. Example 2: $N_+ = N_- = 1$

Chooseibration curvature and background flux

$$F^4 = 2n dx^6 \wedge dx^8; \quad F^5 = 2n dx^7 \wedge dx^8; \quad (6.21a)$$

$$F_{(3)}^{bg1} = (2)^2 \cdot 0 = 2m^4 \wedge dx^7 \wedge dx^5 \wedge dx^6 \wedge dx^9; \quad (6.21b)$$

Again, m , n , and M are constrained by Eq. (6.3). The complex three-form flux is

$$\frac{1}{(2)^2 \cdot 0} G_{(3)} = 2n_{dil}^4 \wedge dx^7 \wedge dx^5 \wedge dx^6 \wedge dx^8 + (m=n)(1=dil) dx^9; \quad (6.22)$$

Define the complex structure as in Eq. (6.5). Then, the type (2,1) condition gives

$$J_{2,2} = \text{diag}(J_2; J_3); \quad (6.23a)$$

$$J_2 = J; \quad (6.23b)$$

$$J_3 = (m=n)(1=dil); \quad (6.23c)$$

So, the base factorizes into $T_{fx^6; x^7}^2 \times T_{fx^8; x^9}^2$ with respect to complex structure. Primitivity implies that g_{Bij} factorizes in the same way. Therefore, the base metric takes the form

$$ds_B^2 = \frac{V_2}{j \text{Im}_2 j} dz^{2,2} + \frac{V_3}{\text{Im}_3} dz^{3,2}; \quad (6.24)$$

Here, we have removed the absolute value bars from Im_3 since the nonnegativity of Im_{dil} and $m=n$ implies nonnegativity of $\text{Im}(1=3)$ and Im_3 .

From Eqs. (3.52a;b), the constraints on NSB-eld moduli are

$$\begin{aligned} 0 &= h_{(3)} = b_{(2)4m} F^4 + b_{(2)5m} F^5 \wedge dx^m; \\ &= 2n b_{(2)4m} dx^6 + b_{(2)5m} dx^7 \wedge dx^8 \wedge dx^m \end{aligned} \quad (6.25a)$$

and

$$\begin{aligned} 0 &= b_{(2)4m} dx^m \wedge F_{(3)5}^{bg} + b_{(2)5m} dx^m \wedge F_{(3)4}^{bg}; \\ &= 2m dx^m \wedge b_{(2)4m} dx^6 + b_{(2)5m} dx^7 \wedge dx^9; \end{aligned} \quad (6.25b)$$

These equations are equivalent to

$$b_{(2)49} = b_{(2)59} = 0; \quad b_{(2)56} = b_{(2)47}; \quad (6.26a)$$

$$b_{(2)48} = b_{(2)58} = 0; \quad b_{(2)56} = b_{(2)47}; \quad (6.26b)$$

respectively. So, there are three unconstrained NS B - eld m oduli,

$$b_{(2) 46}; b_{(2) 57}; \text{ and } b_{(2) 56} + b_{(2) 47}; \quad (6:27)$$

In the RR sector, the uneaten m oduli that follow from the kinetic term s (3:54) are the uncharged scalars

$$e_{(2) 67}; e_{(2) 89}; e_{(4) 4689}; e_{(4) 5789}; e_{(4) 4789} + e_{(4) 5689}; \text{ and } e_{(2)} \text{ \& } e_{(6) 456789}; \quad (6:28)$$

The massless gauge bosons are those that do not couple to RR scalars. From (3:54), they are the three linear combinations of $b_{(2) 8}$, $b_{(2) 9}$, $e_{(4) 458}$, and $e_{(4) 459}$ orthogonal to

$$(2)^2 \text{ }^0(m=n)b_{(2) 8} \quad e_{(4) 459}; \quad (6:29)$$

where orthogonality is defined by the metric appearing in the gauge kinetic term s. (See Sec. 6.4 for further discussion in the context of the dual O 3 orientifold).

Finally, there are $6M$ massless scalars A_I , I^m and M massless gauge bosons A_I from the D 5 branes.

In summary, as in Ex. 1, the massless bosonic elds are one graviton, $3 + M$ vectors, and $16 + 6M$ moduli. However, the moduli now consist of

$$V_b, V_2, V_3, 2 \text{ indep } , 3 \text{ indep } b_{(2) m}, \\ 3 e_{(2)} \text{ scalars, } 3 e_{(4)} \text{ scalars, and } 6M \text{ D 5 scalars.}$$

These elds combine to form one 4D $N = 2$ gravity multiplet, $2 + M$ vector multiplets, and $3 + M$ hypermultiplets.

The amount of unbroken supersymmetry is

$$N_+ = 1; N_- = 1; \quad (6:30)$$

in the notation of Sec. 4.2. Due to the factorization $B = T_{fz^2g}^2 \quad T_{fz^3g}^2$ and the earlier observation that $\text{Im } \tau_3 > 0$, we can write

$$= b \quad B; \text{ where } B = \text{}^{(2)} \quad \text{}^{(3)}; \quad (6:31)$$

for some e_b , $\text{}^{(2)}$ and $\text{}^{(3)}$. Here, $\text{}^{(i)}$ is a spinor on $T_{fz^i g}^2$, and the chirality of $\text{}^{(3)}$ is fixed by the condition $\text{Im } \tau_3 > 0$. The two 6D Killing spinors $+$ and $-$ are related by complex conjugation of T_b^2 and $T_{fz^2g}^2$,

$$1; 2; 3! \quad 1; 2; 3; \quad (6:32)$$

that is,

$$+_b = b; \quad \text{}^{(2)}_+ = \text{}^{(2)}_-; \quad (6:33)$$

6.3. Example 3: More General Flux

Choose bration curvature and background flux

$$F^5 = 2n dx^6 \wedge dx^9 \quad (6:34a)$$

$$F_{(1)} = 2m dx^6 \quad (6:34b)$$

$$F_{(3)}^{bg1} = (2)^2 \cdot 0 = 2m^5 \wedge dx^7 \wedge dx^8 \quad (6:34c)$$

$$H_{(3)}^{bg} = (2)^2 \cdot 0 = 2n dx^7 \wedge dx^8 \wedge dx^9 \quad (6:34d)$$

Again, m , n , and M are constrained by Eq. (6.3). The flux $G_{(3)}$ is

$$\frac{1}{(2)^2 \cdot 0} G_{(3)} = 2m^5 \wedge dx^7 \wedge dx^8 - 2n dx^4 \wedge dx^6 \wedge dx^9 + b_{(2)} \wedge 2m dx^6: \quad (6:35)$$

From the moduli constraints, it is possible to show that $b_{(2)}$, J , and the complex structure all decompose as (49)–(5678). We omit the proof here, since it is tedious and uninteresting; however, this factorization is dual to an analogous factorization for the O3 orientifold discussed in the next subsection. Due to the factorization, we have

$$ds_{X_6}^2 = ds_{5678}^2 + ds_{49}^2; \quad (6:36)$$

where the most general metric in the 5678 directions can be written as

$$ds_{5678}^2 = \frac{V_1}{j\text{Im}_{1j}} \cdot 5 + {}_1Z^{1=2} e^8 \cdot 2 + Z \frac{V_2}{j\text{Im}_{2j}} dx^6 + {}_2dx^7 \cdot 2; \quad (6:37a)$$

with

$$e^8 = dx^8 + e^8_m dx^m; \quad m = 6, 7; \quad (6:37c)$$

parametrizing a S^1 bration, and the most general metric in the 49 directions can be written

$$ds_{49}^2 = \frac{V_3}{j\text{Im}_{3j}} dx^4 + {}_3Z^{1=2} dx^9 \cdot 2: \quad (6:38)$$

Here ${}_1$ and ${}_3$ are pure imaginary, but ${}_2$ can have both real and imaginary parts.

In this parametrization, the (3,0) form and Kähler form are

$$= \frac{V_1 V_2 V_3}{j\text{Im}_{1j} j\text{Im}_{2j} j\text{Im}_{3j}} \cdot 1=2 \cdot 5 + {}_1Z^{1=2} e^8 \wedge Z^{1=2} dx^6 + {}_2dx^7 \wedge dx^4 + {}_3Z^{1=2} dx^9; \quad (6:39a)$$

This takes care of the condition on $G_{(1)}$. The remaining supersymmetry condition (4:42b) on $\mathcal{G}_{(3)}$ is most easily imposed by first demanding the weaker condition

$$Z^{1=2} J_Y G_{(3)} = iG_{(1)} \quad (6:44)$$

(cf. Eq. (4:62)). From the above expressions for J and $G_{(3)}$, we obtain

$$Z^{1=2} J_Y G_{(3)} = 2m \frac{(2)^2 \cdot 0}{v_1} dx^7 + \frac{v_1}{v_3} \frac{n}{m} dx^{11} + \frac{b_{(2)49}}{(2)^2 \cdot 0} + \frac{b_{(2)58}}{(2)^2 \cdot 0} dx^6 \quad ! \quad (6:45)$$

Identifying this with $iG_{(1)}$ gives the constraint

$$(n=m) dx^{11} = (b_{(2)49} + iv_3) = (2)^2 \cdot 0 \quad (6:46a)$$

This constraint allows us to simplify Eq. (6:42a) to

$$1=2 = (b_{(2)58} + iv_1) = (2)^2 \cdot 0 \quad (6:46b)$$

The tilted three-form flux is then

$$\mathcal{G}_{(3)} = (2)^2 \cdot 0 m = (v_1 \cdot 2) v_1^5 \wedge e^8 \wedge v_3 dx^4 \wedge dx^9 \wedge (dx^6 + 2 dx^7) \quad (6:47)$$

The first factor is (1,1) and primitive, and the second is dz^2 . So, $\mathcal{G}_{(3)}$ is (2,1) and primitive, and no further constraints arise from condition (4:42b).

From the kinetic terms (3:54), the massless RR scalars are the uncharged fields

$$e_{(2)29}; e_{(2)89}; e_{(4)5789} + (2)^2 \cdot 0 e_{(2)29}; e_{(4)5689}; e_{(4)5689}; \text{ and } e_{(4)4678} \quad (6:48)$$

The massless gauge bosons are those that do not couple to RR scalars. From (3:54), they are the three linear combinations of v^4 , $b_{(2)9}$, $c_{(2)5}$, and $e_{(4)459}$ orthogonal to

$$(m=n)b_{(2)9} \quad c_{(2)5} \quad (6:49)$$

where orthogonality is defined by the metric appearing in the gauge kinetic terms. (See Sec. 6.4 for further discussion in the context of the dual O3 orientifold).

Finally, there are $6M$ massless scalars A_I , I^m and M massless gauge bosons A_I from the D5 branes.

In summary, as in the previous two examples, the massless bosonic fields are one graviton, $3 + M$ vectors, and $16 + 6M$ moduli. However, in this case the moduli can be taken to be the independent fields

$$\begin{aligned} & 1, 2, 3, v_2, \text{dil}, a^8_6, a^8_7, b_{(2)5}, \\ & 6 e_{(p)} \text{ scalars, and } 6M \text{ D5 scalars,} \end{aligned}$$

with 1 and 3 pure imaginary. These fields combine to form one 4D $N = 2$ gravity multiplet, $2 + M$ vector multiplets, and $3 + M$ hypermultiplets.

The amount of unbroken supersymmetry is

$$N = 2; \tag{6:50}$$

due to the existence of an antiholomorphic involution on 2 compatible with the moduli constraints,

$$2;_{11} ! \quad 2;_{(11)} ; \quad \text{where } \quad_{11} = (b_{(2)58} + iv_1) = (2)^2 \quad^0 : \tag{6:51}$$

In contrast to the previous two examples, there is no decomposition of the $N = 2$ algebra into N_+ and N_- algebras generated by 6D spinors of definite ber chirality. For any $N = 1$ subalgebra of the $N = 2$, the corresponding 6D spinor \wedge contains components of both positive and negative ber chirality (cf. Sec. 4.4).

Nonintegrability of the Almost Complex Structure

In the example just discussed, the ACS selected by the supersymmetry conditions is nonintegrable. The simplest way to verify this is via the torsion classes W_i of Eq. (1.5). This is the one place in the paper in which we will find it useful to compute any of the W_i . The criterion that we will use is [45,32,29]:

$$\text{The ACS is integrable if and only if } W_1 = W_2 = 0:$$

In the example, the ACS is such that the three $(1,0)$ -forms of X_6 are the three factors appearing in Eq. (6:39a). Consequently, the three terms in J of Eq. (6:39b) are each of type $(1,1)$, and we have the following Hodge decomposition of dJ :

$$dJ_{(2;1)+(1;2)} = dZ \wedge \frac{1}{2} Z^{1=2} v_1^5 \wedge e^8 + v_2 dx^6 \wedge dx^7 + \frac{1}{2} Z^{1=2} v_3 dx^4 \wedge dx^9 ; \tag{6:52a}$$

$$dJ_{(3;0)+(0;3)} = 2nZ^{1=2} v_1 dx^6 \wedge dx^9 \wedge e^8 : \tag{6:52b}$$

From the second equation, $W_1 \neq 0$. It is straightforward to compute

$$W_1 = \frac{n_2}{3Z} \frac{v_1 = (v_2 v_3)^{1=2}}{\text{Im}_1 \text{Im}_2 \text{Im}_3} : \quad (6.53)$$

Therefore, the ACS is nonintegrable.

Note, however, that there is no topological obstruction to defining an integrable complex structure. For example, the fibration (6.34b) is compatible with the complex structure

$$w^1 = x^4 + ix^5; \quad w^2 = x^6 + ix^9; \quad \text{and} \quad w^3 = x^7 + ix^8 : \quad (6.54)$$

In this complex structure, $F^{w^1} = 2ndw^2 \wedge dw^2$, with no (0,2) component, so that the fibration is indeed holomorphic. This can also be seen from the fact that X_6 is just a warped version of one of the complex nilmanifolds classified in Ref. [76]. On the other hand, the complex structure (6.54) is not the one selected by the supersymmetry conditions, and is incompatible with the physical metric (6.36) except at certain points in moduli space.

6.4. Dual $O(3)$ orientifold

Consider the $O(3)$ orientifold with internal T^6 and flux [11]

$$F_{(3)}^0 = (2^-)^2 \cdot 0 = 2m \, dx^4 \wedge dx^6 + dx^5 \wedge dx^7 \wedge dx^9 \quad (6.55a)$$

$$H_{(3)}^0 = (2^-)^2 \cdot 0 = 2n \, dx^4 \wedge dx^6 + dx^5 \wedge dx^7 \wedge dx^8 : \quad (6.55b)$$

The Gauss's law constraint (5.1e) is Eq. (6.3). The complex flux $G_{(3)}^0$ is

$$\frac{1}{(2^-)^2} G_{(3)}^0 = 2n_{\text{dil}} \cdot 0 \, dx^4 \wedge dx^6 + dx^5 \wedge dx^7 \wedge dx^8 \quad (m=n) \, (1 = \text{dil}) \cdot 0 \, dx^9 : \quad (6.56)$$

If we parametrize the complex structure as in Eq. (5.8), then the supersymmetry conditions imply a factorization $T^6 \rightarrow T_{f4567g}^4 \times T_{f89g}^2$ with respect to both complex and Kähler structure. The condition that $G_{(3)}^0$ be of type (2,1) implies

$$0 = \frac{0}{T^4} \cdot \frac{0}{T^2}; \quad \text{with} \quad \det \frac{0}{T^4} = 1; \quad \text{dil} \frac{0}{T^2} = m=n; \quad (6.57)$$

where $\frac{0}{T^4}$ is a complex 2×2 matrix and $\frac{0}{T^2}$ is a complex number. The condition that $G_{(3)}^0$ be primitive then gives

$$J^0 = J_{T^4}^0 + J_{T^2}^0 :$$

Working in terms of g_{ij}^0 and g_{ij}^1 has been convenient thus far for deriving the T^6 ! T^4 T^2 factorization, but is bad for describing moduli. As discussed in Sec. 5.1 and in App. E, the parametrization is redundant. For this reason, we will adopt a different parametrization shortly. First, however, note that the supersymmetry constraints on the T^4 factor can be phrased in a parametrization independent way as follows:

$$dx^4 \wedge dx^6 + dx^5 \wedge dx^7 \quad (1,1) \text{ and primitive on } T^4. \quad (6:58)$$

Starting from this condition, it is not hard to show that there is $N = 2$ supersymmetry. For example, at $g_{mn}^{0(T^4)} = \delta_{mn}$, there is an S^2 of complex structures such that the condition is satisfied. The corresponding Kähler forms and holomorphic $(2,0)$ forms are

$$J_{T^4}^0 = n^A J^{(A)}; \quad \hat{J}_{T^4}^0 = n^A \hat{J}^{(A)}; \quad (6:59)$$

where $n = (n^1; n^2; n^3)$ is a unit vector on S^2 , and

$$J^{(1)} = dx^4 \wedge dx^5 + dx^6 \wedge dx^7; \quad J^{(2)} = dx^4 \wedge dx^7 + dx^5 \wedge dx^6; \quad J^{(3)} = dx^4 \wedge dx^6 - dx^5 \wedge dx^7; \\ J^{(1)} = J^{(2)} + iJ^{(3)}; \quad J^{(2)} = J^{(1)} - iJ^{(3)}; \quad J^{(3)} = \text{cyc. perms.} \quad (6:60)$$

The S^2 of complex structures defines a single hyperKähler structure, and corresponds to two 6D negative chirality Weyl spinors $\hat{\chi}_{1;2}^0$ via

$$J^{0(A)}_{ab} = i \hat{\chi}_a^{0y} \hat{\chi}_b^{(A)}; \quad \hat{\chi}^0 = \begin{pmatrix} \hat{\chi}_1^0 \\ \hat{\chi}_2^0 \end{pmatrix}; \quad (6:61)$$

Here, the primes indicate that we are discussing the $O(3)$ orientifold, and the hats indicate that $\hat{\chi}^0$ is a normalized spinor, rescaled relative to the χ appearing in Eq. (4.12). One can also see the $N = 2$ supersymmetry from the fact that there exists exactly one antiholomorphic involution of X_6^0 that preserves the moduli constraints on g_{ij}^0 and g_{dil}^0 ,

$$\hat{\chi}_{T^4}^0; \hat{\chi}_{T^2}^0 \rightarrow -\hat{\chi}_{T^4}^0; \hat{\chi}_{T^2}^0; \quad (6:62)$$

This involution implies that for every Killing spinor such that $\hat{\chi}_1^0 = \hat{\chi}_{T^4}^0 - \hat{\chi}_{T^2}^0$, there is an independent Killing spinor such that $\hat{\chi}_2^0 = \hat{\chi}_{T^4}^0 + \hat{\chi}_{T^2}^0$. The condition $\text{Im} \int_{T^2} \hat{\chi}_2^0 = (m=n) \text{Im} \int_{dil} \hat{\chi}_2^0 > 0$, fixes the chirality of $\hat{\chi}_{T^2}^0$ to be positive. Since $\hat{\chi}_1^0$ has negative chirality by definition, this means that $\hat{\chi}_{T^4}^0$ also has negative chirality. On T^4 , there are exactly two negative chirality spinors, complex conjugate to one another up to an overall

phase that can be absorbed into the definition of the spinors. Therefore, once we know that this example preserves any supersymmetry, we know that it preserves $N = 2$.

A nonredundant parametrization of the metric moduli is obtained by writing the T^4 as a fibration over T^2_{f67g} :

$$ds_{T^4}^2 = \frac{V_1^0}{\text{Im } \tau_1^0} j^{\alpha 4} + \tau_1^{\alpha 5} j^{\beta} + \frac{V_2^0}{\text{Im } \tau_2^0} j dx^6 + \tau_2^0 dx^7 j^{\beta}; \quad (6:63a)$$

$$ds_{T^2}^2 = \frac{V_3^0}{\text{Im } \tau_3^0} j dx^8 + \tau_3^0 dx^9 j^{\beta}; \quad (6:63b)$$

where

$$\tau_1^{\alpha 4} = dx^4 + a^{\alpha 4}_6 dx^6 + a^{\alpha 4}_7 dx^7; \quad (6:64a)$$

$$\tau_1^{\alpha 5} = dx^5 + a^{\alpha 5}_6 dx^6 + a^{\alpha 5}_7 dx^7; \quad (6:64b)$$

with $a^{\alpha m}_n$ constant on T^6 . That is, $a^{\alpha m}$ defines a trivial fibration $F^{\alpha m} = da^{\alpha m} = 0$ over the base $T^2_{f6;7g}$. The natural holomorphic one-forms associated with this parametrization are

$$\omega^1 = \tau_1^{\alpha 4} + \tau_1^{\alpha 5}; \quad dz^2 = dx^6 + \tau_2^0 dx^7; \quad \text{and} \quad dz^3 = dx^8 + \tau_3^0 dx^9; \quad (6:65)$$

Therefore, the (2,1) condition on $G_{(3)}$ becomes

$$\tau_1^0 \tau_2^0 = 1; \quad (6:66a)$$

$$\tau_3^0 \text{dil}^0 = m = n; \quad (6:66b)$$

and the primitivity condition becomes

$$a^{\alpha 4}_7 = a^{\alpha 5}_6; \quad (6:67)$$

From the kinetic terms (5.15), the massless $c_{(4)abcd}$ scalars are the uncharged scalars

$$c_{(4)6789}^0; \quad c_{(4)5689}^0; \quad c_{(4)4789}^0; \quad c_{(4)4589}^0; \quad c_{(4)4689}^0; \quad c_{(4)5789}^0; \quad \text{and} \quad c_{(4)4567}^0; \quad (6:68)$$

The massless gauge bosons are those that do not couple to RR scalars. From (5.15), they are the three linear combinations of $b_{(2)8}^0$, $b_{(2)9}^0$, $c_{(2)8}^0$, and $c_{(2)9}^0$ orthogonal to

$$(m = n)b_{(2)8}^0 + c_{(2)9}^0; \quad (6:69)$$

Here, orthogonality is defined by the metric appearing in the gauge kinetic terms. In terms of the complex gauge bosons (5.16), the metric on the 89 subspace is proportional to

$$\frac{1}{2} \begin{pmatrix} 0 & 1 & d^0 \\ 1 & 0 & d^0 \end{pmatrix} \quad \frac{1}{2} \begin{pmatrix} 0 & 1 & z^3 \\ 1 & 0 & z^3 \end{pmatrix} ; \quad (6:70)$$

The massive linear combination (6.69) is proportional to $\text{Re}(1 = d^0) d_{z^3}^0$.²⁹ Therefore, the 3D space of massless gauge bosons is spanned by

$$\text{Im}(1 = d^0) d_{z^3}^0 ; \quad \text{Re} d_{z^3}^0 ; \quad \text{and} \quad \text{Im} d_{z^3}^0 ; \quad (6:71)$$

Finally, there are $6M$ massless scalars \mathbb{I}^m and M massless gauge bosons $A_{\mathbb{I}}^0$ from the D3 branes.

In summary, the massless bosonic fields are one graviton, $3 + M$ vectors, and $16 + M$ moduli. The moduli consist of

$$V_1^0; V_2^0; V_3^0, 2 \text{ independent } \mathbb{I}^0, 3 \text{ independent } a_{\mathbb{I}^n}^m, \\ 6 c_{(4)}^0 \text{ scalars, and } 6M \text{ D3 scalars.}$$

From the supersymmetry-breaking mass spectra given in Ref. [17], this corresponds to $N = 4 ! N = 2$ with $m_1 = m_2$ for the two massive gravitini. (This is a third way to see that the background preserves $N = 2$ supersymmetry). The moduli space is $M = M_{\mathbb{H}} \times M_{\mathbb{V}}$, where in the approximation that the warp factor is set to unity,

$$M_{\mathbb{H}} = \frac{SO(4; 3 + M)}{SO(4) \times SO(3 + M)} ; \quad (6:72a)$$

$$M_{\mathbb{V}} = \frac{SO(2; 1 + M)}{SO(2) \times SO(1 + M)} \times \frac{SU(1; 1)}{U(1)} ; \quad (6:72b)$$

up to discrete identifications. See App. G for a discussion of the metric on moduli space.

6.5. T-Duality Map

The T-duality map relating the O3 orientifold of Sec. 6.4 to the O5 orientifolds of Secs. 6.1{6.3 is exactly as described in Sec. 5.2, once we perform the following relabelings of coordinates in the O3 orientifold:

$$\text{Sec. 6.1: } x^4; x^5; x^6; x^7; x^8; x^9 \xrightarrow{\text{Sec. 6.4}} x^6; x^7; x^8; x^9; x^4; x^5_{\text{new}} ; \quad (6:73a)$$

$$\text{Sec. 6.2: no relabeling,} \quad (6:73b)$$

$$\text{Sec. 6.3: } x^4; x^5; x^6; x^7; x^8; x^9 \xrightarrow{\text{Sec. 6.4}} x^5; x^8; x^6; x^7; x^9; x^4_{\text{new}} ; \quad (6:73c)$$

²⁹ Note that from $dz^3 = dx^8 + d^0 dx^9$, we have $d_{z^3}^0 = (d_9^0 \quad d_8^0) = (d^0 \quad d^0)$.

Note that in all three cases, the relabeling is a permutation of positive Jacobian. Therefore, the image of the ISD flux (6.55) is again ISD, rather than IA SD. In terms of the $T^6 \rightarrow T^4 \times T^2$ factorization of Sec. 6.4, the three relabelings correspond to T-dualizing along the T^2 factor, along a T^2 in the T^4 factor, and along $S^1 \times T^4$ times $S^1 \times T^2$, respectively.

It is straightforward to follow all moduli and gauge fields through the T-duality map, using the relations in Sec. 5.2. The constraints (6.66a;b) and (6.67) map to (6.13b;a) and (6.14) in Sec. 6.1, to (6.23a;b) and $b_{(2)56} = b_{(2)47}$ in Sec. 6.2, and to (6.46b;a) and (6.42b) in Sec. 6.3. The volumes $V_1^0; V_2^0; V_3^0$ are equal to $V_2; V_3; (2)^4 \Omega = V_b$ in Sec. 6.1, to $(2)^4 \Omega = V_b; V_2; V_3$ in Sec. 6.2, and to $(2)^2 \Omega_{j_1 j_2}; V_2; (2)^2 \Omega_{j_3 j_4}$ in Sec. 6.3. The RR moduli (6.68) map to (6.16), (6.28), and (6.48). The massive gauge bosons (6.69) map to (6.17), (6.29), and (6.49).

7. $N = 3$ Example: The Warped Iwasawa Manifold

7.1. Example 4: $N_+ = 1, N_- = 2$

Consider the choice of fibration curvature and background flux

$$F^4 = 2h dx^6 \wedge dx^8 - m dx^6 \wedge dx^9 - m dx^7 \wedge dx^8 + (m^2 - n) dx^7 \wedge dx^9; \quad (7.1b)$$

$$F^5 = 2h dx^6 \wedge dx^9 + dx^7 \wedge dx^8 - m dx^7 \wedge dx^9; \quad (7.1a)$$

$$F_{(3)}^{bg1} = (2)^2 \Omega = 2f dx^4 \wedge dx^6 \wedge dx^8 - n dx^7 \wedge dx^9 + h dx^5 \wedge n dx^6 \wedge dx^9 - n dx^7 \wedge dx^8 + m n dx^7 \wedge dx^9; \quad (7.1c)$$

with $F_{(1)} = H_{(3)} = 0$. Here, $f; h; n = 1; 2$ and $m = 0; 1$. This choice gives a contribution to the Gauss's law constraint,

$$N_{ux} = 4fh - 4n - m^2; \quad (7.2)$$

so that Eq. (3.15) becomes

$$2M + 4fh - 4n - m^2 = 32; \quad (7.3)$$

The condition $0 \leq N_{ux} \leq 32$ puts restrictions on the allowed combinations of $f; h; m; n$. The discussion of the dual $O3$ theory in Sec. 7.2 shows why this is in some sense a natural class of backgrounds to consider.

For the choice (7:1) to preserve $N = 3$ supersymmetry, we require that the complex (4:28d) be (2,1) and primitive with respect to three independent complex structures. This is equivalent to demanding that $G_{(3)}$ be of type (0,3) with respect a single complex structure, which is then distinct from the previous three [60]. (See Sec. 7.2 for further discussion).

It is convenient to parametrize the metric and almost complex structure as in Ex. 1:

$$w^1 = z^4 + w^1 z^5; \quad (7:4)$$

$$dw^i = dz^{2i+2} + w^i_j dz^{2j+3}; \quad i, j = 2, 3:$$

Here, we have used the symbols w^i instead of z^i , reserving the latter for complex coordinates in which $G_{(3)}$ is of type (2,1) and primitive.

The condition that $G_{(3)}$ be of type (0,3) results in the following constraints:

$$w^i_j = \delta^i_j; \quad \text{where} \quad (h=f)_{dil} = w^1 = \rho; \quad \text{and} \quad (7:5a)$$

$$2 + m + n = 0; \quad \text{or equivalently,} \quad \rho = \frac{1}{2} \sqrt{m + i 4n - m^2}; \quad (7:5b)$$

Here, we have chosen the root ρ such that $\text{Im} \rho > 0$, for agreement of Eq. (7:5a) with $\text{Im} \rho_{dil} = 1 = g_s > 0$. When Eqs. (7:5a;b) are satisfied,

$$\frac{1}{(2\rho)^2} G_{(3)} = 2f w^1 \wedge dw^2 \wedge dw^3; \quad (7:6)$$

which is indeed of type (0,3). Since this statement is independent of the Kähler moduli, the unfluxed metric moduli are the real Kähler moduli V_b and $g_{B w^i w^j}$ ($i, j = 2, 3$), with $g_{B w^i w^j} = g_{B w^j w^i}$. The constraints (7:5a;b) also imply that

$$F^{w^1} = 2hdw^2 \wedge dw^3; \quad (7:7)$$

where $F^{w^1} = F^4 + w^1 F^5$.

Now let us return to the supersymmetry conditions as originally formulated. The three independent complex structures satisfying conditions (4:28a;b) are related to the one just described via

$$\text{complex structure 1:} \quad z^1; dz^2; dz^3 = w^1; dw^2; dw^3; \quad (7:8a)$$

$$\text{complex structure 2:} \quad z^1; dz^2; dz^3 = w^1; dw^2; dw^3; \quad (7:8b)$$

$$\text{complex structure 3:} \quad z^1; dz^2; dz^3 = w^1; dw^2; dw^3; \quad (7:8c)$$

In these three complex structures,

$$F^{z^1} = 2hdz^2 \wedge dz^3; \quad F^{z^2} = 2hdz^2 \wedge dz^3; \quad \text{and} \quad F^{z^3} = 2hdz^2 \wedge dz^3; \quad (7:9a;b;c)$$

respectively. In all three cases, F^{z^i} has no $(0,2)$ component. Therefore, the corresponding foliations are holomorphic, and the complex structures are integrable, in agreement with Sec. 4.2. In contrast, the almost complex structure of the w coordinates is not integrable.

Specialize to complex structure 1. Then we can write Eq. (7:9a) as

$$d^{z^1} = N dz^2 \wedge dz^3; \quad \text{where} \quad N = 2h: \quad (7:10)$$

This is the familiar relation between the three left-invariant one-forms

$$z^1 = z^1 + N z^2 dz^3; \quad dz^2; \quad \text{and} \quad dz^3;$$

on the Iwasawa manifold.

The Iwasawa manifold can be defined as a coset of upper triangular matrices as follows.³⁰ Consider C^3 , presented as the space of upper triangular 3×3 matrices with ones along the diagonal,

$$g_N(z^1; z^2; z^3) = \begin{pmatrix} 0 & & & 1 \\ 1 & z^3 & & z^1 - N \\ 0 & 1 & & z^2 \\ 0 & 0 & & 1 \end{pmatrix} A; \quad \text{where} \quad z^1; z^2; z^3 \in C: \quad (7:11)$$

Here, N is a fixed positive integer. Let $G_3^N(C)$ denote the corresponding group under matrix multiplication. For any N , this group is isomorphic to the complex three-dimensional Heisenberg group $H_3(C)$. We can also define a discrete subgroup $G_3^N(\mathbb{Z})$ containing the matrices

$$g_N(a; b; c) = \begin{pmatrix} 0 & & & 1 \\ 1 & c & & a - N \\ 0 & 1 & & b \\ 0 & 0 & & 1 \end{pmatrix} A; \quad \text{where} \quad a; b; c \in \mathbb{Z}; \quad (7:12)$$

with $\mathbb{Z} = \mathbb{Z} + \mathbb{Z} \in C$ a lattice parametrized by the complex modulus \mathbb{Z} . This subgroup has a natural action on $G_3^N(C)$ by matrix multiplication. Consider the right-coset $M^N = G_3^N(C) = G_3^N(\mathbb{Z})$. The resulting identifications are

$$(z^1; z^2; z^3) = (z^1 + a - Nbz^3; z^2 + b; z^3 + c): \quad (7:13)$$

³⁰ This description is taken directly from the twisted T^3 example in Sec. 2.2 of Ref. [4], with R^3 replaced by C^3 .

This quotient defines the N th Iwasawa manifold.

The standard metric and Kahler form on the Iwasawa manifold are

$$ds_{\text{Iwasawa}}^2 = |z^1|^2 + |dz^2|^2 + |dz^3|^2; \quad (7.14a)$$

$$J_{\text{Iwasawa}} = \frac{i}{2} (z^1 \wedge \bar{z}^1 + dz^2 \wedge d\bar{z}^2 + dz^3 \wedge d\bar{z}^3); \quad (7.14b)$$

However, in the orientifold example that we are considering, the metric and Kahler form are given by³¹

$$ds_6^2 = \frac{V_b}{\text{Im}} |z^1|^2 + 2Z g_{B_{z^i z^j}} dz^i dz^j; \quad (7.15a)$$

$$J = \frac{iV_b}{2 \text{Im}} (z^1 \wedge \bar{z}^1 + iZ g_{B_{z^i z^j}} dz^i \wedge d\bar{z}^j); \quad (7.15b)$$

a generalization of (7.14a;b) that includes arbitrary fiber and base Kahler structure, and also the warp factor Z .

Now let us turn to the NSB-eld moduli. The constraints that follow from Eq. (3.52a) are

$$\begin{aligned} m b_{(2)46} + b_{(2)47} + b_{(2)56} &= 0; & n b_{(2)46} - b_{(2)57} &= 0; \\ m b_{(2)48} + b_{(2)49} + b_{(2)58} &= 0; & n b_{(2)48} - b_{(2)59} &= 0; \end{aligned}$$

or equivalently,

$$b_{(2)w^1 w^i} = b_{(2)w^i w^1} = 0; \quad i = 2;3; \quad (7.16)$$

Eq. (3.52b) gives the same constraints. The unlifted NSB-eld moduli are the orthogonal components

$$b_{(2)w^1 w^i}; \quad b_{(2)w^i w^1}; \quad i = 2;3; \quad (7.17)$$

From the kinetic terms (3.54a;b;c), the massless RR sector scalars are the nine uncharged elds

$$\begin{aligned} e_{(2)67}; \quad e_{(2)89}; \quad e_{(2)69} - e_{(2)78}; \quad e_{(2)67} + e_{(2)456789}; \\ n e_{(2)68} + (m=2) e_{(2)69} + e_{(2)78} + e_{(2)79}; \\ e_{(4)5678} - e_{(4)4679}; \quad e_{(4)4789} - e_{(4)5689}; \\ n e_{(4)4678} + (m=2) e_{(4)4679} + e_{(4)5678} + e_{(4)5679}; \\ \text{and } n e_{(4)4689} + (m=2) e_{(4)4789} + e_{(4)5689} + e_{(4)5789}; \end{aligned} \quad (7.18)$$

³¹ In the conventions of this paper, $ds^2 = g_{ij} dz^i dz^j + g_{\bar{i}\bar{j}} d\bar{z}^i d\bar{z}^j = 2g_{i\bar{j}} dz^i d\bar{z}^j$ (cf. App. A).

The massive gauge bosons are those that couple to the other RR scalars. From (3.54a;b;c), they lie in the 6D space spanned by

$$\begin{aligned}
(2)^2 \text{ }^0(f=h)v^4 & \quad q_{(2)4} \text{ }^0 ; \quad (2)^2 \text{ }^0(f=h)v^5 & \quad m q_{(2)4} \text{ }^0 \quad q_{(2)5} \text{ }^0 ; \\
(2)^2 \text{ }^0(f=h)b_{(2)7} & \quad e_{(4)456} \text{ }^0 ; \quad (2)^2 \text{ }^0(f=h)b_{(2)6} & \quad + m e_{(4)456} \text{ }^0 + e_{(4)457} \text{ }^0 ; \quad (7.19) \\
(2)^2 \text{ }^0(f=h)b_{(2)9} & \quad e_{(4)458} \text{ }^0 ; \quad (2)^2 \text{ }^0(f=h)b_{(2)8} & \quad + m e_{(4)458} \text{ }^0 + e_{(4)459} \text{ }^0 :
\end{aligned}$$

The massless gauge bosons span the orthogonal 6D space, where orthogonality is defined with respect to the metric appearing in the gauge kinetic terms. (See Sec. 7.2 for further discussion in the dual $O3$ orientifold).

Finally, there are $6M$ massless scalars A_I , I^m , and M massless gauge bosons A_I from the D5 branes.

In summary, the massless bosonic fields are one graviton, $6+M$ vectors, and $18+6M$ moduli. The moduli consist of

$$\begin{aligned}
V_b, 4 g_{Bz^i z^j}, 4 \text{ indep } b_{(2)m}, \\
9 e_{(p)} \text{ scalars, and } 6M \text{ D5 scalars.}
\end{aligned}$$

These fields combine to form one 4D $N = 3$ gravity multiplet, and $3+M$ vector multiplets.

The amount of unbroken supersymmetry is

$$N_+ = 1; \quad N_- = 2; \quad (7.20)$$

in the notation of Sec. 4.2. The $N_+ = 1$ supersymmetry corresponds to complex structure 1 (7.8a), with the modulus $\tau_1 = \text{Im } \tau_1 > 0$ appearing in z^1 (cf. Eq. (4.26a)). The $N_- = 2$ supersymmetries correspond to complex structures 2 and 3 (7.8b;c), with the modulus $\tau_1 = \text{Im } \tau_1 < 0$ appearing in z^1 .

7.2. Dual $O3$ orientifold

For the $O3$ orientifold with internal T^6 , choices of flux preserving 4D $N = 3$ supersymmetry were first discussed in Ref. [60]. The requirement for $N = 3$ supersymmetry is

$$G_{(3)}^0 \text{ is } (2,1) \text{ and primitive w.r.t. to three independent complex structures.} \quad (7.21)$$

In contrast, from the equations of motion alone, without imposing any supersymmetry conditions, one obtains the condition that the flux be ISD: ${}_{(3)}G^0 = iG_{(3)}^0$. As has already

been mentioned in Footnote 21, the space of ISD forms includes not only primitive (2,1) forms, but also nonprimitive (1,2) forms $J \wedge !^{(0;1)}$, where $!^{(0;1)}$ is a (0,1) form, as well as (0,3) forms. The condition (7.21) is equivalent to [60]

$$G_{(3)}^0 \quad (0,3) \text{ for some choice of complex structure.} \quad (7.22)$$

The complex structure in this last condition is a fourth complex structure, independent of the three complex structures of the previous condition.³²

To construct a large class of $N = 3$ backgrounds, let us write

$$\frac{1}{(2\ell)^2} G_{(3)}^0 = 2 \, dw^1 \wedge dw^2 \wedge dw^3; \quad (7.23)$$

where ℓ is a positive real number, and where the T^6 factorizes as $T^2 \times T^2 \times T^2$ with respect to complex structure in the following way:

$$w^i = x^i + i y^i; \quad i = 1;2;3; \quad (7.24a)$$

$$(h=f)_{\text{dil}} = 0; \quad f; h \in \mathbb{R}; \quad (7.24b)$$

The coordinates $x^i; y^i$ are related to those used elsewhere in this paper by

$$(x^1; y^1; x^2; y^2; x^3; y^3)_{\text{here}} = (x^5; x^4; x^6; x^7; x^8; x^9)_{\text{rest of paper}}; \quad (7.25)$$

We adopt this notation here and similar notation in Sec. 8.2 since it allows equations involving cyclic permutation of the $x^i; y^i$ to be written succinctly.

When expanded in real coordinates, the complex flux becomes

$$\begin{aligned} \frac{1}{(2\ell)^2} G_{(3)}^0 = & 2 \, dx^1 \wedge dx^2 \wedge dx^3 + i \, dx^1 \wedge dx^2 \wedge dy^3 + \text{cyc. perm s. of 123} \\ & + i^2 \, dx^1 \wedge dy^2 \wedge dy^3 + \text{cyc. perm s. of 123} + i^3 \, dy^1 \wedge dy^2 \wedge dy^3; \end{aligned} \quad (7.26)$$

³² In fact, we could have also rephrased the supersymmetry conditions for the $N = 2$ examples. In each example of Sec. 6, there exists a complex structure such that $G_{(3)}$, $\mathcal{E}_{(3)}$, or $G_{(3)}^0$ (whichever is appropriate) is of the form $J \wedge !^{(0;1)}$. However, unlike the $N = 4 \neq N = 3$ case, where the conditions (7.21) and (7.22) are truly equivalent, in the $N = 4 \neq N = 2$ case, this rephrasing of the supersymmetry conditions seems to involve the assumption that $m_1 = m_2$ for the two massive gravitini.

If, in addition, we assume as part of our ansatz that τ^0 satisfies the quadratic equation

$$P(\tau^0) = l\tau^2 + m\tau + n = 0; \quad l, m, n \in \mathbb{Z}; \quad l > 0; \quad (7.27)$$

then Eq. (7.26) reduces to an expression linear in τ^0 . From Eqs. (7.24a;b), we can then read off the RR and NS fluxes. It is possible to show from the quantization condition (5.4) that by redefinitions of l, m, n , and τ^0 , we can take $l = 1$. Then,

$$\begin{aligned} \frac{1}{(2\pi)^2} F_{(3)}^0 &= 2f \, l dx^1 \wedge dx^2 \wedge dx^3 - n dx^1 \wedge dy^2 \wedge dy^3 + \text{cyc. perm s. of } 123 \\ &+ (m - n) dy^1 \wedge dy^2 \wedge dy^3; \end{aligned} \quad (7.28a)$$

$$\begin{aligned} \frac{1}{(2\pi)^2} H_{(3)}^0 &= 2h \, l dx^1 \wedge dx^2 \wedge dy^3 + \text{cyc. perm s. of } 123 \\ &- m dx^1 \wedge dy^2 \wedge dy^3 + \text{cyc. perm s. of } 123 + (m^2 - 1 - n) dy^1 \wedge dy^2 \wedge dy^3; \end{aligned} \quad (7.28b)$$

The contribution to the Gauss's law constraint (3.15) from this choice of flux is

$$N_{ux} = 4fh(4ln - m^2); \quad (7.29)$$

Not all choices of $(l; m; n)$ are inequivalent. From change of lattice basis on the T^6 , there is an $SL(2; \mathbb{Z})$ identification of the complex structure modulus τ^0 . Modulo identifications, we can assume that τ^0 lies in the fundamental domain of $SL(2; \mathbb{C})$,

$$0 < \text{Re } \tau^0 < \frac{1}{2}; \quad \text{Im } \tau^0 > 0; \quad (7.30)$$

From Eq. (7.27), we have

$$\tau^0 = \frac{1}{2l} \left(m + i \sqrt{4ln - m^2} \right); \quad j^0_j = \frac{p}{n-l}; \quad (7.31)$$

so the condition (7.30) becomes

$$n > 1 - |m| \quad \text{and} \quad 4ln > m^2; \quad (7.32)$$

The possible values of $(f; h; l; m; n)$ are strongly constrained by $0 < N_{ux} < 32$ and the fact that $m^2 \equiv 0; 1 \pmod{4}$. For $(f; h) = (1; 1)$, in addition to the $N_{ux} = 4$ solution without flux, there are just four choices of $(l; m; n)$ satisfying the inequalities (7.32):

$$\begin{aligned} (l; m; n) &= (1; 1; 1); & N_{ux} &= 12; & 2M &= 20; & \tau^0 &= e^{2i\pi/3}; \\ (l; m; n) &= (1; 0; 1); & N_{ux} &= 16; & 2M &= 16; & \tau^0 &= i; \\ (l; m; n) &= (1; 1; 2); & N_{ux} &= 28; & 2M &= 4; & \tau^0 &= \frac{1}{2} \left(1 + i\sqrt{7} \right); \\ (l; m; n) &= (1; 0; 2); & N_{ux} &= 32; & 2M &= 0; & \tau^0 &= i\sqrt{2}; \end{aligned} \quad (7.33a)$$

Allowing arbitrary values of $(f;h)$, there are four additional choices. For $(f;h) = (2;1)$ or $(1,2)$, we can take

$$\begin{aligned} (l;m;n) &= (1;1;1); & N_{ux} &= 24; & 2M &= 8; & \phi^0 &= e^{2i\pi/3}; \\ (l;m;n) &= (1;0;1); & N_{ux} &= 32; & 2M &= 0; & \phi^0 &= i; \end{aligned} \quad (7:33b)$$

Note that the ux (7:28a;b) is correctly quantized with $F_{(3)}^0; H_{(3)}^0 \in (2,2)^0 \oplus H^3(T^6; 2Z)$ in all of four cases.

If the complex structure and axion-dilaton are deformed from $i_j = \phi^0 i_j$ and $(h=f)_{dil}^0 = \phi^0$, with ϕ^0 given by the values in (7:33a;b), then the complex ux is no longer of type $(0,3)$. On the other hand, the ux is still $(0,3)$ for arbitrary choice of Kähler moduli. So, the metric moduli are the nine real degrees of freedom $g_{w^i w^j}^0$.

From the kinetic terms (5.15), the space of massless RR scalars is spanned by the nine uncharged scalars

$$\begin{aligned} & n c_{(4)x^1 y^1 x^2 x^3}^0 + c_{(4)x^1 y^1 y^2 y^3}^0 + (m=2) c_{(4)x^1 y^1 x^2 y^3}^0 + c_{(4)x^1 y^1 y^2 x^3}^0; \\ & c_{(4)x^1 y^1 x^2 y^2}^0; c_{(4)x^1 y^1 x^2 y^3}^0 \quad c_{(4)x^1 y^1 y^2 x^3}^0; \text{ and cyc. perm s. of } 123: \end{aligned} \quad (7:34)$$

This is equivalent to the space spanned by the RR scalars of Hodge type $(2,2)$,

$$c_{(4)w^i w^j w^k w^l}^0; \quad i; j; k; l = 1; 2; 3;$$

The massive gauge bosons are those that couple to the other RR scalars. From (5.15), they lie in the 6D space spanned by

$$(f=h) n b_{(2)x^i}^0 \quad m c_{(2)x^i}^0 \quad c_{(2)y^i}^0; \quad \text{and} \quad (f=h) b_{(2)y^i}^0 + c_{(2)x^i}^0; \quad i = 1; 2; 3: \quad (7:35)$$

In terms of the complex gauge bosons (5.16), this is the space spanned by $d_{w^i}^0$, for $i = 1; 2; 3$. In fact, we can see directly from (5.17) that $c_{(4)w^1 w^2 w^3 w^i}^0$ is the axion eaten by $d_{w^i}^0$ [60]. The massless bosons are the linear combinations orthogonal to this space, where orthogonality is defined with respect to the metric appearing in the gauge kinetic terms. In the $d_{w^i}^0; d_{w^i}^0; d_{w^i}^0; d_{w^i}^0$ basis of complex gauge bosons (5.16), the latter is proportional to

$$\begin{pmatrix} \frac{1}{2} & 0 & 1 & d^0 & 0 & g_6^{0w^i w^j} & w^i \\ 2 & 1 & 0 & d^0 & g_6^{0w^i w^j} & 0 & w^i \end{pmatrix} : \quad (7:36)$$

Therefore, the six massless gauge bosons are (the real and imaginary parts of) $d_{w^i}^0$, for $i = 1; 2; 3$ [60].

Finally, there are $6M$ massless scalars ϕ_I^0 , and M massless gauge bosons A_I^0 from the D3 branes.

In summary, the massless fields are one graviton, $6+M$ vectors, and $18+6M$ moduli. The moduli consist of

$$9 g_{B w^i w^j}, \quad 9 e_{(p)} \text{ scalars, and } 6M \text{ D3 scalars.}$$

These fields combine to form one 4D $N = 3$ gravity multiplet, and $3+M$ vector multiplets.

From Ref. [77], the moduli space of the $N = 3$ theory is completely determined by the number n_V of vector multiplets. It is the coset

$$M = \frac{U(3; n_V)}{U(3) \times U(n_V)}; \quad (7:37)$$

up to discrete identifications, where, in this example, $n_V = 3+M$. For the parametrization of the coset in terms of the above moduli, see Ref. [60].

7.3. T-Duality Map

For the O3 orientifold of Sec. 7.2, the metric is

$$ds_6^2 = 2g_{w^i w^j} dw^i dw^j; \quad (7:38)$$

with $g_{w^i w^j}$ constrained only to be Hermitian. We can alternatively write this metric in the bration form (5.18),

$$ds_6^2 = \frac{V_b^0}{\text{Im } b^0} dw^1 + a^{0w^1 2} + 2g_{B w^i w^j} dw^i dw^j; \quad (7:39)$$

where the components of a^0 are holomorphically constrained,

$$a^{0w^1} = a^{0w^1}_{w^i} dw^i; \quad \text{and} \quad a^{0w^1} = a^{0w^1}_{w^i} dw^i; \quad i=2;3; \quad (7:40a)$$

and where

$$V_b^0 = 2g_{w^1 w^1} \text{Im } b^0; \quad (7:40b)$$

$$g_{B w^i w^j} = g_{w^i w^j} - g_{w^1 w^1} a^{0w^1}_{w^i} a^{0w^1}_{w^j}; \quad i;j=2;3; \quad (7:40c)$$

Here, from Eqs. (7.24a) and (7.25), the complex coordinates in the O3 orientifold are

$$w^1 = x^5 + i x^4 \quad \text{and} \quad w^i = x^{2i+2} + i x^{2i+3}; \quad i=2;3; \quad (7:41)$$

Applying the T-duality map of Sec. 5.2, we find agreement between Secs. 7.1 and 7.2, provided that we identify $\alpha = 0$. For example, the map between complex components of a^0 and $b_{(2)}$ is

$$b_{(2)w^1} = a^{0w^1} = \begin{pmatrix} 0 & 0 \end{pmatrix} \quad \text{and} \quad b_{(2)w^1} = \bar{a}^{0w^1} = \begin{pmatrix} 0 & 0 \end{pmatrix}; \quad (7.42)$$

so, the constraints (7.16) and (7.40a) agree. Here, $w^1 = 4 + 5$ for the O5 orientifold, from which the relation between $b_{(2)w^1}; b_{(2)w^1}$ and $b_{(2)4}; b_{(2)5}$ is

$$b_{(2)w^1} = \frac{1}{2} (b_{(2)5} - b_{(2)4}) \quad \text{and} \quad b_{(2)w^1} = \frac{1}{2} (b_{(2)5} + b_{(2)4}); \quad (7.43)$$

Similarly, using the underscore notation of Sec. 5.2,

$$\begin{aligned} F_{m n}^{w^1} &= F_{m n}^4 + F_{m n}^5 = H_{(3)\underline{4m n}}^0 + {}^0H_{(3)\underline{5m n}}^0 \\ &= \begin{pmatrix} 0 & 0 \end{pmatrix} H_{(3)\underline{w^1 m n}}^0 = (h=f)G_{(3)\underline{w^1 m n}}^0; \end{aligned} \quad (7.44)$$

from which Eq. (7.7) follows.

8. $N = 1$ Example

8.1. Example 5: $N_+ = 1, N_- = 0$

Consider the choice of brane curvature and background flux

$$F^4 = 2h \, dx^6 \wedge dx^8 - dx^6 \wedge dx^9 - dx^7 \wedge dx^8 + dx^7 \wedge dx^9; \quad (8.1a)$$

$$F^5 = 2h \, dx^6 \wedge dx^8 - dx^6 \wedge dx^9 - dx^7 \wedge dx^8 + 2dx^7 \wedge dx^9; \quad (8.1b)$$

$$\frac{1}{(2)^2} F_{(3)}^{bg1} = 2f \, dx^5 \wedge dx^6 \wedge dx^8 - 2f \, dx^4 \wedge dx^7 \wedge dx^9; \quad (8.1c)$$

with $F_{(1)} = H_{(3)} = 0$. This choice gives a contribution to the Gauss's law constraint

$$N_{ux} = 12fh; \quad (8.2)$$

so that Eq. (3.15) becomes

$$2M + 12fh = 32; \quad (8.3)$$

As discussed in Sec. 3.2, N_{ux} is nonnegative. So, f and h have the same sign, and the possible values of $(f;h)$ are $(1,1)$, $(2,1)$, and $(1,2)$, corresponding to $2M = 20, 8$, and $8D5$ branes, respectively.

Define an almost complex structure as in Eq. (6.5). Then, using the decomposition (4.21), it is possible to show that the base T^4 decomposes as $T^4 \simeq T^2 \times T^2$ with respect to complex structure:

$$e^i_j = \delta^i_j; \quad i, j = 2, 3; \quad (8.4)$$

where

$$e^1_1 = e^2_2 = e^3_3 = e^4_4 = (h=f)_{dil} = e^{2i\alpha} : \quad (8.5)$$

When these constraints are satisfied, the complex three-form flux (4.28d) is

$$G_{(3)} = \frac{1}{2} \frac{P}{\text{Im} \tau} (z^1 \wedge dz^2 \wedge dz^3 + \bar{z}^1 \wedge dz^2 \wedge dz^3 + z^1 \wedge dz^2 \wedge dz^3) : \quad (8.6)$$

The primitivity condition (4.28b) then constrains J to be of the form

$$J = \frac{1}{2 \text{Im} \tau} V_b (z^1 \wedge \bar{z}^1 + \bar{z}^1 \wedge z^1) + \frac{1}{2 \text{Im} \tau} (iV_2 dz^2 \wedge dz^2 + iV_3 dz^3 \wedge dz^3 - 2V_{(23)} \text{Im} dz^2 \wedge dz^3) ; \quad (8.7)$$

corresponding to the base metric

$$ds_B^2 = \frac{1}{\text{Im} \tau} (V_2 dz^2 dz^2 + V_3 dz^3 dz^3 + V_{(23)} \text{Re} dz^2 dz^3) : \quad (8.8)$$

From Eqs. (3.52a;b), the constraints on NS B - field moduli are

$$0 = 2h (b_{(2)4m} - b_{(2)5m}) dx^m \wedge (dx^6 \wedge dx^8 - dx^6 \wedge dx^9 - dx^7 \wedge dx^8 + dx^7 \wedge dx^9) + 2hb_{(2)5m} dx^m \wedge dx^7 \wedge dx^9; \quad (8.9a)$$

$$0 = 2fb_{(2)4m} dx^m \wedge dx^7 \wedge dx^9 + 2fb_{(2)5m} dx^m \wedge dx^6 \wedge dx^8 : \quad (8.9b)$$

Together, these constraints imply

$$b_{(2)47} = b_{(2)49} = b_{(2)56} = b_{(2)58} = 0; \quad (8.10a)$$

$$b_{(2)46} = b_{(2)57}; \quad \text{and} \quad b_{(2)48} = b_{(2)59} : \quad (8.10b)$$

So, there are two unconstrained NS B - field moduli,

$$b_{(2)46} + b_{(2)57} \quad \text{and} \quad b_{(2)48} + b_{(2)59} : \quad (8.11)$$

From the kinetic terms (3.54a;b;c), the massless RR sector scalars are the uncharged fields

$$e_{(2)67}; \quad e_{(2)89}; \quad e_{(2)69} - e_{(2)78}; \quad e_{(2)} \text{ } \S \quad e_{(6)456789}; \quad (8.12)$$

$$e_{(4)4678} + e_{(4)5679}; \quad \text{and} \quad e_{(4)5789} + e_{(4)4689} :$$

The massive gauge bosons are those that couple to the other RR scalars. From (3.54a;b;c), they lie in the 9D space spanned by

$$\begin{aligned}
(2)^2 \oplus (f=h)v^4 & \quad Q_{(2)5} ; \quad v^5 ; \quad C_{(2)4} \quad Q_{(2)5} ; \\
(2)^2 \oplus (f=h)b_{(2)6} & \quad e_{(4)456} ; \quad b_{(2)7} ; \quad e_{(4)456} + e_{(4)457} ; \\
(2)^2 \oplus (f=h)b_{(2)8} & \quad e_{(4)458} ; \quad b_{(2)9} ; \quad e_{(4)458} + e_{(4)459} :
\end{aligned} \tag{8.13}$$

The massless bosons are the three linear combinations orthogonal to this space, where orthogonality is defined with respect to the metric appearing in the gauge kinetic terms. (See Sec. 8.2 for further discussion in the dual O3 orientifold).

Finally, there are $6M$ massless scalars A_I , I^m , and M massless gauge bosons A_I from the D5 branes.

In summary, the massless fields are one graviton, $3 + M$ vectors, and $6 + 6M$ moduli. The moduli consist of

$$\begin{aligned}
V_b, V_2, V_3, V_{(23)}, 2 \text{ indep } b_{(2)m}, \\
6 e_{(p)} \text{ scalars, and } 6M \text{ D5 scalars.}
\end{aligned}$$

These fields combine to form one 4D $N = 1$ gravity multiplet, $3 + 3M$ chiral multiplets, and $3 + M$ vector multiplets.

The amount of unbroken supersymmetry is

$$N_+ = 1; \quad N_- = 0; \tag{8.14}$$

in the notation of Sec. 4.2. To verify this, first note that the complex structure modulus of the fiber is τ_1 , with $\text{Im } \tau_1 > 0$ from (8.5). Therefore, the supersymmetry is of + type. Since the base factorizes as $T_{f67g}^2 \times T_{f89g}^2$ with respect to complex structure, we can define chirality operators $e_{(2)} = e_2^2$ on T_{f67g}^2 and $e_{(3)} = e_3^3$ on T_{f89g}^2 , where the indices 2 and 3 are holomorphic z^2 and z^3 indices.³³ Then, since complex structure moduli of T_{f67g}^2 and T_{f89g}^2 are τ_2, τ_3 , with $\text{Im } \tau_i > 0$ from (8.5), there is one independent negative chirality 6D Killing spinor ϵ_- . It is of the form

$$\epsilon_- = \epsilon_+^{(b)} \epsilon_{(2)}^{(2)} \epsilon_{(3)}^{(3)}; \tag{8.15}$$

where $e_{(2)}^{(2)} = \epsilon_{(2)}^{(2)}$ and $e_{(3)}^{(3)} = \epsilon_{(3)}^{(3)}$. The three factors in Eq. (8.15) are on equal footing in the sense that e_i^i (no sum) = +1 in all three cases.

³³ Here, the relative minus sign is necessary for chiralities to be multiplicative, that is, for $e_B = e_{(2)} e_{(3)}$. In the conventions of this paper, $e_B = g_B^{1=2} e^{6789}$ (cf. Eq. (A.17)), which equivalent to $e_B = e_2^2 e_3^3$ when expressed in terms of holomorphic coordinates.

8.2. Dual O3 Orientifold

In this subsection, it is convenient to use the notation

$$(x^1; y^1; x^2; y^2; x^3; y^3)_{\text{here}} = (x^4; x^5; x^6; x^7; x^8; x^9)_{\text{rest of paper}} \quad (8:16)$$

In this notation, consider the O3 orientifold with internal T^6 and flux

$$\begin{aligned} \frac{1}{(2^-)^2} F_{(3)}^0 &= a^0 dx^1 \wedge dx^2 \wedge dx^3 + a dx^1 \wedge dx^2 \wedge dy^3 + \text{cyc. perm s. of 123} \\ &\quad b dx^1 \wedge dy^2 \wedge dy^3 + \text{cyc. perm s. of 123} + b_0 dy^1 \wedge dy^2 \wedge dy^3; \end{aligned} \quad (8:17a)$$

$$\begin{aligned} \frac{1}{(2^-)^2} H_{(3)}^0 &= c^0 dx^1 \wedge dx^2 \wedge dx^3 + c dx^1 \wedge dx^2 \wedge dy^3 + \text{cyc. perm s. of 123} \\ &\quad d dx^1 \wedge dy^2 \wedge dy^3 + \text{cyc. perm s. of 123} + d_0 dy^1 \wedge dy^2 \wedge dy^3; \end{aligned} \quad (8:17b)$$

This class of fluxes was analyzed in Sec. 4.1 of Ref. [11]. Let us quickly review that analysis. The $(2,1)$ constraint on the complex flux $G_{(3)}^0$ implies that

$$G_{ij}^0 = G_{ji}^0; \quad (8:18)$$

that is, $T^6 \simeq T^2 \oplus T^2 \oplus T^2$ with respect to complex structure, where G^0 and d_{dil}^0 satisfy

$$P_1(G^0) = a^0 \otimes 3a^{\otimes 2} - 3b^0 \otimes b_0 = 0; \quad (8:19a)$$

$$P_2(G^0) = c^0 \otimes 3c^{\otimes 2} - 3d^0 \otimes d_0 = 0; \quad (8:19b)$$

and

$$a^0 \otimes d_{\text{dil}}^0 c^0 \otimes c^{\otimes 2} - 2(a \otimes d_{\text{dil}}^0 c)^0 \otimes (b \otimes d_{\text{dil}}^0 d) = 0; \quad (8:19c)$$

A solution exists only if

$$P_1(G^0) = 2(f^0 + g)P(G^0); \quad P_2(G^0) = 2(h^0 + k)P(G^0); \quad (8:20)$$

where $P(G^0)$ is a quadratic polynomial of the form (7.27), and $f; g; h; k \in \mathbb{Z}$. In Eq. (8.20), we have added factors of 2 compared to the corresponding equation in Ref. [11], in order to automatically take into account the $2\mathbb{Z}$ quantization condition on the fluxes. The relations

$$\begin{aligned} 2(fm + gl) &= 3a; & 2(hm + kl) &= 3c; \\ 2(fn + gm) &= 3b; & 2(hn + km) &= 3d; \end{aligned} \quad (8:21)$$

give consistency conditions modulo 3, and the ux contribution to the Gauss's law constraint (5.1e) is

$$N_{ux} = \frac{4}{3} (fk - gh) (m^2 - 4ln) : \quad (8.22)$$

It can be shown from Eq. (8.21) that N_{ux} is divisible by 12.

Ref. [11] went on to consider the particular example

$$a^0; a; b; b_0 = (2; 0; 0; 2); \quad c^0; c; d; d_0 = (2; 2; 2; 4); \quad (8.23a)$$

$$(f; g; h; k; l; m; n) = (1; 1; 1; 2; 1; 1; 1); \quad (8.23b)$$

In App. H, we prove that this is the unique choice of supersymmetric ux in the class (8.17a;b) with the minimum value $N_{ux} = 12$ (and $2M = 20$ D3 branes), modulo $SL(2; Z) \times SL(2; Z)_{\text{dilat}}$ equivalences.³⁴ This choice gives

$$F_{\text{dilat}}^0 = e^{2i\pi/3} : \quad (8.23c)$$

The only other value of $N_{ux} \leq 32$ such that 12 divides N_{ux} is $N_{ux} = 24$ (with $2M = 8$ D3 branes). By arguments analogous to those in App. H, one can show that in this case there are two distinct choices of ux modulo equivalences: one choice differs from Eqs. (8.23a;b;c) by $f; g; \text{dilat}^0 \rightarrow 2f; 2g; 2 \text{dilat}^0$, the other differs by $h; k; \text{dilat}^0 \rightarrow 2h; 2k; \text{dilat}^0 = 2$. So, in the entire class (8.17a;b), there are just three inequivalent choices of supersymmetric ux .

The three possibilities are summarized by

$$\frac{1}{(2)^2} F_{(3)}^0 = 2f dx^1 \wedge dx^2 \wedge dx^3 + dy^1 \wedge dy^2 \wedge dy^3 ; \quad (8.24a)$$

$$\begin{aligned} \frac{1}{(2)^2} H_{(3)}^0 &= 2h dx^1 \wedge dx^2 \wedge dx^3 - 2h dx^1 \wedge dx^2 \wedge dy^3 + \text{cyc. perm s. of 123} \\ &+ 2h dx^1 \wedge dy^2 \wedge dy^3 + \text{cyc. perm s. of 123} - 4h dy^1 \wedge dy^2 \wedge dy^3 : \end{aligned} \quad (8.24b)$$

$$N_{ux} = 12fh; \quad F_{\text{dilat}}^0 = (h=f) \text{dilat}^0 = e^{2i\pi/3}; \quad (8.24c)$$

where $(f; h) = (1; 1), (2, 1),$ or $(1, 2)$.

The complex three-form ux is

$$G_{(3)}^0 = 2ife^{-2i\pi/3} \frac{P}{3} dz^1 \wedge dz^2 \wedge dz^3 + dz^1 \wedge dz^2 \wedge dz^3 + dz^1 \wedge dz^2 \wedge dz^3 : \quad (8.25)$$

³⁴ This was suggested, but not proven in Ref. [11].

The primitivity condition (5.7) implies that J^0 is of the form

$$J^0 = \frac{1}{2j\text{Im}^0j} iV_1^0 dz^1 \wedge dz^1 + iV_2^0 dz^2 \wedge dz^2 + iV_3^0 dz^3 \wedge dz^3 \\ + 2V_{(12)}^0 \text{Im} (dz^1 \wedge dz^2) + 2V_{(23)}^0 \text{Im} (dz^2 \wedge dz^3) + 2V_{(31)}^0 \text{Im} (dz^3 \wedge dz^1) ; \quad (8:26)$$

where $z^i = x^i + iy^i$. In real coordinates, this becomes

$$J^0 = v_1^0 dx^1 \wedge dy^1 + v_2^0 dx^2 \wedge dy^2 + v_3^0 dx^3 \wedge dy^3 \\ + v_{(12)}^0 (dx^1 \wedge dy^2 - dy^1 \wedge dx^2) + (12 \leftrightarrow 23) + (12 \leftrightarrow 31) ; \quad (8:27)$$

where $v_i^0 = V_i^0 \text{sign}^0$ and $v_{(ij)}^0 = V_{(ij)}^0 \text{sign}^0$. The metric is

$$ds_6^2 = \frac{1}{j\text{Im}^0j} \sum_{i=1}^3 V_i^0 dz^i dz^i + \sum_{\substack{i,j=1 \\ i \neq j}}^3 V_{(ij)}^0 dz^i dz^j ; \quad (8:28)$$

where $V_{(ij)}^0 = V_{(ji)}^0$.

From the kinetic terms (5.15), the massless RR scalars are the uncharged scalars

$$c_{(4)x^2y^2x^3y^3}^0 ; c_{(4)x^3y^3x^1y^1}^0 ; c_{(4)x^1y^1x^2y^2}^0 ; c_{(4)x^1y^1x^2y^3}^0 + c_{(4)x^1y^1x^3y^2}^0 ; \\ c_{(4)x^2y^2x^3y^1}^0 + c_{(4)x^2y^2x^1y^3}^0 ; \text{ and } c_{(4)x^3y^3x^1y^2}^0 + c_{(4)x^3y^3x^2y^1}^0 ; \quad (8:29)$$

which are the axionic partners of $V_1^0, V_2^0, V_3^0, V_{(23)}^0, V_{(31)}^0$, and $V_{(12)}^0$, respectively, in $N = 1$ chiral multiplets.

The massive gauge bosons are those that couple to the other RR scalars. From (5.15), they lie in the 9D space spanned by

$$(f=h)b_{(2)x^i}^0 + c_{(2)x^i}^0 ; b_{(2)y^i}^0 ; \text{ and } c_{(2)x^i}^0 + c_{(2)y^i}^0 ; \quad i = 1;2;3 ; \quad (8:30)$$

In terms of the complex gauge bosons (5.16), this is the space spanned by

$$\text{Im} (1 = \text{Im}^0) d_{z^i}^0 ; \text{Re} d_{z^i}^0 ; \text{ and } \text{Im} d_{z^i}^0 ; \quad i = 1;2;3 ; \quad (8:31)$$

The massless bosons are the linear combinations orthogonal to this space, where orthogonality is defined with respect to the metric appearing in the gauge kinetic terms. In the

$d_{z^i}^0 ; d_{z^i}^0 ; d_{z^i}^0 ; d_{z^i}^0$ basis, the latter is proportional to

$$\begin{pmatrix} 1 & 0 & 1 & d^0 & 0 & g_6^0 z^i z^j & z^i \\ 2 & 1 & 0 & d^0 & g_6^0 z^i z^j & 0 & z^i \end{pmatrix} ; \quad (8:32)$$

Therefore, the three massless gauge bosons are

$$\text{Re}(f) d_{z^i}^0 / (f=h) b_{(2)x^i}^0 + b_{(2)y^i}^0 + c_{(2)y^i}^0 ; \quad i=1;2;3: \quad (8.33)$$

Finally, there are $6M$ moduli τ_i and M massless gauge bosons A_I from the $D3$ branes.

In summary, the massless fields are one graviton, $3+M$ vectors, and $6+6M$ moduli. The moduli consist of

$$V_1^0, V_2^0, V_3^0, V_{(23)}^0, V_{(31)}^0, V_{(12)}^0, \\ 6 e_{(p)} \text{ scalars, and } 6M \text{ D5 scalars.}$$

These fields combine to form one 4D $N=1$ gravity multiplet, $3+3M$ chiral multiplets, and $3+M$ vector multiplets.

From the supersymmetry-breaking mass spectra given in Ref. [17], this number of massless moduli and gauge bosons corresponds to $N=4 \rightarrow N=1$ with $m_1 = m_2 = m_3$ for the three massive gravitini. There are $3+3M$ massless chiral multiplets and $3+M$ massless vector multiplets. The metric on moduli space is discussed in App. G. Since the complex structure modulus for each T^2 factor satisfies $\text{Im} \tau^0 > 0$, the 6D spinor ϵ^0 generating the supersymmetries can be written as the product of three 2D spinors, each of definite chirality along a T^2 (cf. the discussion at the end of Sec. 8.1).

8.3. T-Duality Map

If we return from the notation of Sec. 8.2 to the conventional labeling of coordinates via (8.16), then it is straightforward to show that the metric (8.28) can be written as a flat T^2 fibration (5.18), with

$$V_b^0 = V_1^0; \quad (8.34a)$$

$$a^{04} = V_1^0 \tau^1 V_{(12)}^0 dx^6 + V_{(31)}^0 dx^8; \quad (8.34b)$$

$$a^{05} = V_1^0 \tau^1 V_{(12)}^0 dx^7 + V_{(31)}^0 dx^9; \quad (8.34c)$$

$$ds_B^2 = \frac{1}{j \text{Im} \tau^0} V_2^0 (V_{(12)}^0)^2 = V_1^0 dz^2 dz^2 + V_3^0 (V_{(31)}^0)^2 = V_1^0 dz^3 dz^3 \\ + V_{23}^0 V_{(12)}^0 V_{(31)}^0 = V_1^0 dz^2 dz^3 + dz^3 dz^2 : \quad (8.34d)$$

The base metrics (8.8) and (8.34d) agree, provided that we make the identifications

$$V_2 = V_2^0 (V_{(12)}^0)^2 = V_1^0; \quad V_3 = V_3^0 (V_{(31)}^0)^2 = V_1^0; \\ V_{(23)} = V_{(23)}^0 V_{(12)}^0 V_{(31)}^0 = V_1^0; \quad \tau^0 = \tau^0: \quad (8.35)$$

All other quantities map exactly as described in Sec. 5.2.

9. Preview : $N = 2$ Calabi-Yau Duality without Flux³⁵

So far, we have said nothing about g_s or α' corrections. We know that there are at least some such corrections, due to the following pathology in the leading $g_s; \alpha'$ supergravity description of the orientifold.³⁶ As $x \rightarrow x^0$, the Green's function (3.31) behaves like a Green's function on R^4 :

$$G_B(x; x^0) = \frac{1}{8^2} \frac{1}{|x - x^0|^2} + c; \quad (9.1)$$

where c is a constant. So, near an $O5$ plane (with $Q_{O5} = 2$),

$$Z(x) = 1 - g^0 \frac{1}{|x - x_{O5}|^2} + c; \quad (9.2)$$

Consequently, as we let x approach x_{O5} , we encounter a locus $Z = 0$ where the metric is singular and $e^{-2\phi} = g_s^2 Z^{-1}$ diverges. Beyond this locus, we enter the region $Z < 0$, where the metric and $e^{-2\phi}$ are imaginary. A similar pathology affects the leading supergravity description of all orientifolds. On the other hand, orientifold planes are not singular objects in string theory. Therefore, the 10D effective field theory descending from string theory must get corrected in a way that resolves the pathology. In the $N = 2$ and $N = 4$ cases, it is easy to understand this resolution.

In the $N = 2$ example of Sec. 6.2, there is an isometry in the x^9 direction (spontaneously broken by the positions of the eight $O5$ planes and $2M-D5$ branes). If we T-dualize in this direction, we obtain an $O6$ orientifold, with $O6$ planes and $D6$ branes wrapping the $x^4; x^5; x^9$ directions. Since there was no NS flux before the T-duality, there is no new $S^1_{fx^9/g}$ bration introduced by the T-duality. The only flux after the T-duality is $F_{(2)}$, which is (the pullback to 6D of) a two-form in the $T^3_{fx^6; x^7; x^8/g}$ directions. As noted in Ref. [4], this IIA background lifts to a purely geometrical M theory background. The $F_{(2)}$ flux gives the bration of the M theory circle $S^1_{fx^{10}/g}$ over the IIA geometry. The dilaton gives the size of the fiber. Finally, the $D6$ branes and $O6$ planes, the only objects that are singular in the IIA supergravity description, lift locally to smooth Taub-NUT and smooth

³⁵ I am grateful to P. Berglund, V. Braun, B. Florea, C. Johnson, R. Reinbacher, N. Warner, and C. Vafa for comments and suggestions related to this section. In particular, P. Berglund and N. Warner first suggested studying the lift to M theory.

³⁶ Here, leading order in g_s means $\alpha' \rightarrow 0 + \frac{1}{2}$ loop order, that is, tree level in the closed string sector plus tree level in the open string sector.

Atiyah-Hitchin spaces times $R^{6;1}$, respectively [78]. (The Atiyah-Hitchin space is the Z_2 quotient of negative mass Taub-NUT space in which the region too close to the center has been smoothly excised. The excised region corresponds roughly to the unphysical region $Z < 0$ of the orientifold.) Thus, the IIA orientifold lifts to

$$M \text{ theory on } Y_7; \quad \text{where } Y_7 = S^1_{fx^9g} \text{ smooth } Y_6:$$

There is no flux in M theory after the lift. The manifold Y_7 is truly a product, with no discrete identifications mixing the two factors. The orientifold Z_2 operation lifts to a geometrical Z_2 that inverts $x^6; x^7; x^8; x^{10}$. So, it goes into the definition of Y_6 , but does not act on $S^1_{fx^9g}$. Since the M theory compactification preserves 4D $N = 2$ supersymmetry, it follows that

$$Y_6 \text{ is a smooth Calabi-Yau threefold.}$$

Compactifying on $S^1_{fx^9g}$ then gives a standard $N = 2$ Calabi-Yau compactification of IIA. Compactifying on $S^1_{fx^{10}g} Y_6$ (with Y_6 viewed as a fibration) gives the O6 orientifold. Alternatively, F theory on $Y_6 \times T^2$ gives the original O5 orientifold of Sec. 6.2, provided that we identify the F theory torus with $S^1_{fx^{10}g} Y_6$ times $S^1 \times T^2$. Since this F theory T^2 fibration has no global section, there is automatically $F_{(3)}$ flux in the IIB orientifold [79].

Since the geometry $R^{3;1} \times S^1 \times Y_6$, with no flux, solves the equations of motion of 11D supergravity, the only essential ingredient that is missing in the leading IIA description but present in the M theory dual description is the dependence of the 11D background on the x^{10} direction. The 10D IIA supergravity theory is the dimensional reduction of 11D supergravity truncated to lowest Fourier modes along the x^{10} circle. The full Kaluza-Klein reduction of 11D supergravity includes not only the IIA supergravity fields, but also the entire tower of their D0-charged massive cousins from the bound states of N D0 branes, for all $N \geq 1$. Away from the O6 planes, if g_s is tuned to be small, the D0 bound states are heavy, and it is a good approximation to ignore the massive tower. By tuning g_s smaller and smaller, this approximation becomes valid closer and closer to the O6 planes. However, e always diverges at some locus near the O6 planes. As we approach this locus, the D0 bound states become massless and cannot be neglected.

It is natural to ask what class of Calabi-Yau threefolds arises from the duality just described. Immediately, we know at least one piece of topological data. In Sec. 6.2 we determined that there were $n_V = 2 + M$ vector multiplets and $n_H = 3 + M$ hypermultiplets

in addition to the gravity multiplet in the massless spectrum. On the other hand, for Calabi-Yau compactifications of type IIA, $n_V = h^{1;1}$ and $n_H = h^{2;1} + 1$. Therefore,

$$h^{1;1}(Y_6) = h^{2;1}(Y_6) = 2 + M; \quad (9:3)$$

where the possible values of M are $M = 0, 4, 8$, and 12 . In the case that $M = 16$, there is $N = 4$ supersymmetry and Y_6 is $T^2_{\mathbb{F}x^4, \mathbb{F}x^5g} \times K3_{\mathbb{F}x^6, \mathbb{F}x^7, \mathbb{F}x^8, \mathbb{F}x^{10}g}$ [80].

Beyond this, it is possible to determine the intersection numbers from the special Kähler geometry of the moduli space (6:72b). Furthermore, for each M , there is not one Calabi-Yau, but topologically distinct geometries distinguished by the integers $m; n$ of Sec. 6 such that $4m + n = 16 - M$:

M	$(h^{1;1}; h^{2;1})$	$(m; n)$	
0	(2;2)	(4;1); (2;2); (1;4)	(9:4)
4	(6;6)	(3;1); (1;3)	
8	(10;10)	(2;1); (1;2)	
12	(14;14)	(1;1)	

The integer n tells us about $H_1(Y_6; \mathbb{Z})$, which is pure torsion (and subsequently about $H_1(Y_6)$, whose abelianization is $H_1(Y_6; \mathbb{Z})$). In the dual $O(3)$ orientifold of Sec. 6.4, S-duality interchanges the integers m and n .³⁷ Therefore, compactification of type IIA on $Y_6(m; n)$ and $Y_6(n; m)$ gives the same low energy effective field theory. Since both compactifications are of type IIA, the relation between $Y_6(m; n)$ and $Y_6(n; m)$ is something different from Calabi-Yau mirror symmetry.

The details of this duality will appear in a separate paper [56]. There, we will derive the further topological data mentioned in the previous paragraph. In addition, from the standard identifications between 11D supergravity and 10D type IIA supergravity, we will obtain an approximate metric for Y_6 , together with its corresponding closed Kähler form and closed (3,0) form.³⁸

³⁷ To be precise, m and n are interchanged under S-duality followed by the coordinate relabeling $(x^8; x^9)_{\text{new}} = (x^9; x^8)_{\text{old}}$.

³⁸ The approximate metric is Calabi-Yau on a noncompact space that excludes the singular region $Z^0 = 0$, where Z^0 is the warp factor of the $O(6)$ orientifold. This metric is the lowest Fourier mode of the exact Calabi-Yau metric on the compact manifold Y_6 , in a Fourier expansion along the x^{10} direction. It can in principle be systematically extended to the exact Calabi-Yau metric by including all higher Fourier modes, that is, by solving the 6D Einstein equations order by order in an x^{10} Fourier expansion.

An added incentive for studying this duality is that it could help us to compute other g_s and α' corrections in the original orientifold. For corrections that map to worldsheet and D instantons in the Calabi-Yau duals, we can hope to take advantage of the existing results for Calabi-Yau compactifications and then map the results back to the orientifold.

10. Conclusions and Outlook

Type IIB orientifolds with wrapped O5 planes and internal flux yield a class of 4D vacua that remains largely unexplored compared to its O3 and O7 counterparts. We have studied the simplest such vacua here. We have seen that these vacua, in which O5 planes and D5 branes wrap the T^2 fiber over a T^4 base, provide a computable toy model for studying moduli stabilization in a torsionful compactification. The supersymmetry conditions can be expressed in terms of a complex three-form flux $\mathbb{G}_{(3)}$ just as for D3/D7 or heterotic compactifications, and in addition a complex one-form flux $G_{(1)}$. This was illustrated in Secs. 6–8 through five different examples preserving 4D $\mathcal{N} = 1, 2,$ and 3 supersymmetry. As a check, we have also seen that the results of each example agree with the T-dual analysis in the corresponding $T^6 = \mathbb{Z}_2$ orientifold. By providing an intrinsic description in the O5 orientifold, we have generalized the results of Ref. [4], which relied solely on the T-duality map at a special locus in moduli space.

One might have hoped that by giving a description of the O5 orientifold not grounded in T-duality, we would have found new solutions that do not T-dualize to the $T^6 = \mathbb{Z}_2$ O3 orientifold. However, as observed in Sec. 3.1, the condition for the T-duality to fail to exist is that $H_{(3) m}$, with two fiber indices, is nonzero, and this is exactly the case in which the analysis of the O5 orientifold qualitatively changes: the pseudo-BPS conditions of Sec. 3.3 no longer hold; moreover, this component of flux can be thought of as a transversely varying NS B-field oriented parallel to the D5 brane worldvolumes, which gives rise to noncommutativity. It would be interesting to prove that $\mathcal{N} > 1$ supersymmetry demands that $H_{(3) m} = 0$.

There are a number of other questions left unanswered by this paper. In Sec. 3.9, we stated the flux quantization conditions in the O5 orientifold, but offered no proof. It is a significant omission that we were not able to derive these conditions directly in the O5 orientifold. The quantization conditions agree with those of the dual O3 orientifold via

Eqs. (5.31). However, it should be possible, and would be more satisfactory, to instead derive the conditions directly from the appropriate twisted K-theory analysis.³⁹

In our description of the massless spectra, we were more explicit than in Ref. [4] for the O5 orientifold or Ref. [11] for the dual O3 orientifold. As a result, one feature that we noticed is that in all examples, the partial supersymmetry breaking from $N = 4$ to $N = 1, 2,$ or 3 is of the "nongeneric" type in which the massive gravitini have the same mass. A natural question is whether this is required. Do there exist supersymmetric $T^6 = \mathbb{Z}_2$ flux vacua, with quantized flux satisfying the Gauss's law constraint, in which the superHiggs mechanism gives different masses to the gravitini? Also, our analysis of the massless spectrum proceeded on a case by case basis. Is there a more elegant way of computing the massless spectrum through the cohomology of a suitable operator?

Other than in the introduction, we mentioned SU(3) torsion classes in this paper only in Sec. 6.3, where the criterion $W_1 \neq 0$ was used to prove that the almost complex structure of Ex. 3 was nonintegrable. We did not employ SU(2) torsion classes at all. The reason is that the torsion classes simply were not essential for our purposes of analyzing the supersymmetry conditions and moduli stabilization in this particular class of O5 vacua. Nevertheless, in recent months there has been considerable effort devoted to developing a framework that can be used to describe the most general 4D Minkowski vacua of type IIA and IIB string theory. (In the case of noncompact internal geometry this is closely related to studying type II or M theory solutions with an AdS_5 factor [36].) The correct framework for this analysis is in terms of SU(2) structures and torsion classes [41,36,42]. For the example of Sec. 6.3, in which $\mathbb{F}_{(3)}, F_{(1)},$ and $H_{(3)}$ are all nonzero, we were able to avoid talking about SU(2) structures by instead working in terms of the right-SU(3) structures, and the volume form on the fiber. However, this is the one example of the paper in which supersymmetry does not provide us with a single 6D spinor, but two linearly independent spinors χ_R and $\chi_L = \gamma_5 \chi_R$. The two spinors canonically determine an SU(2) structure rather than an SU(3) structure. Therefore, this example should provide a useful testing ground for the IIB supersymmetry conditions as formulated in Ref. [42] in terms of SU(2) torsion classes. One just needs to compute the torsion classes corresponding to the example. (In the other examples, the SU(2) torsion collapses to SU(3) torsion.) Working in terms of these torsion classes might have the added benefit of proving that the almost complex structure is always nonintegrable for the subclass of O5 backgrounds of the type

³⁹ I am indebted to G. Moore for a helpful discussion on flux quantization.

discussed here when there is NS flux, a result that we proved only for the specific example of Sec. 6.3.

Finally, perhaps the most interesting direction to pursue is to make more precise the duality of Sec. 9 relating $N = 2$ orientifold vacua to standard type IIA Calabi-Yau vacua. We have described a number of computable features of the dual Calabi-Yau threefolds and are optimistic about the prospects for identifying these manifolds. As one application of the identification, note that although it is common to speak of connected webs of $N = 2$ string vacua, given a pair of $N = 2$ vacua, it is not necessarily known whether the moduli spaces of the two are connected. For the class of $N = 2$ Calabi-Yau vacua, what is known is that the subspace of smooth Calabi-Yau threefolds that are hypersurfaces in toric varieties is connected. This subspace has trivial fundamental group, and has been completely described by Kreuzer and Skarke, who tabulated all 473,800,776 reflexive polyhedra in four dimensions [81]. In at least the case that $(M; m; n) = (12; 1; 1)$ and $h^{1,1} = h^{2,1} = 14$, the Calabi-Yau Y_6 of Sec. 9 is conceivably contained in this subspace, and the dual orientifold contained in the corresponding web of known connected $N = 2$ vacua. This intriguing duality between $N = 2$ orientifold vacua and standard type IIA Calabi-Yau vacua is currently under investigation [56].

Acknowledgements

It is a pleasure to thank Allan Adams, Per Berglund, Andreas Brandhuber, Ram Brustein, Richard Corrado, Jerome Gauntlett, Jaume Gomis, Fawad Hassan, Anton Kapustin, Albion Lawrence, Christian Romelsberger, and especially Andrew Frey for helpful discussions and useful references. In addition, I thank S. Kachru, P. Tripathy, and S. Trivedi for the enjoyable collaboration from which this investigation is a continuous outgrowth. Finally, I am grateful to Stockholm University, the Aspen Center for Physics, the Kavli Institute for Theoretical Physics, the University of Pennsylvania, and Harvard University for hospitality during the course of this work. This work was supported in part by the DOE under contract DE-FC03-92-ER40701 and by the National Science Foundation under Grant No. PHY 99-07949.

Appendix A . C onventions

The conventions for coordinate indices are as follow s:

$M ; N ; \dots$	denote 10D spacetime indices in the range $0; \dots; 9$,
$\mu ; \nu ; \dots$	denote 4D spacetime indices in the range $0; \dots; 3$,
$a ; b ; \dots$	denote 6D internal indices in the range $4; \dots; 9$,
$\alpha ; \beta ; \dots$	denote T^2 fiber indices in the range $4; 5$,
$m ; n ; \dots$	denote 4D base indices in the range $6; \dots; 9$,
$i ; j ; \dots$ and $\{i ; j ; \dots\}$	denote complex 6D internal indices in the range $1; 2; 3$.

When 10D tensors (such as R_{MN} or the fluxes) are decomposed into $\mu ; \nu$ components, the decomposition is with respect to the basis $dx^\mu ; dx^\nu ; dx^m$, not $dx^\mu ; dx^\nu ; dx^m$, with one exception. In the context of the $O(3)$ orientifold, underscored indices $\underline{\mu} ; \underline{\nu} ; \underline{m}$ denote components in the basis $dx^\mu ; dx^\nu ; dx^m$.

The squares and partially contracted squares of 10D tensors are defined by

$$\begin{aligned} A_{(p)}^2 &= A_{(p)M_1 \dots M_p} A_{(p)}^{M_1 \dots M_p} ; \\ A_{(p)MN}^2 &= A_{(p)M Q_2 \dots Q_p} A_{(p)N}^{Q_2 \dots Q_p} ; \end{aligned} \quad (\text{A } 1)$$

with the metric (2.4) used for raising and lowering indices. This metric is also used to define the square of the bration curvature,

$$F^2 = g_{mn} g^{pq} F_{mn} F_{pq} ; \quad (\text{A } 2)$$

We follow the differential geometry conventions of Nakahara [82] for normalization of differential forms, exterior differentiation, Hodge duality, and curvature. (In the terminology of Misner, Thorne, and Wheeler [83], we employ + + + sign conventions.) In particular, on the 6D internal manifold,

$$\star \wedge^6 = \frac{1}{p!} \epsilon_{a_1 \dots a_p} \epsilon^{a_1 \dots a_p} \text{Vol}_6 ; \quad (\text{A } 3a)$$

$$\frac{1}{3!} J \wedge J \wedge J = \frac{i}{8} \star = \text{Vol}_6 ; \quad (\text{A } 3b)$$

Compared to earlier work [9,11,4], the Hodge star operator maps the same forms to one another, but Vol_6 here is minus Vol_6 there. Therefore, the two possible selfduality conditions on middle dimensional forms,

$$\begin{aligned} \star \wedge^3 &= + i \wedge^3 \quad \text{imaginary-selfduality (ISD)} ; \\ \star \wedge^3 &= - i \wedge^3 \quad \text{imaginary-antselfduality (IASD)} ; \end{aligned} \quad (\text{A } 4)$$

are the same here as in the earlier work [9,11,4].

Our normalization conventions for 6D Hermitian metrics in complex coordinates $z^i; \bar{z}^j$ follow by compatibility with the expression for the metric in arbitrary 6D coordinates $x^a; \bar{x}^b$:

$$ds^2 = g_{ab} dx^a dx^b = g_{i\bar{j}} dz^i d\bar{z}^j + g_{\bar{j}i} d\bar{z}^j dz^i = 2g_{i\bar{j}} dz^i d\bar{z}^j \quad (\text{A.5})$$

Note the factor of 2 on the RHS of the last equality.

The notation $!_{(p)} = !_{(p)}^0 + !_{(p)}^1 + !_{(p)}^2$ is defined in Sec. 3.2, and denotes the decomposition of an internal 6D p-form $!_{(p)}$ into components $!_{(p)}^i$ of rank i on the T^2 fiber.

Slashes denote contraction of tensors with 10D Dirac matrices,

$$\cancel{A}_{(p)M_1 \dots M_k} = \frac{1}{(p-k)!} A_{(p)M_1 \dots M_k} \gamma^{M_{k+1} \dots M_p} \quad (\text{A.6})$$

Antisymmetrized products of Dirac matrices are

$$\begin{aligned} \gamma^{M_1 \dots M_n} &= [\gamma^{M_1} \gamma^{M_2} \dots \gamma^{M_n}], \\ a_1 \dots a_n &= [a_1 a_2 \dots a_n]. \end{aligned} \quad (\text{A.7})$$

The antisymmetrized product of all ten γ^M gives the 10D chirality operator,

$$\gamma^{(10)} = \gamma^{1=2} \gamma^{0123456789}; \quad (\text{A.8})$$

where g_{MN} is the 10D metric (2.4). Uppercase Dirac matrices satisfy

$$\gamma^M \gamma^N + \gamma^N \gamma^M = 2g_{MN} \quad (\text{A.9})$$

Lowercase Dirac matrices satisfy

$$\gamma^a \gamma^b + \gamma^b \gamma^a = 2g_{ab}; \quad (\text{A.10})$$

where $\eta = \text{diag}(1;1;1;1)$ and g_b is the 6D internal metric (2.7).

The relation between uppercase and lowercase Dirac matrices is

$$\gamma^M = Z^{1=4} \gamma^m; \quad \gamma_a = Z^{1=4} \gamma^a; \quad (\text{A.11})$$

where

$$Z^{(4)} = i \gamma^{0123}; \quad (\text{A.12})$$

Here, aside from the overall warp factor (cf. Eq. (2.8)), we follow Candele, Horowitz, Strominger, and Witten [5]. We also follow the latter in employing a Majorana representation. The γ_M ; are real and Hermitian, except for γ_0 ; γ_0 , which are real and antiHermitian. The γ_a are imaginary and Hermitian.

In addition, we define

$$\begin{aligned} \gamma^{(4)} &= i\gamma^{1\ 0123}; \\ \gamma^{(6)} &= i\gamma^{1=2\ 1=2\ 456789}; \\ \gamma_b &= i\gamma^{1=2\ g^{(T_{\text{fib}}^2)}\ 1=2\ 45}; \\ \gamma_B &= \gamma^{1=2\ 6789}; \end{aligned} \tag{A.13}$$

and similarly,

$$\begin{aligned} \gamma^{(6)} &= i\gamma_6^{1=2\ 456789}; \\ \gamma_b &= i\gamma^{(T_{\text{fib}}^2)}\ 1=2\ 45}; \\ \gamma_B &= \gamma_B^{1=2\ 6789}; \end{aligned} \tag{A.14}$$

Since $\gamma^{(4)2} = 1$, we then have the relations

$$\begin{aligned} \gamma^{(4)} &= \gamma^{(4)}\ 1; \quad \gamma_b = 1\ \gamma_b; \quad \text{and} \quad \gamma_B = 1\ \gamma_B; \\ \gamma^{(6)} &= \gamma_b\ \gamma_B = 1\ \gamma^{(6)}; \quad \gamma^{(10)} = \gamma^{(4)}\ \gamma^{(6)} = \gamma^{(4)}\ \gamma^{(6)}; \end{aligned} \tag{A.15}$$

as well as

$$\gamma_{a_1 \dots a_{2n}} = \gamma^{n=2}\ 1\ \gamma_{a_1 \dots a_{2n}}; \quad \gamma_{a_1 \dots a_{2n+1}} = \gamma^{(2n+1)=4}\ \gamma^{(4)}\ \gamma_{a_1 \dots a_{2n}}; \tag{A.16}$$

In our representation conventions, $\gamma^{(4)}$, $\gamma^{(6)}$, and γ_b are imaginary and Hermitian, while γ_B is real and Hermitian. Thus, complex conjugation \dagger of a 6D internal spinor reverses its $\gamma^{(6)}$ and γ_b chirality, but leaves its γ_B chirality unchanged. Complex conjugation $u \dagger u$ of a 4D spacetime spinor u reverses its $\gamma^{(4)}$ chirality.

We can further decompose the lowercase Dirac matrices as

$$\gamma_m = e\ 1; \quad \gamma_m = \gamma^{1=2}\ e_b\ \gamma_m; \tag{A.17}$$

where

$$e_b = i\gamma^{(T_{\text{fib}}^2)}\ e^{45}; \quad e_B = \gamma_B^{1=2}\ e^{6789}; \tag{A.18}$$

and

$$f e; e g = 2\gamma^{(T_{\text{fib}}^2)}; \quad f e_m; e_n g = 2\gamma_{B\ m\ n}; \tag{A.19}$$

Since $b^2 = 1$, we then have the relations

$$b = e_b \quad 1; \quad B = 1 \quad e; \quad (A 20)$$

and finally

$$= e \quad 1; \quad m^n = 1 \quad e_m^n; \quad m_n = Z e \quad e_m n; \quad (A 21)$$

which are needed in Sec. 4.2.

The convention for subscripts on 6D, 4D, and 2D spinors is that the subscripts always indicate the fiber chirality and/or minus the base chirality. Therefore, the only 6D spinors that we refer to directly are of negative ⁽⁶⁾ chirality.

Appendix B. Coordinate Identifications

The fiber coordinates are periodically identified via

$$x = x + 1 \quad \text{at} \quad x \in \mathbb{Z}; \quad x \in \mathbb{Z} : \quad (B 1)$$

In the case that the base B is a torus, Eq. (2.5) implies that

$$A = \frac{1}{2} F_{mn} x^m dx^n; \quad F_{mn} = \text{const} \in \mathbb{Z}; \quad (B 2)$$

up to a fiber coordinate redefinition

$$x_{\text{new}} = x + \quad ; \quad A_{\text{new}} = A \quad d \quad ; \quad \text{where} \quad = \quad f x^m g : \quad (B 3)$$

In the gauge (B 2), the base coordinates have a periodic identification that also acts on the fiber coordinates:

$$(x^m; x) \rightarrow (x^m + 1; x - \frac{1}{2} F_{mn} x^n) \quad \text{at} \quad x \in \mathbb{Z}; \quad x \in \mathbb{Z} : \quad (B 4)$$

This twisted identification is the unique coordinate identification that both projects to $x^m \rightarrow x^m + 1$ on the base and is compatible with

$$= (dx + A)_{x^m} = (dx + A)_{x^m + 1} \quad \text{at} \quad x \in \mathbb{Z}; \quad x \in \mathbb{Z} : \quad (B 5)$$

Other gauge choices lead to similar coordinate identifications.

Appendix C . Derivation of Pseudo-BPS Constraints

Let

$$\begin{aligned} \mathcal{L}_1 &= g_{RR} + 2r_r \dot{r} \quad ; \\ \mathcal{L}_2 &= g_{RR} + 2r_r \dot{r} + \frac{1}{4}F^2; \\ \mathcal{L}_3 &= r^2 + 2(r_r \dot{r})^2; \end{aligned} \tag{C.1}$$

and define

$$\mathcal{L} = \mathcal{L}_1 + \mathcal{L}_2 + \mathcal{L}_3 : \tag{C.2}$$

The warp factor Z and dilaton ϕ are assumed to depend only on the base coordinates. For such a function f , we have

$$\begin{aligned} r_r f &= Z^{-1} g_B^{mm} (\log Z)_{;m} \dot{r} \\ r_r f &= Z^{-1} g_B^{mm} (\log Z)_{;m} \dot{r} \\ r_m r_n f &= r_{Bm} r_{Bn} \left(\frac{1}{4} Z^{-1} Z_{;k} g_B^{k^0 k} g_{Bmn} + Z_{;m} \dot{r}_{;n} + Z_{;m} \dot{r}_{;n} f_{;k} \right) : \end{aligned} \tag{C.3}$$

Using Eqs. (3.2) and (C.3), it is straightforward to show that

$$\begin{aligned} r_B Z^{-1} e^{-2\phi} r_B Z^{-3=2} e^{-2\phi} &= Z^{1=2} Z^{-1} e^{-2\phi} Z^{-3=2} e^{-2\phi} \\ &+ Z^{-1} e^{-2\phi} Z^{-3=2} e^{-2\phi} r_B Z^{-3=2} e^{-2\phi} \quad ; \end{aligned} \tag{C.4}$$

where contractions are performed using base metric g_{Bmn} . On the other hand, from Eqs. (3.1) and (3.4), we have⁴⁰

$$\begin{aligned} \mathcal{L}_1 &= e^2 F_{(1)}^2 + \frac{1}{3!} \mathbb{F}_{(3)}^2 + \frac{1}{2} \frac{1}{5!} \mathbb{F}_{(5)}^2 + (2_r \dot{r})^2 e^{-X} Q_i \frac{4(x_r x_i)}{Z^4 g_B} ; \\ \mathcal{L}_2 \quad \frac{1}{2} \mathcal{L}_1 &= \frac{1}{2} e^2 \left(\frac{1}{3!} (\mathbb{F}_{(3)}^1)^2 + \frac{1}{5!} (\mathbb{F}_{(5)}^1)^2 + \frac{2}{5!} (\mathbb{F}_{(5)}^2)^2 + \frac{1}{4} F^2 \right) ; \\ \mathcal{L}_3 &= e^2 F_{(1)}^2 + \frac{1}{2} \frac{1}{3!} \mathbb{H}_{(3)}^2 + \frac{1}{2} \frac{1}{3!} \mathbb{H}_{(3)}^2 + \frac{1}{2} (2_r \dot{r})^2 e^{-X} Q_i \frac{4(x_r x_i)}{Z^4 g_B} : \end{aligned} \tag{C.5}$$

Therefore,

$$\begin{aligned} &= \frac{1}{2} e^2 F_{(1)}^2 + \frac{2}{3!} (\mathbb{F}_{(3)}^0)^2 + \frac{1}{3!} (\mathbb{F}_{(3)}^1)^2 + \frac{2}{5!} (\mathbb{F}_{(5)}^1)^2 + \frac{1}{5!} (\mathbb{F}_{(5)}^2)^2 \\ &+ \frac{1}{2} \frac{1}{3!} (\mathbb{H}_{(3)}^0)^2 + \frac{1}{2} \frac{1}{2!} F^2 + (2_r \dot{r})^2 e^{-X} Q_i \frac{4(x_r x_i)}{Z^4 g_B} : \end{aligned} \tag{C.6}$$

⁴⁰ Here, $\mathbb{F}_{(5)} = (1 - \mathbb{F}_{(5)}^{\text{int}})$, where $\mathbb{F}_{(5)}^{\text{int}} = \sum^P \mathbb{F}_{(5)}^i$ denotes the purely internal part of $\mathbb{F}_{(5)}$, with no 4D spacetime indices. Note that this implies a relative factor of 2 in $\frac{1}{2} \frac{1}{5!} \mathbb{F}_{(5)}^2 = \frac{1}{5!} \mathbb{F}_{(5)}^{\text{int}^2}$.

By substituting Eq. (C.6) into Eq. (C.4) and adding to the result $Z^3 e^2$ times Eq. (3.14a), we obtain the desired pseudo-BPS condition Eq. (3.18).

In writing Eqs. (C.5) and (C.6), we have assumed that $H_{(3)}^2 = 0$. If this is not the case, then there is an additional term

$$-\frac{1}{2} \int \mathbb{F}_{(3)}^2 = -\frac{e^2}{3!} (\mathbb{F}_{(3)}^2)^2 - \frac{1}{5!} (H_{(3)}^2)^2 \quad (\text{C.7})$$

on the RHS of the second equation (C.5), and additional terms

$$= \frac{3}{2} e^2 \frac{1}{3!} (\mathbb{F}_{(3)}^2)^2 - \frac{1}{2} \frac{1}{3!} (H_{(3)}^2)^2 \quad (\text{C.8})$$

on the RHS of Eq. (C.6).

Here $\mathbb{F}_{(3)}^2 = C_0 H_{(3)}^2$, where $C_{(0)}$ is a local section of a $U(1)$ bundle (such that $F_{(1)} = dC_{(0)}$) and cannot be written as a single-valued function. Since $\mathbb{F}_{(3)}^2$ and $H_{(3)}^2$ are gauge-invariant globally-defined three-forms, this implies that $F_{(1)}$ and $H_{(3)}^2$ cannot simultaneously be nonzero. Therefore, $F_{(3)}^2 = 0$. However, even when $F_{(1)} = 0$ and $H_{(3)}^2 \neq 0$, the second term in Eq. (C.8) still presents a problem; since it is negative, we obtain an additional negative term on the RHS of Eq. (3.18), and can no longer deduce the Hodge duality relations (3.20). Therefore, we restrict to the case $H_{(3)}^2 = 0$, in which we know how to proceed.

Appendix D. IIB Fermion Variations

The string frame fermion variations to linear order in the spinors are

$$\begin{aligned} \delta_{L;R} \psi_{L;R} &= \frac{1}{2} \not{\partial} \psi_{L;R} - \frac{1}{2} \not{H}_{(3)} \psi_{L;R}; \\ \delta_{R;L} \psi_{L;R} &= \frac{1}{2} e^{-\frac{1}{2} \not{F}_{(1)}} \not{H}_{(3)} \psi_{R;L} + \frac{1}{2} \not{F}_{(3)} \psi_{R;L}; \\ \delta_{L;R} \psi_{L;R M} &= \psi_{L;R M} - \frac{1}{4} \not{H}_{(3)M} \psi_{L;R}; \\ \delta_{R;L} \psi_{L;R M} &= \frac{1}{8} e^{-\frac{1}{2} \not{F}_{(1)}} \not{F}_{(3)} \not{F}_{(5) M} \psi_{R;L} : \end{aligned} \quad (\text{D.1})$$

Here, subscripts to the left (right) of a gamma are correlated with upper (lower) sign choices. The subscripts L and R indicate whether the spin content of a fermion comes from the left-moving or right-moving sector of the worldsheet conformal field theory. In

this appendix, we follow Hassan [75], except for a sign difference in the chirality of all of the 10D spinors, due to opposite conventions for $\epsilon^{(10)}$. We take

$$\epsilon^{(10)}_{L;R} = \epsilon_{L;R}; \quad \epsilon^{(10)} = +; \quad \text{and} \quad \epsilon^{(10)}_{L;R M} = \epsilon_{L;R M}; \quad (\text{D } 2)$$

where $\epsilon^{(10)}$ is given by Eq. (A.8). Slashes denote contraction with Dirac matrices as defined in Eq. (A.6). Also,

$$r_M{}_{L;R} = \mathbb{Q}_M + \frac{1}{4} w_{M ab}{}^{ab}{}_{L;R}; \quad (\text{D } 3)$$

where $w_{M ab}$ is the spin connection, and

$$\mathbb{F}^{(n)M_1 \dots M_n} = n \mathbb{Q}_{M_1} C^{(n-1)M_2 \dots M_n]} \frac{n!}{3!(n-3)!} H^{(3)M_1 M_2 M_3} C^{(n-3)M_4 \dots M_n]}; \quad (\text{D } 4)$$

The last equation means that

$$\begin{aligned} \mathbb{F}^{(n)} &= F^{(n)} \wedge H^{(3)} \wedge C^{(n-3)}; \\ d\mathbb{F}^{(n)} &= H^{(3)} \wedge F^{(n-2)} + \text{local sources}; \end{aligned} \quad (\text{D } 5)$$

By defining the complex quantities

$$\mathbb{F} = F_L + i F_R; \quad \mathbb{Q} = Q_L + i Q_R; \quad \mathbb{H} = H_L + i H_R; \quad \text{and} \quad \mathbb{M} = M_L + i M_R; \quad (\text{D } 6)$$

we can equivalently write the fermion variations as

$$\begin{aligned} \delta \psi &= \frac{1}{2} i \not{\mathbb{Q}} \not{\epsilon} + \frac{1}{4} e \not{\mathbb{F}}^{(1)} + \frac{1}{4} e \not{\mathbb{F}}^{(3)} \not{\mathbb{H}}^{(3)}; \\ \delta \chi_M &= r_M + \frac{i}{8} e \not{\mathbb{F}}^{(1)} \not{M} + \frac{1}{2} \not{\mathbb{F}}^{(5)} \not{M} + \frac{i}{8} e \not{\mathbb{F}}^{(3)} \not{M} - 2 \not{\mathbb{H}}^{(3)} \not{M}; \end{aligned} \quad (\text{D } 7)$$

In terms of the axion-dilaton and complex flux,⁴¹

$$\begin{aligned} d\text{dil} &= C_{(0)} + i e; \\ G_{(3)} &= \mathbb{F}_{(3)} - i e H_{(3)} = F_{(3)} \wedge H_{(3)}; \end{aligned} \quad (\text{D } 8)$$

⁴¹ Here, we have given the axion-dilaton for type IIB supergravity with no orientifold projection. This is also the axion-dilaton for orientifolds with 03 or 07 planes and Becker-type spinor constraints. However, as discussed in Sec. 4.2, a different quantity plays the role of the axion-dilaton for 05 orientifolds, since the zero mode of $C_{(0)}$ is projected out. For the type I orientifold with 09 planes, $d\text{dil} = a + i e$, where a is the scalar dual to $C_{(2)}$.

this becomes

$$\begin{aligned}
e^{\mu\nu} &= \frac{1}{2} \mathcal{G}^{\text{dil}} + \frac{1}{4} \mathcal{G}^{(3)} \\
e^{\left(\begin{smallmatrix} \mu \\ M \end{smallmatrix} \right. \left. \begin{smallmatrix} \nu \\ \frac{1}{4} M \end{smallmatrix} \right)} &= e^{\left(\begin{smallmatrix} \mu \\ r_M \end{smallmatrix} \right. \left. \begin{smallmatrix} \nu \\ \frac{1}{8} M \end{smallmatrix} \right)} + \frac{1}{4} F^{(1)M} + \frac{1}{16} \mathcal{F}^{(5)M} \\
&\quad + \frac{1}{8} \mathcal{G}^{(3)M} + \frac{1}{16} M \mathcal{G}^{(3)} :
\end{aligned} \tag{D.9}$$

Here, we have assumed a real representation of the Dirac matrices. We have also used the relation

$$\mathcal{H}^{(3)}; M \mathcal{G} = 2\mathcal{H}^{(3)M} \tag{D.10}$$

in deriving (D.9).

When expressed in terms of the complex quantities \mathcal{G}^{dil} and $\mathcal{G}^{(3)}$, the equations simplify in Einstein frame. Using a prime to indicate the latter, the relations between string frame and Einstein frame are

$$\begin{aligned}
g_{MN}^0 &= e^{-2} g_{MN}; & \mathcal{G}_M^0 &= e^{-4} \mathcal{G}_M; & r_M^0 &= r_M + \frac{1}{8} M^N \mathcal{G}_N; \\
\mathcal{F}^0 &= e^{-8}; & \mathcal{H}^0 &= e^{-8}; & \mathcal{G}^0 &= e^{-8} \left(\mathcal{G} + \frac{1}{4} M \right);
\end{aligned} \tag{D.11}$$

Therefore, the Einstein frame analog of Eqs. (D.9) is

$$\begin{aligned}
\mathcal{G}^0 &= \frac{1}{2} e^{\mathcal{G}^{\text{dil}0}} + \frac{1}{4} e^{-2} \mathcal{G}^{(3)0}; \\
\mathcal{G}_M^0 &= r_M^0 + \frac{1}{4} e^{F^{(1)M}} + \frac{1}{16} \mathcal{F}^{(5)M} + e^{-2} \left(\frac{1}{8} \mathcal{G}^{(3)M} + \frac{1}{16} M \mathcal{G}^{(3)0} \right);
\end{aligned} \tag{D.12}$$

where a slash and prime denotes the analog of Eq. (A.6) with $\mathcal{G}_M \rightarrow \mathcal{G}_M^0$.

Eqs. (D.12) give the Einstein frame fermion variations in the usual string theory conventions, in which $F^{(p)}$, $\mathcal{H}^{(3)}$, and $\mathcal{G}^{(3)}$ are the same in string frame and Einstein frame. In the supergravity literature, the quantity $\mathcal{G}^{(3)}$ is given a slightly different definition. The relation between that definition and the one given in Eqs. (D.8) has been discussed elsewhere [6,75].

Appendix E. The Meaning of W_{GVW} in the $O3$ Orientifold with Internal T^6

As mentioned in Sec. 5.1, one subtlety in our description of metric moduli stabilization for the $O3$ orientifold based on internal T^6 is that not all of the (18 real) g_{ij} and (9 real) g_{ij} correspond to the (21 real) physical metric moduli.

In contrast to the case for a proper Calabi-Yau threefold, there is not a one-to-one correspondence

$$\begin{aligned} \int_j D_{i_j} &= \int^{(1;2)} \quad (\text{sum on } i; j), \\ \mathfrak{g}_1 &= \int_{(k_j)}^{k_1} \int^{(1;2)}_{k_1 j_1} \quad ; \end{aligned} \tag{E 1}$$

between deformations of complex structure ((1,2) forms or $T^{(0;1)}$ -valued (1,0) forms) and deformations of g_{ij} . The reason is that for a manifold such as T^6 with nontrivial $H^{(0;1)}$, there exist nonprimitive (1,2) forms

$$\int^{(1;2)} = J \wedge \int^{(0;1)}; \tag{E 2}$$

that generate vanishing metric deformations. For a T^6 , the three such forms are $J \wedge dz^i$. Correspondingly, there are three complex unphysical degrees of freedom in \int_j that do not correspond to metric degrees of freedom.

This can be understood in the supergravity theory as follows. In the absence of flux, the theory has $N = 4$ supersymmetry. The fluxes break this to $N < 4$ supersymmetry, but the breaking should still be described within the formalism of $N = 4$ gauged supergravity. Unlike global $N = 4$ supersymmetry, $N = 1$ supergravity cannot be cleanly described in $N = 1$ language, due to the lack of a simple on-shell formulation for $N = 1$ gravitino multiplets with a single auxiliary field analogous to the D or F fields for vector or chiral multiplets.⁴² Nevertheless, upon choosing an $N = 1$ subalgebra of $N = 4$, one can still organize all fields into $N = 1$ multiplets and decompose the bosonic parts of all supersymmetry variations into D-terms, F-terms, and "other-terms." In this sense, we can still define an $N = 1$ superpotential, whose variation gives the F-terms. However, this superpotential is part of a family of superpotentials parametrized by a choice of the $N = 1$ subalgebra. Equivalently, it is parametrized by a choice of a point in the space

$$Y = \frac{SU(4)_R}{U(1)_R \times SU(3)}; \quad \dim_{\mathbb{C}} Y = 3; \tag{E 3}$$

of embeddings of the $U(1)_R$ -symmetry of $N = 1$ in the $SU(4)_R$ -symmetry of $N = 4$. Here, the $SU(3)$ is the commutant of $U(1)_R$ in $SU(4)_R$. For T^6 , W_{GVW} is exactly this type of family of superpotentials, parametrized by the three complex unphysical degrees

⁴² Here, we have in mind an on-shell multiplet whose on-shell dynamical degrees of freedom are a spin 3/2 field and a spin 1 field. For work on on-shell gravitino multiplets, see Ref. [84].

of freedom in i_j . To obtain the F-term s, W_{GVW} should only be varied with respect to the physical degrees of freedom. The F-term s impose the conditions

$$G_{(3)} = (1,2)\text{-nonprimitive} + (2;1); \quad (\text{E.4})$$

where the (1,2)-nonprimitive component is of the form (E.2).

On the other hand, in an operational sense, varying W_{GVW} with respect to d_{il} and all of the i_j , including unphysical degrees of freedom, is a convenient thing to do: it is more easily implemented in practice than varying with respect to only physical deformations, and this seemingly unphysical procedure imposes exactly the (2,1) condition on $G_{(3)}$, which is still a subset of the supersymmetry conditions. The philosophy of Ref. [11] was simply to use W_{GVW} in this seemingly unphysical way, as a convenient tool for imposing the (2,1) condition.

One disadvantage of this approach is that it is not always clear how many physical moduli there are among the redundant set of all i_j and $g_{i\bar{j}}$ unixed by the supersymmetry conditions. For the examples that we present, this is an issue only in the $N = 2$ case. (For our $N = 1$ and $N = 3$ examples, d_{il} and all complex structure moduli are fixed, and all of the unixed Kahler moduli are physical.) We avoid the problem by choosing a noncanonical decomposition of unixed metric moduli into Kahler and complex structure moduli in Sec. 6.4. The choice is natural from the point of view of the dual 05 orientifold of Sec. 6.2. The choice corresponds exactly to the choice of $\epsilon_+ = 0$ or $\epsilon_- = 0$ in Sec 4.2, which leads to an integrable complex structure possessing a fiber base decomposition as in Eq. (4.21).

Appendix F. T-Duality Map for RR Potentials

The T-duality map for the RR potentials $C_{(p)}$ is analogous to Eq. (5.28). The relations between components in the $\epsilon; dx^m$ basis of the 05 orientifold and the $\epsilon^0; dx^m$ basis of the 03 orientifold are

$$\begin{aligned} C_{(p)}^0 &= \frac{1}{2} C_{(p+2)}^{02} = (2^-)^2 \epsilon^0; \\ C_{(p)}^1 &= C_{(p)}^{01}; \\ C_{(p)}^2 &= C_{(p-2)}^{00} (2^-)^2 \epsilon^0; \end{aligned} \quad (\text{F.1})$$

We assume that $H_{(3)\underline{m}}^0 = 0$ so that the T-duality exists, but retain $F_{(3)\underline{mnr}}^0$. (The equations of motion ultimately require that $F_{(3)\underline{mnr}}^0 = 0$.)

In the $O(3)$ orientifold, a choice of gauge for the potentials leading to the quantized flux (5.4) is

$$\begin{aligned} B_{(2)}^{bg} &= \frac{1}{2} H_{(3)\underline{nr}}^0 x^r dx^{\wedge} dx^n + \frac{1}{3!} H_{(3)\underline{mnr}}^0 x^r dx^m \wedge dx^n; \\ C_{(2)}^{bg} &= \frac{1}{2} F_{(3)\underline{r}}^0 x^r dx^{\wedge} dx + \frac{1}{2} F_{(3)\underline{nr}}^0 x^r dx^{\wedge} dx^n + \frac{1}{3!} F_{(3)\underline{mnr}}^0 x^r dx^m \wedge dx^n; \end{aligned} \quad (F.2)$$

In the $\omega^0; dx^m$ basis (where $\omega^0 = dx + a^0_m$), this becomes

$$\begin{aligned} B_{(2)}^{bg} &= \frac{1}{2} H_{(3)\underline{nr}}^0 x^r \omega^0 \wedge dx^n + \frac{1}{3!} H_{(3)\underline{mnr}}^0 \frac{1}{2} H_{(3)\underline{nr}}^0 a^0_m x^r dx^m \wedge dx^n; \\ C_{(2)}^{bg} &= \frac{1}{2} F_{(3)\underline{r}}^0 x^r \omega^0 \wedge \omega^0 + \frac{1}{2} F_{(3)\underline{nr}}^0 F_{(3)\underline{r}}^0 a^0_n x^r \omega^0 \wedge dx^n \\ &\quad + \frac{1}{3!} F_{(3)\underline{mnr}}^0 \frac{1}{2} F_{(3)\underline{nr}}^0 a^0_m + \frac{1}{2} F_{(3)\underline{r}}^0 \omega^0_m a^0_n x^r dx^m \wedge dx^n; \end{aligned} \quad (F.3)$$

In addition, there are background components of $C_{(4)}^{bg}$ with one and two ω^0 indices, such that

$$F_{(5)sm}^0 e^{ared} = dC_{(4)}^{bg} - C_{(2)}^{bg} \wedge H_{(3)}^0; \quad (F.4)$$

with $F_{(5)}^0$ given by Eq. (5.30).

The moduli are $c_{(0)}^0$ and $c_{(4)}^0 = C_{(4)}^0 - C_{(4)}^{bg}$. The latter are given by

$$\begin{aligned} C_{(4)}^0 &= \frac{1}{2!2!} C_{(4)\underline{rs}}^0 dx^{\wedge} dx^{\wedge} dx^r \wedge dx^s + \frac{1}{3!} C_{(4)\underline{nrs}}^0 dx^{\wedge} dx^n \wedge dx^r \wedge dx^s \\ &\quad + \frac{1}{4!} C_{(4)m nrs}^0 dx^m \wedge dx^n \wedge dx^r \wedge dx^s; \end{aligned} \quad (F.5)$$

In the $\omega^0; dx^m$ basis, this becomes

$$\begin{aligned} C_{(4)}^0 &= \frac{1}{2!2!} C_{(4)\underline{rs}}^0 \omega^0 \wedge \omega^0 \wedge dx^r \wedge dx^s \\ &\quad + \frac{1}{3!} C_{(4)\underline{nrs}}^0 \frac{1}{2} C_{(4)\underline{rs}}^0 a^0_n \omega^0 \wedge dx^n \wedge dx^r \wedge dx^s \\ &\quad + \frac{1}{4!} C_{(4)m nrs}^0 \frac{1}{3!} C_{(4)\underline{nrs}}^0 a^0_n + \frac{1}{2!2!} C_{(4)\underline{rs}}^0 a^0_m a^0_n dx^m \wedge dx^n \wedge dx^r \wedge dx^s; \end{aligned} \quad (F.6)$$

Applying the T-duality map (F.1), replacing $a_{(2)}^0$ with $b_{(2)}$ via Eq. (5.25a), and at the same time using the flux relations (5.31), we obtain

$$\begin{aligned} C_{(0)} &= F_{(1)m} x^m; \\ C_{(2)45} &= e_{(2)45}; \\ C_{(2)n} &= \frac{1}{2} F_{(3)nr}^{bg} x^r + b_n C_{(0)}; \\ C_{(2)mn} &= C_{(2)mn}^{bg} + e_{(2)mn}; \\ C_{(4)nrs} &= C_{(4)nrs}^{bg} + e_{(4)nrs} \wedge b_{(2)} \wedge e_{(2)nrs}; \\ C_{(4)45mn} &= F_{(5)45mnr}^{bg} x^r + b_{(2)} \wedge \frac{1}{2} F_{(3)5qp} x^p dx^q_{45mn} + \frac{1}{2} b_{(2)} \wedge b_{(2)} C_{(0)45mn}; \\ C_{(6)45m nrs} &= e_{(6)45m nrs} + b_{(2)} \wedge e_{(4)45m nrs} + \frac{1}{2} b_{(2)} \wedge b_{(2)} \wedge e_{(2)45m nrs}; \end{aligned} \quad (F.7)$$

where

$$C_{(2)rs}^{bg} = C_{(4)45rs}^{bg}; \quad C_{(4)nrs}^{bg} = C_{(4)nrs}^{bg}; \quad (F.8)$$

and where the T-duality map of RR moduli is

$$\begin{aligned} e_{(2)45} &= C_{(0)}^0; \quad e_{(2)rs} = C_{(4)45rs}^0; \\ e_{(4)nrs} &= C_{(4)nrs}^0; \quad e_{(6)456789} = C_{(4)6789}^0; \end{aligned} \quad (F.9)$$

Eq. (F.7) is of the same form as Eqs. (3.43) and (3.44), with the gauge choice

$$C_{(0)} = F_{(1)m} x^m; \quad C_{(2)n}^{bg} = \frac{1}{2} F_{(3)nr}^{bg} x^r; \quad C_{(4)45m n}^{bg} = F_{(5)45m nr}^{bg} x^r; \quad (F.10)$$

Similarly, from Eqs. (5.25b;c), the gauge choices for $A_{(2)}$ and $B_{(2)}^{bg}$ resulting from Eq. (F.2) and the T-duality map are

$$A_{(2)} = \frac{1}{2} F_{mn} x^m dx^n; \quad B_{(2)}^{bg} = \frac{1}{3!} H_{(3)mnr}^{bg} x^m dx^n \wedge dx^r; \quad (F.11)$$

One can check that the potentials (F.7) do indeed give rise to the fluxes (3.20a), (3.47), and (3.50).

Appendix G. Moduli Space Metrics

In this appendix, we discuss the kinetic terms for the moduli of the $T^6 = Z_2 \times O3$ orientifold. We restrict to the $N = 2$ flux (6.55) of Sec. 6.4 and the $N = 1$ flux (8.24) of Sec. 8.2, and work to leading order in Z .⁴³ For the $N = 3$ case, the kinetic terms were derived by Frey and Polchinski [60], and shown to describe a sigma model with target (7.37), as required by Ref. [77]. Here, we follow Frey and Polchinski, except for a minor difference in conventions,⁴⁴ and the inclusion of the axion-dilaton modulus.

⁴³ Properly treating the warp factor in the 4D kinetic terms for the 6D metric moduli is a problem that we do not attempt to address here. It was partially studied in Ref. [13], however, the treatment there neglected terms involving $\partial Z = (\partial Z = \partial g_{ab}) \partial g_{ab}$. It is currently under investigation [61].

⁴⁴ We take $x^a = x^a + 1$ on the T^6 , whereas Frey and Polchinski take $x^a = x^a + 2$. So, some coefficients differ by powers of 2. In addition, we do not absorb factors $g_s = e^{h-1}$ into the normalizations of the 4D fields, since, in the $N = 2$ case, is an unlifted modulus whose vev is not a priori determined.

The 10D string frame action for the metric, dilaton, and axion $C_{(0)}$ is

$$S_{G; C_{(0)}}^{10S} = \frac{1}{2} \int_{Z_2} d^{10}x \sqrt{-G_S} e^{-2\phi} R_{10S} + 4(\partial\phi)^2 - \frac{1}{2} (\partial C_{(0)})^2; \quad (G.1)$$

where $Z_2 = (\mathbb{Z})^6$ and $(G_S)_{MN}$ denotes the metric (5.1a). The integral runs over the Z_2 covering space $\mathbb{R}^{3;1} \times T^6$, so there is an overall factor of $1=2$ multiplying the action. In the 10D Einstein frame,

$$(G_E)_{MN} = e^{-2\phi} (G_S)_{MN}; \quad (G.2)$$

the action becomes

$$S_{G; \text{dil}}^{10E} = \frac{1}{4} \int_{Z_2} d^{10}x \sqrt{-G_E} R_{10E} - \frac{\partial_M \text{dil} \partial^M \text{dil}}{2 \text{Im}_{\text{dil}}}; \quad (G.3)$$

where now the contractions are performed using the new metric G_E . The dimensional reduction down to 4D is

$$S_{G; \text{dil}}^4 = \frac{1}{2} \int_0^{2\pi} d^4x \sqrt{-G_4} R_4 - \frac{\partial_{\text{dil}} \partial_{\text{dil}}}{2 \text{Im}_{\text{dil}}^2} - \frac{\partial_a \partial^a}{2} - \frac{1}{4} G^{ac} G^{bd} \partial_{ab} \partial_{cd}; \quad (G.4)$$

Here, $(2\pi)^6 = (\det G_6)^{1/2} = 2$, with G_4 and G_6 the restrictions of G_E to 4D and 6D, respectively. Finally, in terms of the 4D Einstein frame metric $(g_E)_{ab} = (G_4)_{ab}$ and rescaled 6D metric $\tilde{g}_{ab} = (G_6)_{ab} = (2\pi)^2 \delta_{ab}$, this becomes

$$S_{g; \text{dil}}^{4E} = \frac{1}{2} \int_0^{2\pi} d^4x \sqrt{-g} R_E - \frac{\partial_{\text{dil}} \partial_{\text{dil}}}{2 \text{Im}_{\text{dil}}^2} - \frac{1}{4} \tilde{g}^{ac} \tilde{g}^{bd} \partial_{ab} \partial_{cd}; \quad (G.5)$$

As in Ref. [60], once the moduli $c_{(4)abcd}$ and I^a are included, the total 4D Einstein-Hilbert plus moduli action is

$$S_{g; \text{dil}; c; I}^{4E} = S_g^{4E} + S_{\text{dil}}^{4E} + S_{c; I}^{4E}; \quad \text{where} \quad (G.6a)$$

$$S_g^{4E} = \frac{1}{2} \int_0^{2\pi} d^4x \sqrt{-g} R_E; \quad (G.6b)$$

$$S_{\text{dil}}^{4E} = \frac{1}{2} \int_0^{2\pi} d^4x \sqrt{-g} \frac{\partial_{\text{dil}} \partial_{\text{dil}}}{2 \text{Im}_{\text{dil}}^2}; \quad (G.6c)$$

$$S_{c; I}^{4E} = \frac{1}{2} \int_0^{2\pi} d^4x \sqrt{-g} \left[\partial_{ab} c_{abcd} \partial^a I^b + \frac{1}{4} \tilde{g}^{ac} \tilde{g}^{bd} D^{ab} D^{cd} + \partial_{ab} c_{abcd} \partial^a c^b \right]; \quad (G.6d)$$

In Eq. (G.6c),

$$I^a = I^a = (2 \dots); \quad (G.7a)$$

$$C_{(4)abcd} = (2 \dots)^4 \omega_{abcdef} \epsilon^{ef}; \quad \text{where } \epsilon_{456789} = 1; \quad (G.7b)$$

and

$$D^{ab} = \partial^{ab} + I^{[a} \partial_{I^{b]}}; \quad (G.7c)$$

The second term in D^{ab} is the analog of the gauge Chern-Simons term in $F_{(3)}$ of type I or in $H_{(3)}$ of the heterotic theories. It is well known that the sigma model (G.6d) with a, b running over D values and I running over M values parametrizes the coset $T_{D, D+M}$, where $T_{m, n}$ is the Grassmannian

$$T_{m, n} = \frac{SO(m; n)}{SO(m) \times SO(n)}; \quad (G.8)$$

up to discrete identifications [85].

Thus, for $N = 4$, with $2M = 32 D3$ branes and no flux, the scalar manifold of the low energy effective field theory is

$$M_{N=4} = \frac{SU(1;1)}{U(1)} \times \frac{SO(6;22)}{SO(6) \times SO(22)}; \quad (G.9)$$

up to discrete identifications, where the first factor is from d_{11} and the second from g_{22} . When there is nonvanishing flux, the supersymmetry is reduced, and the moduli constraints reduce this moduli space to a proper submanifold that is also a product of homogeneous spaces.

For $N = 3$ flux, the axion-dilaton and some moduli are fixed. In the w -coordinates of Sec. 8.2, the unlifted components of g_{ij} are $I^i, I^{\{, i\} = |^i, i\} = |^i$, and the above sigma model parametrizes the coset $U(3; 3+M) = U(3) \times U(3+M)$, as discussed in Refs. [77,60].

For the $N = 1$ flux of Sec. 8.2, the moduli constraints are similar. The axion-dilaton is again fixed. In the z -coordinates of Sec. 8.2, the nonvanishing components of g_{ij} are as in the $N = 3$ case in w -coordinates, but with the addition restriction that $i\}$ and $i\}$ be symmetric (imaginary and real, respectively). That is, $i\} = j\}$ and $i\} = j\}$. It would be interesting to describe this $6 + 6M$ dimensional real hypersurface in $U(3; 3+M) = U(3) \times U(3+M)$ as a coset, perhaps $U(3; 1+M) = U(3) \times U(1+M)$.

Finally, for the $N = 2$ case of Sec. 6.4, the moduli constraints are such that the scalar manifold factorizes into an $a = 4;5;6;7$ part and an $a = 8;9$ part. If there were no further constraints, the moduli space would be $SU(1;1) = U(1) \times T_{2;2+M} \times T_{4;4+M}$. The first two factors form the manifold $ST_{2;2+M}$ in the special Kahler series

$$ST_{2;m} = \frac{SU(1;1)}{U(1)} \times \frac{SO(2;n)}{SO(2) \times SO(n)} : \quad (G.10)$$

The third factor is quaternionic. Therefore, we would identify the vector multiplet moduli space with $ST_{2;2+M}$ and the hypermultiplet moduli space with $T_{4;4+M}$. However, there are additional moduli constraints that further reduce these special Kahler and quaternionic manifolds. First, there is the constraint $(m=n)(1=d_{il}) = \frac{0}{3}$, where $\frac{0}{3}$ is the complex structure modulus of ab in the $8;9$ directions. Therefore, $ST_{2;2+M}$ is further lifted to⁴⁵

$$M_V = ST_{2;1+M} : \quad (G.11)$$

If, in the $4;5;6;7$ directions, we write (cf. Eqs. (6.63a;b))

$$ab dx^a dx^b = \frac{1}{\text{Im} \frac{0}{1}} \theta^4 + \frac{0}{1} \theta^5{}^2 + \frac{2}{\text{Im} \frac{0}{2}} dx^6 + \frac{0}{2} dx^7{}^2 ; \quad (G.12)$$

with $\theta^4; \theta^5$ given by Eqs. (6.64a;b), then the remaining moduli constraints are

$$\frac{0}{1} \frac{0}{2} = 1; \quad a^4{}_7 = a^5{}_6; \quad \text{and} \quad \theta^4{}_6 = \theta^5{}_7 : \quad (G.13)$$

The hypermultiplet moduli space M_H is the hypersurface (G.13) in $T_{4;4+M}$. Since it must be quaternionic, we conclude that⁴⁶

$$M_H = T_{4;3+M} : \quad (G.14)$$

As we have already observed in Secs. 6.4 and 8.2, for classes of flux discussed in this paper, the partial breaking of $N = 4$ to $N = 1;2$ supersymmetry is such that the resulting

⁴⁵ The only other homogeneous symmetric space G/H of dimension $2(2+M)$, with M divisible by four, that is special Kahler is $U(1;2+M) = U(1) \times U(2+M)$. However, this is not a hypersurface in $ST_{2;2+M}$. For a nice review of special Kahler and hyperKahler/quaternionic geometry in connection with both locally and globally $N = 2$ supersymmetric field theories, see Ref. [86].

⁴⁶ Likewise, the only other homogeneous symmetric quaternionic manifolds of dimension $4(3+M)$ are $Sp(2M+8) = Sp(2) \times Sp(2M+6)$ and $U(2;3+M) = U(2) \times U(3+M)$. Neither of these is a hypersurface in $T_{4;4+M}$.

massive gravitini all have equal mass. It is noteworthy that for the $T^6 = Z_2 \times O_3$ orientifold, there does not exist a single proper example in the literature (that is, a choice of flux consistent with Dirac quantization and the Gauss's law constraint) in which the breaking involves massive gravitini of unequal mass. The "generic" case of partial supersymmetry breaking with unequal gravitino mass is apparently not as generic as one might have thought. It is not clear that such a case is possible. However, if it is, then the resulting mass spectra and superHiggs mechanism would be governed by the results of Ref. [17], in which the reduced moduli spaces for this case have also been worked out.

Appendix H. Uniqueness of Flux in Sec. 8.2

In this appendix we prove the claim that the flux (8.23) is the unique choice of supersymmetric flux in the class (8.17a;b) with the minimal value $N_{\text{flux}} = 12$, modulo $SL(2;Z) \times SL(2;Z)_{\text{dual}}$ equivalences. (Here $SL(2;Z)$ is the diagonal subgroup of $SL(2;Z)^3 \times SL(6;Z)$, where $SL(6;Z)$ is the duality group of T^6 , and $SL(2;Z)^3$ is the product of the duality groups of the T^2 factors in the $T^6 \rightarrow T^2 \times T^2 \times T^2$ factorization of the complex structure for this example.) For notational simplicity, we drop the primes on \hat{P} and \hat{M} that were used in Sec. 8.2 solely to denote that O_3 from O_5 quantities. Instead, primes will indicate $SL(2;Z)$ transformed quantities below. It is convenient to rewrite Eq. (7.27) in matrix form as

$$P(\tau) = \frac{1}{2} \begin{pmatrix} 1 & \tau \\ \tau & 1 \end{pmatrix} \hat{P}^{-1}; \quad \text{where } \hat{P} = \begin{pmatrix} 2l & m \\ m & 2n \end{pmatrix}; \quad (\text{H } 1)$$

Then,

$$\det \hat{P} = 4ln - m^2; \quad (\text{H } 2)$$

and under $SL(2;Z)$ transformations,

$$\hat{P} \rightarrow \hat{P}^0; \quad \text{where } \hat{P}^{-1} = \frac{1}{\det \hat{P}} \begin{pmatrix} 2n & -m \\ m & 2l \end{pmatrix}; \quad (\text{H } 3a)$$

$$\hat{P} \rightarrow \hat{P}^0; \quad \text{where } \hat{P}^0 = M^T \hat{P} M; \quad (\text{H } 3b)$$

with

$$M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2;Z); \quad (\text{H } 3c)$$

For $N_{ux} = 12$, we have $(fk - gh)(4ln - m^2) = 9$. Since $4ln - m^2 \equiv 1 \pmod{4}$, the only possible factorization is

$$fk - gh = 3; \quad 4ln - m^2 = 3: \quad (H.4)$$

Here we have used the fact that $4ln - m^2 > 0$ in order for $P(\lambda)$ to have two complex conjugate roots with nonzero imaginary part. This is required for the T^6 to be nondegenerate.

It is well known that \mathcal{H} can also be mapped into the fundamental domain (7.30) by $SL(2;Z)$ transformations. For $2F_0$, the conditions (7.32) combined with the second equality in Eq. (H.4) imply that $3 \mid 4l(n-1)$, with $n-1 > 0$. Therefore, $l = n$. From Eq. (H.4), we then have $3 = (2l + jn - j)(2l - jn - j)$. So, $l = m = n = 1$, corresponding to $\tau = \frac{1}{2} + i\frac{\sqrt{3}}{2}$ from Eq. (7.32). The two possibilities are dual via $\tau \rightarrow -\bar{\tau}$. (Only the $m = +1$ solution is actually in the fundamental domain as defined in (7.30)). Thus,

$$(l; m; n) = (1; 1; 1) \quad \text{and} \quad \tau = e^{2i\pi/3}; \quad \text{up to } SL(2;Z) \text{ duality.} \quad (H.5)$$

Now consider the equivalences on $(f; g; h; k)$. Under $SL(2;Z)_{d_{il}}$ duality,

$$A = \begin{pmatrix} f & g \\ h & k \end{pmatrix} \quad ; \quad A^0 = N A; \quad \text{where } N \in SL(2;Z)_{d_{il}}: \quad (H.6)$$

From $a = b = \frac{2}{3}(f + g)$, $c = d = \frac{2}{3}(h + k)$, together with $fk - gh = 3$, we conclude that $\gcd(f; h) = 1$. (Assume, to the contrary, that $\gcd(f; h) = 3$. Then, demanding that $a; b; c; d \in 2\mathbb{Z}$ implies that $\gcd(g; k) = 3$ and subsequently that 9 divides $fk - gh$. This contradicts $fk - gh = 3$.) Therefore, $f + h = 1$ for some $e; \nu$, so that the $SL(2;Z)_{d_{il}}$ duality $A^0 = \begin{pmatrix} f^0 & g^0 \\ h^0 & k^0 \end{pmatrix} = \begin{pmatrix} h & f \\ f & g \\ h & k \end{pmatrix}$ gives $h^0 = 1$. Next, the duality $A^{00} = \begin{pmatrix} f^{00} & g^{00} \\ h^{00} & k^{00} \end{pmatrix} = \begin{pmatrix} 0 & 1 & f^0 & g^0 \\ 1 & f^0 & 1 & k^0 \end{pmatrix}$ gives $h^{00} = 0$. Finally, the $SL(2;Z)_{d_{il}}$ duality $A^{000} = \begin{pmatrix} f^{000} & g^{000} \\ 0 & k^{000} \end{pmatrix} = \begin{pmatrix} 1 & f^0 & g^0 \\ 0 & 1 & 0 & k^0 \end{pmatrix}$ allows us to replace g^{00} with any $g^{000} + g^0 \pmod{k^{00}}$. From $\det A^{000} = \det A^{00} = \det A^0 = \det A = 3$, we have $f^{000}k^{000} = 3$. We can assume that $f^{000}; k^{000} > 0$ (via $SL(2;Z)$ duality by $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$, if necessary). There are two possible factorizations: $f^{000} = 1$ and $k^{000} = 3$, or $f^{000} = 3$ and $k^{000} = 1$. The second case is excluded by $c = d = 2k^{000} = 3 \notin 2\mathbb{Z}$. In the first case, the congruence $3a = 2(f^{000} + g^{000}) \equiv 0 \pmod{3}$ gives $g^{000} \equiv 1 \pmod{3} \equiv 1 \pmod{k^{00}}$. So, $g^{000} = 1$ and $\begin{pmatrix} f & g \\ h & k \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 0 & 3 \end{pmatrix}$ up to $SL(2;Z)_{d_{il}}$ duality. This is equivalent to

$$\begin{pmatrix} f & g \\ h & k \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \quad \text{up to } SL(2;Z)_{d_{il}} \text{ duality.} \quad (H.7)$$

Together, Eqs. (H.5) and (H.7) prove the desired uniqueness of the choice (8.23b).

References

- [1] C.M. Hull, \Superstring Compactifications With Torsion And Space-Time Supersymmetry," *Print-86-0251 (CAMBRIDGE)*.
- [2] A. Strominger, \Superstrings With Torsion," *Nucl. Phys. B 274, 253 (1986)*.
- [3] K. Dasgupta, G. Rajesh and S. Sethi, \M theory, orientifolds and G-ux," *JHEP 9908, 023 (1999) [arXiv:hep-th/9908088]*.
- [4] S. Kachru, M.B. Schulz, P.K. Tripathy and S.P. Trivedi, \New supersymmetric string compactifications," *JHEP 0303, 061 (2003) [arXiv:hep-th/0211182]*.
- [5] P. Candelas, G.T. Horowitz, A. Strominger and E. Witten, \Vacuum Configurations For Superstrings," *Nucl. Phys. B 258, 46 (1985)*.
- [6] M. Graña and J. Polchinski, \Gauge/gravity duals with holomorphic dilaton," *Phys. Rev. D 65, 126005 (2002) [arXiv:hep-th/0106014]*.
- [7] H. Verlinde, \Holography and compactification," *Nucl. Phys. B 580, 264 (2000) [arXiv:hep-th/9906182]*;
C.S. Chan, P.L. Paul and H. Verlinde, \A note on warped string compactification," *Nucl. Phys. B 581, 156 (2000) [arXiv:hep-th/0003236]*.
- [8] I.R. Klebanov and M.J. Strassler, \Supergravity and a coning gauge theory: Duality cascades and SB-resolution of naked singularities," *JHEP 0008, 052 (2000) [arXiv:hep-th/0007191]*.
- [9] S.B. Giddings, S. Kachru and J. Polchinski, \Hierarchies from fluxes in string compactifications," *Phys. Rev. D 66, 106006 (2002) [arXiv:hep-th/0105097]*.
- [10] L. Randall and R. Sundrum, \A large mass hierarchy from a small extra dimension," *Phys. Rev. Lett. 83, 3370 (1999) [arXiv:hep-ph/9905221]*;
L. Randall and R. Sundrum, \An alternative to compactification," *Phys. Rev. Lett. 83, 4690 (1999) [arXiv:hep-th/9906064]*.
- [11] S. Kachru, M.B. Schulz and S. Trivedi, \Moduli stabilization from fluxes in a simple IIB orientifold," *JHEP 0310, 007 (2003) [arXiv:hep-th/0201028]*.
- [12] P.K. Tripathy and S.P. Trivedi, \Compactification with flux on K3 and tori," *JHEP 0303, 028 (2003) [arXiv:hep-th/0301139]*;
A. Giryavets, S. Kachru, P.K. Tripathy and S.P. Trivedi, \Flux compactifications on Calabi-Yau threefolds," *JHEP 0404, 003 (2004) [arXiv:hep-th/0312104]*.
- [13] O. DeWolfe and S.B. Giddings, \Scales and hierarchies in warped compactifications and brane worlds," *Phys. Rev. D 67, 066008 (2003) [arXiv:hep-th/0208123]*.
- [14] J. Polchinski and A. Strominger, \New Vacua for Type II String Theory," *Phys. Lett. B 388, 736 (1996) [arXiv:hep-th/9510227]*;
T.R. Taylor and C. Vafa, \RR flux on Calabi-Yau and partial supersymmetry breaking," *Phys. Lett. B 474, 130 (2000) [arXiv:hep-th/9912152]*;
P. Mayr, \On supersymmetry breaking in string theory and its realization in brane

- worlds," Nucl. Phys. B 593, 99 (2001) [arXiv:hep-th/0003198];
- B. R. Greene, K. Schalm and G. Shiu, "Warped compactifications in M and F theory," Nucl. Phys. B 584, 480 (2000) [arXiv:hep-th/0004103];
- G. Curio, A. Klemm, D. Lust and S. Theisen, "On the vacuum structure of type II string compactifications on Calabi-Yau spaces with H-fluxes," Nucl. Phys. B 609, 3 (2001) [arXiv:hep-th/0012213];
- G. Curio, A. Klemm, B. Kors and D. Lust, "Fluxes in heterotic and type II string compactifications," Nucl. Phys. B 620, 237 (2002) [arXiv:hep-th/0106155];
- G. Curio, B. Kors and D. Lust, "Fluxes and branes in type II vacua and M-theory geometry with G_2 and Spin(7) holonomy," Nucl. Phys. B 636, 197 (2002) [arXiv:hep-th/0111165];
- J. Louis and A. Micu, "Type II theories compactified on Calabi-Yau threefolds in the presence of background fluxes," Nucl. Phys. B 635, 395 (2002) [arXiv:hep-th/0202168];
- K. Becker, M. Becker, M. Haack and J. Louis, "Supersymmetry breaking and α' -corrections to flux induced potentials," JHEP 0206, 060 (2002) [arXiv:hep-th/0204254];
- A. R. Frey and A. Mazumdar, "3-form induced potentials, dilaton stabilization, and running moduli," Phys. Rev. D 67, 046006 (2003) [arXiv:hep-th/0210254];
- R. Blumenhagen, D. Lust and T. R. Taylor, "Moduli stabilization in chiral type IIB orientifold models with fluxes," Nucl. Phys. B 663, 319 (2003) [arXiv:hep-th/0303016];
- M. Berg, M. Haack and B. Kors, "An orientifold with fluxes and branes via T-duality," Nucl. Phys. B 669, 3 (2003) [arXiv:hep-th/0305183];
- S. P. de Alwis, "On potentials from fluxes," Phys. Rev. D 68, 126001 (2003) [arXiv:hep-th/0307084];
- J. F. G. Cascales and A. M. Uranga, "Chiral 4d string vacua with D-branes and moduli stabilization," arXiv:hep-th/0311250;
- J. F. G. Cascales, M. P. Garcia del Moral, F. Quevedo and A. M. Uranga, "Realistic D-brane models on warped throats: Fluxes, hierarchies and moduli stabilization," JHEP 0402, 031 (2004) [arXiv:hep-th/0312051];
- D. Lust, S. Reert and S. Stieberger, "Flux-induced Soft Supersymmetry Breaking in Chiral Type IIB Orientifolds with D3/D7-Branes," arXiv:hep-th/0406092.
- [15] A. R. Frey, "Warped strings: Self-dual flux and contemporary compactifications," arXiv:hep-th/0308156.
- [16] G. Dall'Agata, "Type IIB supergravity compactified on a Calabi-Yau manifold with H-fluxes," JHEP 0111, 005 (2001) [arXiv:hep-th/0107264];
- L. Andrianopoli, R. D'Auria, S. Ferrara and M. A. Lledo, "Super Higgs effect in extended supergravity," Nucl. Phys. B 640, 46 (2002) [arXiv:hep-th/0202116];
- L. Andrianopoli, R. D'Auria, S. Ferrara and M. A. Lledo, "Gauging of flat groups in four dimensional supergravity," JHEP 0207, 010 (2002) [arXiv:hep-th/0203206];

- L. Andrianopoli, R. D'Auria, S. Ferrara and M. A. Lledo, "Duality and spontaneously broken supergravity in flat backgrounds," Nucl. Phys. B 640, 63 (2002) [arXiv:hep-th/0204145];
- R. D'Auria, S. Ferrara and S. Vaula, "N = 4 gauged supergravity and a IIB orientifold with fluxes," New J. Phys. 4, 71 (2002) [arXiv:hep-th/0206241];
- R. D'Auria, S. Ferrara, M. A. Lledo and S. Vaula, "No-scale N = 4 supergravity coupled to Yang-Mills: The scalar potential and super Higgs effect," Phys. Lett. B 557, 278 (2003) [arXiv:hep-th/0211027];
- S. Ferrara, "Duality, gauging and super Higgs effect in string and M-theory," arXiv:hep-th/0211116;
- L. Andrianopoli, R. D'Auria, S. Ferrara and M. A. Lledo, "N = 2 super-Higgs, N = 1 Poincare vacua and quaternionic geometry," JHEP 0301, 045 (2003) [arXiv:hep-th/0212236];
- C. Angelantonj, R. D'Auria, S. Ferrara and M. Trigiante, "K3 T²=Z₂ orientifolds with fluxes, open string moduli and critical points," Phys. Lett. B 583, 331 (2004) [arXiv:hep-th/0312019];
- M. Grana, T. W. Grimm, H. Jockers and J. Louis, "Soft supersymmetry breaking in Calabi-Yau orientifolds with D-branes and fluxes," arXiv:hep-th/0312232;
- T. W. Grimm and J. Louis, "The effective action of N = 1 Calabi-Yau orientifolds," arXiv:hep-th/0403067.
- [17] R. D'Auria, S. Ferrara, F. Gargiulo, M. Trigiante and S. Vaula, "N = 4 supergravity Lagrangian for type IIB on T⁶=Z₂ in presence of fluxes and D3-branes," JHEP 0306, 045 (2003) [arXiv:hep-th/0303049].
- [18] S. Ferrara and M. Porrati, "N = 1 no-scale supergravity from IIB orientifolds," Phys. Lett. B 545, 411 (2002) [arXiv:hep-th/0207135];
- L. Andrianopoli, R. D'Auria, S. Ferrara and M. A. Lledo, "Gauged extended supergravity without cosmological constant: No-scale structure and supersymmetry breaking," Mod. Phys. Lett. A 18, 1001 (2003) [arXiv:hep-th/0212141].
- [19] A. Lawrence and J. McGreevy, "Local string models of soft supersymmetry breaking," arXiv:hep-th/0401034.
- [20] L. Susskind, "The anthropic landscape of string theory," arXiv:hep-th/0302219;
- M. R. Douglas, "The statistics of string/M theory vacua," JHEP 0305, 046 (2003) [arXiv:hep-th/0303194];
- S. Ashok and M. R. Douglas, "Counting flux vacua," arXiv:hep-th/0307049;
- T. Banks, M. Dine and E. Gorbatov, "Is there a string theory landscape?," arXiv:hep-th/0309170;
- M. R. Douglas, "Statistics of string vacua," arXiv:hep-ph/0401004;
- M. Dine, "Is there a string theory landscape: Some cautionary notes," arXiv:hep-th/0402101;

- M . R . D ouglas, B . Shi m an and S . Zelditch, \C ritical points and supersym m etric vacua," arX iv:m ath .cv/0402326;
- F . D enef and M . R . D ouglas, \D istributions of flux vacua," arX iv:hep-th/0404116;
- A . G iryavets, S . K achru and P . K . Tripathy, \O n the taxonomy of flux vacua," arX iv:hep-th/0404243;
- L . Susskind, \Supersym m etry breaking in the anthropic landscape," arX iv:hep-th/0405189;
- M . R . D ouglas, \Statistical analysis of the supersym m etry breaking scale," arX iv:hep-th/0405279.
- [21] S . K achru, R . K allos, A . Linde and S . P . Trivedi, \D e Sitter vacua in string theory," Phys.Rev.D 68, 046005 (2003) [arX iv:hep-th/0301240].
- [22] F . D enef, M . R . D ouglas and B . Florea, \B uilding a better racetrack," arX iv:hep-th/0404257.
- [23] D . Robbins and S . Sethi, \A barren landscape," arX iv:hep-th/0405011.
- [24] A . Saltm an and E . Silverstein, \The scaling of the no-scale potential and de Sitter m odel building," arX iv:hep-th/0402135.
- [25] K . D asgupta, C . H erdeiro, S . H irano and R . K allos, \D 3/D 7 in ationary m odel and M -theory," Phys.Rev.D 65, 126002 (2002) [arX iv:hep-th/0203019];
- M . Fabinger and E . Silverstein, \D -Sitter space: Causal structure, therm odynamics, and entropy," arX iv:hep-th/0304220;
- A . R . Frey, M . Lippert and B . W illiam s, \The fall of stringy de Sitter," Phys.Rev.D 68, 046008 (2003) [arX iv:hep-th/0305018];
- S . K achru, R . K allos, A . Linde, J . M aldacena, L . M cAllister and S . P . Trivedi, \T owards in ation in string theory," JCAP 0310, 013 (2003) [arX iv:hep-th/0308055];
- C . P . Burgess, R . K allos and F . Q uevedo, \de Sitter string vacua from supersym m etric D -term s," JHEP 0310, 056 (2003) [arX iv:hep-th/0309187];
- E . Silverstein and D . Tong, \Scalar speed lim its and cosm ology: A cceleration from D -cceleration," arX iv:hep-th/0310221;
- J . P . H su, R . K allos and S . P rokushkin, \O n brane in ation w ith volum e stabiliza-tion," JCAP 0312, 009 (2003) [arX iv:hep-th/0311077];
- A . Buchel and R . Roiban, \In ation in warped geom etries," arX iv:hep-th/0311154;
- F . K oyam a, Y . Tachikawa and T . W atari, \Supergravity analysis of hybrid in ation m odel from D 3-D 7 system ," Phys.Rev.D 69, 106001 (2004) [arX iv:hep-th/0311191];
- P . G . C am ara, L . E . Ibanez and A . M . U ranga, \F lux-induced SU SY -breaking soft term s," arX iv:hep-th/0311241;
- H . F irouzjahi and S . H . H . Tye, \C loser towards in ation in string theory," Phys.Lett.B 584, 147 (2004) [arX iv:hep-th/0312020];
- A . Buchel, \O n e ffective action of string theory flux com pactifications," Phys.Rev.D 69, 106004 (2004) [arX iv:hep-th/0312076];

- R. Brustein and S. P. de Alwis, "Moduli potentials in string compactifications with fluxes: Mapping the discretuum," *arXiv:hep-th/0402088*;
- L. Kofman, A. Linde, X. Liu, A. Maloney, L. McAllister and E. Silverstein, "Beauty is attractive: Moduli trapping at enhanced symmetry points," *arXiv:hep-th/0403001*;
- O. DeWolfe, S. Kachru and H. Verlinde, "The giant in AdS," *arXiv:hep-th/0403123*;
- N. Iizuka and S. P. Trivedi, "An inflationary model in string theory," *arXiv:hep-th/0403203*;
- M. Alishahiha, E. Silverstein and D. Tong, "DBI in the sky," *arXiv:hep-th/0404084*;
- M. Berg, M. Haack and B. Kors, "Loop corrections to volume moduli and inflation in string theory," *arXiv:hep-th/0404087*;
- A. Buchel and A. Ghodsi, "Brane world inflation," *arXiv:hep-th/0404151*.
- [26] B. de Wit, D. J. Raine and N. D. Hari Dass, "Residual Supersymmetry of Compactified $D = 10$ Supergravity," *Nucl. Phys. B* 283, 165 (1987);
- J. Maldacena and C. Nunez, "Supergravity description of field theories on curved manifolds and a no go theorem," *Int. J. Mod. Phys. A* 16, 822 (2001) [*arXiv:hep-th/0007018*];
- S. Ivanov and G. Papadopoulos, "A no-go theorem for string warped compactifications," *Phys. Lett. B* 497, 309 (2001) [*arXiv:hep-th/0008232*].
- [27] J. P. Gauntlett, D. Martelli and D. Waldram, "Superstrings with intrinsic torsion," *arXiv:hep-th/0302158*.
- [28] J. P. Gauntlett, N. Kim, D. Martelli and D. Waldram, "Fivebranes wrapped on SLAG three-cycles and related geometry," *JHEP* 0111, 018 (2001) [*arXiv:hep-th/0110034*];
- J. P. Gauntlett, D. Martelli, S. Pakis and D. Waldram, "G-structures and wrapped NS5-branes," *arXiv:hep-th/0205050*.
- [29] G. L. Cardoso, G. Curio, G. Dall'Agata, D. Lust, P. Manousselis and G. Zoupanos, "Non-Kaehler string backgrounds and their ve torsion classes," *Nucl. Phys. B* 652, 5 (2003) [*arXiv:hep-th/0211118*].
- [30] K. Becker and M. Becker, "M-Theory on Eight-Manifolds," *Nucl. Phys. B* 477, 155 (1996) [*arXiv:hep-th/9605053*].
- [31] C. Vafa, "Evidence for F-Theory," *Nucl. Phys. B* 469, 403 (1996) [*arXiv:hep-th/9602022*];
- S. Sethi, C. Vafa and E. Witten, "Constraints on low-dimensional string compactifications," *Nucl. Phys. B* 480, 213 (1996) [*arXiv:hep-th/9606122*];
- K. Dasgupta and S. Mukhi, "A note on low-dimensional string compactifications," *Phys. Lett. B* 398, 285 (1997) [*arXiv:hep-th/9612188*].
- [32] S. Gurrieri, J. Louis, A. Micu and D. Waldram, "Mirror symmetry in generalized Calabi-Yau compactifications," *Nucl. Phys. B* 654, 61 (2003) [*arXiv:hep-th/0211102*].
- [33] J. P. Gauntlett and S. Pakis, "The geometry of $D = 11$ Killing spinors," *JHEP* 0304, 039 (2003) [*arXiv:hep-th/0212008*].

- [34] D. Martelli and J. Sparks, "G-structures, fluxes and calibrations in M-theory," *Phys. Rev. D* **68**, 085014 (2003) [[arXiv:hep-th/0306225](#)].
- [35] J. P. Gauntlett, J. B. Gutowski and S. Pakis, "The geometry of D = 11 null Killing spinors," *JHEP* **0312**, 049 (2003) [[arXiv:hep-th/0311112](#)].
- [36] J. P. Gauntlett, D. Martelli, J. Sparks and D. Waldram, "Supersymmetric AdS₅ solutions of M-theory," [arXiv:hep-th/0402153](#).
- [37] P. Kaste, R. Minasian, M. Petrini and A. Tomasiello, "Kaluza-Klein bundles and manifolds of exceptional holonomy," *JHEP* **0209**, 033 (2002) [[arXiv:hep-th/0206213](#)];
P. Kaste, R. Minasian, M. Petrini and A. Tomasiello, "Nontrivial RR two-form field strength and SU(3)-structure," *Fortsch. Phys.* **51**, 764 (2003) [[arXiv:hep-th/0301063](#)];
P. Kaste, R. Minasian and A. Tomasiello, "Supersymmetric M-theory compactifications with fluxes on seven-manifolds and G-structures," *JHEP* **0307**, 004 (2003) [[arXiv:hep-th/0303127](#)].
- [38] S. Fidanza, R. Minasian and A. Tomasiello, "Mirror symmetric SU(3)-structure manifolds with NS fluxes," [arXiv:hep-th/0311122](#).
- [39] T. Z. Husain, "M2-branes wrapped on holomorphic curves," *JHEP* **0312**, 037 (2003) [[arXiv:hep-th/0211030](#)];
T. Z. Husain, "That's a wrap!," *JHEP* **0304**, 053 (2003) [[arXiv:hep-th/0302071](#)].
- [40] M. J. Du and J. T. Liu, "Hidden spacetime symmetries and generalized holonomy in M-theory," *Nucl. Phys. B* **674**, 217 (2003) [[arXiv:hep-th/0303140](#)];
C. Hull, "Holonomy and symmetry in M-theory," [arXiv:hep-th/0305039](#);
G. Papadopoulos and D. Tsimpis, "The holonomy of the supercovariant connection and Killing spinors," *JHEP* **0307**, 018 (2003) [[arXiv:hep-th/0306117](#)];
G. Papadopoulos and D. Tsimpis, "The holonomy of IIB supercovariant connection," *Class. Quant. Grav.* **20**, L253 (2003) [[arXiv:hep-th/0307127](#)];
A. Batrachenko, M. J. Du, J. T. Liu and W. Y. Wen, "Generalized holonomy of M-theory vacua," [arXiv:hep-th/0312165](#);
A. Batrachenko and W. Y. Wen, "Generalized holonomy of supergravities with 8 real supercharges," [arXiv:hep-th/0402141](#).
- [41] G. Dall'Agata and N. Prezas, "N = 1 geometries for M-theory and type IIA strings with fluxes," *Phys. Rev. D* **69**, 066004 (2004) [[arXiv:hep-th/0311146](#)].
- [42] G. Dall'Agata, "On supersymmetric solutions of type IIB supergravity with general fluxes," [arXiv:hep-th/0403220](#).
- [43] K. Behrndt and C. Jeschek, "Fluxes in M-theory on 7-manifolds and G-structures," *JHEP* **0304**, 002 (2003) [[arXiv:hep-th/0302047](#)];
K. Behrndt and M. Cvetič, "Supersymmetric intersecting D6-branes and fluxes in massive type IIA string theory," *Nucl. Phys. B* **676**, 149 (2004) [[arXiv:hep-th/0308045](#)];
K. Behrndt and C. Jeschek, "Fluxes in M-theory on 7-manifolds: G-structures and superpotential," [arXiv:hep-th/0311119](#);

- K. Behrndt and M. Cvetič, "General $N = 1$ supersymmetric flux vacua of (massive) type IIA string theory," arXiv:hep-th/0403049.
- [44] A. R. Frey, "Notes on $SU(3)$ structures in type IIB supergravity," arXiv:hep-th/0404107.
- [45] A. Gray and L. M. Hervella, "The sixteen classes of almost Hermitian manifolds and their linear invariants," *Ann. Math. Pura Appl.* 123 35 (1980);
S. Chiossi and S. Salamon, "The Intrinsic Torsion of $SU(3)$ and G_2 Structures," in *Differential Geometry, Valencia 2001*, World Sci. Publishing, River Edge, NJ, 115 (2002) [arXiv:math.DG/0202282].
- [46] J. Gutowski, G. Papadopoulos and P. K. Townsend, "Supersymmetry and generalized calibrations," *Phys. Rev. D* 60, 106006 (1999) [arXiv:hep-th/9905156];
J. Gutowski and G. Papadopoulos, "AdS calibrations," *Phys. Lett. B* 462, 81 (1999) [arXiv:hep-th/9902034];
J. Gutowski, "Generalized calibrations," arXiv:hep-th/9909096.
- [47] G. L. Cardoso, G. Curio, G. Dall'Agata and D. Lust, "BPS action and superpotential for heterotic string compactifications with fluxes," *JHEP* 0310, 004 (2003) [arXiv:hep-th/0306088].
- [48] K. Becker and K. Dasgupta, "Heterotic strings with torsion," *JHEP* 0211, 006 (2002) [arXiv:hep-th/0209077];
K. Becker, M. Becker, K. Dasgupta and P. S. Green, "Compactifications of heterotic theory on non-Kähler complex manifolds. I," *JHEP* 0304, 007 (2003) [arXiv:hep-th/0301161];
K. Becker, M. Becker, K. Dasgupta and S. Prokushkin, "Properties of heterotic vacua from superpotentials," *Nucl. Phys. B* 666, 144 (2003) [arXiv:hep-th/0304001].
- [49] K. Becker, M. Becker, P. S. Green, K. Dasgupta and E. Sharpe, "Compactifications of heterotic strings on non-Kähler complex manifolds. II," arXiv:hep-th/0310058.
- [50] E. Goldstein and S. Prokushkin, "Geometric model for complex non-Kähler manifolds with $SU(3)$ structure," arXiv:hep-th/0212307.
- [51] R. Gopakumar and C. Vafa, "On the gauge theory/geometry correspondence," *Adv. Theor. Math. Phys.* 3, 1415 (1999) [arXiv:hep-th/9811131];
R. Dijkgraaf and C. Vafa, "Matrix models, topological strings, and supersymmetric gauge theories," *Nucl. Phys. B* 644, 3 (2002) [arXiv:hep-th/0206255].
- [52] M. Becker, K. Dasgupta, A. Knauf and R. Tatar, "Geometric transitions, flops and non-Kähler manifolds. I," arXiv:hep-th/0403288.
- [53] J. F. G. Cascales and A. M. Uranga, "M5-brane geometries, T-duality and fluxes," *JHEP* 0401, 021 (2004) [arXiv:hep-th/0307156];
M. Serone and M. Trappetti, "String vacua with flux from freely-acting orbifolds," *JHEP* 0401, 012 (2004) [arXiv:hep-th/0310245].

- K. Pilch and N. P. Warner, "Generalizing the $N = 2$ supersymmetric RG flow solution of IIB supergravity," Nucl. Phys. B 675, 99 (2003) [arXiv:hep-th/0306098];
- K. Pilch and N. P. Warner, "N = 1 supersymmetric solutions of IIB supergravity from Killing spinors," arXiv:hep-th/0403005;
- D. Nemeschansky and N. P. Warner, "A family of M-theory flows with four supersymmetries," arXiv:hep-th/0403006.
- [68] K. Becker, M. Becker and R. Sriharsha, "PP-waves, M-theory and fluxes," Nucl. Phys. B 676, 172 (2004) [arXiv:hep-th/0308014].
- [69] G. L. Cardoso, G. Curio, G. Dall'Agata and D. Lust, work in progress.
- [70] A. Hanany and B. Kol, "On orientifolds, discrete torsion, branes and M theory," JHEP 0006, 013 (2000) [arXiv:hep-th/0003025].
- [71] P. Candelas, "Lectures On Complex Manifolds," in Superstrings 87, proceedings of the Trieste Spring School (1987) 1-88;
- M. Marino, R. Minasian, G. W. Moore and A. Strominger, "Nonlinear instantons from supersymmetric p-branes," JHEP 0001, 005 (2000) [arXiv:hep-th/9911206].
- [72] P. Bouwknegt, J. Evslin and V. Mathai, "T-duality: Topology change from flux," arXiv:hep-th/0306062.
- [73] P. Bouwknegt, J. Evslin and V. Mathai, "On the topology and flux of T-dual manifolds," arXiv:hep-th/0312052;
- P. Bouwknegt and D. Ridout, "A note on the equality of algebraic and geometric D-brane charges in WZW models," arXiv:hep-th/0312259;
- P. Bouwknegt, K. Hannabuss and V. Mathai, "T-duality for principal torus bundles," arXiv:hep-th/0312284;
- V. Mathai and J. Rosenberg, "T-duality for torus bundles via noncommutative topology," arXiv:hep-th/0401168.
- [74] E. Bergshoeff, C. M. Hull and T. Ortin, "Duality in the type II superstring effective action," Nucl. Phys. B 451, 547 (1995) [arXiv:hep-th/9504081].
- [75] S. F. Hassan, "T-duality, spacetime spinors and RR fields in curved backgrounds," Nucl. Phys. B 568, 145 (2000) [arXiv:hep-th/9907152].
- [76] S. Salamon, "Complex structures on nilpotent Lie algebras," arXiv:math.DG/9808025;
- E. Abbena, S. Garbiero, S. Salamon, "Almost Hermitian Geometry on Six Dimensional Nilmanifolds," arXiv:math.DG/0007066.
- [77] L. Castellani, A. Ceresole, S. Ferrara, R. D'Auria, P. Fre and E. Maina, "The Complete $N = 3$ Matter Coupled Supergravity," Nucl. Phys. B 268, 317 (1986).
- [78] A. Sen, "A note on enhanced gauge symmetries in M- and string theory," JHEP 9709, 001 (1997) [arXiv:hep-th/9707123].
- [79] E. Witten, "Non-Perturbative Superpotentials In String Theory," Nucl. Phys. B 474, 343 (1996) [arXiv:hep-th/9604030].

- [80] N. Seiberg, "IR dynamics on branes and space-time geometry," *Phys. Lett. B* 384, 81 (1996) [[arXiv:hep-th/9606017](#)];
 N. Seiberg and E. Witten, "Gauge dynamics and compactification to three dimensions," [arXiv:hep-th/9607163](#).
- [81] M. Kreuzer and H. Skarke, "Relexive polyhedra, weights and toric Calabi-Yau bractions," *Rev. Math. Phys.* 14, 343 (2002) [[arXiv:math-ag/0001106](#)];
 M. Kreuzer and H. Skarke, "Complete classification of relexive polyhedra in four dimensions," *Adv. Theor. Math. Phys.* 4, 1209 (2002) [[arXiv:hep-th/0002240](#)].
- [82] M. Nakahara, "Geometry, Topology and Physics," (Graduate student series in physics), Institute of Physics Publishing, Bristol, UK (1990).
- [83] C. W. Misner, K. S. Thorne and J. A. Wheeler, "Gravitation," W. H. Freeman and Company, New York (2000).
- [84] S. J. Gates and V. A. Kostelecky, "Supersymmetric Matter Gravitino Multiplets," *Nucl. Phys. B* 248, 570 (1984).
- [85] J. Maharana and J. H. Schwarz, "Noncompact Symmetries In String Theory," *Nucl. Phys. B* 390, 3 (1993) [[arXiv:hep-th/9207016](#)].
- [86] P. Frey, "Lectures on Special Kahler Geometry and Electric-Magnetic Duality Rotations," *Nucl. Phys. Proc. Suppl.* 45B C, 59 (1996) [[arXiv:hep-th/9512043](#)].