

Noncommutative models in patch cosmology

Gianluca Calcagni

Dipartimento di Fisica, Università di Parma,
Parco Area delle Scienze 7/A, I-43100 Parma, Italy
andINFN - Gruppo collegato di Parma, Parco Area delle Scienze 7/A, I-43100 Parma, Italy
(Dated: June 1, 2004)

We consider several classes of noncommutative inflationary models within an extended version of patch cosmological braneworlds, starting from a maximally invariant generalization of the action for scalar and tensor perturbations to a noncommutative brane embedded in a commutative bulk. Slow-roll expressions and consistency relations for the cosmological observables are provided, both in the UV and IR region of the spectrum; the inflaton field is assumed to be either an ordinary scalar field or a Born-Infeld tachyon. The effects of noncommutativity are then analyzed in a number of ways and energy regimes.

PACS numbers: 98.80.Cq, 04.50.+h, 11.10.Nx

I. INTRODUCTION

The idea that the early Universe experienced a phase of accelerated expansion has come to a crucial point. Born as a panacea for some problems of the standard big bang scenario, the inflationary paradigm has been developed and refined during these years, always successfully explaining the available observational data. The upcoming generation of high-precision cosmological experiments such as WMAP (Wilkinson Microwave Anisotropy Probe) [1, 2, 3] and Planck [4] might definitely operate a selection on the great amount of inflation-inspired models. On the other hand, new theoretical scenarios in which the high-energy physics grows more and more in importance have produced a set of interesting research fields implementing the traditional 4D cosmology: therefore we have string cosmology, braneworld cosmology, noncommutative cosmology, and so on.

In their seminal paper [5], Brandenberger and Ho presented a model of large-scale perturbation spectra, in which a noncommutative geometrical structure is generated by the stringy spacetime uncertainty relation (SSUR)

$$\Delta x_p \geq \frac{\ell_s}{2}; \quad (1)$$

where ℓ_s is the string length scale and $x_p = a(t)x$ is a physical space coordinate. It has been argued that this is a universal property for string and brane theory [6, 7, 8]. This picture (henceforth "BH") has then been further explored in [9, 10, 11, 12, 13, 14, 15] and presents many common features with trans-Planckian scenarios with a modified dispersion relation [16, 17, 18, 19, 20, 21, 22].

In this paper we construct another noncommutative model based on the same philosophy of [5] and confront it with BH in its two versions, the first one with the Friedmann-Robertson-Walker (FRW) 2-sphere factored

out in the action measure and the second one with a unique effective scale factor. Scalar and tensor amplitudes and indices as well as consistency equations are obtained through the slow-roll (SR) formalism, both for an ordinary scalar field and a Born-Infeld cosmological tachyon. All the observables turn out to be functions of a noncommutative parameter, called β , measuring the magnitude of the Hubble energy H at the time of horizon crossing in comparison with the fundamental string mass $M_s = \ell_s^{-1}$.

Some works have studied the inflationary perturbations treating β on either almost [12, 15] or exactly [13, 14] the same ground of the SR parameters, computing UV amplitudes and indices via a double or SR expansion for small parameters, respectively. Here we will follow a different approach and consider β as a distinct object with respect to the SR tower; we will keep only the lowest-SR-order part of the observables and regard any β -term as pertaining to these leading-SR-order quantities. We stress that, while the parameter β accounts for nonlocal effects coming from the string scale ℓ_s , the SR tower is determined by the dynamics of the cosmological inflationary expansion. Therefore, they describe quite distinct physical phenomena. In fact, there is no connection between β and the recursively defined SR tower, although even the first SR parameter is introduced by hand; the elements of the tower are built up of time derivatives of H and they all vanish in a de Sitter background, while β , which contains only the Hubble parameter and the string scale, does not. In particular, the lowest-SR-order spectral amplitudes, equivalent to those obtained in a quasi-de Sitter model, will depend on β . Besides this motivation, such a procedure has additional advantages. For example, we can study regimes with not-so-small β within the SR approximation; secondly, if one keeps the magnitude of β unconstrained, one can also explore the IR region of the spectrum, $\beta \ll 1$, through appropriate techniques.

These effective noncommutative models can be extended to braneworld scenarios in which the 3-brane ex-

Electronic address: calcagni@sunipr.it

periences a cosmological expansion governed by an effective Friedmann equation. The precise theoretical setup is highly nontrivial even in the commutative case, because of the number of requirements to impose on the background forms and spacetime geometry in order to have a cosmological four-dimensional variety. We will phenomenologically assume to have a 3-brane in which the SSUR (1) holds for all the braneworld coordinates $fx, g, = 0; 1; 2; 3$, while the extra dimension y along the bulk remains decoupled from the \ast -algebra.

A very qualitative way to see how such a noncommutative scenario might emerge is the following. One of the most promising braneworlds is the Randall-Sundrum (RS) model [23, 24] or its Gauss-Bonnet (GB) generalization (e.g., [25, 26], and references therein), motivated by M-theory as low-energy products of a dimensionally reduced 11D supergravity to a 10D string theory, down to a 5D effective gravity [27, 28, 29, 30, 31, 32, 33] (see also [34]). The resulting 11D manifold is $AdS_5 \times X_{CY}$, where the brane is located at the fixed point $y = y_b$ of the Z_2 symmetry in the 5D anti-de Sitter bulk and the other six dimensions are compactified on a Calabi-Yau 3-fold X_{CY} . The 5D gravitational coupling is related to the 11D one by $\frac{2}{5} = \frac{2}{11} V_{CY}$, where V_{CY} is the internal volume of the Calabi-Yau space and $\frac{2}{11} = M_s^{-9}$; thus, we will identify the noncommutative string mass as the fundamental energy scale of the full theory. To diagonalize the noncommutative algebra and induce a pure 4D SSUR on the brane one might expect the expectation values of the 11D background fields such that the extra direction commutes, $[y; x] = 0$. Some other subtleties to deal with are discussed in Sec. IIIC.

A useful approach to study perturbation spectra in braneworld scenarios is patch cosmology, which makes use of a nonstandard Friedmann equation, coupled to the slow-roll formalism; despite all the shortcomings of this approximated treatment of extra-dimensional physics, it gives several important first-in-pact information. The four-dimensional scenario is automatically included.

We will not provide a full derivation of standard results and leave the reader to consult the available literature. For an introduction to inflation and perturbation theory, see [35, 36]. For a review on braneworld, see [37]. For patch cosmology, scalar and tachyon inflation, and a more complete list of assumptions, technicalities, comments, and references, see [26].¹

The plan of the paper is as follows. The general setup is established in Sec. II, further developing the formalism of [26]; in Sec. III we review the BH models and introduce a new prescription for the noncommutative action, describing then the UV and the IR spectral regions. Sec. IV fully develops these models; in Sec. V a detailed anal-

ysis summarizes the main results and Sec. VI is devoted to concluding remarks.

II. GENERAL SETUP

We will keep the general framework of a noncommutative 3-brane in which, either in a limited time interval during its evolution or in a given energy patch, the cosmological expansion satisfies an effective Friedmann equation

$$H^2 = \frac{2}{q} q; \quad (2)$$

where q is constant and $\frac{2}{q} > 0$ is some factor with energy dimension $[q] = E^{1-2q}$. Often we will use the more convenient parameter $2(1-q^1)$. Table I reports the characteristics of the 4D and braneworld cosmologies we will consider. There, $\frac{2}{5} = 8 m_5^{-3} = M_5^{-3}$ is the five-dimensional gravitational coupling, $\frac{2}{5}$ is the Gauss-Bonnet coupling and $\frac{2}{5}$ is the brane tension. Gravity experiments impose $M_5 > 10^8$ GeV and $\frac{2}{5} > 10^3$ GeV; best-fit analyses of BH noncommutative models gives estimates for the string scale $M_s = 10^{11} - 10^{17}$ GeV [9, 10]. In typical Horava-Witten scenarios, the fundamental scale is of order of the GUT scale, $M_s = 10^6$ GeV.

TABLE I: The energy regimes described in the text.

Regime	q	$\frac{2}{q}$
GB	2=3	1
RS	2	1
4D	1	0

We neglect any contribution from both the Weyl tensor and the brane-bulk exchange; assuming there is some conservation mechanism for a perfect fluid with equation of state $p = w$, the continuity equation on the brane reads $\dot{\rho} + 3H(\rho + p) = 0$. Let us consider an inflationary four-dimensional flat universe filled with an homogeneous inflation field. If this is an ordinary scalar field $\phi(t)$ with potential V , then the energy density and pressure are

$$\rho = \frac{1}{2} \dot{\phi}^2 + V(\phi) = p + 2V(\phi); \quad (3)$$

and the effective equation of motion is

$$\ddot{\phi} + 3H\dot{\phi} - V'(\phi) = 0; \quad (4)$$

Another choice is to consider a homogeneous tachyon field $T(t)$, that is a scalar, causal field satisfying the Dirac-Born-Infeld action [53, 54]

$$S_{DBI} = \int d^4x \int da^3 V(T) \sqrt{1 - \dot{T}^2}; \quad (5)$$

¹ To the reference list of [26] we add [38, 39, 40, 41, 42, 43, 44, 45] for higher derivative and Gauss-Bonnet gravity models and [46, 47, 48, 49, 50, 51, 52] for the cosmological tachyon.

energy density and pressure read

$$\rho = \frac{V(T)}{c_s}; \quad (6)$$

$$p = -\frac{V(T)}{c_s} = -\frac{V(T)^2}{c_s}; \quad (7)$$

$$\text{where } c_s = \frac{p}{\rho} = \frac{p}{1 - T^2}.$$

A. Slow-roll parameters and commutative observables

According to the inflationary paradigm, an early-Universe period of accelerated expansion is driven by a scalar field slowly "rolling" down its potential toward a local minimum. Thanks to this idea, one can construct a set of useful quantities (the SR parameters) which govern the dynamics of the system and make the computational task easier through suitable SR expansions. The first parameter is the time variation of the Hubble length H^{-1} ,

$$\frac{\dot{H}}{H^2} = \frac{3q(1+w)}{2}; \quad (8)$$

Full SR towers involving either the Hubble parameter or the inflation potential can be defined from dynamical considerations within the Hamilton-Jacobi formulation of the problem. Here we will need only the first three bricks of the Hubble tower. For a scalar field, these are

$$= \frac{3q}{2} \frac{1}{H^2}; \quad (9)$$

$$\frac{d \ln \rho}{d \ln a} = \frac{1}{H^2}; \quad (10)$$

$$\frac{1}{H^2} = \frac{1}{H^2}; \quad (11)$$

The evolution equations of the parameters are second-SR-order expressions,

$$\dot{\rho} = H^2 \rho [2 - \frac{\dot{H}}{H^2}]; \quad (12a)$$

$$\dot{p} = H^2 p; \quad (12b)$$

further time derivatives raise the SR order by one at each step. For the tachyon field, the SR parameters are

$$\tau = \frac{3q}{2} T^2; \quad (13)$$

$$\tau = \frac{T}{H T}; \quad (14)$$

$$\frac{1}{H^2} = \frac{T}{H^2 T} = \frac{\ddot{T}}{H^2 T^2}; \quad (15)$$

with time variation

$$\dot{\tau} = 2H \tau; \quad (16a)$$

$$\dot{\tau} = H \tau; \quad (16b)$$

Note that one can compute second-SR-order tachyon expressions by going to the form $\tau \approx 2$ in Eq. (12a).

A derivation of the perturbation amplitudes will be seen in the noncommutative case. Here, we just quote the results for the commutative observables, denoted by a superscript (c). The perturbation amplitudes can be written as

$$A^{(c)} = \frac{aH}{5z}; \quad (17)$$

to lowest SR order (equivalently, in a quaside Sitter 4D spacetime). In general, the squared function z is

$$z^2 = \frac{a^2(1+p)}{q H^2} = \frac{a^2(1+w)}{q \frac{2}{H}}; \quad (18)$$

where q is a proportionality coefficient; for an ordinary and tachyon scalar on the brane,

$$z(\tau) = \frac{a}{H}; \quad (19)$$

$$z(T) = \frac{aT}{c_s \frac{1}{q} H^2}; \quad (20)$$

with $q(\tau) = 1$ and $q(T) = 1/c_s$. The scalar amplitude is, to lowest SR order,

$$A_s^{(c)2} = \frac{3q}{25} \frac{1}{H^2}; \quad (21)$$

where q is given by either Eq. (9) or (13), according to the model. The spectral index and its running are, to lowest SR order,

$$n_s^{(c)} - 1 = \frac{d \ln A_s^{(c)2}}{d \ln k} = 2 - 4; \quad (22)$$

$$\frac{dn_s^{(c)}}{d \ln k} = \frac{h}{5(n_s^{(c)} - 1) + 4(3 - \tau)}; \quad (23)$$

while for the tachyon

$$n_s^{(c)} - 1 = 2 - \frac{(2 + \tau)}{h}; \quad (24)$$

$$\frac{dn_s^{(c)}}{d \ln k} = \frac{(3 - \tau)}{5(n_s^{(c)} - 1) + (2 + \tau)}; \quad (25)$$

These equations can be obtained through Eq. (12) and the lowest-order relation

$$\frac{d}{d \ln k} = \frac{d}{H dt}; \quad (26)$$

In Eqs. (23) and (25), the parameter τ^2 has been dropped (see the discussion in [26]).

Let α be the inverse of the bulk curvature scale; the effective 4D Newton constant is $\frac{2}{4} = \frac{2}{5} = (1 + 4^{-2})$, which in RS gives $\frac{2}{4} = \frac{2}{4} = 6$. The gravitational spectrum in RS and GB scenarios has been investigated in [25, 55] for a de Sitter brane. It turns out that $A_t^{(c)2} =$

$A_{t(4D)}^{(c)2} F^2(H=1)$, where $A_{t(4D)}^{(c)2}$ is the commutative 4D amplitude with $z_{4D} = \frac{P}{2a} = \frac{1}{4} F$ is a complicated function of η and $H=1$ determined by the normalization of the y -dependent part $\phi_0(y)$ of the graviton zero mode calculated on the brane position, $F^2 = \frac{2}{3} (y_b)^2 = \frac{2}{3}$. Writing $A_t^2 = A_{t(4D)}^2 z_{4D}^2 = z^2$, one has

$$z(\eta) = \frac{P}{4F} \frac{2a}{F} : \quad (27)$$

This remarkably simple generalization of the 4D function z is possible only because of the maximally symmetry of the de Sitter brane, which permits a variable separation of the wave equation for the Kaluza-Klein gravity modes, $h(x; y) = h^{(m)}(x) \phi_m(y)$. To be consistent with the patch solution (2), we consider the approximated version F_q^2 of F^2 in the proper energy limits: $F_1^2 = 1$, $F^2(H=1)$ in 4D, $F_{2=3}^2 = 3H(2)$, $F_0^2(H=1)$ in RS, and $F_{2=3}^2 = (1+4)^2 = (8-H)$, $F^2(H=1)$ in GB [25]. We can write down the patch version of Eq. (27) by noting that in four dimensions the graviton background can be described by Eq. (18) with $\phi_1(\eta) = 1$ and a perfect fluid $p_h = \rho_h/3$ which does not contribute to the cosmic acceleration $a = aH^2(1)$, being $\eta = 1$ in Eq. (8). Generalizing this stationary-solution trick one has $w(\eta) = 2 = (3q) - 1$ and

$$z(\eta) = \frac{P}{4F_q} \frac{2a}{F_q} ; \quad (28a)$$

$$F_q^2 = \frac{3q}{q(\eta)} \frac{2}{4} \frac{H}{4} ; \quad (28b)$$

where the coefficient $q(\eta)$ is determined by the gravity model and is $\phi_1(\eta) = 1 = \phi_{2=3}(\eta)$ and $\phi_2(\eta) = 2/3$. The commutative tensor amplitude is then

$$A_t^{(c)2} = \frac{3q}{25} \frac{2}{4} \frac{H}{2} \frac{2+}{q(\eta)} ; \quad (29)$$

while the spectral index and its running are

$$n_t^{(c)} = \frac{d \ln A_t^{(c)2}}{d \ln k} = (2 +) ; \quad (30)$$

$$n_t^{(c)} = (2 +) (n_s^{(c)} - 1) + (2 +) ; \quad (31)$$

both for the ordinary scalar and the tachyon field. The tensor-to-scalar ratio is

$$r^{(c)} = \frac{A_t^{(c)2}}{A_s^{(c)2}} = \frac{1}{q(\eta)} ; \quad (32)$$

B. Leading-order noncommutative observables

Let A denote a lowest-order perturbation amplitude, $A^2 = f A_t; A_s g$; in general, it can be written as

$$A(; H ;) = A^{(c)}(H ;) () ; \quad (33)$$

where θ is a noncommutative parameter to be defined later, $A^{(c)} = A(\theta = 1)$ is the amplitude in the commutative limit, and $()$ is a function encoding leading-SR-order noncommutative effects. It will turn out that, up to $O(\theta^2)$ terms,

$$\frac{d \ln A^2}{d \ln k} = ; \quad (34)$$

where $\theta = ()$ is a function of θ such that $\theta = 0()$. The spectral index is

$$n = \frac{d \ln A^2}{d \ln k} = n^{(c)} + ; \quad (35)$$

for the scalar spectrum, $n = n_s - 1$. The index running is

$$\frac{dn}{d \ln k} = \theta^{(c)} + \frac{d^2 \ln A^2}{d \ln k^2} ; \quad (36)$$

In the scalar field case, the last term can be written as

$$\frac{d^2 \ln A^2}{d \ln k^2} = [(2) - 2] ; \quad (37)$$

[here $\theta = \theta(H)$ to first SR order] with $\theta \neq 2$ in the tachyon case. Because of Eq. (33), the tensor-to-scalar ratio is $r = r^{(c)}$ and the consistency equations read

$$\begin{aligned} s(\theta) &= r_q f(5) (n_s - 1) \\ &+ [4(3 +) - (7 + +)] \theta g ; \quad (38) \\ s(\Gamma) &= r_q f(3 +) (n_s - 1) \\ &+ [(2 +) (3 +) - (5 + 2 +)] \theta g ; \quad (39) \end{aligned}$$

The lowest-SR-order consistency equation for the tensor index is

$$n_t = [(2 +)] r ; \quad (40)$$

and its running is

$$n_t = r_q [(2 +) (n_s - 1) + (2 +)] r_q ; \quad (41)$$

There is also a next-to-leading order version of Eq. (40), which we will not consider here.

III. NONCOMMUTATIVE MODELS

Let us introduce the new time variable $\tau = 2R^+$, $d\tau = da/H$. With a constant SR parameter θ , an integration by parts with respect to a gives

$$= \frac{a}{H} \frac{1}{1 + } - \frac{a}{H} ; \quad (42)$$

Inequality (1) can be rewritten in terms of comoving coordinates as

$$x \leq 1_s^2 ; \quad (43)$$

and the corresponding algebra of noncommutative spacetime is time independent,

$$[\cdot, \cdot] = i\ell_s^2 : \quad (44)$$

The \star -product realizing Eq. (44) is defined as

$$(f \star g)(x; y) = e^{(i\ell_s^2=2)(\partial_x \otimes \partial_y - \partial_y \otimes \partial_x)} f(x; y) g(x; y) \Big|_{x^0=y^0=x} : \quad (45)$$

This realization of noncommutativity is in contrast with

$$[x, x] = i\ell_s^2 ; \quad (46)$$

where ℓ_s is the noncommutative parameter. This type of noncommutative cosmology, which does not preserve the FRW symmetries, has been studied in [56, 57, 58]. Other implementations can be found in [59, 60, 61, 62, 63, 64, 65].

A. BH models

In the following we will adopt the short notation $a = a(\cdot)$ and $a = a(\cdot) \Big|_{k_0}$. For the skipped details, see [5]. Consider now the action of a free scalar field living in a (1+1)-dimensional FRW space. In the noncommutative models we will study, each conventional product is replaced by the \star -product (45); thus, the gravitational sector of the theory is not a completely passive spectator but is involved via the \star -coupling of the metric with the matter content. The new 2D action reads, noting that $a^2 = a \star a$ [5],

$$S_{BH} = \int d^2x \frac{1}{2} \partial_\mu a \star \partial^\mu a + \int d^2x \frac{1}{2} \partial_\mu a \star \partial^\mu a \Big|_{k_0} : \quad (47)$$

In the comoving momentum space,

$$(x; y) = V^{1/2} \int_{k < k_0} \frac{dk}{2} \phi(k) e^{ikx} ; \quad (48)$$

where V is the total spatial coordinate volume and

$$k_0 = M_s a_{eff} \quad (49)$$

is a cutoff realizing the stringy uncertainty relation. The most convenient way to recast the action is to write the scale factor as a Fourier integral, $a^2(\cdot) = \int d! a^2(!) e^{i! \cdot}$, and perform the \star -products of the complex exponentials in the integrand, removing the cutoff in the limit $k_0 \rightarrow \infty$ when absorbing the δ -integrals in momentum spaces. The result is

$$S = V \int_{k < k_0} d^2k \frac{1}{2} \partial_k a \star \partial_k a + \int_{k < k_0} d^2k \frac{1}{2} \partial_k a \star \partial_k a \Big|_{k_0} ; \quad (50)$$

where

$$k = \frac{1}{2} (a_+^2 + a_-^2) ; \quad (51)$$

Defining two new objects

$$a_{eff}^2 = \frac{1}{k} a_+ a_- ; \quad (52)$$

$$y^2 = \frac{1}{k} \frac{a_+^2 + a_-^2}{2a_+ a_-} ; \quad (53)$$

and the effective conformal time coordinate

$$\frac{d\tau}{d} = a_{eff}^2 ; \quad (54)$$

the scalar action becomes

$$S = V \int_{k < k_0} d\tau dk \frac{1}{2} y^2 \partial_k a_{eff}^2 \partial_k a_{eff}^2 + \int_{k < k_0} d\tau dk \frac{1}{2} y^2 \partial_k a_{eff}^2 \partial_k a_{eff}^2 ; \quad (55)$$

where the primes are derivatives with respect to τ .

B. A new prescription for noncommutativity

Cyclic permutations of the \star -product inside the integral (47) leave the action invariant. Therefore, it is natural to see whether a different noncyclic ordering of the factors gives a theory with interestingly new predictions. The other nontrivial noncommutative action one can obtain is

$$S_{new} = V \int d^2x \frac{1}{2} \partial_\mu a \star \partial^\mu a + \int d^2x \frac{1}{2} \partial_\mu a \star \partial^\mu a \Big|_{k_0} : \quad (56)$$

The same computational pattern of the previous section leads to Eq. (55) with k given by

$$k = \frac{1}{2} (a_+^2 + a_-^2) ; \quad (57)$$

and

$$a_{eff}^2 = a_+^p a_-^p ; \quad (58)$$

$$y^2 = \frac{a_+^2 + a_-^2}{2a_+ a_-} ; \quad (59)$$

In this case there is only a partial smearing of the product of scale factors and one might guess that the resulting noncommutative phenomenology would be less pronounced than that of BH model. In the UV limit it will turn out that, within a given variation of the noncommutative parameter and in some region in the space of parameters, the range of the quantity $j_s(\cdot) = j_s(T)$ is slightly smaller than in the BH model but always of the same order of magnitude. In the infrared region, however, the two models are almost undistinguishable; see Sec. V.

C. Four-dimensional effective actions and amplitudes

When going to 3+1 dimensions, the measure z_k^2 of the integral will contain the nonlocal effect coming from the SSUR:

$$S = \int_{k < k_0} d^3k \frac{1}{2} z_k^2 \dot{u}_k^2 : \quad (60)$$

Here we will consider two classes of models. In the first one, we suppose the total measure to be given by the product of the noncommutative (1+1)-measure and the commutative one:

$$z_k = zy; \quad (61)$$

then, as we are going to show in a moment,

$$= \frac{a_{\text{eff}}^2}{a^2 y} \quad (\text{class 1}); \quad (62)$$

These models, in which the FRW 2-sphere is factored out, will be dubbed as "1." Another interesting prescription consists in replacing the commutative scale factor in z_k with the effective one; then, say a_{eff} ,

$$z_k = z \frac{a_{\text{eff}}}{a}; \quad (63)$$

and

$$= \frac{a_{\text{eff}}}{a} \quad (\text{class 2}); \quad (64)$$

models with this will be named "2."

Let us now look at cosmological perturbations coming from an inflationary era and assume, as it is the case, that is a generic perturbation satisfying the action (60). The spectral amplitude coming from the k -th mode of the perturbation is

$$A^2 = \frac{2k^3}{25^{\frac{3}{2}}} \langle \dot{u}_k^2 \rangle; \quad (65)$$

where angle brackets denote the vacuum expectation value and the expression is evaluated at the reference time t_* to be discussed in a while. V is a change of variable,

$$u_k = \sqrt{z_k} \chi_k; \quad (66)$$

the action (60) gives the Mukhanov equation

$$u_k'' + k^2 \frac{z_k''}{z_k} u_k = 0; \quad (67)$$

Noting that $d\sim = d = (a=a_{\text{eff}})^2$, we get the useful relation

$$\sim = \frac{1}{aH} \frac{a}{a_{\text{eff}}}^2; \quad (68)$$

in the lowest SR approximation. If the SR parameters are small, then they are constant to leading order because their derivatives are higher order. It is then possible to solve the Mukhanov equation with exactly constant SR parameters and perturb the obtained solution. Such cosmological solutions do exist and can be constructed in a variety of situations; among them, a particularly important one is power-law inflation, which we will use when considering the infrared region of the spectrum. Therefore,

$$\frac{1}{z_k} \frac{d^2 z_k}{d\sim^2} = \frac{a_{\text{eff}}^4}{a} \frac{1}{z} \frac{d^2 z}{d\sim^2} = \frac{a_{\text{eff}}^4}{a} \frac{1}{2} \frac{1}{\sim^2} \frac{1}{\sim^2}; \quad (69)$$

where $\sim = 3/2 + O(\epsilon)$. With constant \sim , the solution of this equation is the same as that of the commutative case, namely $j_{\sim}(\sim)/(\sim)^{3/2} H^{(1)}(\sim/k)$, where $H^{(1)}$ is the Hankel function of the first kind of order \sim . In the long wavelength limit $k \rightarrow 0$, when the mode with comoving wave number k is well outside the horizon, the appropriately normalized solution becomes, from Eq. (68),

$$j_{\sim}^2 = \frac{1}{2k} \frac{1}{k\sim}^2 = \frac{1}{2k} \frac{aH}{k}^2 \frac{a_{\text{eff}}^4}{a}; \quad (70)$$

nally, one gets Eq. (33) by inserting either definition (61) or (63) in Eq. (65).

Given a noncommutative brane in a commutative bulk, the nonlocal smearing will only affect the pure four-dimensional part of the graviton-zero mode action, while leaving the pure transversal normalization unchanged; from the discussion in Sec. II, it is then clear that the noncommutative tensor spectral amplitude will be $A_t^2 = A_t^{(c)2} / \int_0^2 (dy) A_{t(4D)}^2$. Therefore, for the gravitational spectrum, \mathcal{P}_t denotes the coefficient functions of the noncommutative 4D polarization tensor $h^{(0)}(\mathbf{x})$ and z is given by Eq. (28).

In the case of the inflation field, $\sim = R$ is the curvature perturbation on comoving hypersurfaces, generated by quantum fluctuations of the field inflating the early Universe.

The action and Mukhanov equation for a perturbation generated by a tachyon field has an additional factor in front of k^2 in Eqs. (60) and (67), namely the speed of sound for the perturbation: $k^2 \rightarrow c_s^2 k^2$ [66, 67]. Since the SSUR does not affect products of homogeneous quantities, the noncommutative generalization of the tachyonic scalar amplitude is straightforward [68]. Now, one may ask how the inhomogeneous version of the original Born-Infeld action (5) is modified when inserting the $*$ -products. Let us recall that noncommutativity naturally arises in string theory when a Neveu-Schwarz (NS-NS) B-field is switched on in the low-energy tree-level action. However, this results in a linearization of the tachyonic action and, on the other hand, a large noncommutative parameter may trigger brane decay processes [69]; therefore, the simple noncommutative version of the cosmological tachyon might seem too naive.

Anyhow, tachyon scenarios are not new to counterintuitive behaviors. In the slow-roll approximation, $\dot{\phi}^2 \ll 1$, the action (5) can be linearized and the rescaled field $\bar{\phi} = \sqrt{V} \phi$ behaves like an ordinary scalar; nevertheless, the theoretical prediction encoded in the consistency relations is different with respect to that of the genuine scalar scenario [see Eqs. (38) and (39)]. Here, something similar happens, in aging to turn on and increase the B-field smoothly, and the final result differs from the scalar case indeed.

Moreover, the stringy linearization is a feature of realization (46) rather than (44) and the former may give rise to a different cosmological model in which FRW isotropy is not preserved [57]; also, a priori it would be highly nontrivial to construct a Lorentz-violating cosmological brane model (in fact, in the case of a de Sitter brane, maximal symmetry is crucial for coordinate-separating the graviton wave equation [25, 55]).

To further understand the difficulties lying in a full implementation of noncommutative string theory in cosmology, it is important to stress that all that has been said about the algebra (46) (i.e. instability and cosmological scenarios) is true only in a purely spatial \star -product, $\phi_i = 0$. When trying to introduce noncommutativity in both space and time, as is the case of realization (44), it may be difficult to achieve a coherent, well-defined theory. In fact, in the Seiberg-Witten limit reproducing the noncommutative geometry, α' and θ are kept fixed while $B \rightarrow 1$ and the Regge slope $\alpha' \rightarrow 0$. Let $E_i = B_{0i}$ be the electric part of the NS 2-form and assume $E = E_i dx^i \neq 0$. Then, while the B-field goes to infinity and approaches the critical value $E_{cr} = (2\alpha')^{-1}$, a classical instability develops and the rate of open string pair production diverges [70]; heuristically, the string is tore apart by the increasing electric field strength. For these reasons we regard algebra (44) as the starting point of the cosmological setup rather than the ultimate product of some high-energy theory, for the moment leaving the details of the latter aside.

D. The UV region

In order to correctly evaluate the perturbation spectra, one must determine the time τ_0 when the k -th mode is generated and, later, when it crosses the Hubble horizon. Because of the momentum cutoff (49), the analysis for the noncommutative case must be conducted separately in the mildly and strongly noncommutative regions.

From the very beginning, one can define the time τ when a perturbation with wave number k crosses the horizon by the formula $k = k(\tau) = a(\tau)H(\tau)$. This relation provides an operative definition of the number of e-foldings ($k/H \exp N$) and the time variation of k , Eq. (26). Of course, this is valid for any cosmology in which time definitions have zero uncertainty, that is, for commutative cosmologies and noncommutative cosmologies in the range far from the upper bound (49), in

the so-called ultraviolet region, where $k \gg k_0$. In fact, the time of horizon crossing is different from its commutative counterpart τ_c , since $\tau_c < \tau$ and the crossing mode is delayed [5]. In [14] this effect is quantified as $k_c = k = \exp[(\tau - \tau_c)]$.

On the contrary, one might define the horizon crossing through the z function as

$$k^2 = \frac{z_k^{(0)}}{z_k} 2(aH)^2; \quad (71)$$

and get an extra factor of 2; due to the structure of the Mukhanov equation, this approach would be valid in any case, let it be the commutative or the noncommutative one.

In the UV region, the cosmological energy scale when the perturbation is generated is much smaller than the stringy scale, $H(\tau > \tau_0) = H(\tau_0) = M_s$, and noncommutative effects are soft; thus, the smeared versions of α' can be approximated by α since

$$k_s^2 \ll 1 \quad (\text{UV region}); \quad (72)$$

from Eq. (42). It is convenient to define the noncommutative parameter

$$\frac{kH}{aM_s^2} \ll 1; \quad (73)$$

whose time derivative is

$$\dot{\epsilon} = 4H \quad : \quad (74)$$

Note that this relation states that ϵ is almost constant in a rapidly accelerating background, regardless of its magnitude. The analogy with the evolution equations of the SR tower, e.g., Eq. (12), suggested the authors of [13, 14] treat ϵ as a sort of SR parameter, keeping all the parameters at the same truncation level in the expressions of the UV observables.

At horizon crossing,

$$\epsilon = \frac{\dot{\phi}}{k} \frac{p}{2aH} = 2 \frac{H}{M_s} \ll 1; \quad (75)$$

and Eq. (74) is valid for ϵ , too. The ultraviolet region is by definition the region in which $H = M_s \ll 1$,² it is characterized by long wavelength perturbations generated inside the Hubble radius and, in a cosmological microwave background (CMB) spectrum, this would correspond to the portion of the Sachs-Wolfe (inflationary) plateau with not-too-small spherical modes, $10^{-1} < l < 100$.

In the commutative case, to use one pivot scale instead of the other amounts to different next-to-lowest-order expansions in the SR parameters; the 4D consistency equations are thus unaffected, since the introduction of the

² Without risk of confusion, we will continue to use the symbol ϵ to indicate the ratio $H = M_s$ when discussing the UV limit (1) of spectral quantities.

optimized pivot scale (71) results in a rescaled coefficient $C \rightarrow C + \ln \frac{H}{M_s}$ and this one is not present in them (see, e.g., [14] and references therein for details). This is also true in the RS scenario [71] as well as in general patch cosmology [72].

In the noncommutative case, the change of the pivot scale doubles the magnitude of the parameter (75). The resulting models will display the same theoretical features of the $k = aH$ models, but shifted backward along the energy scale determined by the ratio $H = M_s$. Observational constraints should take the rescaling of the string mass into account, when changing the pivot scale.

In the limit (72), we can Taylor expand the scale factors a around τ_0 for small k . To first order in the SR parameters and to all orders in ϵ , the nonlocal dependence of the scale factor is

$$a(k) = a(\tau_0) \left[1 - \frac{1}{2} \frac{H^2}{M_s^2} \tau_0^2 + \frac{1}{6} \frac{H^4}{M_s^4} \tau_0^4 - \dots \right] \quad (76)$$

where the factor in front of τ_0 comes from a series whose radius of convergence is τ_0 . More precisely, when $\tau_0 \ll 1$ then $H = M_s < 0.8$: Since we are interested in lowest-SR-order amplitudes, we can neglect the SR tower and find

$$a(k) = a(\tau_0) \left[1 - \frac{1}{2} \frac{H^2}{M_s^2} \tau_0^2 + \dots \right] \quad (77)$$

The concrete procedure to compute the spectral amplitudes will be to use the horizon crossing formula (71) at τ_0 in the UV region, and the saturation time τ_0 in the IR region. In [5] and other papers these instants are dubbed τ_k and τ_k^0 , respectively, to highlight the dependence on the wave number.

E. BH model IR region

In the IR region things are quite different: the wave modes are generated outside the horizon and, since they are frozen until they cross the horizon, their magnitude depends on the time when they were generated. This corresponds to the (k -dependent) time τ_0 when the SSUR is saturated, $k(\tau_0) = k_0(\tau_0)$, and quantum fluctuations start out with their vacuum amplitude. The effective and smeared scale factors must be evaluated at this instant; the expansion (77) is no longer valid since $H = M_s$ in the infrared. To proceed one can explicitly use the exact solution around which the equation of motion for the perturbation has been expanded. The power-law solution corresponds to a constant index w , when the scale factor is $a(\tau) = \tau^{n/(n+1)}$, and $H = n \tau^{-1/(n+1)} = (n+1)$. For an exponential scale factor (de Sitter expansion, $n \rightarrow 1$), $a(\tau) = H^{-1}$, in accordance with Eq. 42). From equations (49) and (52),

$$a(k) = k^{-1} \tau_0^{1/(n+1)} \quad (78)$$

where $\tau_0 = \tau_0(k)$ and

$$\frac{2}{H} = \frac{M_s}{H} \quad (79)$$

In the infrared region,

$$k \tau_0^2 \ll 1 \quad (\text{IR region}); \quad (80)$$

and

$$a = H^{-1} k^{-1} \tau_0^{1/(n+1)} \quad (81)$$

$$a = H^{-1} k^{-1} \tau_0^{1/(n+1)} \quad (82)$$

where evaluation at τ_0 is understood. When $\tau_0 \rightarrow 1$, we recover the UV or quasicommutative region since $k \tau_0^2 \rightarrow 1$. Actually, the UV and IR spectra may be joined together in an intermediate region, as it was shown in [10]; in particular, see their Eq. (12), corresponding in the de Sitter limit to $\tau_0^2 = (1 - 3\epsilon/2)$. We will not be able to recover this spectrum within our formalism; however, we will describe other hybrid regimes by using the methods adopted in the IR region ($2 - 10$) for 1 . For future reference, note that

$$- = 2 H \quad (83)$$

F. New model IR region

In the "New" model, the effective scale factor is given by Eq. (58). From Eq. (49),

$$a(k) = k^{-1} \tau_0^{1/(n+1)} \quad (84)$$

where

$$\frac{1}{2} \frac{H^2}{M_s^2} \tau_0^2 = 1 \quad (85)$$

With this definition, the new expressions for a and a are identical to Eqs. (81) and (82), with τ_0 replaced by τ_0 . Equation (83) is replaced by

$$- = \frac{4}{1+2} H \quad (86)$$

In the far IR region, $\tau_0^2 \rightarrow 1$, while in the UV limit $\tau_0 \rightarrow 0$.

Without further justifications, the IR region of the spectrum, $H = M_s$, may be not very satisfactory from a string-theoretical point of view, both because we are above the fundamental energy scale³ and due to the above-mentioned classical instabilities. As it is done in many other occasions in early-Universe cosmology, we will turn a blind eye to this point and seek what are the observational consequences of the extreme regime of the present noncommutative models.

³ However, the space-momentum stringy uncertainty relation, implying $\Delta x \Delta p \geq \frac{1}{2}$, is not a universal property of the theory.

IV . NONCOMMUTATIVE ZOOLOGY

We are ready to collect all the machineries developed so far and inspect the noncommutative models at hand.

A . BH 1

In the BH 1 case,

$$^2 = \frac{2(a_+ a_-)^3}{a^4(a_+^2 + a_-^2)} ; \quad (87)$$

In the UV region,

$$^2 = \frac{(1 + \frac{1}{2})^3}{1 + \frac{1}{2}} ; \quad (88a)$$

$$= \frac{8(2 + \frac{1}{2})}{1 + \frac{1}{2}} ; \quad (88b)$$

$$= \frac{8(2 + \frac{1}{2} + 1)}{(2 + \frac{1}{2})(1 + \frac{1}{2})} ; \quad (88c)$$

For $\frac{1}{2} \rightarrow 1$ one recovers the nearly commutative, $\frac{1}{2}$ -expanded behavior⁴

$$^2 = \frac{1}{16} ; \quad (89a)$$

$$= \frac{1}{4} ; \quad (89b)$$

$$= \frac{1}{4} ; \quad (89c)$$

In the IR region,

$$^2 = \frac{3}{(2 + \frac{1}{2})(1 + \frac{1}{2})} ; \quad (90a)$$

$$= \frac{4(2 + \frac{1}{2})}{(2 + \frac{1}{2})(1 + \frac{1}{2})} ; \quad (90b)$$

$$= \frac{2(2^2 + 6 + 5)}{(3 + 2)(2 + \frac{1}{2})(1 + \frac{1}{2})} ; \quad (90c)$$

In the commutative limit ($\frac{1}{2} \rightarrow 1$), $^2 = \frac{1}{4}$, while in the strongly noncommutative regime ($\frac{1}{2} \rightarrow 0$), $^2 = \frac{3}{4}$ and $\frac{1}{2} = \frac{1}{4}$, in agreement with [10].⁵

B . BH 2

From equations (52) and (64),

$$^2 = \frac{a_+ a_-}{a^2} ; \quad (91)$$

In the UV,

$$^2 = \frac{1}{4} ; \quad (92a)$$

$$= \frac{4}{1} ; \quad (92b)$$

$$= \frac{4}{1} ; \quad (92c)$$

For $\frac{1}{2} \rightarrow 1$, $\frac{1}{2} \rightarrow 4$ and $\frac{1}{2} \rightarrow 4$. In the IR,

$$^2 = \frac{2}{1 + \frac{1}{2}} ; \quad (93a)$$

$$= \frac{2}{1 + \frac{1}{2}} ; \quad (93b)$$

$$= \frac{2}{1 + \frac{1}{2}} ; \quad (93c)$$

When $\frac{1}{2} \rightarrow 1$, $\frac{1}{2} \rightarrow 2(1 + \frac{1}{2})$.

C . New 1

The correction to the commutative amplitude reads

$$^2 = \frac{2(a_+ a_-)^{3=2}}{a^2(a_+ + a_-)} ; \quad (94)$$

In the UV region,

$$^2 = (1 + \frac{1}{2})^{3=2} ; \quad (95a)$$

$$= \frac{6}{1} ; \quad (95b)$$

$$= \frac{4}{1} ; \quad (95c)$$

In the IR limit,

$$^2 = \frac{3=2}{1 + \frac{1}{2}} ; \quad (96a)$$

$$= \frac{6}{1 + 2} ; \quad (96b)$$

$$= \frac{8(1 + \frac{1}{2})}{(1 + 2)^2} ; \quad (96c)$$

In the strongly non-commutative limit ($\frac{1}{2} \rightarrow 0$), $^2 = \frac{3}{4}$ and $\frac{1}{2} = 6 + O(\frac{1}{2})$.

D . New 2

From Eqs. (58) and (64),

$$^2 = \frac{p \frac{1}{a_+ a_-}}{a} ; \quad (97)$$

The UV limit gives

$$^2 = \frac{p}{1} ; \quad (98a)$$

$$= \frac{2}{1} ; \quad (98b)$$

$$= \frac{4}{1} ; \quad (98c)$$

⁴ Throughout the paper we will keep only the leading-order term in the approximated $\frac{1}{2}$ since there is a $\frac{1}{2}$ factor in front of it in Eqs. (38) and (39).

⁵ Eqs. (44)–(47) of [5] are not correct, due to a missing power of y in the inserted z_k^2 ; in Eqs. (23)–(25) of [10] the correct amplitude is recovered.

In the IR region,

$$^2 = \frac{1}{1+1} ; \quad (99a)$$

$$= \frac{2}{1+2} ; \quad (99b)$$

$$= \frac{8}{(1+2)^2} ; \quad (99c)$$

For $1, = 2 + O(2)$.

V. DISCUSSION

To summarize, we can compare the considered models in the perturbative limits, that is, the UV commutative limit (1) and the IR noncommutative limit (1). Trivially, in the nonperturbative or commutative IR region (1), a and one recovers the standard spectrum, $^2 = 1$ and $= 0$; also, by construction, the noncommutative UV region is ill-defined.

In general, we can write the UV commutative limit of the relevant quantities as

$$^2 = 1 - b ; \quad (100a)$$

$$4b ; \quad (100b)$$

$$4 ; \quad (100c)$$

where b is a constant. As anticipated, the structure of the IR amplitudes also permits a perturbative expansion around 1 ; in this case, spectral amplitudes are evaluated at $k < k_0$ via the power-law solution. The IR commutative limit is then

$$^2 = 1 - \frac{b}{2} = 2 ; \quad (101a)$$

$$2b = 2 ; \quad (101b)$$

$$2 ; \quad (101c)$$

from the previous discussions, it is natural to interpret this as an intermediate momentum region at the edge of the UV regime, around < 1 where Eq. (77) ceases to be valid, and corresponding to perturbations generated across the Hubble horizon. In fact, what one does is hit this region starting from the low-momentum IR side. The above-mentioned junction spectrum of [10] is located somewhere closer to the infrared.

Table II shows that all the models display similar asymptotic limits toward different numerical coefficients, the BH ones being larger than the New ones; the coefficient of BH 1 is 4 times that of model 2 within each region (UV or IR), while this ratio is reduced to $b_1 = b_2 = 3$ in the New model. Thus, there is less difference between model New 1 and model New 2 with respect to that occurring between BH 1 and BH 2, further confirming that the "half-sneering" of the new scenario somehow softens noncommutative effects.

The intermediate spectrum (101) breaks down when $^2 < 0$, that is when $H = M_s > 0.5$ (BH 1), 0.8 (New 1), 1

TABLE II: The commutative limit, to lowest order in 1 .

Model	b (commutative limit)
BH 1	4
BH 2	1
New 1	3/2
New 2	1/2

(BH 2) and 1.4 (New 2); therefore Eq. (101) well describes class 2 models at the UV boundary < 1 while it is not particularly reliable for class 1 models.

In the deep UV or commutative limit, the linear approximation (100) properly encodes all the phenomenology of the models; however, the exact noncommutative amplitude better describes the behavior of the cosmological observables in the full span of the UV region. To see this, let us compare the function $\Delta\sigma$, governing the energy dependence of the spectral index (35), with its approximated version $\Delta\sigma_{\text{appr}}$ given by Eq. (100b); we plot the quantity $(\Delta\sigma_{\text{appr}}) =$ for the UV models in Fig. 1. The BH 2, New 1 and New 2 models display the same linear trend in $\Delta\sigma$, while the BH 1 curve is a little below the bisector; the approximation error is up to 50% for < 0.5 , corresponding to $H = M_s < 0.7$, and goes below 10% when $H = M_s < 0.5$. An analogous treatment of Eqs.




FIG. 1: The relative approximation error $(\Delta\sigma_{\text{appr}}) =$ vs in the UV sector. The thin line is for BH 1, the thick line is a superposition of BH 2, New 1 and New 2.

(38) and (39) shows that the difference between the exact and the approximated scalar running may be even greater than the WMAP experimental error for this observable, $s_{\text{appr}} > 10^{-2}$, for any and suitable values for n_s and r in the allowed range. Therefore, the following analysis has been conducted with the full nonlinear amplitude.

Table III reports the noncommutative high-energy limit in the IR region. In particular, the spectral amplitude of New 1 is twice the amplitude of BH 1; however, within each class (1 and 2) a unique set of consistency relations is generated. In the perturbative noncommutative limit, 1 , the IR version of $(\Delta\sigma_{\text{appr}}) =$ is shown

in Fig. 2. The relative approximation error is up to 20% for the BH models and < 0.5 , while it is up to 40% for the New models. The curves of New 1 and New 2 models coincide.

TABLE III: Noncommutative zoology in the IR high-energy limit, to lowest order in δ .

Model	IR noncommutative limit
BH 1	$\delta^2 = 2$
New 1	δ^3
BH 2	2
New 2	2




FIG. 2: The relative approximation error $(\Delta\sigma/\sigma)_{\text{appr}}$ vs δ in the IR sector. The thin solid line is for BH 1, the thin dashed line is for BH 2 and the thick line is a superposition of New 1 and New 2.

In standard cosmology, the consistency equation relating the tensor index n_t and r is adopted in order to reduce the space of parameters. Until now, this has been done only for the 4D and RS cases, both displaying the same 4D degenerate version of Eq. (40). The function

contains a new theoretical parameter, the string energy scale M_s , which enlarges the standard space of cosmological variables. In principle, this might pose some problems if one wanted a reasonably stringent constraint on the observables, facing an uncertainty similar to that one gets when keeping n_t unfixed [73]. In the UV commutative region $\delta \ll 1$, however, one can use the known results for the 4D and RS likelihood analysis in order to compare the consistency equations in the allowed range [74]. For the Gauss-Bonnet case one should rely on the results found in [25, 75].

The IR noncommutative limit is easier to deal with since the asymptotic form of Eq. (40) is independent of the string scale, as it is shown in Table IV. Some features are particularly interesting: (1) The infrared RS-2 models are the only ones with a negative tensor tilt, other noncommutative realizations giving a tilt sign opposite to that of the commutative case; (2) 4D class 2 models predict an exactly scale-invariant tensor spectrum to

lowest order in SR, setting $n_t = 0$ (2); (3) The highest proportionality coefficient is provided by GB class 1 models, allowing a greater tilt given the same tensor-to-scalar ratio.

TABLE IV: The consistency equation (40) in the commutative UV and noncommutative IR limit.

(N on)commutative models	$n_t=r$		
	GB	RS	4D
Commutative UV ($\delta = 0$)	1	2	2
Class 1 IR ($\delta = 6$)	5	2	4
Class 2 IR ($\delta = 2$)	1	$\frac{2}{3}$	0

Although there are $3 \times 2 = 48$ models at hand and a great amount of information to deal with, some preliminary considerations will permit us to simplify such an intricate taxonomy and draw theoretical curves in a reasonable region in the n_s - r plane. Let us first compare the BH scenario with the New one and define j_j ($j_{\text{BH}} + j_{\text{New}} = 2$ and $(j_{\text{BH}} - j_{\text{New}}) = j_j$). Figure 3(a) shows that in the commutative region BH and New models are considerably different, being $\frac{U^V}{1} = 2(7+5) = 10 = 11$ when $\delta \rightarrow 0$ and $\frac{U^V}{2} = 2 = 3$. In the limit $\delta \rightarrow 1$, $\frac{U^V}{1} \rightarrow \frac{U^V}{2}$; this is a spurious effect due to the breaking of the Taylor expansion (77), as one can see by considering the commutative limit of the IR spectra in Fig. 3(b). In fact, $\frac{IR}{1} \neq \frac{IR}{2}$ when $\delta \rightarrow 2$ and, as expected, $\frac{IR}{1} \rightarrow 10 = 11$ and $\frac{IR}{2} \rightarrow 2 = 3$ when $\delta \rightarrow 1$. All this is in accordance with Table II. However, in the IR noncommutative limit there is little difference between BH and New models, being $< 10\%$. Therefore, we will only show the results of New in the infrared and skip the almost identical counterparts in BH. A similar inspection shows that class-1 and class-2 models are quantitatively nondegenerate, getting $\frac{1}{1} = \frac{3}{2}$ for New and BH-IR, and $\frac{1}{1} = \frac{4}{2}$ for BH-UV, in agreement with Tables II and III. Note that these results are independent of the bulk physics.

The versatility of the patch formalism allows coupling it to a noncommutative background in a great number of ways. For example, a realistic picture of the cosmological evolution would be to adopt one particular patch regime in a time interval when a given region of the (non)commutative spectrum is generated; one may then associate the IR region of extra-horizon-generated perturbations with the early-Universe high-energy period, when the extra dimension opens up and the Friedmann equation suffers either GB and/or RS modifications. The consequent evolution is GB-IR \rightarrow RS-IR/UV \rightarrow 4D-UV. Another possibility is to consider pure energy patches and study the noncommutative spectrum in GB, RS, and 4D separately.

Let us compare the running of the scalar index of ordinary-inflation and tachyon-inflation fields,

$$s = s(\delta) = s(T): \quad (102)$$

Since the graphic material is very abundant, we give just

a selection of it; the full set of bi- and three-dimensional figures of this and other combined analyses are available upon request to the author. In Fig. 4 the relative running r_s ($n_s = 1; r; \gamma$) is presented for 4D noncommutative models in the ultraviolet. Two-dimensional slices are then displayed in Figs. 5 and 6. Figure 5 shows that the relative running in Randall-Sundrum is rather modest; on the contrary, in GB and 4D noncommutativity may conspire to bias Eq. (102) and, in particular, the scalar running above the current WMAP uncertainty estimates, $O(10^{-2})$. Braneworld effects, if any, should become more apparent in Planck data, for which the forecasted error is one order of magnitude smaller, $r_s \sim O(10^{-3})$ [76]. In each 2D plot we keep the commutative model as a reference. Note that to increase either n_s or (γ^{-1}) pushes r_s toward positive values. Finally, Figs. 7 and 8 show some features of the New scenarios in the infrared region.

VI. CONCLUDING REMARKS

In this paper we have considered several classes of noncommutative inflationary models within an extended version of patch cosmological braneworlds, starting from a maximally invariant \ast -generalization of the action for scalar and tensor perturbations. Observables and consistency relations are provided via a SR approximation. The main results are:

Class 1 and class 2 models are appreciably distinct from each other in the full span of the spectrum.

BH and New models give almost the same predictions in the IR region of the spectrum.

The relative running r_s (102) is generally more pronounced in the GB scenario than in 4D, while in RS the effect is less evident.

Either increasing η or going to the commutative limit, $H \rightarrow M_{\text{pl}} \rightarrow 0$, the relative running r_s tends toward positive values.

The consistency relation η / r , Eq. (40), greatly differs from one noncommutative model to another.

These models are far from being fully explored. For instance, one could impose also the extra dimension(s) to be noncommutative and extend the algebra (44) or other realizations to the transverse direction(s). A brane with finite thickness would emerge because of the minimum length scale l_s ; in this case our analysis could be thought as performed on mean-valued quantities along the brane thickness. For example, $\int_{\text{brane}} dy, \int_{\text{brane}} d\eta$, and so on. The subject requires further attention and a good starting point might be the cosmological thick brane setup [77, 78, 79, 80, 81, 82, 83, 84, 85, 86].

An interesting possibility is to choose another vacuum state rather than the adiabatic vacuum with which the perturbation spectrum is usually calculated. This scheme has been outlined in [16] and developed in [15, 20].

Other important aspects might be the subject of future studies. First, the use of the gravitational version of the function $z(\sim)$, Eq. (27), would permit one to compute next-to-leading-order expressions both for the tensor amplitude and the consistency equation for the tensor index [71, 72]. Secondly, a numerical simulation of the CMB spectrum as well as a likelihood analysis involving the consistency equation (40), or its next-order version, in the IR limits of Table IV are required in order to constrain the space of cosmological parameters in the low-momentum region of the perturbation spectra. Third, different analyses would point out other important aspects of the models; one may set his/her fancy free by looking at cross comparisons like in [26] and define general relative runnings $r_s^{(\gamma)}(\gamma) = r_s^{(0)}(\gamma)$. Also, stand-alone analyses with explicit inflationary models would constrain the inflation potential according to the predictions for the cosmological observables obtained from the SR expressions for n_t , n_s , and r_s . All these topics are currently under investigation.

Acknowledgments

I am grateful to L.G. Rigobol for his invaluable suggestions and advice during the completion of this work. I also thank Y.S.M. Yung for useful discussions.

-
- [1] C.L. Bennett et al., *Astrophys. J., Suppl. Ser.* 148, 1 (2003).
 - [2] D.N. Spergel et al., *Astrophys. J., Suppl. Ser.* 148, 175 (2003).
 - [3] S.L. Bridle, A.M. Lewis, J. Weller, and G. Efstathiou, *Mon. Not. R. Astron. Soc.* 342, L72 (2003).
 - [4] <http://astro.estec.esa.nl/SA-general/Projects/Planck/>
 - [5] R. Brandenberger and P.-M. Ho, *Phys. Rev. D* 66, 023517 (2002).
 - [6] T. Yoneya, in *Wandering in the Fields*, edited by K. Kawarabayashi and A. Ukawa (World Scientific, Singapore, 1987), p.419.
 - [7] M. Li and T. Yoneya, *Phys. Rev. Lett.* 78, 1219 (1997).
 - [8] T. Yoneya, *Prog. Theor. Phys.* 103, 1081 (2000).
 - [9] Q.-G. Huang and M. Li, *J. High Energy Phys.* 06, 014 (2003).
 - [10] S. Tsujikawa, R. Maartens, and R. Brandenberger, *Phys. Lett. B* 574, 141 (2003).
 - [11] Q.-G. Huang and M. Li, *J. Cosmol. Astropart. Phys.* 11, 001 (2003).
 - [12] Q.-G. Huang and M. Li, *astro-ph/0311378*.
 - [13] H. Kim, G.S. Lee, and Y.S.M. Yung, *hep-th/0402018*.
 - [14] H. Kim, G.S. Lee, H.W. Lee, and Y.S.M. Yung, *Phys. Rev. D* 70, 043521 (2004).

- [15] R.-G. Cai, Phys. Lett. B 593, 1 (2004).
- [16] U. H. Danielsson, Phys. Rev. D 66, 023511 (2002).
- [17] R. Easter, B. R. Greene, W. H. Kinney, and G. Shiu, Phys. Rev. D 66, 023518 (2002).
- [18] J. Martin and R. H. Brandenberger, Phys. Rev. D 68, 063513 (2003).
- [19] S. Cramonini, Phys. Rev. D 68, 063514 (2003).
- [20] G. L. Alberghi, R. Casadio, and A. Tronconi, Phys. Lett. B 579, 1 (2004).
- [21] S. Shankaranarayanan and L. Sriram Kumar, hep-th/0403236.
- [22] S. Koh, S. P. Kim, and D. J. Song, gr-qc/0402065.
- [23] L. Randall and R. Sundrum, Phys. Rev. Lett. 83, 3370 (1999).
- [24] L. Randall and R. Sundrum, Phys. Rev. Lett. 83, 4690 (1999).
- [25] J. F. Dufaux, J. E. Lidsey, R. M. Aartens, and M. Sami, Phys. Rev. D 70, 083525 (2004).
- [26] G. Calcagni, Phys. Rev. D 69, 103508 (2004).
- [27] P. Horava and E. Witten, Nucl. Phys. B 460, 506 (1996).
- [28] P. Horava and E. Witten, Nucl. Phys. B 475, 94 (1996).
- [29] A. Lukas, B. A. Ovrut, K. S. Stelle, and D. Walram, Phys. Rev. D 59, 086001 (1999).
- [30] A. Lukas, B. A. Ovrut, and D. Walram, Phys. Rev. D 60, 086001 (1999).
- [31] A. Lukas, B. A. Ovrut, and D. Walram, Phys. Rev. D 61, 023506 (2000).
- [32] S. Kobayashi and K. Koyama, J. High Energy Phys. 02, 056 (2002).
- [33] R. A. Mowitt, J. Dent, and B. Dutta, hep-th/0405050.
- [34] K. Maeda and N. Ohta, Phys. Lett. B 597, 400 (2004).
- [35] J. E. Lidsey, A. R. Liddle, E. W. Kolb, E. J. Copeland, T. Barreiro, and M. Abney, Rev. Mod. Phys. 69, 373 (1997).
- [36] V. F. Mukhanov, H. A. Feldman, and R. H. Brandenberger, Phys. Rep. 215, 203 (1992).
- [37] R. M. Aartens, Living Rev. Relativity 7, 1 (2004).
- [38] S. Nojiri and S. D. Odintsov, J. High Energy Phys. 07, 049 (2000).
- [39] I. P. Neupane, J. High Energy Phys. 09, 040 (2000).
- [40] Y. M. Cho, I. P. Neupane, and P. S. Wesson, Nucl. Phys. B 621, 388 (2002).
- [41] S. Nojiri, S. D. Odintsov, and S. Ogushi, Phys. Rev. D 65, 023521 (2002).
- [42] J. E. Lidsey, S. Nojiri, and S. D. Odintsov, J. High Energy Phys. 06, 026 (2002).
- [43] S. Nojiri, S. D. Odintsov, and S. Ogushi, Int. J. Mod. Phys. A 17, 4809 (2002).
- [44] M. H. Dehghani, Phys. Rev. D 70, 064009 (2004).
- [45] M. Minamitsuji and M. Sasaki, hep-th/0404166.
- [46] Y.-S. Piao, R.-G. Cai, X. Zhang, and Y.-Z. Zhang, Phys. Rev. D 66, 121301(R) (2002).
- [47] Y.-S. Piao, Q.-G. Huang, X. Zhang, and Y.-Z. Zhang, Phys. Lett. B 570, 1 (2003).
- [48] G. Felder and L. Kofman, Phys. Rev. D 70, 046004 (2004).
- [49] M. R. Garousi, M. Sami, and S. Tsujikawa, Phys. Rev. D 70, 043536 (2004).
- [50] Z.-K. Guo and Y.-Z. Zhang, J. Cosmol. Astropart. Phys. 08, 010 (2004).
- [51] C. Kim, H. B. Kim, Y. Kim, O.-K. Kwon, and C. O. Lee, hep-th/0404242.
- [52] A. DeBenedictis, A. Das, and S. K. Koster, gr-qc/0402047.
- [53] M. R. Garousi, Nucl. Phys. B 584, 284 (2000).
- [54] G. W. Gibbons, Phys. Lett. B 537, 1 (2002).
- [55] D. Langlois, R. M. Aartens, and D. Wands, Phys. Lett. B 489, 259 (2000).
- [56] C.-S. Chu, B. R. Greene, and G. Shiu, Mod. Phys. Lett. A 16, 2231 (2001).
- [57] F. Lizzi, G. Mangano, G. Miele, and M. Peloso, J. High Energy Phys. 06, 049 (2002).
- [58] O. Bertolami and L. Guisado, Phys. Rev. D 67, 025001 (2003).
- [59] F. Lizzi, G. Mangano, G. Miele, and G. Sparano, Int. J. Mod. Phys. A 11, 2907 (1996).
- [60] F. Lizzi, G. Mangano, and G. Miele, Mod. Phys. Lett. A 16, 1 (2001).
- [61] S. Alexander, R. Brandenberger, and J. Magueijo, Phys. Rev. D 67, 081301 (2003).
- [62] M. Fukuma, Y. Kono, and A. Miwa, Nucl. Phys. B 682, 377 (2004).
- [63] M. Fukuma, Y. Kono, and A. Miwa, hep-th/0312298.
- [64] H. Garcia-Compean, O. Obregon, and C. Ramirez, Phys. Rev. Lett. 88, 161301 (2002).
- [65] G. D. Barbosa and N. Pinto-Neto, hep-th/0407111.
- [66] J. Garriga and V. F. Mukhanov, Phys. Lett. B 458, 219 (1999).
- [67] A. Frolov, L. Kofman, and A. Starobinsky, Phys. Lett. B 545, 8 (2002).
- [68] D.-J. Liu and X.-Z. Li, astro-ph/0402063.
- [69] K. Dasgupta, G. Rajesh, and S. Mukhi, J. High Energy Phys. 06, 022 (2000).
- [70] C. Bachas and M. Porrati, Phys. Lett. B 296, 77 (1992).
- [71] G. Calcagni, J. Cosmol. Astropart. Phys. 06, 002 (2004).
- [72] G. Calcagni, hep-ph/0406057.
- [73] G. Efsthathiou, Mon. Not. R. Astron. Soc. 332, 193 (2002).
- [74] S. Tsujikawa and A. R. Liddle, J. Cosmol. Astropart. Phys. 03, 001 (2004).
- [75] S. Tsujikawa, M. Sami, and R. M. Aartens, Phys. Rev. D 70, 063525 (2004).
- [76] J. R. Bond, C. Contaldi, A. Lewis, and D. Pogosyan, astro-ph/0406195.
- [77] P. Kanti, I. L. Kogan, K. A. Olive, and M. Pospelov, Phys. Lett. B 468, 31 (1999).
- [78] P. Kanti, I. L. Kogan, K. A. Olive, and M. Pospelov, Phys. Rev. D 61, 106004 (2000).
- [79] C. Csaki, J. Erlich, T. J. Hollowood, and Y. Shim, Nucl. Phys. B 581, 309 (2000).
- [80] S. Kobayashi, K. Koyama, and J. Soda, Phys. Rev. D 65, 064014 (2002).
- [81] P. Mounaix and D. Langlois, Phys. Rev. D 65, 103523 (2002).
- [82] A. Wang, Phys. Rev. D 66, 024024 (2002).
- [83] C. Barcelo, C. Gemani, and C. F. Sopuerta, Phys. Rev. D 68, 104007 (2003).
- [84] K. A. Bronnikov and B. E. Meierovich, Gravitation Cosmol. 9, 313 (2003).
- [85] K. Ghoroku and M. Yahiro, hep-th/0305150.
- [86] S. Kanno and J. Soda, J. Cosmol. Astropart. Phys. 07, 002 (2004).

This figure "fig3.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/hep-th/0406006v3>

This figure "fig4.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/hep-th/0406006v3>

This figure "fig5.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/hep-th/0406006v3>

This figure "fig6.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/hep-th/0406006v3>

This figure "fig7.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/hep-th/0406006v3>

This figure "fig8.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/hep-th/0406006v3>