

A B C of  $N = 8, d = 1$  supermultipletsS. Bellucci<sup>a</sup>, E. Ivanov<sup>b</sup>, S. Krivonos<sup>b</sup>, O. Lechtenfeld<sup>c</sup><sup>a</sup> INFN-Laboratori Nazionali di Frascati, C.P. 13, 00044 Frascati, Italy  
bellucci@lnf.infn.it<sup>b</sup> Bogoliubov Laboratory of Theoretical Physics, JINR, 141980 Dubna, Russia  
eivanov, krivonos@thsun1.jinr.ru<sup>c</sup> Institut für Theoretische Physik, Universität Hannover, 30167 Hannover, Germany  
lechtenf@itp.uni-hannover.de

## Abstract

We construct a variety of on-shell  $N = 8; d = 1$  supermultiplets with finite numbers of component fields as direct sums of properly constrained  $N = 4; d = 1$  superfields. We also show how these multiplets can be described in  $N = 8; d = 1$  superspace where the whole amount of supersymmetry is manifest. Some of these multiplets can be obtained by dimensional reduction from  $N = 2$  multiplets in  $d = 4$ , whereas others cannot. We give examples of invariant superfield actions for the multiplets constructed, including  $N = 8$  superconformally invariant ones.

# 1 Introduction

Supersymmetric quantum mechanics (SQM) [1] is relevant to a vast bunch of phenomena associated with supersymmetric theories in diverse dimensions and string theory (see e.g. [2, 3] and references therein). In particular, superconformal quantum mechanics (SCQM) [4]–[9], [3] has profound implications in the AdS/CFT [10] and black holes arena (see e.g. [11]–[18]). It is also closely related to the integrable Calogero–Moser-type systems [19, 20]. Taking into account these and some other uses, the construction and analysis of new SQM models (including SCQM models) is an urgent and interesting task with many potentially important outcomes.

Most of the SQM models explored to date possess  $N = 4; d = 1$  supersymmetries. Much less is known about higher- $N$  SQM models. Some of them were addressed many years ago in the seminal paper [21] within an on-shell Hamiltonian approach, some others (with  $N = 8$ ) received attention lately in connection with branes and black holes [22, 23, 34], and as effective theories describing low-energy dynamics of BPS monopoles in  $N = 4$  super Yang–Mills theory [24].  $N = 4$  and  $N = 8$  SQM models were also used to describe  $N = 1$  and  $N = 2$  four-dimensional supersymmetric gauge theories in a small spatial volume (see [25] and references therein).

The natural formalism for dealing with supersymmetric models is the on-shell superfield approach. Thus for the construction of new SQM models with extended  $d = 1$  supersymmetry, one needs, first of all, the complete list of the corresponding on-shell  $d = 1$  supermultiplets and the superfields which comprise these multiplets. One of the peculiarities of  $d = 1$  supersymmetry is that some of its on-shell multiplets cannot be obtained via direct dimensional reduction from the multiplets of higher- $d$  supersymmetries with the same number of spinorial charges. Another peculiarity is that some on-shell multiplets of the latter have on-shell  $d = 1$  counterparts. Taking into account these and some other specific features of  $d = 1$  supersymmetry, it is desirable to have convenient self-consistent methods of deducing  $d = 1$  on-shell multiplets and relevant superfields directly within the  $d = 1$  setting, without resorting to dimensional reduction.

One of such methods was proposed in [6] and further advanced in recent papers [7, 8, 9]. It is based on nonlinear realizations of the finite-dimensional superconformal groups in  $d = 1$  (see also [11]). The irreducible superfields representing one or another on-shell  $d = 1$  supermultiplet come out as the Goldstone superfields parametrizing one or another coset manifold of the proper  $d = 1$  superconformal group. The superfield irreducibility constraints naturally emerge as a part of manifestly covariant inverse Higgs [26] conditions on the relevant Cartan superforms. This method is advantageous in that it automatically specifies the superconformal properties of the involved supermultiplets, which are of importance, e.g. when constructing the SCQM models associated with these multiplets.<sup>1</sup>

In [8], the full set of on-shell  $N = 4$  supermultiplets with 4 physical fermions (and a finite number of auxiliary fields) was deduced, proceeding from nonlinear realizations of the

---

<sup>1</sup> Sometimes, there is no need to proceed just from superconformal algebras for deducing the correct superfield irreducibility conditions; one can make use of much simpler superalgebras which are related to the superconformal ones via a contraction and include as subalgebras the corresponding  $d = 1$  Poincaré superalgebras (see [8] for an instructive example).

most general  $N = 4; d = 1$  superconformal group  $D(2; 1; \epsilon)$  in its various coset superspaces.<sup>2</sup> Besides already known  $N = 4$  multiplets, some new ‘on-shell’ multiplets were found within this setting, viz. nonlinear ‘tensor’ and nonlinear chiral  $N = 4$  multiplets. The application of the same method to the less studied case of  $N = 8; d = 1$  supersymmetry was initiated in [9]. There we considered nonlinear realizations of the  $N = 8; d = 1$  superconformal group  $OSp(4^2|4)$  in its two different cosets and showed that two interesting  $N = 8; d = 1$  multiplets, with on-shell field contents  $(3, 8, 5)$  and  $(5, 8, 3)$ , naturally come out as the corresponding Goldstone multiplets. We constructed superconformally invariant actions for these multiplets in  $N = 4; d = 1$  superspace and revealed a surprising phenomenon which consists in the following. The same  $N = 8; d = 1$  supermultiplet admits a few non-equivalent splittings into pairs of irreducible on-shell  $N = 4; d = 1$  multiplets, such that different  $N = 4$  superconformal subgroups of  $OSp(4^2|4)$ , viz.  $SU(1; 1|2)$  and  $OSp(4^2|2)$ , are manifest for different splittings. Respectively, the on-shell component action of the given  $N = 8$  multiplet in general admits several different representations in terms of  $N = 4; d = 1$  superfields.

The basic aim of the present paper is to give a superfield description of all other linear on-shell  $N = 8; d = 1$  supermultiplets with 8 fermions, in both  $N = 8$  and  $N = 4$  superspaces.

Towards deriving an exhaustive list of on-shell  $N = 8$  supermultiplets and the relevant constrained  $N = 8; d = 1$  superfields, we could proceed in the same way as in the case of  $N = 4$  supermultiplets in [8], i.e. by considering nonlinear realizations of all known  $N = 8$  superconformal groups in their various cosets. However, this task is more complicated as compared to the  $N = 4$  case, in view of the existence of many non-equivalent  $N = 8$  superconformal groups ( $OSp(4^2|4)$ ,  $OSp(8|2)$ ,  $F(4)$  and  $SU(1; 1|4)$ , see e.g. [28]), with numerous coset manifolds.

In order to avoid these complications, we take advantage of two fortunate circumstances. Firstly, from the results of [29, 30] it follows that the field contents of linear on-shell multiplets of  $N = 8; d = 1$  supersymmetry with 8 physical fermions range from  $(8, 8, 0)$  to  $(0, 8, 8)$ , with the intermediate multiplets corresponding to all possible divisions of 8 bosonic fields into physical and auxiliary ones. Thus we are aware of the full list of such multiplets independently of the issue of their interpretation as the Goldstone ones parametrizing the proper superconformal cosets. Besides the two extreme possibilities,

$$(a) (8; 8; 0); \quad (b) (0; 8; 8); \quad (1.1)$$

we have seven further ones,

$$(c) (7; 8; 1); \quad (d) (6; 8; 2); \quad (e) (5; 8; 3); \quad (f) (4; 8; 4); \\ (g) (3; 8; 5); \quad (h) (2; 8; 6); \quad (i) (1; 8; 7); \quad (1.2)$$

The superfield description of the multiplets (1.2e) and (1.2g) was given in [9]. The superfield description of the remaining multiplets is just the subject of the present paper.

The second circumstance allowing us to advance without resorting to the nonlinear realizations techniques is the aforesaid existence of various splittings of  $N = 8$  multiplets into pairs of irreducible  $N = 4$  supermultiplets. We know how to represent the latter

---

<sup>2</sup>Two other  $N = 4; d = 1$  superconformal groups,  $SU(1; 1|2)$  and  $OSp(4^2|2)$ , can be treated as special cases of  $D(2; 1; \epsilon)$ , with  $\epsilon = 0$  or  $\epsilon = 1$  and  $\epsilon = 1$ , respectively [27, 28].

in terms of constrained  $N = 4$  super elds, so it proves to be a matter of simple algebra to guess the form of the four extra supersymmetries mixing the  $N = 4$  super elds inside each pair and extending the manifest  $N = 4$  supersymmetry to  $N = 8$ . After xing such pairs, it is again rather easy to embed them into the appropriately constrained  $N = 8; d = 1$  super elds. Beyond this, one can wonder how to reproduce these super elds within the superconformal coset techniques, as it was done for the multiplets (1.2e) and (1.2g) in [9]. We hope to return to the analysis of this important question elsewhere.

The paper is organized as follows. In the preparatory Section 2 we list all the known finite-dimensional o-shell multiplets of  $N = 4; d = 1$  Poincare supersymmetry and recall their  $N = 4$  super eld formulations, with the corresponding constraints [8]. In Section 3 we sketch the construction of the  $N = 8, d = 1$  superspace most suitable for our purposes. Section 4 is split into nine subsections related to the nine di erent  $N = 8$  o-shell multiplets we are considering. There we describe the structure of the multiplets in terms of  $N = 4$  and  $N = 8$  super elds. In particular, we construct the corresponding irreducibility constraints in  $N = 4$  superspace, as well as the implicit  $N = 4$  Poincare supersymmetry transformations, completing the manifest ones to the full  $N = 8$  supersymmetry. Invariant free actions are also constructed for all  $N = 8$  supermultiplets, and interaction terms are given for a few of them.

## 2 $N = 4, d = 1$ supermultiplets

In this Section, based upon Ref. [8], we tabulate all possible o-shell multiplets of  $N = 4; d = 1$  supersymmetry with finite number of component elds and the corresponding super elds as Goldstone multiplets of the nonlinearly realized most general  $N = 4; d = 1$  superconformal group  $D(2;1; )$  (for the generic or for some special values of this parameter).

We use the following definition of the  $N = 4; d = 1$  spinor derivatives

$$D^i = \frac{\partial}{\partial \theta_i} + i^i \partial_t; \quad \bar{D}_i = \frac{\partial}{\partial \bar{\theta}_i} + i_i \partial_t; \quad D^i; \bar{D}_j = 2i^i_j \partial_t : \quad (2.1)$$

Sometimes it will be more convenient to use the quartet form of the same derivatives:

$$D^{ia} = \frac{\partial}{\partial \theta_{ia}} + i^{ia} \partial_t \quad (D^i; D^i); \quad \quad (i; i) : \quad (2.2)$$

Respectively, there are two equivalent parametrizations of the  $N = 4; d = 1$  superspace

$$R^{(1|4)} = (t; i;^k) = (t;^{ia}) : \quad (2.3)$$

In the Table below, besides the eld content of multiplets and the corresponding super elds, we also indicate to which coset of the internal  $R$ -symmetry  $SU(2)$  of the  $N = 4; d = 1$  Poincare superalgebra the physical bosonic elds (or a subset of them) can belong and whether one of these elds can be identified with the Goldstone coset eld (dilaton) associated with the dilatation generator  $D(2;1; )$ . On all these multiplets, except for the chiral one, the supergroup  $D(2;1; )$  can be realized at any value of the parameter . The  $N = 4; d = 1$  chirality is compatible only with the choice of the supergroup  $SU(1;1|2)$  as the superconformal one.

### N = 4 supermultiplets

multiplet	content	R symm.	dilaton		SF
old tensor	(1, 4, 3)	{	yes	any	u
chiral	(2, 4, 2)	c.charge	yes	0; 1	;
nl. chiral	(2, 4, 2)	su(2)=u(1)	no	any	;
tensor	(3, 4, 1)	su(2)=u(1)	yes	any	$V^{ij}$
nonlinear	(3, 4, 1)	su(2)	no	any	$N^{ia}$
hyper	(4, 4, 0)	su(2)	yes	any	$q^i; q_i$
ferm. hyper	(0, 4, 4)	{	no	any	$^a;_a$

The constraints read as follows:

Old Tensor:  $D^i D_i u = \bar{D}^i \bar{D}_i u = 0$   
 (Note:  $D^i; \bar{D}_i u = \text{const}$ ; generic is compatible only with  $\text{const} = 0$ , otherwise  $= 0$  or  $-1$ )

Chiral:  $D^i = \bar{D}_i = 0$   
 (Note: it exists only for  $D(2;1;-1)$  or  $D(2;1;0)$   $\rightarrow$   $su(1;1|2) \rightarrow su(2)$ )

Nonlinear Chiral:  $D^1 = D^2; \bar{D}_2 = \bar{D}_1$  (and c.c.)

Tensor:  $D^{(i} V^{jk)} = \bar{D}^{(i} V^{jk)} = 0$

Nonlinear:  $N^a (D^i D^j N_a^k) = N^a (\bar{D}^i \bar{D}^j N_a^k) = 0; N^{ai} N_{ai} = 2$

Hypermultiplet:  $D^{(i} q^{j)} = \bar{D}^{(i} q^{j)} = 0$  (and c.c.)

Fermionic Hypermultiplet:  $D^{(i} \eta^{j)} = \bar{D}^{(i} \eta^{j)} = 0$  (and c.c.)

### 3 N = 8, d=1 superspace

The maximal automorphism group of  $N = 8; d=1$  super Poincare algebra (without central charges) is  $SO(8)$  and so eight real Grassmann coordinates of  $N = 8; d=1$  superspace  $R^{(1|8)}$  can be arranged into one of three 8-dimensional real irreps of  $SO(8)$ . The constraints defining the irreducible  $N = 8$  supermultiplets in general break this  $SO(8)$  symmetry. So, it is preferable to split the 8 coordinates into two real quartets

$$R^{(1|8)} = (t;_{ia}; \#_A); \quad \overline{(\#_{ia})} = ^{ia}; \overline{(\#_A)} = \#^A; \quad i; a; \quad ; A = 1; 2; \quad (3.1)$$

in terms of which only four commuting automorphism  $SU(2)$  groups will be explicit. The further symmetry breaking can be understood as identification of some of these  $SU(2)$ , while extra symmetries, if exist, mix different  $SU(2)$  indices. The corresponding covariant derivatives are defined by

$$D^{ia} = \frac{\partial}{\partial_{ia}} + i^{ia} \partial_t; \quad r^A = \frac{\partial}{\partial \#_A} + i \#^A \partial_t; \quad (3.2)$$

By construction, they obey the algebra:<sup>3</sup>

$$D^{ia}; D^{jb} = 2i^{ijab} \partial_t; \quad r^A; r^B = 2i^{AB} \partial_t : \quad (3.3)$$

Note that the dimensional reduction from  $N=2; d=4$  superspace to  $d=1$  yields a superspace with two manifest  $SU(2)$  automorphism groups realized on the Grassmann coordinates: one of them is just the  $R$ -symmetry of  $N=2; d=4$  Poincare superalgebra and the other originates from the  $d=4$  Lorentz group  $SL(2; C)$ . These two  $SU(2)$  correspond to the appropriate identifying of the doublet indices of two groups  $SU(2)$  in the above product of four such groups. In what follows we shall not be bound to any sort of dimensional reduction; the differential constraints on the relevant  $N=8; d=1$  super elds will be deduced directly within the  $d=1$  setting, in terms of the spinor covariant derivatives (3.2).

## 4 $N=8, d=1$ supermultiplets

As mentioned in Introduction, our real strategy of deducing a super eld description of the  $N=8; d=1$  supermultiplets (1.1), (1.2) consisted in selecting the appropriate pair of the constrained  $N=4; d=1$  super elds from those listed in Section 2 and then in guessing the constrained  $N=8$  super eld which accommodate this pair in a most economical way. In the Subsections below, just to make the presentation more coherent, we turn the argument around and start with postulating the  $N=8; d=1$  constraints. The  $N=4$  super eld formulations will be deduced from the  $N=8$  ones.

### 4.1 Supermultiplet $(0, 8, 8)$

The on-shell  $N=8; d=1$  supermultiplet  $(0, 8, 8)$  is carried out by two real fermionic  $N=8$  super elds  $\psi^{aA}; \chi^i$  subjected to the following constraints:

$$D^{(ia} \chi^{j)} = 0; D^{i(a} \psi_{A}^{b)} = 0; r^{(A} \chi_{i}^{B)} = 0; r^{(A} \psi_{a}^{B)} = 0; \quad (4.1)$$

$$r^A \psi_a^A = D^{ia} \chi_i; \quad r^A \chi_i = D^{ia} \psi_a^A : \quad (4.2)$$

In order to understand the structure of this supermultiplet in terms of  $N=4$  super elds and to prove that the above constrained  $N=8$  super elds indeed carry the irreducible on-shell eld content  $(0, 8, 8)$ , we proceed as follows. As the first step, let us single out the  $N=4$  subspace in the  $N=8$  superspace  $R^{(1\mathcal{B})}$  as the set of coordinates

$$R^{(1\mathcal{A})} = (t; \psi_{ia}) \quad R^{(1\mathcal{B})}; \quad (4.3)$$

and expand the  $N=8$  super elds over the extra Grassmann coordinate  $\theta_A$ . Then we observe that the constraints (4.2) imply that the spinor derivatives of all involved super elds with respect to  $\theta_A$  can be expressed in terms of spinor derivatives with respect to  $\psi_{ia}$ .

---

<sup>3</sup>We use the following convention for the skew-symmetric tensor :  $\epsilon_{ij}{}^{jk} = \delta_i^k$ ;  $\epsilon_{12} = \epsilon^{21} = 1$ .

This means that the only essential  $N = 4$  super eld components of  ${}^{aA}$  and  ${}_i$  in their  $\#$ -expansion are the first ones

$${}^{aA} = {}^{aA}{}_{j=0} ; \quad {}_i = {}_i{}_{j=0} : \quad (4.4)$$

These fermionic  $N = 4$  super elds are subjected, in virtue of eqs. (4.1) and (4.2), to the irreducibility constraints in  $N = 4$  superspace

$$D^{a(i}{}_j) = 0 ; D^{i(a}{}_b)A = 0 : \quad (4.5)$$

As follows from the list of Section 2, these super elds are just two fermionic  $N = 4$  hypermultiplets, each carrying  $(0, 4, 4)$  independent component elds. So, being combined together, they accommodate the whole on-shell component content of the  $N = 8$  multiplet  $(0, 8, 8)$ , which proves that the  $N = 8$  constraints (4.1), (4.2) are the true choice.

Thus, from the  $N = 4$  superspace perspective, the  $N = 8$  supermultiplet  $(0, 8, 8)$  amounts to the sum of two  $N = 4; d = 1$  fermionic hypermultiplets with the on-shell component content  $(0, 4, 4) + (0, 4, 4)$ .

The transformations of the implicit  $N = 4$  Poincare supersymmetry, completing the manifest one to the full  $N = 8$  supersymmetry, have the following form in terms of the  $N = 4$  super elds defined above:

$${}^{aA} = \frac{1}{2} A D^{ia}{}_i ; \quad {}_i = \frac{1}{2} A D_a{}^i {}^{aA} : \quad (4.6)$$

The compatibility of (4.6) with the constraints (4.5) is verified using the following corollaries of these constraints:

$$D^{ia}D^{jb}{}_c A = 2i^{ij}{}_c{}^b {}^{aA} ; D^{ia}D^{jb}{}_k = 2i^{ab}{}_k{}^j {}_i : \quad (4.7)$$

The invariant free action can be written as

$$S = \int dt d^4x \left( \frac{1}{2} \frac{1}{A} D^{ia}{}_i D_a{}^j {}^{aA} + \frac{1}{2} \frac{1}{A} D_a{}^i {}^{aA} D^{ia}{}_i \right) : \quad (4.8)$$

Because of the presence of explicit theta's in the action (4.8), the latter is not manifestly invariant even with respect to the manifest  $N = 4$  supersymmetry. Nevertheless, one can check that (4.8) is invariant under this supersymmetry which is realized on the super elds as

$${}^{aA} = \delta^{aA} Q^{jb}{}_j {}^{aA} ; \quad {}_i = \delta_i{}^j Q^{jb}{}_j {}_i ; \quad (4.9)$$

where

$$Q^{ia} = \frac{\partial}{\partial \theta_{ia}} \quad {}_i {}^{ia} \partial_t ; \quad (4.10)$$

$\theta_{ia}$  is the supertranslation parameter and  $\delta$  means the 'active' variation (taken at a fixed point of the  $N = 4$  superspace). Actually, the action (4.8) can be given the manifestly  $N = 4$  supersymmetric form as a sum of two integrals over the appropriate harmonic analytic subspaces of  $N = 4; d = 1$  superspace (see [31]). Below we will employ this equivalent harmonic superspace form of the on-shell action of the fermionic hypermultiplet (Subsections 4.2 and 4.5), as well as one more its form, as an integral over the chiral  $N = 4; d = 1$  superspace (Subsection 4.3).

## 4.2 Supermultiplet (1, 8, 7)

This supermultiplet can be described by a single scalar  $N = 8$  superfield  $U$  which obeys the following irreducibility conditions:

$$D^{ia} D_a^j U = r^j r^i U; \quad (4.11)$$

$$r^{(i} r^{j)} U = 0; \quad D^{i(a} D^{j b)} U = 0; \quad (4.12)$$

Let us note that the constraints (4.11) reduce the manifest  $R$ -symmetry to  $[SU(2)]^3$  due to the identification of the indices  $i$  and  $A$  of the covariant derivatives  $D^{ia}$  and  $r^A$ .

This supermultiplet possesses a unique decomposition into the pair of the  $N = 4$  supermultiplets as  $(1, 8, 7) = (1, 4, 3) + (0, 4, 4)$ . The corresponding  $N = 4$  superfield projections can be defined as

$$u = U|_{\theta=0}; \quad \psi^i = r^i U|_{\theta=0}; \quad (4.13)$$

and they obey the standard constraints

$$D^{(ia} D^{j)} \psi^i = 0; \quad D^{i(a} D^{j b)} u = 0; \quad (4.14)$$

The second constraint directly follows from (4.12), while the first one is implied by the relation

$$\frac{\partial}{\partial t} D_a^{(i} r^{j)} U = 0; \quad (4.15)$$

which can be proven by applying the differential operator  $D^{kb} r^i$  to the  $N = 8$  superfield constraint (4.11) and making use of the algebra of covariant derivatives.

The additional implicit  $N = 4$  supersymmetry is realized on these  $N = 4$  superfields as follows:

$$u = \psi_i \psi^i; \quad \psi^i = \frac{1}{2} \psi_j D^{ia} D_a^j u; \quad (4.16)$$

The simplest way to deal with the action for this supermultiplet is to use harmonic superspace [32, 33, 31], at least for the spinor superfields  $\psi^i$ .

We use the definitions and conventions of Ref. [31]. The harmonic variables parametrizing the coset  $SU(2)_R/U(1)_R$  are defined by the relations

$$u^+ u_i = 1, \quad u_i^+ u_j = u_j^+ u_i = \delta_{ij}; \quad \overline{(u^+ i)} = u_i; \quad (4.17)$$

The harmonic projections of  $\psi^i = \psi^i$ ;  $\psi_i$  are defined by

$$\psi^+ = \psi_i^+; \quad \psi^- = \psi_i^+; \quad (4.18)$$

and the constraints (4.14) are rewritten as

$$D^+ D^- u = \overline{D^+} \overline{D^-} u = 0; \quad D^+ \overline{D^-} u + \overline{D^+} D^- u = 0; \quad (4.19)$$

$$D^+ \psi^+ = \overline{D^+} \psi^+ = 0; \quad D^{++} \psi^+ = \overline{D^{++}} \psi^+ = 0; \quad (4.20)$$

Here  $D^i = D^i$ ;  $\overline{D^i} = \overline{D^i}$  and  $D^- = D^i u_i$ ;  $\overline{D^-} = \overline{D^i} u_i$ ,  $D^+ = u^i \partial/\partial u^i$  (in the central basis of the harmonic superspace), with  $D^{ia}$  given in (2.2). The relations (4.20) imply



that  $\psi^+; \bar{\psi}^+$  are analytic harmonic  $N=4; d=1$  super elds living on the analytic subspace  $(t_A; \psi^+; \bar{\psi}^+; u_i)$  which is closed under  $N=4$  supersymmetry. In this setting, the transformations of the hidden  $N=4$  supersymmetry (4.16) are rewritten as

$$\begin{aligned}\psi^+ &= \frac{1}{2} \psi^+ D^+ \bar{D}^+ - \bar{\psi}^+ \bar{D}^+ D - \frac{1}{2} D^+ \bar{D}^+ u; \\ \bar{\psi}^+ &= \frac{1}{2} \bar{\psi}^+ D^+ \bar{D}^+ - \psi^+ \bar{D}^+ D - \frac{1}{2} \bar{D}^+ D^+ u; \\ u &= \psi^+ \psi^+ + \bar{\psi}^+ \bar{\psi}^+ + \psi^+ \bar{\psi}^+ : \end{aligned} \quad (4.21)$$

The action is given by

$$S = \frac{1}{2} \int dtd^4 u^2 + \int dud \psi^+ \bar{\psi}^+ \quad (4.22)$$

where  $dud = dtd_A d^+ d^+$  is the measure of integration over the analytic superspace. The action (4.22) is manifestly  $N=4$  supersymmetric since it is written in terms of  $N=4$  super elds. However, its invariance with respect to the hidden  $N=4$  supersymmetry (4.21) must be explicitly checked. The variation of the first term in (4.22) can be represented as (we explicitly write only the terms involving the parameters)

$$\int dtd^4 \frac{u^2}{2} = \int dtd^4 du \psi^+ \bar{\psi}^+ u = 2 \int dtd^4 du \psi^+ u; \quad (4.23)$$

while the variation of the second term reads

$$\int dud \psi^+ \bar{\psi}^+ = 2 \int dud D^+ \bar{D}^+ \psi^+ u; \quad (4.24)$$

Keeping in mind that

$$\int dud D^+ \bar{D}^+ = \int dtd^4 du; \quad (4.25)$$

we see that the action (4.22) is indeed invariant under  $N=8$  supersymmetry.

### 4.3 Supermultiplet $(2, 8, 6)$

The  $N=8$  super eld formulation of this supermultiplet involves two scalar bosonic super elds  $U; \bar{U}$  obeying the constraints

$$r^{(ai} r^{b)j} U = 0; \quad r^{a(i} r^{bj)} \bar{U} = 0; \quad (4.26)$$

$$r^{ai} U = D^{ia} \bar{U}; \quad r^{ai} \bar{U} = D^{ia} U \quad (4.27)$$

where we have identified indices  $i$  and  $A$ ,  $a$  and  $\alpha$  of the covariant derivatives, thus retaining only two manifest  $SU(2)$  automorphism groups. From (4.26), (4.27) some useful corollaries follow:

$$D^{ia} D_a^j U + r^{aj} r_a^i U = 0; \quad D^{i(a} D^{j)b} U = 0; \quad (4.28)$$

$$D^{ia} D_i^b \bar{U} + r^{bi} r_a^i \bar{U} = 0; \quad D^{(ia} D^{j)b} \bar{U} = 0; \quad (4.29)$$

Comparing (4.28), (4.29) and (4.26) with (4.11), (4.12), we observe that the  $N = 8$  supermultiplet with the eld content  $(2, 8, 6)$  can be obtained by combining two  $(1, 8, 7)$  supermultiplets and imposing the additional relations (4.27) on the corresponding  $N = 8$  super elds.

In order to construct the invariant actions and to prove that the above  $N = 8$  constraints indeed yield the multiplet  $(2, 8, 6)$ , we should reveal the structure of this supermultiplet in terms of  $N = 4$  super elds, as we did in the previous cases. However, in the case at hand, we have two different choices to split the  $(2, 8, 6)$  supermultiplet

1.  $(2, 8, 6) = (1, 4, 3) \quad (1, 4, 3)$
2.  $(2, 8, 6) = (2, 4, 2) \quad (0, 4, 4)$

As already mentioned, the possibility to have a few different on-shell  $N = 4$  decompositions of the same  $N = 8$  multiplet has been observed in our earlier paper [9]. It is related to different choices of the manifest  $N = 4$  supersymmetries as subgroups of the  $N = 8$  super Poincare group. We shall treat both options.

1.  $(2, 8, 6) = (1, 4, 3) \quad (1, 4, 3)$

In order to describe the  $N = 8$   $(2, 8, 6)$  multiplet in terms of  $N = 4$  super elds we should choose the appropriate  $N = 4$  superspace. The first (evident) possibility is to choose the  $N = 4$  superspace with the coordinates  $(t; \text{ia})$  i.e. the same as in Subsections 4.1 and 4.2 (see eq. (4.3)). In this superspace one  $N = 4$  Poincare supergroup is naturally realized, while the second one mixes two irreducible  $N = 4$  super elds which comprise the  $N = 8$   $(2, 8, 6)$  supermultiplet in question. Expanding the  $N = 8$  super elds  $U; \text{ in } \#^{\text{ia}}$ , one finds that the constraints (4.26), (4.27) leave in  $U$  and as independent  $N = 4$  projections only those of zeroth order in  $\#^{\text{ia}}$

$$u = U|_{\#^{\text{ia}}=0}; \quad = |_{\#^{\text{ia}}=0} : \quad (4.30)$$

Each  $N = 4$  super eld proves to be subjected, in virtue of (4.26), (4.27), to the additional constraint:

$$D^{i(a} D^{j)b} u = 0; \quad D^{(ia} D^{j)b} = 0 : \quad (4.31)$$

Thus we conclude that our  $N = 8$  multiplet  $U; \text{ , when rewritten in terms of } N = 4$  super elds, amounts to a direct sum of two  $N = 4$  multiplets  $u$  and  $\text{ , both having the same on-shell eld contents } (1, 4, 3)$ .

The transformations of the implicit  $N = 4$  Poincare supersymmetry completing the manifest one to the full  $N = 8$  Poincare supersymmetry have the following form in terms of these  $N = 4$  super elds:

$$u = \text{ia} D^{\text{ia}} ; \quad = \text{ia} D^{\text{ia}} u : \quad (4.32)$$

It is rather easy to construct the action in terms of  $N = 4$  super elds  $u$  and  $\text{ , such that it is invariant with respect to the implicit } N = 4$  supersymmetry (4.32). The generic action has the form

$$S = \int d^4x \text{ F}(u; \text{ );} \quad (4.33)$$

where the function  $F$  obeys the Laplace equation

$$F_{uu} + F = 0 : \quad (4.34)$$

The simplest example of such an action is supplied by the free action

$$S_{\text{free}} = \frac{1}{2} \int dtd^4 \left( u^2 - \bar{u}^2 \right) : \quad (4.35)$$

One may wonder whether the latter action is positively defined or not. After passing to the component fields we find that, with the following definition of the superspace integration measure

$$\int dtd^4 = \frac{1}{24} \int dt D^{ia} D_{ja} D^{jb} D_{ib} ; \quad (4.36)$$

and with the auxiliary fields eliminated by their equations of motion, the action (4.35) yields the correct kinetic terms for the physical bosons

$$S_{\text{free}} = \int dt \left( \frac{u^2}{2} + \frac{\bar{u}^2}{2} + \text{fermions} \right) : \quad (4.37)$$

The sign minus between two terms in (4.35) is actually related to the difference between the defining constraints (4.31) for the superfields  $u$  and  $\bar{u}$  and has no impact on the positive definiteness of the component action. Below we shall meet more examples of such "fake \non-positive definiteness" of the superfield actions. The corresponding component actions are correct in all cases.

An interesting subclass of the actions (4.33) is provided by those of the form :

$$S = \int dtd^4 \left( f(u) + \sum_{n=1}^{\infty} g_n(u) \bar{u}^{2n} \right) ; \quad (4.38)$$

with the additional conditions (which follow from (4.34))

$$2g_1 = f^{(0)} ; \quad g_{n+1} = \frac{g_n^{(0)}}{2(2n+1)(n+1)} ; \quad (4.39)$$

where the primes denote the derivatives with respect to  $u$ . The immediate corollary is that any action written in terms of the  $N=4$  superfield  $u$  can be promoted to an invariant of  $N=8$  supersymmetry by adding the appropriate interaction with the superfield  $\bar{u}$ .<sup>4</sup> For example, the action of  $N=4$  SCQM [6]

$$S_{N=4} = \int dtd^4 u \log u$$

---

<sup>4</sup>One can equally start with an action of the superfield  $\bar{u}$  and promote it to an  $N=8$  supersymmetric action by turning on the appropriate interactions with  $u$ .

can be generalized to have  $N = 8$  supersymmetry as

$$\begin{aligned} S_{N=8} &= \int d^4x \left[ \frac{1}{2} u \log(u^2 + v^2) - \arctan \frac{v}{u} \right] \\ &= \frac{1}{2} \int d^4x [(u + i v) \log(u + i v) + (u - i v) \log(u - i v)]: \end{aligned} \quad (4.40)$$

Note that the  $N = 8$  superconformal invariance of this action is not automatic and remains to be checked.

$$2. \quad (2, 8, 6) = (2, 4, 2) \quad (0, 4, 4)$$

There is a more sophisticated choice of a  $N = 4$  subspace in the  $N = 8; d = 1$  superspace, which gives rise to the second possible  $N = 4$  superfield splitting of the considered  $N = 8$  supermultiplet, that is into the multiplets  $(2, 4, 2)$  and  $(0, 4, 4)$ .

First of all, let us define a new set of covariant derivatives

$$D^{ia} = \frac{1}{2} D^{ia} - i r^{ai}; \quad \bar{D}^{ia} = \frac{1}{2} D^{ia} + i r^{ai}; \quad D^{ia}; \bar{D}^{jb} = 2i \epsilon^{ij} \epsilon^{ab} \partial_t; \quad (4.41)$$

and new  $N = 8$  superfields  $V; \bar{V}$  related to the original ones as

$$V = U + i; \quad \bar{V} = U - i; \quad (4.42)$$

In this basis the constraints (4.26), (4.27) read

$$\begin{aligned} D^{ia} V &= 0; \quad \bar{D}^{ia} \bar{V} = 0; \\ D^{i(a} D^{j b)} \bar{V} + \bar{D}^{i(a} \bar{D}^{j b)} V &= 0; \quad D^{(ia} D^{j)b} \bar{V} - \bar{D}^{(ia} \bar{D}^{j)b} V = 0; \end{aligned} \quad (4.43)$$

Now we split the complex quartet covariant derivatives (4.41) into two sets of the doublet  $N = 4$  ones as

$$D^i = D^{i1}; \quad \bar{D}^i = \bar{D}^{i2}; \quad r^i = D^{i2}; \quad \bar{r}^i = \bar{D}^{i1} \quad (4.44)$$

and cast the constraints (4.43) in the form

$$\begin{aligned} D^i V &= 0; \quad r^i V = 0; \quad \bar{D}^i \bar{V} = 0; \quad \bar{r}^i \bar{V} = 0; \\ D^i D^j \bar{V} - \bar{r}^i \bar{r}^j V &= 0; \quad D^i r^j \bar{V} - \bar{D}^i \bar{r}^j V = 0; \end{aligned} \quad (4.45)$$

Next, as an alternative  $N = 4$  superspace, we choose the set of coordinates closed under the action of  $D^i; \bar{D}^i$ , i.e.

$$(t; x_{i1} + i x_{i2}; x_{i2} - i x_{i1}); \quad (4.46)$$

while the  $N = 8$  superfields are expanded with respect to the orthogonal combinations  $x_{i1} - i x_{i2}, x_{i2} + i x_{i1}$  which are annihilated by  $D^i; \bar{D}^i$ .

As a consequence of the constraints (4.45), the quadratic action of the derivatives  $r^i$  and  $\bar{r}^i$  on every  $N = 8$  superfield  $V; \bar{V}$  can be expressed as  $D^i; \bar{D}^i$  of some other superfield. Therefore, only the zeroth and first order components of each  $N = 8$  superfield are

independent  $N = 4$  super eld projections. Thus, we are left with the following set of  $N = 4$  super elds:

$$v = V^j; \quad \bar{v} = \bar{V}^j; \quad v^i = \bar{r}^i v; \quad \bar{v}^i = r^i \bar{v}; \quad (4.47)$$

These  $N = 4$  super elds prove to be subjected to the additional constraints which also follow from (4.45)

$$D^i v = 0; \quad \bar{D}^i \bar{v} = 0; \quad D^{ij} = 0; \quad \bar{D}^{ij} = 0; \quad D^{ij} = \bar{D}^{ij}; \quad (4.48)$$

The  $N = 4$  super elds  $v, \bar{v}$  comprise the standard  $N = 4; d = 1$  chiral multiplet  $(2, 4, 2)$ , while the  $N = 4$  super elds  $v^i, \bar{v}^i$  subjected to (4.48) and both having the on-shell contents  $(0, 4, 4)$  are recognized as the fermionic version of the  $N = 4; d = 1$  hypermultiplet.

The implicit  $N = 4$  supersymmetry is realized by the transformations

$$\begin{aligned} v &= v^i; \quad v^i = \frac{1}{2} D^2 v - 2i v^i; \\ \bar{v} &= \bar{v}^i; \quad \bar{v}^i = \frac{1}{2} \bar{D}^2 \bar{v} + 2i \bar{v}^i. \end{aligned} \quad (4.49)$$

The invariant free action has the following form :

$$S_f = \int d^4x \, v \bar{v} - \frac{1}{2} \int d^4x \, v^i v_i - \frac{1}{2} \int d^4x \, \bar{v}^i \bar{v}_i; \quad (4.50)$$

Let us note that this very simple form of the action for the  $N = 4$   $(0, 4, 4)$  supermultiplet  $v^i, \bar{v}^i$  is related to our choice of the  $N = 4$  superspace. Indeed, with our definition (4.46), the  $(0, 4, 4)$  supermultiplet is described by the doublet of chiral fermions with the additional constraints (4.48) an essential part of which is simply the  $N = 4; d = 1$  chirality conditions. Therefore, the chiral superspace is the best choice to write the on-shell action of the multiplet  $(0, 4, 4)$  in the present case. It is worthwhile to emphasize that all differently looking superspace on-shell actions of the multiplet  $(0, 4, 4)$ , viz. both terms in (4.8), the action (4.22) and the last two terms in (4.50), yield the same component action for this multiplet.

#### 4.4 Supermultiplet $(3, 8, 5)$

This supermultiplet has been discussed in detail in [9] where it was termed as the  $N = 8$  tensor multiplet'. Here we shortly remind its main features.

In the  $N = 8$  superspace this supermultiplet is described by the triplet of bosonic super elds  $V^{ij}$  obeying the irreducibility constraints

$$D_a^{(i} V^{jk)} = 0; \quad r^{(i} V^{jk)} = 0; \quad (4.51)$$

So three out of four original automorphism  $SU(2)$  symmetries remain manifest in this description.

The  $N = 8$  supermultiplet  $(3, 8, 5)$  can be decomposed into  $N = 4$  supermultiplets in the two ways

$$1. \quad (3, 8, 5) = (3, 4, 1) \quad (0, 4, 4)$$

$$2. \quad (3, 8, 5) = (1, 4, 3) \quad (2, 4, 2)$$

As in the previous case, we discuss both these options.

$$1. \quad (3, 8, 5) = (3, 4, 1) \quad (0, 4, 4)$$

This splitting requires choosing the coordinate set (4.3) as the relevant  $N = 4$  superspace. Expanding the  $N = 8$  super elds  $V^{ij}$  in  $\#_i$ , one finds that the constraints (4.51) leave in  $V^{ij}$  the following four bosonic and four fermionic  $N = 4$  projections:

$$v^{ij} = V^{ij} ; \quad \quad \quad r_j V^{ij} ; \quad A \quad r_i r_j V^{ij} \quad (4.52)$$

where  $j$  means restriction to  $\#_i = 0$ . As a consequence of (4.51), these  $N = 4$  super elds prove to obey the constraints

$$D_a^{(i} v^{jk)} = 0 ; \quad D_a^{(i} j) = 0 ;$$

$$A = 6m \quad D_i^a D_{aj} v^{ij} ; m = \text{const} : \quad (4.53)$$

Thus, for the considered splitting, the  $N = 8$  tensor multiplet super eld  $V^{ij}$  amounts to a direct sum of the  $N = 4$  tensor multiplet super eld  $v^{ij}$  with the  $o$ -shell content  $(3; 4; 1)$  and a fermionic  $N = 4$  hypermultiplet  $r_j$  with the  $o$ -shell content  $(0, 4, 4)$ , plus a constant  $m$  of the mass dimension.

The transformations of the implicit  $N = 4$  Poincare supersymmetry which together with the manifest  $N = 4$  supersymmetry constitute the full  $o$ -shell  $N = 8$  Poincare supersymmetry have the following form in terms of the above  $N = 4$  super elds:

$$v^{ij} = \quad (i j) \quad -(i j) ; \quad \quad \quad r_j = \quad 2i \Gamma_j v^{ij} \quad \frac{1}{3} i D_j \overline{D}_k v^{jk} + 6m \quad i \quad (4.54)$$

where

$$i \quad 1i ; \quad -i \quad 2i ; \quad i \quad 1i ; \quad i \quad 2i ; \quad (4.55)$$

$$2. \quad (3, 8, 5) = (1, 4, 3) \quad (2, 4, 2)$$

This option corresponds to another choice of the  $N = 4$  superspace, which amounts to dividing the  $N = 8; d = 1$  Grassmann coordinates into doublets with respect to some other  $SU(2)$  indices. The relevant splitting of  $N = 8$  superspace into the  $N = 4$  subspace and the complement of the latter can be performed as follows. Firstly, we define the new covariant derivatives as

$$D^a = \frac{1}{\sqrt{2}} D^{1a} + i r^{a1} ; \quad \overline{D}_a = \frac{1}{\sqrt{2}} D_a^2 - i r_a^2 ;$$

$$r^a = \frac{i}{\sqrt{2}} D^{2a} + i r^{a2} ; \quad r_a = \frac{i}{\sqrt{2}} D_a^1 - i r_a^1 : \quad (4.56)$$

Then we choose the set of coordinates closed under the action of  $D^a; D_a$ , i.e.

$$t; \quad x_a \quad i\#_{a1}; \quad x^a + i\#^{a1}; \quad (4.57)$$

while the  $N=8$  super elds are expanded with respect to the orthogonal combinations  $i\#_2^a, \quad x_1^a + i\#_1^a$  annihilated by  $D^a; D_a$ .

The basic constraints (4.51), being rewritten in the basis (4.56), take the form

$$\begin{aligned} D^a v &= 0; \quad D^a v \quad r^a = 0; \quad r^a v + D^a = 0; \quad r^a = 0; \\ \bar{r}_a' &= 0; \quad \bar{r}_a v + \bar{D}_a' = 0; \quad \bar{D}_a v \quad \bar{r}_a' = 0; \quad \bar{D}_a' = 0 \end{aligned} \quad (4.58)$$

where

$$v \quad 2iV^{12}; \quad ' \quad V^{11}; \quad ' \quad V^{22}; \quad (4.59)$$

Due to the constraints (4.58), the derivatives  $r^a$  and  $\bar{r}_a$  of every  $N=8$  super eld in the triplet  $(V^{12}; V^{11}; V^{22})$  can be expressed as  $D^a; \bar{D}_a$  of some other super eld. Therefore, only the zeroth order (i.e. taken at  $i\#_2^a = x_1^a + i\#_1^a = 0$ ) components of each  $N=8$  super eld are independent  $N=4$  super eld projections. These  $N=4$  super elds are subjected to the additional constraints which also follow from (4.58)

$$D^a D_a v = \bar{D}_a \bar{D}^a v = 0; \quad D^a' = 0; \quad \bar{D}_a' = 0; \quad (4.60)$$

The  $N=4$  super elds  $' ; ' \quad$  comprise the standard  $N=4; d=1$  chiral multiplet  $(2, 4, 2)$ , while the  $N=4$  super eld  $v$  subjected to (4.60) has the needed on-shell content  $(1, 4, 3)$ .

The implicit  $N=4$  supersymmetry acts on the  $N=4$  super elds  $v; ' ; ' \quad$  as follows:

$$v = x_a D^a' + x^a \bar{D}_a'; \quad ' = x_a D^a v; \quad ' = x^a \bar{D}_a v; \quad (4.61)$$

The invariant  $N=4$  super eld actions for both decompositions of the  $N=8$  multiplet  $(3, 8, 5)$  were presented in Ref. [9].

Finally, we note that the considered multiplet can be treated as a dimensional reduction of the  $N=2; d=4$  tensor multiplet, whence its name  $N=8$  tensor multiplet' used in [9]. Three physical bosons of the  $d=1$  case as compared to four bosonic degrees of freedom of the  $d=4$  case are related to the fact that one such degrees in  $d=4$  is represented by the 'notoph'. After reduction to  $d=1$  the notoph eld strength becomes a constant, and it is just the constant  $m$  appearing in (4.53).

## 4.5 Supermultiplet $(4, 8, 4)$

This supermultiplet can be described by a quartet of  $N=8$  super elds  $Q^a$  obeying the constraints:

$$D_i^{(a} Q^{b)} = 0; \quad r_i^{(} Q_a^{)} = 0; \quad (4.62)$$

Let us note that the constraints (4.62) are manifestly covariant with respect to three  $SU(2)$  subgroups realized on the indices  $i; a$  and  $.$

From (4.62) some important relations follow:

$$D^{ia} D^{jb} Q^c = 2i^{ij} \epsilon^{ab} Q^a; \quad r^i r^j Q^a = 2i^{ij} \quad Q^a; \quad (4.63)$$

Using them, it is possible to show that the super elds  $Q^a$  contain the following independent components:

$$Q^a_j; \quad D^i_a Q^a; \quad r^i Q^a; \quad D^i_a r^j Q^a; \quad (4.64)$$

where  $j$  means now restriction to  $_{ia} = \#_i = 0$ . This directly proves that we deal with the irreducible  $(4, 8, 4)$  supermultiplet.

In order to construct the corresponding action, we pass to  $N = 4$  super elds. There are three different possibilities to split this  $N = 8$  multiplet into the  $N = 4$  ones:

1.  $(4, 8, 4) = (4, 4, 0) \quad (0, 4, 4)$
2.  $(4, 8, 4) = (3, 4, 1) \quad (1, 4, 3)$
3.  $(4, 8, 4) = (2, 4, 2) \quad (2, 4, 2)$

Once again, we shall consider all these three cases separately.

1.  $(4, 8, 4) = (4, 4, 0) \quad (0, 4, 4)$

This case implies the choice of the  $N = 4$  superspace (4.3). Expanding the  $N = 8$  super elds  $Q^a$  in  $\#_i$ , one may easily see that the constraints (4.62) leave in  $Q^a$  the following four bosonic and four fermionic  $N = 4$  super eld projections:

$$q^a = Q^a_j; \quad _{ia} \quad r^i Q^a : \quad (4.65)$$

Each  $N = 4$  super eld is subjected, in virtue of (4.62), to an additional constraint

$$D^{i(a} q^{b)} = 0; \quad D^{i(a} _{b)i} = 0: \quad (4.66)$$

Consulting Section 1, we come to the conclusion that these are just the hypermultiplet  $q^i$  with the on-shell eld content  $(4, 4, 0)$  and a fermionic analog of the  $N = 4$  hypermultiplet  $_{ia}$  with the eld content  $(0, 4, 4)$ .

The transformations of the implicit  $N = 4$  Poincare supersymmetry have the following form in terms of these  $N = 4$  super elds:

$$q^a = \frac{1}{2} \epsilon^i \epsilon^a_i; \quad _{ia} = 2i \epsilon^i q^a_i : \quad (4.67)$$

Harmonic superspace provides the most adequate framework for constructing the action for this splitting. We introduce the harmonic variables parametrizing the coset  $SU(2)/U(1)$  as in eq. (4.17) and define the harmonic projections of  $q^a; \quad _{ai}$  by

$$q^+ = q^a u^+_a; \quad _{+i} = _{ia} u^+_a; : \quad (4.68)$$

Now the constraints (4.66) take the standard form of the Grassmann harmonic analyticity conditions

$$D^{i+} q^+ = 0; \quad D^{i+} _{+j} = 0 \quad (4.69)$$



where  $D^{i+} = D^{ia}u_a^+$ . Thus, the super elds  $q^+$  and  $\bar{q}^{i+}$  live in the analytic harmonic  $N=4; d=1$  superspace  $(t_a; u_a^+)$ . In this setting, the transformations of the hidden  $N=4$  supersymmetry (4.67) are rewritten as

$$q^+ = \frac{1}{2} \bar{q}^{i+} u_a^+; \quad \bar{q}^{i+} = 2i \bar{q}^{i+} q^+ : \quad (4.70)$$

The free action is

$$S = \frac{i}{2} \int d^4x \quad q^+ q^+ \bar{q}^{i+} u_a^+ : \quad (4.71)$$

where  $d^4x = d^4t_a d^4u_a^+$  is the measure of integration over the analytic superspace.

$$2. (4, 8, 4) = (3, 4, 1) + (1, 4, 3)$$

In order to describe this  $N=4$  super eld realization of the  $N=8$  supermultiplet  $(4; 8; 4)$ , we introduce the  $N=8$  super elds  $V^{ab}; V$  as

$$Q^a = \bar{Q}^{ab} V^{ab}; \quad V^{ab} = V^{ba}; \quad (4.72)$$

and use the covariant derivatives (4.41) to rewrite the basic constraints (4.62) as

$$D^{i(a} V^{bc)} = 0; \quad \bar{D}^{i(a} V^{bc)} = 0; \quad (4.73)$$

$$D^{ia} V = \frac{1}{2} \bar{D}^{ib} V^{ab}; \quad \bar{D}^{ia} V = \frac{1}{2} D^{ib} V^{ab} : \quad (4.74)$$

The constraints (4.73) define  $V^{ab}$  as the  $N=8$  super eld comprising the on-shell multiplet  $(3, 8, 5)$ , while, as one can deduce from (4.73), (4.74), the  $N=8$  super eld  $V$  has the content  $(1, 8, 7)$ . Then the constraints (4.74) establish relations between the fermions in these two super elds and reduce the number of independent auxiliary elds to four, so that we once again end up with the irreducible  $N=8$  multiplet  $(4, 8, 4)$ .

Two sets of  $N=4$  covariant derivatives

$$D^a; \bar{D}^a \quad D^{1a}; \bar{D}^{2a} \quad \text{and} \quad r^a; r^a \quad D^{2a}; \bar{D}^{1a}$$

are naturally realized in terms of the  $N=4$  superspaces  $(t; u_a + i\#_{1a}; u_a - i\#_{2a})$  and  $(t; u_a + i\#_{2a}; u_a - i\#_{1a})$ . In terms of the new derivatives the constraints (4.73), (4.74) become

$$D^{(a} V^{bc)} = \bar{D}^{(a} V^{bc)} = r^{(a} V^{bc)} = \bar{r}^{(a} V^{bc)} = 0; \\ D^a V = \frac{1}{2} r^b V^{ab}; \quad \bar{D}^a V = \frac{1}{2} \bar{r}^b V^{ab}; \quad r^a V = \frac{1}{2} D^b V^{ab}; \quad \bar{r}^a V = \frac{1}{2} \bar{D}^b V^{ab} : \quad (4.75)$$

Now we see that the  $r^a; r_a$  derivatives of the super elds  $V; V^{ab}$  are expressed as  $D^a; \bar{D}^a$  of the super elds  $V^{ab}; V$ , respectively. Thus, in the  $(u_a + i\#_{2a}; u_a - i\#_{1a})$  expansions of the super elds  $V; V^{ab}$  only the first components (i.e. those of zero order in the coordinates

( $\epsilon_{2a} + i\epsilon_{2a}; \epsilon_{1a} - i\epsilon_{1a}$ ) will be independent  $N = 4$  super elds. We denote them  $v; v^{ab}$ . The hidden  $N = 4$  supersymmetry is realized on these  $N = 4$  super elds as:

$$v = \frac{1}{2} \epsilon_a D_b v^{ab} + \frac{1}{2} \epsilon_a \bar{D}_b v^{ab}; \quad v^{ab} = \frac{4}{3} \epsilon^{(a} D^{b)} v - \epsilon^{(a} \bar{D}^{b)} v; \quad (4.76)$$

while the super elds themselves obey the constraints

$$D^{(a} v^{bc)} = \bar{D}^{(a} v^{bc)} = 0; \quad D^{(a} \bar{D}^{b)} v = 0; \quad (4.77)$$

which are remnant of the  $N = 8$  super eld constraints (4.75).

The invariant free action reads

$$S = \int d^4x \sqrt{-g} \left( -\frac{1}{2} \partial_\mu v^\mu + \frac{3}{8} v^{ab} \partial_\mu v_{ab} \right); \quad (4.78)$$

### 3. $(4, 8, 4) = (2, 4, 2) + (2, 4, 2)$

This case is a little bit more tricky. First of all we define the new set of  $N = 8$  super elds  $W; \bar{W}$  in terms of  $V^{ij}; \bar{V}^{ij}$  defined earlier in (4.72)

$$W = V^{11}; \bar{W} = \bar{V}^{22}; \quad \frac{2}{3} V + \frac{3}{2} V^{12}; \quad \frac{2}{3} \bar{V} + \frac{3}{2} \bar{V}^{12} \quad (4.79)$$

and construct two new sets of  $N = 4$  derivatives  $D^i; \bar{D}^i$  from those defined in (4.41):

$$D^i = \frac{1}{\sqrt{2}} D^{i1} + \bar{D}^{i1}; \quad \bar{D}^i = \frac{1}{\sqrt{2}} D^{i2} + \bar{D}^{i2}; \\ r^i = \frac{1}{\sqrt{2}} D^{i1} - \bar{D}^{i1}; \quad \bar{r}^i = \frac{1}{\sqrt{2}} D^{i2} - \bar{D}^{i2}; \quad (4.80)$$

The basic constraints (4.73), (4.74) can be rewritten in terms of the super elds  $W; \bar{W}$  and the derivatives  $D^i; \bar{D}^i$  as

$$D^i W = r^i W = 0; \quad \bar{D}^i \bar{W} = \bar{r}^i \bar{W} = 0; \quad D^i = \bar{r}^i = 0; \quad r^i = \bar{D}^i = 0; \\ \bar{D}^i W = D^i \bar{W} = 0; \quad D^i \bar{W} + \bar{D}^i W = 0; \quad \bar{r}^i W = r^i \bar{W} = 0; \quad r^i \bar{W} + \bar{r}^i W = 0; \quad (4.81)$$

The proper  $N = 4$  superspace is defined as the one on which the covariant derivatives  $D^1; \bar{D}^2; r^1; \bar{r}^2$  are naturally realized. The constraints (4.81) imply that the remaining set of covariant derivatives, i.e.  $D^2; \bar{D}^1; r^2; \bar{r}^1$ , when acting on every involved  $N = 8$  super eld, can be expressed as spinor derivatives from the first set acting on some other  $N = 8$  super eld. Thus the first  $N = 4$  super eld components of the  $N = 8$  super elds  $W; \bar{W}$  are the only independent  $N = 4$  super eld projections. The transformations of the implicit  $N = 4$  Poincare supersymmetry have the following form in terms of these  $N = 4$  super elds:

$$w = D^1 + r^1; \quad \bar{w} = \bar{r}^1 W - D^1 \bar{W}; \\ w = \bar{D}_1 + \bar{r}_1; \quad \bar{w} = \bar{D}_1 W - r^1 \bar{W}; \quad (4.82)$$

The free invariant action is

$$S = \int d^4x \, \text{tr} \, \bar{\psi} \gamma^\mu \psi \quad : \quad (4.83)$$

#### 4.6 Supermultiplet (5, 8, 3)

This supermultiplet has been considered in details in Refs. [22, 9]. It was termed there the 'N = 8 vector multiplet'. Here we sketch its main properties.

In order to describe this supermultiplet, one should introduce the bosonic N = 8 superfields  $V_a; U$  obeying the constraints

$$D^{ia} V_a + \gamma^a_{\dot{a}i} r^{\dot{a}} U = 0; \quad r^{\dot{a}} V_a + D^{\dot{a}i} U = 0 : \quad (4.84)$$

It is worth noting that the constraints (4.84) are covariant not only under three SU(2) automorphism groups (realized on the doublet indices  $i, a$  and  $\dot{a}$ ), but also under the SO(5) automorphisms. These SO(5) transformations mix the spinor derivatives  $D^{ia}$  and  $r^{\dot{a}i}$  in the indices  $i$  and  $a$ , while two SU(2) groups realized on these indices constitute SO(4)  $\subset$  SO(5). The superfields  $U; V^a$  form an SO(5) vector: under the SO(5) transformations belonging to the coset SO(5)/SO(4) they transform as

$$V_a = \gamma_a^{\dot{a}i} U; \quad U = \gamma^a_{\dot{a}i} V^{\dot{a}i} : \quad (4.85)$$

As in the previous cases we may consider two different splittings of the N = 8 vector multiplet into irreducible N = 4 superfields

1. (5, 8, 3) = (1, 4, 3) + (4, 4, 0)
2. (5, 8, 3) = (3, 4, 1) + (2, 4, 2)

Once again, they correspond to two different choices of the N = 4; d = 1 superspace as a subspace in the original N = 8; d = 1 superspace.

1. (5, 8, 3) = (1, 4, 3) + (4, 4, 0)

The relevant N = 4 superspace is  $R^{(1|4)}$  parametrized by the coordinates  $(t; \dot{a})$  and defined in (4.3). As in the previous cases, it follows from the constraints (4.84) that the spinor derivatives of all involved superfields with respect to  $\dot{a}$  are expressed in terms of spinor derivatives with respect to  $\dot{a}$ . Thus the only essential N = 4 superfield components of  $V_a$  and  $U$  in their  $\dot{a}$ -expansion are the first ones

$$v_a = V_a|_{\dot{a}=0}; \quad u = U|_{\dot{a}=0} : \quad (4.86)$$

They accommodate the whole on-shell component content of the N = 8 vector multiplet. These the bosonic N = 4 superfields are subjected, in virtue of (4.84), to the irreducibility constraints in N = 4 superspace

$$D^{i(a} v^{b)} = 0; \quad D^{i(a} D^{\dot{b}}_{\dot{a}} u = 0 : \quad (4.87)$$

Thus, from the  $N = 4$  superspace standpoint, the vector  $N = 8$  supermultiplet is the sum of the  $N = 4; d = 1$  hypermultiplet  $v_a$  with the on-shell component contents  $(4, 4, 0)$  and the  $N = 4$  bid' tensor multiplet  $u$  with the contents  $(1, 4, 3)$ .

The transformations of the implicit  $N = 4$  Poincare supersymmetry are

$$v_a = D_a^i u; \quad u = \frac{1}{2} D^{ia} v_a; \quad (4.88)$$

$$2. \quad (5, 8, 3) = (3, 4, 1) + (2, 4, 1)$$

Another interesting  $N = 4$  superfield splitting of the  $N = 8$  vector multiplet can be achieved by passing to the complex parametrization of the  $N = 8$  superspace as

$$t; \quad x_i = x + i\theta^i; \quad \bar{x}^i = \bar{x} - i\theta^i$$

where we have identified the indices  $a$  and  $i$ , thus having reduced the number of manifest  $SU(2)$  automorphism symmetries to just two. In this superspace the covariant derivatives  $D^i, \bar{D}^j$  defined in (4.41) (with the identification of indices just mentioned) are naturally realized. We are also led to define new superfields

$$V = \frac{1}{2} D^i v_i; \quad W = \frac{1}{2} D^i v_i + i D^i u; \quad \bar{W} = \frac{1}{2} D^i v_i - i D^i u; \quad (4.89)$$

In this basis of  $N = 8$  superspace the original constraints (4.84) amount to

$$\begin{aligned} D^i W &= \frac{1}{4} (\bar{D}^i \bar{W} + D^i \bar{W}); \quad \bar{D}^i W = \frac{1}{4} (D^i W + \bar{D}^i W); \\ D^i \bar{W} &= 0; \quad \bar{D}^i W = 0; \quad (D^k D^i) W = (\bar{D}^k \bar{D}^i) \bar{W}; \end{aligned} \quad (4.90)$$

Next, we single out the  $N = 4; d = 1$  superspace as  $t; \quad x_i = x + i\theta^i$ ; and split our  $N = 8$  superfields into the  $N = 4$  ones in the standard way. As in all previous cases, the spinor derivatives of each  $N = 8$  superfield with respect to  $\bar{\theta}^2$  and  $\theta_2$ , as a consequence of the constraints (4.90), are expressed as derivatives of some other superfields with respect to  $\bar{\theta}^2$  and  $\theta_2$ . Therefore, only the rest (i.e. taken at  $\bar{\theta}^{2a} = 0$  and  $\theta_{2a} = 0$ )  $N = 4$  superfield components of the  $N = 8$  superfields really matter. They accommodate the entire on-shell field content of the multiplet. These  $N = 4$  superfields are defined as

$$W_j; \quad \bar{W}^j; \quad w = W; \quad \bar{w} = \bar{W} \quad (4.91)$$

and satisfy the constraints following from (4.90)

$$D^1 w = 0; \quad \bar{D}^1 \bar{w} = 0; \quad D^1 w = \bar{D}^1 \bar{w} = 0; \quad D^1 w = D^1; \quad \bar{D}^1 \bar{w} = \bar{D}^1; \quad (4.92)$$

They tell us that the  $N = 4$  superfields  $w$  and  $\bar{w}$  form the standard  $N = 4$  chiral multiplet  $(2, 4, 2)$ , while the  $N = 4$  superfield  $w$  represents the  $N = 4$  tensor multiplet  $(3, 4, 1)$ .

The implicit  $N = 4$  supersymmetry is realized on  $w$ ,  $\bar{w}$ , and  $\psi$  as

$$w = \frac{1}{2} (\bar{D}^{\dot{a}})^{\alpha} (\bar{D}^{\dot{a}})^{\beta} \psi_{\alpha\beta}; \quad \bar{w} = \frac{4}{3} \bar{D}^{\dot{a}} w; \quad \psi = \frac{4}{3} D^{\dot{a}} w : \quad (4.93)$$

An analysis of  $N = 8$  supersymmetric actions for the  $N = 8$  vector multiplet may be found in [9]. Here we present only two examples of such actions.

The first action

$$S = 2 \int d^4x \frac{1}{\hat{u} + \sqrt{\hat{u}^2 + 2\phi^2}} \quad (4.94)$$

corresponds to the splitting 1 above. It possesses both  $N = 4$  superconformal symmetry and  $N = 8$  Poincare supersymmetry. The closure of these two supersymmetries yields the whole  $N = 8$  conformal supersymmetry associated with the supergroup  $OSp(4^*, 4)$  [9].

The second action is constructed using the splitting 2. It includes the  $N = 4$  tensor  $w$  and the chiral  $\psi$  multiplets. The free action invariant under the implicit  $N = 4$  supersymmetry (4.93) reads

$$S_{\text{free}} = \int d^4x w^2 - \frac{3}{4} \psi^2 : \quad (4.95)$$

The  $N = 4$  superconformally invariant action which is also invariant under (4.93) has the very simple form

$$S_{\text{kin}} = 2 \int d^4x \frac{\log \left( \frac{w^2 + \frac{1}{2}}{w^2} \right)}{w^2} : \quad (4.96)$$

It is also  $OSp(4^*, 4)$  invariant.

Finally, we note that the multiplet  $(5, 8, 3)$  can be regarded as a dimensional reduction of the abelian  $N = 2; d = 4$  gauge multiplet [22] (this is the reason why sometimes it is referred to as  $N = 8$  vector multiplet [9]). Three extra physical scalar fields of the  $d = 1$  case come from the spatial components of the  $d = 4$  gauge vector potential which become gauge invariant after reduction to  $d = 1$ . The description of this multiplet in  $N = 8; d = 1$  harmonic superspace was given in [34, 35].

## 4.7 Supermultiplet $(6, 8, 2)$

This supermultiplet can be described by two  $N = 8$  tensor multiplets  $V^{ij}$  and  $W^{ab}$ ,

$$D_a^{(i} V^{jk)} = 0; \quad r_a^{(i} V^{jk)} = 0; \quad D_i^{(a} W^{bc)} = 0; \quad r_i^{(a} W^{bc)} = 0; \quad (4.97)$$

with the additional constraints

$$D_j^a V^{ij} = r^{bi} W^a_b; \quad r_j^a V^{ij} = D_b^i W^{ab} : \quad (4.98)$$

The role of the latter constraints is to identify the eight fermions which are present in  $V^{ij}$  with the fermions from  $W^{ab}$ , and to reduce the number of independent auxiliary fields in both superfields to two

$$F_1 = D_i^a D_{aj} V^{ij}; \quad F_2 = D_a^i D_{ib} W^{ab}; \quad (4.99)$$

where  $j_m$  means here restriction to  $\epsilon_{ia} = \delta_{ia} = 0$ .

There are two different possibilities to split this  $N = 8$  multiplet into the  $N = 4$  ones

$$1. \quad (6, 8, 2) = (3, 4, 1) \quad (3, 4, 1)$$

$$2. \quad (6, 8, 2) = (4, 4, 0) \quad (2, 4, 2)$$

As before, we discuss peculiarities of both decompositions.

$$1. \quad (6, 8, 2) = (3, 4, 1) \quad (3, 4, 1)$$

The corresponding  $N = 4$  supersubspace is (4.3). The  $N = 8$  constraints imply that the only essential  $N = 4$  super eld components of  $V^{ij}$  and  $W^{ab}$  in their #-expansion are the rst ones

$$v^{ij} = V^{ij}{}_j; \quad w^{ab} = W^{ab}{}_j; \quad (4.100)$$

These six bosonic  $N = 4$  super elds are subjected, in virtue of (4.97), (4.98), to the irreducibility constraints in  $N = 4$  superspace

$$D^a(i V^{jk}) = 0; \quad D^i(a W^{bc}) = 0; \quad (4.101)$$

Thus, the  $N = 8$  supermultiplet  $(6, 8, 2)$  amounts to the sum of two  $N = 4; d = 1$  tensor multiplets  $v^{ij}; w^{ab}$  with the on-shell eld contents  $(3, 4, 1) \quad (3, 4, 1)$ .

The transformations of the implicit  $N = 4$  Poincare supersymmetry are

$$v^{ij} = \frac{2}{3} {}^{(i} D^j) {}_b w^{ab}; \quad w^{ab} = \frac{2}{3} {}^{(a} D^b) {}_i v^{ij}; \quad (4.102)$$

The free  $N = 8$  supersymmetric action has the following form :

$$S = \int dtd^4 \quad v^2 \quad w^2 : \quad (4.103)$$

$$2. \quad (6, 8, 2) = (4, 4, 0) \quad (2, 4, 2)$$

In this case, to describe the  $(6, 8, 2)$  multiplet, we combine two  $N = 4$  super elds: chiral super eld

$$D^i = \overline{D}^i = 0 \quad (4.104)$$

and the hypermultiplet  $q^{ia}$

$$D^{(i} q^{j)a} = \overline{D}^{(i} q^{j)a} = 0; \quad (4.105)$$

The transformations of implicit  $N = 4$  supersymmetry read

$$q^{ja} = {}^a D^i + {}^a \overline{D}^i; \quad = \frac{1}{2} {}^a D^i q_{ia}; \quad = \frac{1}{2} {}^a \overline{D}^i q_{ia}; \quad (4.106)$$

The invariant free action is as follows:

$$S_{free} = \int dtd^4 \quad q^2 \quad 4 : \quad (4.107)$$

## 4.8 Supermultiplet (7, 8, 1)

This supermultiplet has a natural description in terms of two  $N = 8$  superfields  $V^{ij}$  and  $Q^a$  satisfying the constraints

$$D^{(ia}V^{jk)} = 0; \quad r^{(i}V^{jk)} = 0; \quad D^{i(a}Q^{b)} = 0; \quad r^i_{\phantom{i}i} Q^a = 0; \quad (4.108)$$

$$D^a_j V^{ij} = i r^i Q^a; \quad r_j V^{ij} = i D^i_a Q^a : \quad (4.109)$$

The constraints (4.108) leave in the superfields  $V^{ij}$  and  $Q^a$  the sets (3, 8, 5) and (4; 8; 4) of irreducible components, respectively. The role of the constraints (4.109) is to identify the fermions in the superfields  $V^{ij}$  and  $Q^a$  and to reduce the total number of independent auxiliary components in both superfields to just one.

For this supermultiplet there is a unique splitting into  $N = 4$  superfields as

$$(7; 8; 1) = (3; 4; 1) \quad (4; 4; 0):$$

The proper  $N = 4$  superspace is parametrized by the coordinates  $(t; i_a)$ . The constraints (4.108), (4.109) imply that the only essential  $N = 4$  superfield components in the  $\#$ -expansion of  $V^{ij}$  and  $Q^a$  are the first ones

$$v^{ij} = V^{ij}|_{\# = 0}; \quad q^a = Q^a|_{\# = 0} : \quad (4.110)$$

These seven bosonic  $N = 4$  superfields are subjected, as a corollary of (4.108), (4.109), to the irreducibility constraints in  $N = 4$  superspace

$$D^{a(i}v^{jk)} = 0; \quad D^{i(a}q^{b)} = 0 : \quad (4.111)$$

Thus the  $N = 8$  supermultiplet (7, 8, 1) amounts to the sum of the  $N = 4; d = 1$  hypermultiplet  $q^a$  with the (4, 4, 0) on-shell field content and the  $N = 4$  tensor multiplet  $v^{ij}$  with the (3, 4, 1) content.

The implicit  $N = 4$  Poincare supersymmetry is realized by the transformations

$$v^{ij} = \frac{2i}{3} (D^i_j q^a); \quad q^a = \frac{i}{2} D^{ja} v_{ij} : \quad (4.112)$$

The free action can be also easily written

$$S = \int dt d^4x \left( v^2 - \frac{4}{3} q^2 \right) : \quad (4.113)$$

## 4.9 Supermultiplet (8, 8, 0)

This supermultiplet is analogous to the supermultiplet (0, 8, 8): they differ in their overall Grassmann parity. It is described by the two real bosonic  $N = 8$  superfields  $Q^{aA}; \quad i$  subjected to the constraints

$$D^{(ia}Q^{b)A} = 0; \quad D^{i(a}Q^{b)A} = 0; \quad r^{(A}Q^{B)} = 0; \quad r^A Q^{aB} = 0; \quad (4.114)$$

$$r^A Q^a_A = D^{ia} Q^A_i; \quad r^A Q^A_i = D^{ia} Q^A_a : \quad (4.115)$$

Analogously to the case of the supermultiplet  $(8, 8, 0)$ , from the constraints (4.115) it follows that the spinor derivatives of all involved superfields with respect to  $\theta_A$  are expressed in terms of spinor derivatives with respect to  $\theta_{ia}$ . Thus the only essential  $N = 4$  superfield components in the  $\theta$ -expansion of  $Q^{aA}$  and  $\chi^i$  are the first ones

$$Q^{aA} = Q^{aA}|_{\theta=0}; \quad \chi^i = \chi^i|_{\theta=0}: \quad (4.116)$$

They accommodate the whole on-shell component content of the multiplet  $(8, 8, 0)$ . These bosonic  $N = 4$  superfields are subjected, as a consequence of (4.114), (4.115), to the irreducibility constraints in  $N = 4$  superspace

$$D^{a(i}\chi^{j)} = 0; \quad D^{i(a}Q^{b)A} = 0: \quad (4.117)$$

Thus the  $N = 8$  supermultiplet  $(8, 8, 0)$  can be represented as the sum of two  $N = 4; d = 1$  hypermultiplets with the on-shell component contents  $(4, 4, 0) + (4, 4, 0)$ .

The transformations of the implicit  $N = 4$  Poincaré supersymmetry in this last case are as follows:

$$Q^{aA} = \frac{1}{2} \theta^A D^{ia} \chi_i; \quad \chi^i = \frac{1}{2} \theta_A D_a^i Q^{aA}: \quad (4.118)$$

The invariant free action is

$$S = \int d^4x \, d^4\theta \, Q^2 \chi^2: \quad (4.119)$$

The most general action still respecting four  $SU(2)$  automorphism symmetries has the following form:

$$S = \int d^4x \, d^4\theta \, F(Q^2; \chi^2); \quad (4.120)$$

where, as the necessary condition of  $N = 8$  supersymmetry, the function  $F(Q^2; \chi^2)$  should obey the equation

$$\frac{\partial^2}{\partial Q^2 \partial Q^2} Q^2 F(Q^2; \chi^2) + \frac{\partial^2}{\partial \chi^2 \partial \chi^2} \chi^2 F(Q^2; \chi^2) = 0: \quad (4.121)$$

## 5 Summary and conclusions

In this paper, as a further development of findings of our previous paper [9], we presented superfield formulations of the full amount of on-shell  $N = 8; d = 1$  supermultiplets with 8 physical fermions, both in  $N = 8$  and  $N = 4$  superspaces. We listed all possible  $N = 4$  superfield splittings of these multiplets. For each such splitting we gave the  $N = 8$  supersymmetric free  $N = 4$  superfield action, and for some special cases we quoted examples of  $N = 8$  supersymmetric actions with interaction. Only two of these  $d = 1$  multiplets,  $(3, 5, 8)$  and  $(5, 8, 3)$  considered earlier in [9], can be understood in the framework of the dimensional reduction  $N = 2; d = 4 \rightarrow N = 8; d = 1$ . It is far from obvious whether the  $N = 8; d = 1$  on-shell superfield formulations of other multiplets considered here can be



recovered in a similar way.<sup>5</sup> Fortunately, there is no actual need to care about this since our approach is self-consistent in  $d=1$ .

Our results are summarized in the Table below.

$N = 8$  supermultiplets

Multiplet	$N = 8$ Super elds	$N = 8$ Constraints	$N = 4$ splittings
$(0, 8, 8)$	${}^{aA}{}_{\dot{A}}{}^i{}_{\dot{i}}$	$(4.1), (4.2)$	$(0; 4; 4) \quad (0; 4; 4)$
$(1, 8, 7)$	$U$	$(4.11), (4.12)$	$(1; 4; 3) \quad (0; 4; 4)$
$(2, 8, 6)$	$U; \quad$	$(4.26), (4.27)$	$(1; 4; 3) \quad (1; 4; 3)$ $(2; 4; 2) \quad (0; 4; 4)$
$(3, 8, 5)$	$V^{ij}$	$(4.51)$	$(3; 4; 1) \quad (0; 4; 4)$ $(1; 4; 3) \quad (2; 4; 2)$
$(4, 8, 4)$	$Q^a$	$(4.62)$	$(4; 4; 0) \quad (0; 4; 4)$ $(3; 4; 1) \quad (1; 4; 3)$ $(2; 4; 2) \quad (2; 4; 2)$
$(5, 8, 3)$	$U; V^a$	$(4.84)$	$(1; 4; 3) \quad (4; 4; 0)$ $(3; 4; 1) \quad (2; 4; 2)$
$(6, 8, 2)$	$V^{ij}; W^{ab}$	$(4.97), (4.98)$	$(3; 4; 1) \quad (3; 4; 1)$ $(4; 4; 0) \quad (2; 4; 2)$
$(7, 8, 1)$	$V^{ij}; Q^a$	$(4.108), (4.109)$	$(3; 4; 1) \quad (4; 4; 0)$
$(8, 8, 0)$	$Q^{aA}{}_{\dot{A}}{}^i{}_{\dot{i}}$	$(4.114), (4.115)$	$(4; 4; 0) \quad (4; 4; 0)$

These results should be regarded as preparatory for a more detailed study of the  $N = 8$  SQM models associated with the supermultiplets considered. In particular, it would be interesting to reproduce these supermultiplets from nonlinear realizations of  $N = 8; d = 1$  superconformal groups, construct the relevant superconformal actions and reveal a possible relation of the corresponding  $N = 8$  SQM to the physics of branes and black holes. An intriguing question is whether some dynamical models with higher  $N > 8; d = 1$  supersymmetry can be constructed by combining some of the  $N = 8$  multiplets considered in this paper. For deeper geometrical understanding of the  $N = 8$  models proposed above, it would be interesting to consider their Hamiltonian formulation, as it was done in [36] and [18] for  $N = 4$  SQM.

It should be pointed out that in the present work we addressed only those multiplets which satisfy linear constraints in the superspace. As we know, there exist  $N = 4; d = 1$  multiplets with nonlinear defining constraints (e.g. nonlinear versions of the chiral  $(2, 4, 2)$  multiplet [8], as well as of the hypermultiplet  $(4, 4, 0)$  [37]). It would be interesting to construct analogous nonlinear versions of some  $N = 8$  multiplets from the above set. Also, in [31, 8] plenty of nonlinear  $\phi$ -shell relations between the  $N = 4; d = 1$  super elds listed in Section 2 were found, as a generalization of the linear relations described in [29, 30]. These substitutions preserve the number 4 of the fermionic elds, but express some auxiliary elds in terms of the time derivatives of physical bosonic elds. In the

<sup>5</sup>Perhaps one should start from an enlarged set of  $N = 4; d = 4$   $\phi$ -shell super elds, with broken Lorentz covariance.

$N = 8$  case one can expect similar relations, and it is of interest to find the explicit form of them and to explore their possible implications in  $N = 8$  SQM models.

Finally, in this paper we restricted ourselves to  $N = 8; d = 1$  multiplets with a finite number of off-shell components. It still remains to be understood how they are related to multiplets with an infinite number of auxiliary fields, which naturally appear in various versions of harmonic  $N = 8; d = 1$  superspace (see e.g. [34, 38, 35]). Also, the relevance of the latter to the  $N = 8$  SQM model building needs to be explored further.

## Acknowledgements

This work is dedicated to the memory of our friend and colleague Tolya Pashnev.

This research was partially supported by the European Community's Human Potential Programme under contract HPRN-CT-2000-00131 Quantum Spacetime, the INTAS-00-00254 grant, the NATO Collaborative Linkage Grant PST CLG .979389, RFBR-DFG grant No 02-02-04002, grant DFG No 436 RUS 113/669, RFBR grant No 03-02-17440 and a grant of the Heisenberg-Landau programme. E.I. thanks the Institute of Theoretical Physics of the University of Hannover and the Theory Group of the University of Padua for the warm hospitality at the final stage of this work. S.K. thanks INFN - Laboratori Nazionali di Frascati and the Institute of Theoretical Physics of the University of Hannover for the warm hospitality extended to him during the course of this work.

## References

- [1] E. Witten, Nucl. Phys. B 188 (1981) 513.
- [2] R. de Lima Rodrigues, "The quantum mechanics SUSY algebra: an introductory review", hep-th/0205017.
- [3] R. Britto-Pacumio, J. Michelson, A. Strominger, A. Volovich, "Lectures on superconformal quantum mechanics and multi-black hole moduli spaces", hep-th/9911066.
- [4] V. Akulov, A. Pashnev, Teor. Mat. Fiz. 56 (1983) 344.
- [5] S. Fubini, E. Rabinovici, Nucl. Phys. B 245 (1984) 17.
- [6] E. Ivanov, S. Krivonos, V. Leviant, J. Phys. A : Math. Gen. 22 (1989) 4201.
- [7] E. Ivanov, S. Krivonos, O. Lechtenfeld, JHEP 0303 (2003) 014, hep-th/0212303.
- [8] E. Ivanov, S. Krivonos, O. Lechtenfeld, Class. Quantum Grav. 21 (2004) 1031, hep-th/0310299.
- [9] S. Bellucci, E. Ivanov, S. Krivonos, O. Lechtenfeld, Nucl. Phys. B 684 (2004) 321, hep-th/0312322.

- [10] J.M. Maldacena, Adv. Theor. Math. Phys. 2 (1998) 231;  
Int. J. Theor. Phys. 38 (1999) 1113-1133, hep-th/9711200;  
S.S. Gubser, I.R. Klebanov, A.M. Polyakov, Phys. Lett. B 428 (1998) 105,  
hep-th/9802109;  
E. Witten, Adv. Theor. Math. Phys. 2 (1998) 253-291, hep-th/9802150.
- [11] J.A. de Azcarraga, J.M. Izquierdo, J.C. Perez Bueno, P.K. Townsend,  
Phys. Rev. D 59 (1999) 084015, hep-th/9810230.
- [12] P. Claus, M. Derix, R. Kallosh, J. Kumar, P.K. Townsend, A. Van Proeyen,  
Phys. Rev. Lett. 81 (1998) 4553, hep-th/9804177.
- [13] G.W. Gibbons, G. Papadopoulos, K.S. Stelle, Nucl. Phys. B 508 (1997) 623,  
hep-th/9706207.
- [14] G. Papadopoulos, Class. Quantum Grav. 17 (2000) 3715, hep-th/0002007;  
J.M.ichelson, A. Strominger, Commun. Math. Phys. 213 (2000) 1, hep-th/9907191;  
JHEP 9909 (1999) 005, hep-th/9908044.
- [15] A. Maloney, M. Spradlin, A. Strominger, JHEP 0204 (2002) 003, hep-th/9911001.
- [16] G.W. Gibbons, P.K. Townsend, Phys. Lett. B 454 (1999) 187, hep-th/9812034.
- [17] E. Ivanov, S. Kirionos, J. Niederle, Nucl. Phys. B 677 (2004) 485, hep-th/0210196.
- [18] S. Bellucci, A. Galajinsky, E. Ivanov, S. Kirionos, Phys. Lett. B 555 (2003) 99,  
hep-th/0212204.
- [19] F. Calogero, J. Math. Phys. 10 (1969) 2197.
- [20] S. Bellucci, A. Galajinsky, S. Kirionos, Phys. Rev. D 68 (2003) 064010,  
hep-th/0304087.
- [21] M. de Crombrugghe, V. Rittenberg, Ann. Phys. 151 (1983) 99.
- [22] D.-E. Diaconescu, R. Entin, Phys. Rev. D 56 (1997) 8045, hep-th/9706059.
- [23] J. Gomis, P.K. Townsend, M.N.R. Wohlfarth, JHEP 0212 (2002) 027,  
hep-th/0211020.
- [24] D. Bak, K. Lee, P. Yi, Phys. Rev. D 62 (2000) 025009, hep-th/9912083.
- [25] A.V. Smilga, "Low dimensional sisters of Seiberg-Witten effective theory",  
hep-th/0403294.
- [26] E.A. Ivanov, V.I. Ogievetsky, Teor. Mat. Fiz. 25 (1975) 164.
- [27] L. Frappat, P. Sorba, A. Sciarrino, "Dictionary on Lie superalgebras",  
hep-th/9607161.
- [28] A. Van Proeyen, "Tools for supersymmetry", hep-th/9910030.

- [29] S.J. Gates, Jr., L.Rana, Phys. Lett. B 342 (1995) 132, hep-th/9410150.
- [30] A. Pashnev, F. Toppan, J. Math. Phys. 42 (2001) 5257, hep-th/0010135.
- [31] E. Ivanov, O. Lechtenfeld, JHEP 0309 (2003) 073, hep-th/0307111.
- [32] A. Galperin, E. Ivanov, V. Ogievetsky, E. Sokatchev, Pis'ma ZhETF 40 (1984) 155 [JETP Lett. 40 (1984) 912];  
A.S. Galperin, E.A. Ivanov, S. Kalitzin, V.I. Ogievetsky, E.S. Sokatchev, Class. Quantum Grav. 1 (1984) 469.
- [33] A.S. Galperin, E.A. Ivanov, V.I. Ogievetsky, E.S. Sokatchev, Harmonic superspace, Cambridge University Press, Cambridge 2001, 306 p.
- [34] B. Zupnik, Nucl. Phys. B 554 (1999) 365, B 644 (2002) 405E, hep-th/9902038.
- [35] E.A. Ivanov, A.V. Smilga, "Symplectic sigma models in superspace", hep-th/0402041.
- [36] S. Bellucci, A. Nersisyan, Phys. Rev. D 64 (2001) 021702, hep-th/0101065;  
Nucl. Phys. Proc. Suppl. 102 (2001) 227, hep-th/0103005.
- [37] S. Hellerman, J. Polchinski, "Supersymmetric quantum mechanics from light cone quantization", in: M.A. Shifman (ed.), "The many faces of the superworld", hep-th/9908202.
- [38] S. Bellucci, E. Ivanov, S. Krivonos, Phys. Lett. B 460 (1999) 348, hep-th/9811244.