

HU-EP-04/35, D F U P G -2004/98

3D GEORGI-G LASHOW MODEL AND CONFINING STRINGS
AT ZERO AND FINITE TEMPERATURES

DMITRI ANTONOV

Institute of Physics, Humboldt University of Berlin,
Newtonstr. 15, 12489 Berlin, Germany
E-mail: antonov@physik.hu-berlin.de

MARIA CRISTINA DIAMANTINI

Dipartimento di Fisica dell'Università di Perugia and
INFN, sezione di Perugia,
Via A. Pascoli, 06123 Perugia, Italy
E-mail: diamanti@mail.cern.ch

In this review, we discuss confining and finite-temperature properties of the 3D $SU(N)$ Georgi-Glashow model, and of 4D compact QED. At zero temperature, we derive string representations of both theories, thus constructing the $SU(N)$ -version of Polyakov's theory of confining strings. We discuss geometric properties of confining strings, as well as the appearance of the string tension from the field-theoretical one in 4D, and k -string tensions at $N > 2$. In particular, we point out the relevance of negative stiffness for stabilizing confining strings, an effect recently re-discovered in material science. At finite temperature, we present a derivation of the confining-string free energy and show that, at the one-loop level and for a certain class of string models in the large- D limit, it matches that of QCD at large N . This crucial matching is again a consequence of the negative stiffness. In the discussion of the finite-temperature properties of the 3D Georgi-Glashow model, in order to be closer to QCD, we mostly concentrate at the effects produced by some extensions of the model by external matter fields, such as dynamical fundamental quarks or photinos, in the supersymmetric generalization of the model.

Permanent address: ITEP, B. Cherenkovskaya 25, RU-117 218, Moscow, Russia.

2 D. Antonov and M. C. Diamantini

Table of Contents

1	Introduction	3
2	The $SU(N)$ 3D GG model	12
3	String tension of the W Wilson loop in the fundamental representation	16
4	$SU(N)$ confining strings	21
4.1	Fundamental representation	21
4.2	Adjoint representation	27
4.3	k -strings	29
5	Generalization to the $SU(N)$ -analogue of 4D compact QED with the θ -term	36
6	Geometric aspects of confining strings: the physics of negative stiffness	41
7	High temperature behavior of confining strings	45
8	The influence of matter fields to the deconfinement phase transition in the 3D GG model at finite temperature	49
8.1	Introduction	49
8.2	Higgs-inspired corrections to the RG flow in the absence of W bosons	53
8.3	Finite-temperature 3D compact QED with massless fundamental fermions	62
8.4	3D Georgi-Glashow model and its supersymmetric generalization at finite temperature	67
9	Summary	73

Many papers related to this review would have never been written without Ian Kogan's seminal ideas on the finite-temperature phase transitions in string theory and in the 3D Georgi-Glashow model. The sudden death of Ian Kogan was the loss for physics and a personal tragedy for everybody who had known him, in particular for the authors of this manuscript. In the personal sense, Ian was a very bright man with a warm heart, and he will be deeply missed as a good friend. We nevertheless strongly believe that Ian's ideas will survive over decades, inspiring many new generations of physicists.

1. Introduction

During last 30 years, the problem of quark confinement remains as a great challenge not only to QCD theorists, but to the whole theoretical high-energy physics community. In general, confinement can be defined as the absence in the spectrum of a certain field theory of physical finite and finite states of some particles, whose fields are nevertheless present in the Lagrangian of that theory (see e.g. [1,3] for reviews and books on confinement). With regard to QCD, this definition means the absence of asymptotic quark and gluon states, i.e. states which carry color.^a This fact is reflected in the linear growth with the distance of the potential between two color particles, as well as in the logarithmic growth of the strong-coupling constant. The latter makes the standard perturbative diagrammatic techniques inapplicable at the distances larger than $O(\Lambda_{\text{QCD}}^{-1})$. However, such distances are of the primary interest, since only there physically observable colorless states of hadrons are formed, whereas at smaller distances one deals with the unphysical colored states { quarks, gluons, and ghosts. A question, which may naturally be posed at this point, is whether the QCD Lagrangian, operating with these colored states only, yields eventually the correct description of colorless degrees of freedom as well. A reliable indication that this is really the case comes from the simulations of the QCD Lagrangian on a lattice (for recent reviews see e.g. [2,5,6]), which yield a reasonably good description of hadronic spectra. However, as of today, a systematic analytic way of description of large-distance effects in QCD in terms of microscopic (colored) degrees of freedom is unfortunately lacking. The breakdown of the perturbative expansion at large distances under discussion naturally introduces the notion nonperturbative for these effects, as well as for techniques, which

^a An attempt to construct a momentum-space interpretation of this phenomenon has been done (see e.g. Ref. [4]). In this review, we will, however, prefer to deal with the conventional space-time picture, which enables one to directly operate with such notions as the vacuum correlation length or potential between color particles.

4 D. Antonov and M. C. Diamantini

physicists attempt to invent for their description.

It is actually not a great fault that the perturbative expansion is inapplicable at large distances, since, being formulated in terms of colored states, it is anyway not suitable to keep gauge invariance under control. Indeed, an individual diagram is always gauge-dependent, merely because a certain gauge fixing should be performed before one starts to compute diagrams. Contrary to that, hadronic states, being colorless, are gauge-invariant. Moreover, since the QCD vacuum possesses an unbroken gauge symmetry,^b only gauge-invariant amplitudes, being averaged over the vacuum, yield a nonvanishing result.^c Although the gauge dependence disappears when one sums up contributions of individual diagrams to a certain gauge-invariant quantity, it would be more natural to have an expansion operating with such quantities already at any intermediate stage. Such an expansion, which is the expansion in the inverse number of colors, has been proposed by 't Hooft [9] and further developed in the framework of loop equations [10] (see e.g. [11] for reviews). This is a classic example of a nonperturbative approach to QCD.

Another nonperturbative phenomenon in QCD, which is of the same fundamental importance as confinement, and whose complete analytic understanding is also still lacking, is the spontaneous chiral symmetry breaking (SCSB). This is the symmetry of the QCD Lagrangian with N_f massless flavors under the global transformations, which are the $U(N_f) \times U(N_f)$ independent rotations of left- and right-handed quark fields. These are equivalent to the independent vector and axial $U(N_f)$ rotations of the full four-component Dirac spinors, under which the QCD Lagrangian remains invariant as well. At the same time, the axial transformations mix states with different P -parities. Therefore, were the chiral symmetry unbroken, one would observe parity degeneracy of all the states, whose other quantum numbers are the same. The observed splittings between such states

^b We do not discuss in this review the modern line of research, devoted to the so-called color superconductivity [7]. In the color superconducting phase of QCD, which takes place at high baryonic densities, corresponding to the values of the quark chemical potential $\mu \sim 400$ MeV, the gauge symmetry is spontaneously broken.

^c To get a description of large-distance effects in terms of microscopic degrees of freedom, it looks natural to integrate over the high-momentum modes, in the same way as it is usually done in a derivation of renormalization-group (RG) equations in statistical mechanics. For instance, this approach gives a correct RG behavior of the running strong-coupling constant (at the one-loop level) [3]. However, it typically suffers from the above-mentioned problem of violation of the gauge invariance, as far as RG equations for Green functions are concerned. Recent progress in this direction of research is concentrated around the concept of the so-called exact RG flow (see e.g. Ref. [8] for a recent review).

are, however, too large to be explained by the small quark masses. Namely, this splitting is of the order of hundreds MeV, whereas the current masses of light u - and d -quarks are of the order of a few MeV.^d This observation tells us that the chiral symmetry of the QCD Lagrangian is broken down spontaneously. The phenomenon of SC SB naturally leads to the appearance of light pseudoscalar Goldstone bosons, whose role is played by pions, which are indeed the lightest of all the hadrons. The order parameter of SC SB is the chiral quark condensate $\langle \bar{\psi}\psi \rangle \approx (250 \text{ MeV})^3$. Its appearance is quite natural in QCD, where, due to the strong attraction between quarks and antiquarks, it costs very little energy to create a massless quark-antiquark pair. Once having zero total momentum and angular momentum, such pairs should carry net chiral charge, hence is the notion "chiral condensate".

Since the early days of QCD, when it has been realized that its nonperturbative phenomena, confinement and SC SB, are of the fundamental importance, numerous nonperturbative approaches have been proposed in an attempt to describe some of these two phenomena in a controllable way.^e On the side of confinement, these approaches include e.g. the already mentioned expansion in the large number of colors [9], loop equations [10], stochastic vacuum model [1], method of Abelian projections [14].^f On the side of SC SB, the classic approach is the one based on the Nambu {Jona-Lasinio} type models [17], as well as on the related nonlinear chiral meson Lagrangians [18] (the latter approach has further been developed in [19]; for a review see [20]). There exist also microscopic models of SC SB in QCD, based on instantons [21] or dyons [22], which have been put forward in Refs. [23] and [24], respectively.

Although a lot of physical effects have been captured by the above-mentioned methods, none of these methods unfortunately provides the full solution of QCD. The concept of "full solution" has several facets. According to a conventional understanding of a solution of a certain field theory,

^d The current mass of the s -quark, which is around 150 MeV, is still smaller than the typical splitting values at least in a factor of three.

^e Unfortunately, practically no approaches exist, which describe both phenomena. Partially, that is because it is not yet fully accepted that these phenomena are related to each other microscopically, i.e. that the same vacuum configurations are responsible for both phenomena. For instance, as it has been shown in Ref. [12], this is not so in the case of the so-called Faddeev-Niemi effective action [13]. An example of a nonperturbative approach, where the two phenomena are related to each other, is the stochastic vacuum model of QCD [1], which does not refer to a particular microscopic vacuum configuration.

^f A comprehensive lattice analysis of the latter has recently been performed [15] (see Ref. [16] for reviews).

6 D. Antonov and M. C. Diamantini

it means a prescription of how to compute an arbitrary gauge-invariant vacuum amplitude. In QCD, the results should be in an agreement with the experimental and lattice data on the properties of hadrons. The gluonic and quark averages should then give a correct quantitative description of confinement and SC SB, respectively. The values of local averages (condensates) and relations between these quantities should also agree with those known from QCD sum rules [25]. Furthermore, the standard diagrammatic expansion should be reproducible (giving, in particular, the asymptotic freedom), and the nonperturbative dimensional quantities should be made expressible in terms of the only dimensional QCD parameter, Λ_{QCD} . Moreover, some constraints, which are accepted to be rigorous in QCD, should be obeyed. For instance, the above-mentioned expansion in the large number of colors should be correctly reproducible and should respect the large- N loop equation. The obtained solution of QCD should also accommodate classical vacuum configurations of the gluodynamics action, such as instantons.

In this review, we are going to discuss the 3D Georgi-Glashow (GG) model, which is the QCD-related model possessing the property of confinement [26]. Our primary goal will therefore be the study of confinement (and not of the SC SB), as well as of the deconfining phase transition at finite temperatures. The advantage of the 3D GG model with respect to QCD is that confinement in it takes place in the weak-coupling regime. It turns out that, already in this regime, the vacuum (i.e. the ground state) of this model is nonperturbative, being populated by 't Hooft-Polyakov monopoles [27], which provide the permanent confinement of external fundamental charges. As a guiding principle of our analysis we will use the string picture of confinement, therefore let us briefly discuss it.

In QCD, the linearly rising confining interquark potential is associated to a string-like configuration of the gluonic field between quarks, usually called the QCD string. Indeed, the energy of a string grows linearly with its length, $E(R) = \sigma R$.⁹ According to the Regge phenomenology, the string energy density σ , called the string tension, is approximately equal $(440 \text{ MeV})^2$. The string can naturally be called confining (the notion, which is always used in confining gauge theories other than QCD, such as the 3D GG model), since with the increase of the distance R , it stretches and prevents a quark and an antiquark from the separation to macroscopic distances. As any

⁹ In this review, we only briefly discuss the phenomenon of string breaking (see the end of subsection 4.2). The string breaking always happens at a certain distance if dynamical matter fields, transforming by the same representation of the gauge group as the confined external ones, are present.

3D Georgi-Glashow model and confining strings 7

dimensionful quantity in QCD, the string tension should be proportional to the respective power of Λ_{QCD} . Namely,

$$\begin{aligned} \langle \sigma \rangle_{\text{QCD}} &= a^{-2} \exp \left[\int_{g^2(a^{-2})}^{g^2} \frac{dg^2}{g^2 (g^2)} \right], \\ &= a^{-2} \exp \left[\frac{16^{-2}}{\frac{11}{3}N_c - \frac{2}{3}N_f} g^2(a^{-2}) \right]; \end{aligned} \quad (1.1)$$

where $a \neq 0$ stands for the distance UV cutoff (e.g. the lattice spacing). Furthermore, $\langle \sigma \rangle$ means "at the one-loop level", at which

$$\langle \sigma \rangle = \frac{11}{3}N_c - \frac{2}{3}N_f \frac{g^2}{16^{-2}};$$

As it is explicitly seen from Eq. (1.1), all the coefficients in the expansion of $\langle \sigma \rangle$ in (positive) powers of g^2 vanish, which means that the QCD string is indeed an essentially nonperturbative object. The string should therefore be produced by some nonperturbative background fields. On top of these, however, one expects to have some quantum fluctuations of the gauge field, which give rise to the string excitations.

The confining quark-antiquark potential corresponds to the so-called area law of the Wilson loop:^h

$$\langle W(C) \rangle = \frac{1}{N_c} \int_{\mathcal{C}} \text{tr} P \exp \left[ig \int_{\mathcal{C}} A^a T^a dx \right] e^{-\int_{\mathcal{C}} \mathcal{F}^j \cdot \mathcal{F}^j} e^{-\int_{\mathcal{C}} j_{\text{min}}(C) \cdot j};$$

Here, \mathcal{C} is the surface of the minimal area, bounded by the trajectory C of the quark-antiquark pair, and $j:::jm$ means either a length or an area. The confining string, which sweeps out the minimal surface, can naturally be viewed as a product of the above-mentioned strong background fields. Instead, quantum fluctuations around these enable the string to sweep out with a nonvanishing probability any other surface (\mathcal{C}) , different from $\mathcal{C}_{\text{min}}(C)$. To derive the string representation of QCD would then mean to give sense

^h It should be confronted with the perimeter law,

$$\langle W(C) \rangle = \int_{\mathcal{C}} \mathcal{F}^j \cdot \mathcal{F}^j e^{-\text{const} \cdot \int_{\mathcal{C}} \mathcal{F}^j \cdot \mathcal{F}^j}$$

which corresponds to the Coulomb potential and takes place in unconfined theories, e.g. (non-compact) QED.

8 D. Antonov and M. C. Diamantini

to the following formula:

$$\langle W(C) \rangle = \langle W_{\text{min}}(C) \rangle = \sum_{(C)} e^{S[C]} \tag{1.2}$$

Here, $\sum_{(C)}$ and $S[C]$ stand for a certain sum over string world sheets and a string effective action, both of which are yet unknown in QCD.

The first problem one encounters in trying to determine $S[C]$ is that fundamental strings [28] can be quantized only in critical dimensions: a consistent quantum theory describing strings out of critical dimensions has not yet been found. The simplest model, the Nambu-Goto string, can be quantized only in space-time dimension $D = 26$ or $D = 1$ because of the conformal anomaly. It is appropriate to describe an effective string picture for confinement in QCD, but is inappropriate to describe fundamental smooth strings dual to QCD [29], since large Euclidean world-sheets are crumpled. The picture of a fundamental string theory dual to QCD is strongly supported by a recent lattice calculation by Luscher and Weisz [30], where evidence of a string behavior in the static quark-antiquark potential has been found down to distances of 0.5 fm.

In the rigid-string action [31,32], the marginal term proportional to the square of the extrinsic curvature, introduced to avoid crumpling, turns out to be infrared irrelevant and, thus, unable to provide smooth surfaces.

Recent progress in this field is based on a new type of action. In its local formulation [33,34], the string action is induced by an antisymmetric tensor field. This action realizes explicitly the necessary zig-zag invariance of confining strings [34,35]. It can be derived without extra assumptions [36] for the confining phase of compact U(1) gauge theories [3]. An alternative approach to the induced string action was originally proposed in [37], and is based on a d -dimensional, curved space-time string action with the quarks living on a four-dimensional horizon [38]. The formulation of the string theory in the d -dimensional curved space-time is closely related to the AdS/CFT (Anti de Sitter/Conformal Field Theory) correspondence [39]. In fact, with a special choice of the metric in the curved space, one recovers the AdS₅ space, thereby providing a string theory description of a conformal gauge theory [39].

The main characteristic of the effective string action obtained by integrating out the tensor field is a non-local interaction with negative stiffness, that can be expressed as a derivative expansion of the interaction between surface elements. To perform an analytic analysis of the geometric proper-

ties of these strings, this expansion must be truncated: this clearly makes the model non-unitary, but in a spurious way. Moreover, since the stiffness is negative, a stable truncation must, at least, include a sixth-order term in the derivatives. The role of negative stiffness, as first pointed out in [36,40] is crucial. It is in fact the sixth-order term, forced by the negative stiffness, that suppresses the formation of spikes on the surfaces and leads to a smooth surface in the large- D approximation. In fact, in [40,41] it has been shown that, in the large- D approximation, this model has an infrared fixed point at zero stiffness, corresponding to a tensionless smooth string whose world sheet has Hausdorff dimension 2, exactly the desired properties to describe QCD flux tubes. As first noticed in [42,43], the long-range orientational order in this model is due to an antiferromagnetic interaction between normals to the surface, a mechanism confirmed by numerical simulations [44]. The presence of the infrared fixed point does not depend on the truncation [41] and it is present for all ghost- and tachyon-free truncations. Moreover, the effective theory describing the infrared behavior of the coning string is a conformal field theory (CFT) with central charge $c = 1$.

Another important feature of the negative-stiffness model is its high-temperature behavior. Contrary to all previous string models for QCD, it is able to reproduce the large- N QCD behavior, found by Polchinski and Yang in [89], in both sign and reality properties [45].

It is remarkable that the role of negative stiffness, while first discovered in the context of string [36,40] and membranes [47], has been rediscovered, and actually experimentally tested, also in material science [48]. In fact, it has been found that composites with negative stiffness inclusions have higher overall stiffness than that of their constituents. Such composites find applications in which high stiffness and damping are needed, permitting extreme properties not previously anticipated.

Equation (1.2) would clearly be a generalization of a path-integral representation for a propagator of a point-like particle, say boson, $h(x) = \int_{P_{xx^0}} e^{iS[P_{xx^0}]}$, where P_{xx^0} is a path connecting the points x and x^0 . In particular, within this analogy, the role of the classical trajectory of a particle would be played by $\arg \min(C)$. For a point-like particle, the measure in the sum over paths is known and depends only on the dimensionality of the space-time. The world-line action $S[P_{xx^0}]$ can also be evaluated, either analytically [for certain potentials $V(\cdot)$ or external gauge fields, in case if is charged and interacts with any], or using Feynman's variational method. On the string side, a derivation of the measure in the sum over world sheets has been discussed in Ref. [49] for the case of the Abelian Higgs model in

the London limit. As it will be seen below, in a certain case, string effective action derivable in the 3D GG model is the same as that of the London limit of the dual Abelian Higgs model. The summation over string world sheets is, however, realized in these two models in different ways. Namely, in the dual Abelian Higgs model it stems from the integration over singularities of the dual Higgs field (vortex cores), which is already present in the original partition function. Instead, the 3D GG model does not contain any dual Abelian Higgs field, and the summation over world sheets in this model is realized by means of the resulting string effective action itself [34]. Numerous investigations of this action have further been performed (see e.g. Refs. [36, 42, 43]), and some of these will be discussed in this review. A separate section will be devoted to the finite-temperature properties of conning strings [45].

For the purpose of study of the so-called k -strings and merely to be closer to the real QCD, we will deal with the $SU(N)$ -generalization of the standard $SU(2)$ GG model, which will be introduced in the next section. As for the $(N = 2)$ 3D GG model, it is a classic example [26] of a theory, which allows for an analytic description of confinement. As it has already been mentioned, confinement in this model is due to the plasma of point-like magnetic monopoles, which produce random magnetic fluxes through the contour of the Wilson loop. In the weak-coupling regime of the model, this plasma is dilute, and the interaction between monopoles is Coulombic, being induced by the dual-photon exchanges. Since the energy of a single monopole is a quadratic function of its flux, it is energetically favorable for the vacuum to support a configuration of two monopoles of unit charge (in the units of the magnetic coupling constant, g_m), rather than a single, doubly-charged monopole. Owing to this fact, monopoles of unit charge dominate in the vacuum, whereas monopoles of higher charges tend to dissociate to them. Summing over the grand canonical ensemble of monopoles of unit charge, interacting with each other by the Coulomb law, one arrives at an effective low-energy theory, which is a 3D sine-Gordon theory of a dual photon. The latter acquires a mass (visible upon the expansion of the cosine potential) by means of the Debye screening in the Coulomb plasma. The appearance of this (exponentially small) mass and, hence, of a finite (albeit exponentially large) vacuum correlation length is crucial for the generation of the fundamental string tension, i.e. for the confinement of an external fundamental matter. It is worth noting that a physically important interpretation of these ideas in terms of spontaneous breaking of magnetic Z_2 symmetry has been presented in reviews [50] and Refs. therein.

While conning properties of the 3D GG model are well known since the

pioneering Polyakov's paper [26], the finite-temperature properties of this model have been addressed only recently, starting with the papers [51, 52]. It turned out [52] that charged matter fields of W -bosons play the crucial role for the dynamics of the phase transition. Below we will review this issue and also discuss the influence of other matter fields to the finite-temperature properties of the model. Such fields are either already present in the original Lagrangian (e.g. Higgs [53], or dual photons in the supersymmetric generalization of the model [54]), or can be included in the framework of a certain extension of the model (e.g. massless fundamental quarks [55] or a heavy fundamental bosonic matter [56]).

The outline of the review is the following. In the next section, we will introduce the 3D GG model in the general $SU(N)$ case. In section 3, we will find the string tension of the fundamental Wilson loop defined at a flat contour (henceforth called for shortness "flat Wilson loop"). In section 4, we will develop a theory of conining strings based on the Kalb-Ramond field, which enables one to deal with non-flat Wilson loops. Using for concreteness the case of fundamental representation, we will present in subsection 4.1 two methods, by means of which the theory of conining strings can be derived. In subsection 4.2, the case of the adjoint Wilson loop will be considered, and the respective theory of conining strings will be constructed in the large- N limit. In subsection 4.3, we will study the spectrum of k -strings, i.e. strings between sources in (higher) representations with a nonvanishing N -ality. These sources carry a charge k with respect to the center of the gauge group Z_N and can be seen as a superposition of k fundamental charges. Clearly, the spectrum of such strings is an important ingredient for the complete description of the conining dynamics of the 3D GG model. In section 5, the obtained $SU(N)$ -theory of conining strings will be generalized to the $SU(N)$ -version of 4D compact QED (in the continuum limit) with the field-theoretical term. As it has been found in Refs. [33, 43], for the usual compact QED, this term leads to the appearance of the string term. The latter, being proportional to the number of self-intersections of the world sheet, might help in the solution of the problem of crumpling of large world sheets [3, 31]. Critical values of β , at which this happens, will be derived in the general $SU(N)$ -case for fundamental and adjoint representations, as well as for k -strings. In section 6, various geometric features of conining strings will be studied. In section 7, thermodynamics of conining strings will be discussed, and a derivation of the one-loop free energy of a string in the large- D limit will be presented. Again, we will show that it is the presence of negative stiffness that allows one to reproduce the large- N behavior of

12 D. Antonov and M. C. Diamantini

high-temperature QCD.

In section 8, we will pass from the thermodynamics of confining strings to the thermodynamics of the 3D GG model itself. After an introduction into this subject in subsection 8.1, we will pay a particular attention to the influence of various matter fields to the dynamics of the deconfining phase transition. In subsections 8.2 and 8.3, we will consider an approximation where W-bosons are disregarded. Subsection 8.2 will be devoted to the influence of the Higgs field (when it is not infinitely heavy) to the RG flow, while in subsection 8.3 we will consider the model in the presence of external dynamical fundamental quarks. In subsection 8.4, we will first discuss the crucial role of W-bosons to the dynamics of the phase transition in the finite-temperature 3D GG model and then consider the supersymmetric generalization of the model. In the Summary section 9, the main points discussed in the review will be emphasized once again.

2. The SU(N) 3D GG model

The Euclidean action of the 3-d Georgi-Glashow model reads [26]

$$S = \int d^3x \left[\frac{1}{4g^2} F^a{}^2 + \frac{1}{2} (D^a)^2 + \frac{1}{4} (A^a)^2 \right]; \quad (2.1)$$

where the Higgs field ϕ^a transforms by the adjoint representation, i.e. $D^a \phi^a = \partial^a \phi^a + g^{abc} A^b \phi^c$. The weak-coupling regime $g^2 \ll m_W$, which will be assumed henceforth, parallels the requirement that m_W should be large enough to ensure the spontaneous symmetry breaking from SU(2) to U(1). At the perturbative level, the spectrum of the model in the Higgs phase consists of a massless photon, two heavy, charged W_{\pm} -bosons with mass $m_W = g v$, and a neutral Higgs field with mass $m_H = \sqrt{2} g v$.

What is, however, more important is the nonperturbative content of the model, represented by the famous 't Hooft-Polyakov monopole [27]. It is a solution to the classical equations of motion, which has the following Higgs- and vector-field parts:

$$\begin{aligned} \phi^a &= v \frac{x^a}{r}, \quad u(0) = 0, \quad u(r) \xrightarrow{r \rightarrow \infty} \exp(i \mathbf{m} \cdot \mathbf{r}) = (gr); \\ A^i &\xrightarrow{r \rightarrow \infty} \frac{1}{r^2} \epsilon^{ijk} x^j x^k, \\ H &= \frac{1}{r^3} \left[4 - (\mathbf{x} \cdot \mathbf{x}_2) (\mathbf{x}_3) \right]; \end{aligned}$$

as well as the following action $\mathcal{S} = \frac{4\pi}{g^2} \int d^3x$. Here, $g^2 = m_W^2$ is the weak-coupling parameter, $\mathbf{m} = (m_H = m_W)$ is a certain monotonic, slowly varying function, $\mathbf{m} \cdot \mathbf{m} = 1$, $\mathbf{m}(0) = 1$ (BPS-limit) [57],

(1) '1:787 [58].

The following approximate saddle-point solution (which becomes exact in the BPS-limit) has been found in Ref. [26]:

$$S = N S_0 + \frac{g_m^2}{8} \sum_{\substack{a,b=1 \\ a \neq b}}^N \frac{q_a q_b}{r_{ab}} e^{-m_H r_{ab}} + O\left(\frac{1}{m_W R}\right);$$

where $m_W^{-1} R \sum_{a,b=1}^N q_a q_b g_m = 4$, $[g_m] = [\text{mass}]^{-1/2}$. Therefore, while at $m_H \neq 1$, the usual compact QED action is recovered, in the BPS-limit one has

$$S' = N S_0 + \frac{g_m^2}{8} \sum_{\substack{a,b=1 \\ a \neq b}}^N \frac{q_a q_b}{r_{ab}};$$

i.e. the interaction of two monopoles doubles for opposite and vanishes for equal charges. Therefore, in this limit, the standard monopole-antimonopole Coulomb plasma recombines itself into two mutually noninteracting subsystems, consisting of monopoles and antimonopoles. The interaction between objects inside each of these subsystems has the double strength with respect to the interaction in the initial plasma.

When $m_H < 1$, the summation over the grand canonical ensemble of monopoles has been performed in Ref. [59] and reads

$$Z_{\text{mon}} = 1 + \sum_{N=1}^{\infty} \frac{1}{N!} \int \prod_{a=1}^N d^3 z_a \prod_{a=1}^N e^{-S_a} = \int \mathcal{D}D \exp \int d^3 x \left[\frac{1}{2} (\partial_\mu \phi)^2 + \frac{1}{2} (\partial_\mu \chi)^2 + \frac{m_H^2}{2} \phi^2 - 2 g_m \cos(g_m \chi) \right]; \tag{2.2}$$

Here, ϕ is the dual-photon field and χ is the field additional with respect to compact QED, which describes the Higgs boson. Furthermore, the monopole fugacity (i.e. the statistical weight of a single monopole), \mathcal{Z} , has the following form [26]:

$$\mathcal{Z} = \frac{m_W^{7/2}}{g} e^{-S_0}; \tag{2.3}$$

The function $\mathcal{Z} = \mathcal{Z}(m_H = m_W)$ is determined by the loop corrections. It is known [60] that this function grows in the vicinity of the origin (i.e. in the BPS limit). However, the speed of this growth is such that it does not spoil the exponential smallness of \mathcal{Z} in the weak-coupling regime under study.

Our next aim will be to construct the $SU(N)$ -generalization of the partition function (2.2). Let us first introduce the $(N-1)$ -dimensional vector H of the mutually commuting diagonal generators of the group $SU(N)$. Together with certain pairwise linear combinations of the off-diagonal generators which, in analogy with the $SU(2)$ -group, are called step (rising and lowering) generators $E_{\pm i}, i = 1; \dots; \frac{N(N-1)}{2}$, the diagonal generators form the following algebra:

$$[H, E_{\pm i}] = \alpha_{\pm i} E_{\pm i}; \quad [E_i, E_{-i}] = \alpha_i H;$$

Vectors α_i 's here are called root vectors of the group $SU(N)$. The vector potential, $A^a, a = 1; \dots; N^2 - 1$, can be respectively decomposed into photons and W -bosons as follows:

$$A^a = \sum_i \left(W^{+i} E_{-i} + W^{-i} E_i + A^i H \right);$$

where from now on $\sum_i = \sum_{i=1}^{\frac{N(N-1)}{2}}$. Next, embeddings of the $SU(2)$ -monopole into the maximal Abelian subgroup $U(1)^{N-1}$ of the group $SU(N)$ can be characterized by the space-time variations of the Higgs field ϕ^a . Outside the monopole core, ϕ^a can be chosen along the z -axis: $\phi^a(0;0;z) = \tilde{\nu}_a H$. Here, $\tilde{\nu}_a$ is the $(N-1)$ -dimensional vector of v.e.v.'s of the Higgs field ϕ^a . For an arbitrary space-time direction,

$$\phi^a(r; r') = (r; r')^a \phi^a(0;0;z) = \tilde{\nu}_a H(r; r');$$

The matrix $\tilde{\nu}_a$ can be chosen such that

$$\tilde{\nu}_a(r; r') = Y^a + X_i^a(r) T_i \frac{r-r'}{r}; \tag{2.4}$$

Here, the 3-component matrix-valued vector

$$T_i = \frac{E_{i+} + E_{i-}}{2}; \frac{E_i - E_{-i}}{2}; \alpha_i H$$

characterizes the embedding of the $SU(2)$ Lie algebra into the root space of the group $SU(N)$, associated with the root α_i . There exist therefore $\frac{1}{2}N(N-1)$ embeddings, corresponding to the same number of monopoles. The constant $Y^a = (\tilde{\nu}_a \cdot (\tilde{\nu}_a \alpha_i) \alpha_i) H$, which parametrically depends on i , $Y^a = Y_{(i)}^a$, is the hypercharge associated with the i -th embedding. Since the vector $\tilde{\nu}_a \cdot (\tilde{\nu}_a \alpha_i) \alpha_i$ belongs to the plane containing α_i and $\tilde{\nu}_a$ (and is perpendicular to α_i), the first term on the r.h.s. of Eq. (2.4) breaks $SU(N)$

down to $SU(2) \times U(1)^{N-2}$, whereas the second term, with $X_i^a \sim \frac{1}{r} \sim \frac{1}{r} \sim \frac{1}{r}$, breaks further $SU(2)$ down to $U(1)$. Again, like in the $SU(2)$ -case, the only part of the monopole vector potential, which does not vanish exponentially at large distances, is the diagonal (photonic) one. It reads

$$A^i = \frac{1}{g} \nabla^i \frac{x}{r};$$

that corresponds to the magnetic field

$$H^i = g_m \frac{x}{4r^3} \epsilon_{ij} (\delta_{ij} - \frac{x_i x_j}{r^2}) H^j(\mathbf{x}');$$

where again $g_m = 4\pi/g$. Therefore, in the $SU(N)$ -case, monopoles also interact by means of the long-ranged Coulomb forces. This intermonopole interaction, mediated by dual photons, results into the following $SU(N)$ -analogue of the partition function (2.2):

$$Z_{mon}^N = \int \prod_{i=1}^N D\tilde{\phi}_i \exp \left[-\int d^3x \left(\frac{1}{2} (\partial_\mu \tilde{\phi}_i)^2 + \frac{1}{2} (\partial_\mu \tilde{\phi}_i)^2 + \frac{m_H^2}{2} \tilde{\phi}_i^2 - 2g_m \sum_{i=1}^N \cos(g_m \tilde{\phi}_i) \right) \right]; \quad (2.5)$$

Here, the dual-photon field is described by the $(N-1)$ -dimensional vector $\tilde{\phi}$.

Averaging in Eq. (2.5) over the Higgs field by means of the cumulant expansion one gets in the second order of this expansion [64]:

$$Z_{mon}^N \approx \int \prod_{i=1}^N D\tilde{\phi}_i \exp \left[-\int d^3x \left(\frac{1}{2} (\partial_\mu \tilde{\phi}_i)^2 + \frac{m_H^2}{2} \tilde{\phi}_i^2 - 2g_m \sum_{i=1}^N \cos(g_m \tilde{\phi}_i) \right) \right] + 2 \int d^3x d^3y \sum_{i,j} \cos(g_m \tilde{\phi}_i(\mathbf{x})) K(\mathbf{x}-\mathbf{y}) \cos(g_m \tilde{\phi}_j(\mathbf{y})) \quad (2.6)$$

In this equation, $\tilde{D} = \exp \left[-\frac{m_H^2}{2} \int d^3x \tilde{\phi}_i^2 \right]$ is the modified fugacity (which can be shown to remain exponentially small as long as the cumulant expansion is convergent) and $K(\mathbf{x}) = \int \frac{d^3p}{(2\pi)^3} \frac{e^{i\mathbf{p}\cdot\mathbf{x}}}{p^2 + m_H^2} = 1$. The Debye mass of the dual photon, stemming from Eq. (2.6) by virtue of the formula $\langle \tilde{\phi}_i \tilde{\phi}_i \rangle = \frac{N}{2}$, $i = 1, \dots, N-1$, reads

$$m_D = g_m \sqrt{\frac{N-1}{N}} \left[1 + \frac{N(N-1)}{2} \right]; \quad (2.7)$$

16 D. Antonov and M. C. Diamantini

where $I = \int_{\mathbb{R}^3} d^3x K(x)$. At $m_H = m_W$, the following expression for I has been obtained [64]:

$$I \sim \frac{4}{m_H m_W^2} \exp\left[-\frac{4}{m_H = m_W}\right] \quad (2.8)$$

The parameter of the cumulant expansion is $O(N^2)$. By virtue of Eqs. (2.3) and (2.8), one can readily see that the condition for this parameter to be (exponentially) small reads $N < \exp\left[-\frac{1}{e}\right]$. Approximating by its value at infinity, we find $N < e^{8.9}$. Therefore, when one takes into account the propagation of the heavy Higgs boson, the necessary condition for the convergence of the cumulant expansion is that the number of colors may grow not arbitrarily fast, but should rather be bounded from above by some parameter, which is nevertheless exponentially large.ⁱ A similar analysis can be performed in the BPS limit, $m_H = g^2$. There, one readily finds $I \sim (g = m_H)^2$, and

$$I N^2 / N^2 \exp\left[-\frac{1}{2}\right] \quad :$$

Approximating by its value at the origin, we see that the upper bound for N in this limit is smaller than in the vicinity of the compact-QED limit and reads $N < e^{\frac{1}{2}}$.

3. String tension of the W Wilson loop in the fundamental representation

At zero temperature, 3D GG model is a clear analogy of the 2D XY model in its continuum limit (see e.g. Refs. [26, 62]). The analogy is due to the fact that vortices of the 2D XY model correspond to monopoles of the 3D GG model, whereas spin waves correspond to free (non-dual) photons. Disorder in both theories is produced by topological defects, i.e. by vortices or monopoles.^j Instead, spin waves or free photons cannot disorder correlation functions at large distances and affect them only at the distances smaller than the vacuum correlation length. As a result, monopoles lead to the area law of the W Wilson loop, whereas free photons lead to its perimeter law.

ⁱAs for the convergence of the cumulant expansion itself, it is a natural requirement, which should be obeyed by any field theory with a normal stochastic, rather than the coherent, vacuum.

^jAt finite temperatures, disorder is primarily generated by W -bosons [52], which, at zero temperature under discussion, are practically irrelevant due to their heaviness. The quantitative discussion of the role of W -bosons at finite temperature will be presented in subsection 8.4.

Thus, since confinement and string tension, we are interested in, are generated by monopoles, in this section photons will be omitted. The partition function of the grand canonical ensemble of monopoles in the Euclidean space-time reads

$$Z_{m\ on}^N = \sum_{N=0}^{\infty} \frac{g_m^N}{N!} \exp \left[\frac{g_m^2}{2} \int d^3x d^3y \tilde{y}^N(\mathbf{x}) D_0(\mathbf{x}, \mathbf{y}) \tilde{y}^N(\mathbf{y}) \right] : (3.1)$$

Here, $D_0(\mathbf{x}) = 1/(4\pi|\mathbf{x}|)$ is the 3D Coulomb propagator, and the monopole density is defined as $\tilde{y}^N(\mathbf{x}) = \sum_{k=1}^N \mathbf{e}_{i_k}(\mathbf{x} - \mathbf{r}_k)$ at $N \geq 1$ and $\tilde{y}^0(\mathbf{x}) = 0$. Furthermore, the average is defined as follows:

$$\langle \tilde{y}_{i_1} \dots \tilde{y}_{i_N} \rangle = \frac{1}{Z} \int d^3x_1 \dots d^3x_N \tilde{y}_{i_1}(\mathbf{x}_1) \dots \tilde{y}_{i_N}(\mathbf{x}_N) : (3.2)$$

Upon the explicit summation, the partition function (3.1) can be represented in the sine-Gordon type form, which is the large- m_H limit of the partition function (2.5):

$$Z_{m\ on}^N = \int D\tilde{\phi} \exp \left[\int d^3x \left[\frac{1}{2} (\partial_\mu \tilde{\phi})^2 - \sum_i \cos(g_m \tilde{\phi}_i) \right] \right] : (3.3)$$

The Debye mass (2.7) becomes reduced to [64]: $m_D = g_m \sqrt{\frac{p}{N}}$.

The following comment is worth to be made at this point. The description of the grand canonical ensemble of monopoles in terms of the dual-photon field essentially implies the validity of the mean-field approximation. This approximation, which enables one to disregard fluctuations of fields of individual monopoles, is only valid if the number of monopoles contained in the Debye volume, m_D^3 , is large. Up to exponentially small corrections, the mean value of the monopole density, evaluated according to the formula $\rho_{mean} = \frac{1}{V^{(3)}} \frac{\partial \ln Z_{m\ on}}{\partial \ln}$, reads $\rho_{mean} \sim N(N-1)$, where $V^{(3)}$ is the three-volume occupied by the system. Therefore, the number of monopoles contained in the Debye volume is

$$\rho_{mean} m_D^3 \sim \frac{N-1}{g_m^3} :$$

This is indeed an exponentially large quantity, even at $N \geq 1$. With the increase of N , the accuracy of the mean-field approximation is being further enhanced.

18 D. Antonov and M. C. Diamantini

Next, one can introduce the monopole field-strength tensor F^N violating the Bianchi identity as $\frac{1}{2} \epsilon_{\mu\nu\lambda} \partial_\lambda F^N = g_m \tilde{N}$. Since it was argued to omit photons throughout this section, we obtain for F^N the following expression:

$$F^N(\mathbf{x}) = g_m \epsilon_{\mu\nu\lambda} \partial_\lambda \int d^3y D_0(\mathbf{x}-\mathbf{y}) \tilde{N}(\mathbf{y}) \quad (3.4)$$

Then, by virtue of the Stokes' theorem and the formula

$$\text{tr} \exp(i\mathcal{O}H) = \prod_{a=1}^N \exp(i\mathcal{O} \cdot \tilde{a}) ;$$

where \mathcal{O} is an arbitrary $(N-1)$ -component vector and \tilde{a} 's are the weight vectors of the group $SU(N)$, $a = 1; \dots; N$, we obtain the following expression for the Wilson loop defined at the N -monopole configuration:

$$W(C)_{\text{mon}}^N = \frac{1}{N} \text{tr} \exp \left(\frac{ig}{2} \int_C d^3x F^N \right) = \frac{1}{N} \prod_{a=1}^N W_a^N \quad (3.5)$$

Here,

$$W_a^N = \exp \left(\frac{ig}{2} \tilde{a} \int_C d^3x F^N \right) \quad (3.6)$$

and $\mathbf{x}^{(C)} = \int_C d^2x(\mathbf{x}) \otimes \mathbf{x}(\mathbf{x})$ is the vorticity tensor current defined

at a certain surface (C) bounded by the contour C and parametrized by the vector $\mathbf{x}(\mathbf{r})$ with $\mathbf{r} = (r_0; r_1)$ standing for the 2D coordinate. A straightforward calculation with the use of Eq. (3.4) yields

$$W_a^N = \exp \left(i\tilde{a} \int_C d^3x \tilde{N} \right) ; \quad (3.7)$$

where $\mathbf{x}^{(C)} = \int_C d^2x(\mathbf{x}) \otimes \frac{1}{|\mathbf{x}(\mathbf{x})|} \mathbf{x}(\mathbf{x})$ is the solid angle under which the

surface (C) is seen by an observer located at the point \mathbf{x} , $d\Omega = \frac{1}{2} d^2x$. (Note that owing to the Gauss' law, $\int_C d^3x \tilde{N} = 4\pi$ for a closed surface, i.e. when C is shrunk to a point.) The ratio of two W_a^N 's defined at different surfaces,

Ω_1 and Ω_2 , bounded by the contour C , reads

$$\begin{aligned} \frac{W_a^N(\Omega_1)}{W_a^N(\Omega_2)} &= \exp \left(i\tilde{a} \int_{\Omega_1} d^3x \tilde{N}(\mathbf{x}) \otimes \frac{1}{|\mathbf{x}(\mathbf{x})|} \right) = \\ &= \prod_{k=1}^N \exp \left(i\tilde{a} \otimes \mathbf{e}_k \int_{\Omega_1} d^3x \tilde{N}(\mathbf{x}) \otimes \frac{1}{|\mathbf{x}(\mathbf{x})|} \right) ; \quad (3.8) \end{aligned}$$

Due to the Gauss' law, the last integral in this equation is equal either to 4 or to 0, depending on whether the point \mathbf{x} lies inside or outside the volume bounded by the surface S_1 [2]. Since the product $\sim_a \mathbf{e}_{ik}$ is equal either to $\frac{1}{2}$ or to 0, we conclude that $\frac{W_a^N(1)}{W_a^N(2)} = 1$. This fact proves the independence of W_a^N of the choice of the surface in the definition (3.6). Clearly, this is the consequence of the quantization condition $gg_m = 4$ used in the derivation of Eq. (3.7).

A representation of the partition function (3.1), alternative to Eq. (3.3) and more appropriate for the investigation of the Wilson loop, is the one in terms of the dynamical monopole densities [64]. It can be obtained by multiplying Eq. (3.1) by the following unity:

$$\int_{\mathbb{R}^3} \delta(\mathbf{x} \sim \mathbf{y}) \exp(i g_m \int_{\mathbf{x}}^{\mathbf{y}} \mathbf{A} \cdot d\mathbf{x}) = \int_{\mathbb{R}^3} \delta(\mathbf{x} \sim \mathbf{y}) \exp(i g_m \int_{\mathbf{x}}^{\mathbf{y}} \mathbf{A} \cdot d\mathbf{x}) ;$$

so that the field \mathbf{A} plays the role of the Lagrange multiplier. We obtain for the partition function:

$$Z_{mon}^N = \int_{\mathbb{R}^3} \delta(\mathbf{x} \sim \mathbf{y}) \exp\left(\frac{g_m^2}{2} \int_{\mathbb{R}^3} d^3x d^3y \sim(\mathbf{x}) D_0(\mathbf{x} \sim \mathbf{y}) \sim(\mathbf{y}) + i g_m \int_{\mathbb{R}^3} d^3x \sim\right) \\ \sum_{N=0}^{\infty} \frac{1}{N!} \exp(i g_m \int_{\mathbb{R}^3} d^3x \sim^N) ; \quad (3.9)$$

where the last sum is equal to

$$\exp\left(2 \int_{\mathbb{R}^3} d^3x \sum_i \cos(g_m \mathbf{e}_{i1} \sim)\right) ; \quad (3.10)$$

Accordingly, Eq. (3.7) becomes replaced by

$$W_a^N = \frac{1}{Z_{mon}^N} \int_{\mathbb{R}^3} \delta(\mathbf{x} \sim \mathbf{y}) \exp\left(\frac{g_m^2}{2} \int_{\mathbb{R}^3} d^3x d^3y \sim(\mathbf{x}) D_0(\mathbf{x} \sim \mathbf{y}) \sim(\mathbf{y}) + \right. \\ \left. + \int_{\mathbb{R}^3} d^3x i g_m \sim + 2 \int_{\mathbb{R}^3} \cos(g_m \mathbf{e}_{i1} \sim) + i \mathbf{r}_a \sim\right) ;$$

and the full expression for the monopole contribution to the Wilson loop [instead of Eq. (3.5)] reads $W(C)_{mon} = \frac{1}{N} \sum_{a=1}^N W_a$.

Let us next introduce the magnetic field according to the formulae $\mathbf{e} \cdot \mathbf{B} = \sim, \mathbf{e} \cdot \mathbf{B} = 0$. This yields

20 D. Antonov and M. C. Diamantini

$$W_a = \frac{1}{Z_{mon}^N} \int_{\mathcal{C}} \mathcal{D}\tilde{B} \exp \left(\int_{\mathcal{C}} d^3x \left(\frac{g_m^2}{2} \tilde{B}^2 + \right. \right. \\ \left. \left. + i g_m \tilde{\omega} \tilde{B} + 2 \sum_i \cos(g_m \varphi_i \tilde{\omega}) + 4 i \tilde{\omega}_a \right) \right); \quad (3.11)$$

(C)

where, as it has been discussed, the surface (C) is arbitrary. The magnetic and the dual-photon fields can be integrated out of Eq. (3.11) by solving the respective saddle-point equations. From now on in this section, we will consider the contour C located in the (x;y)-plane. This naturally leads to the following Ansatz for the fields to be inserted into the saddle-point equations: $\tilde{B} = \tilde{B}(z)$, $\tilde{\omega} = \tilde{\omega}(z)$. For the points (x;y) lying inside the contour C, these equations then read:

$$i g_m \tilde{\omega}^0 + g_m^2 \tilde{B} - 4 i \tilde{\omega}_a(z) = 0; \quad (3.12)$$

$$i \tilde{B}^0 - 2 \sum_i \varphi_i \sin(g_m \varphi_i \tilde{\omega}) = 0; \quad (3.13)$$

where $dz = dx + iy dy$. Note that differentiating Eq. (3.12) and substituting the result into Eq. (3.13), we obtain the equation

$$\tilde{\omega}^0 - 2 \sum_i \varphi_i \sin(g_m \varphi_i \tilde{\omega}) = g \tilde{\omega}_a^0(z);$$

which is the SU(N)-generalization of the respective saddle-point equation obtained in Ref. [26]. These equations can be solved by using one more natural Ansatz for the saddle points of the fields, $\tilde{B}(z) = \tilde{\omega}_a(z)$, $\tilde{\omega}(z) = \tilde{\omega}_a(z)$, that, due to the formula

$$\tilde{\omega}_a \tilde{\omega}_b = \frac{1}{2} \delta_{ab} - \frac{1}{N} \delta_{ab}; \quad (3.14)$$

it makes W_a a-independent. Let us next multiply Eq. (3.13) by $\tilde{\omega}_a$, taking into account that for any a, (N-1) positive roots yield the scalar product with $\tilde{\omega}_a$ equal to 1/2, while the others are orthogonal to $\tilde{\omega}_a$. Equations (3.12), (3.13) then go over to

$$2i \tilde{\omega}^0 + g_m^2 \tilde{B} = 4 i \tilde{\omega}_a(z); \quad \tilde{B}^0 + 2i N \sin \tilde{\omega} = 0; \quad (3.15)$$

where $g_m = 2$. One can readily check that the solution to this system of equations has the following form:

$$B(z) = i \frac{8m_D}{g_m^2} \frac{e^{m_D |z|}}{1 + e^{2m_D |z|}}; \quad (z) = 4 \operatorname{sgn} z \arctan e^{m_D |z|} : \quad (3.16)$$

Taking the value $B(0)$ for the evaluation of the Wilson loop, we obtain for the string tension:

$$= 4 \frac{N}{2N} \frac{14m_D}{g_m^2} = \frac{8}{g_m} \frac{N}{N} \frac{1}{N} : \quad (3.17)$$

This result demonstrates explicitly the nonanalytic dependence of on g , but does not yield the correct overall numerical factor. That is because $B(z)$, necessary for this calculation, is defined ambiguously. The ambiguity originates from the exponentially large thickness of the string: $|z| \sim d m_D^{-1}$. It is therefore unclear which value of $B(z)$ we should take: either $B(0)$, as it was done, or the one averaged over some range of z . It turns out that, in the weak-field (low-density) limit, the problem can be solved for an arbitrarily-shaped surface, by using the representation in terms of the Kalb-Ramond field, which will be described in the next section.

4. SU(N) con ning strings

4.1. Fundamental representation

The dualphoton can alternatively be described in terms of the Kalb-Ramond field. One of the ways to introduce this field is to consider it as the field-strength tensor corresponding to the field B^0 , namely $B^0 = \frac{1}{2g_m} \epsilon^{ijk} \partial_j \partial_k B^0$. The Wilson loop (3.11) then takes the form^k

$$W_a = \frac{1}{Z_{mon}^N} \int \mathcal{D}\tilde{h} \int \mathcal{D}\tilde{B}^0 \exp \left(\int d^3x \left[\frac{1}{4} \tilde{h}^2 + \frac{i}{2} \tilde{h} \epsilon^{ijk} \partial_j \partial_k \tilde{B}^0 + 2 \sum_i \cos(g_m \tilde{h} \cdot \tilde{e}_i) + \frac{ig}{2} \tilde{h} \cdot \tilde{a} \right] \right) : \quad (4.1)$$

The field \tilde{h} can further be integrated out by solving the saddle-point equation of the form (3.13), where B^0 is replaced by $\frac{1}{2g_m} \epsilon^{ijk} \partial_j \partial_k \tilde{B}^0$ (and \tilde{h} depends

^k Note the following correspondence between the Coulomb interaction of monopole densities and the actions of the magnetic- and Kalb-Ramond fields:

$$\frac{1}{4} \int d^3x \tilde{h}^2 = \frac{g_m^2}{2} \int d^3x B^2 = \frac{g_m^2}{2} \int d^3x d^3y \tilde{h}(\mathbf{x}) D_0(\mathbf{x} - \mathbf{y}) \tilde{h}(\mathbf{y}) :$$

such an extremization is equivalent to the definition of the extremal surface through the equation $H[\mathbf{x}(\sigma)] = 0$. Due to Eqs. (4.5), (4.7), H indeed vanishes on both sides of the stationary surface. The reason for that is the following distinguished property of the extremal surface: $\mathbf{j} \cdot \mathbf{j}$ is smaller than $\langle \mathbf{j} \cdot \mathbf{j} \rangle$ on one side and larger than $\langle \mathbf{j} \cdot \mathbf{j} \rangle$ on the other side. Therefore, according to Eq. (4.5), on both sides of the extremal surface, $\nabla_{\perp} h = 0$. This is equivalent to the equation $\nabla_{\perp} H = 0$ and hence (upon the multiplication of both sides by $\nabla_{\perp} \cdot \nabla_{\perp}$) to $H = 0$. [In particular, for a flat contour the extremal surface is the flat surface inside this contour. This can explicitly be seen from the solutions (3.16) of Eqs. (3.15). Namely, at any z such that $|\mathbf{j} \cdot \mathbf{j}| \ll \langle \mathbf{j} \cdot \mathbf{j} \rangle$, $B^0(z) \approx \pm \text{sgn} z \sqrt{2\langle \mathbf{j} \cdot \mathbf{j} \rangle} e^{-|z|/\lambda}$, where both equalities hold with the exponential accuracy. The equation $B^0(z) = 0$ is equivalent to the condition of vanishing of H on both sides of the flat surface.]

Next, the general form of an antisymmetric rank-2 tensor field \mathbf{h} is

$$\mathbf{h} = \alpha \mathbf{A} + \beta \mathbf{A} + \gamma \mathbf{C} \quad (4.8)$$

The constraint $\alpha \mathbf{h} = 0$, imposed in Eq. (4.1), is equivalent to setting \mathbf{A} equal zero. From now on, we will promote \mathbf{h} to include also the fields of free photons, \mathbf{A} , by abolishing this constraint. Let us further go into the weak-field limit, $|\mathbf{h}| \ll 1$. Using the Cauchy inequality, we have

$$|\mathbf{h}| \leq \frac{|\mathbf{j} \cdot \mathbf{j}|}{(N-1)} = \frac{|\mathbf{j} \cdot \mathbf{j}|}{2N(N-1)}; \quad (4.9)$$

so that the weak-field limit is equivalent to the following low-density approximation: $|\mathbf{j} \cdot \mathbf{j}| \ll \frac{1}{2N(N-1)}$. To understand in which sense this inequality implies the low-density approximation, recall the mean monopole density we had from Eq. (3.3):

$$\langle \mathbf{j} \cdot \mathbf{j} \rangle_{\text{mean}} \approx N(N-1); \quad (4.10)$$

Therefore, at large- N , the low-density approximation implies that $|\mathbf{j} \cdot \mathbf{j}|$ should be of the order N^{-2} times smaller than its mean value.

In the weak-field limit, we now have the following expression for the total Wilson loop, where the constraint $\alpha \mathbf{h} = 0$ is removed:

$$\exp \int d^3x \left[\frac{1}{12m_D^2} \mathbf{H}^2 + \frac{1}{4} \mathbf{h}^2 + \frac{ig}{2} \mathbf{A} \cdot \mathbf{h} \right]; \quad (4.11)$$

24 D. Antonov and M. C. Diamanti

Here, Z^{tot} is given by the same integral over $\tilde{\mathfrak{h}}_{\text{eld}}$, but with \mathfrak{h} set to zero, and $\tilde{H} = @ \tilde{\mathfrak{h}} + @ \tilde{\mathfrak{h}} + @ \tilde{\mathfrak{h}}$ is the Kalb-Ramond eld-strength tensor. [The apparent \mathfrak{h} -dependence of the r.h.s. of Eq. (4.11) is due to the fact that in course of the $\tilde{\mathfrak{h}}$ -expansion of $V(\tilde{\mathfrak{h}})$, only one branch of this potential has been taken into account. As it was discussed above, the \mathfrak{h} -dependence disappears upon the summation over all the branches.] Owing to Eq. (4.8), Eq. (4.11) is obviously factorized as $W(C;)_{\text{weak}}^{\text{tot}; a}_{\text{eld}} = W_a W_a^{\text{phot}}$, where the free photonic contribution to the Wilson loop reads

$$W_a^{\text{phot}} = \exp \left[- \frac{g^2 N}{4N} \int_C dx^I dx^0 D_0(x) \right] \quad (4.12)$$

Doing the integration over $\tilde{\mathfrak{h}}$ in Eq. (4.11), we obtain

$$W(C;)_{\text{weak}}^{\text{tot}; a}_{\text{eld}} = \exp \left[- \frac{g^2 N}{4N} \int_C dx^I dx^0 D_{m_D}(x) \right] + \frac{m_D^2}{2} \int_C d(x^I) \int_C d(x^0) D_{m_D}(x) \quad (4.13)$$

where $D_{m_D}(x) = e^{m_D x^i} (4 - x^j)$ is the 3D Yukawa propagator. Note that the free photonic contribution is completely canceled out of this expression, i.e. it is only the dualphoton (of the mass m_D) which mediates the $C-C$ - and \mathfrak{h} -interactions.

One can further expand the nonlocal string effective action

$$S_{\text{str}} = (g m_D)^2 \frac{N}{8N} \int_C d(x^I) \int_C d(x^0) D_{m_D}(x) \quad (4.14)$$

in the powers of derivatives with respect to the world-sheet coordinates σ^a 's. Note that the actual parameter of this expansion is $1 = (m_D R)^2$, where R

$\text{Area}(\Sigma)$ is the size of Σ (see the discussion below). The resulting quasi-local action reads (cf. Refs. [36, 65]):

$$S_{\text{str}} = \int_C d^2 \sigma \sqrt{-g} + \frac{1}{m_D^2} \int_C d^2 \sigma \sqrt{-g} g^{ab} (\partial_a t) (\partial_b t) + \int_C d^2 \sigma \sqrt{-g} R + O \left(\frac{1}{m_D^4 R^2} \right) \quad (4.15)$$

Here, $\partial_a = @ = @^a$, and the following quantities characterize Σ : $g_{ab}(\sigma) = (@_a x^i(\sigma)) (@_b x^j(\sigma))$ is the induced metric tensor, $g = \det k g^{ab} k$,

$t_{ab} = \nabla_a \nabla_b x^c - \nabla_b \nabla_a x^c = R^c{}_{ab} x^a$ is the extrinsic-curvature tensor, $R = \nabla^a \nabla_a \ln \sqrt{g} = \nabla^a \bar{g}^a$ is the expression for the scalar (or extrinsic) curvature in the conformal gauge $g_{ab} = \bar{g}_{ab}$.

The third term on the r.h.s. of Eq. (4.15) is known to be a full derivative, and therefore it does not actually contribute to the string effective action, while the second term describes the so-called rigidity (or stiffness) of the string [31,32]. The reason for the notation μ^2 , introduced in Ref. [31], is that, as it has been shown in that paper, it is μ^2 , which is asymptotically free. This asymptotic freedom then indicates that the rigidity term can only be infrared relevant, provided the respective β -function has a zero in the infrared region. However, such a zero has not been found. Due to this fact, the rigidity term is not a good candidate to solve the old-standing problem of crumpling of large world sheets in the Euclidean space-time. This necessitates to seek other possible solutions of this problem. One of such solutions, based on the string β -term, has been proposed already in Ref. [31]. A possible derivation of such a term, within the $SU(N)$ -analogue of the 4D compact QED will be presented below.

The string coupling constants in Eq. (4.15) read

$$g_m^2 = 2 \frac{N}{\mu^2} \frac{1}{g_m^2} ; \tag{4.16}$$

$$g_3^2 = g_2^2 = \frac{2(N-1)}{4g_m^3 N^{3-2}} : \tag{4.17}$$

A comment is in order regarding the negative sign of μ^2 . Up to a total derivative, the rigidity term reads

$$\int d^2x \sqrt{g} g^{ab} (\partial_a x^c) (\partial_b x^c) = \int d^2x \sqrt{g} (\Delta x^c)^2 ;$$

where the Laplacian, associated to the metric g^{ab} , acts onto x^c as $\Delta x^c = \nabla_a \nabla^a x^c = \nabla_a \bar{g}^{ab} \partial_b x^c$. In the conformal gauge, one thus readily gets for the rigidity term $\int d^2x \sqrt{g} (\Delta x^c)^2$. For a nearly flat surface, $g \approx c^2$, and we have for the string propagator (in the 2D Euclidean space):

$$\langle x^c(x) x^c(0) \rangle = \int d^2p \frac{e^{ipx}}{p^2 + \frac{1}{c^2} p^2} ;$$

For $c < 0$, this propagator has a tachyonic pole, whereas an unphysical mass pole shows up otherwise. We therefore conclude that the negative sign of μ^2 is important for the stability of strings.

In the weak-field (or low-density) approximation, we can thus fix the proportionality factor 2^{-2} at \mathcal{L} , that is close to the factor 8 of Eq. (3.17) which, however, could have not been fixed within the method of section 3. Another important fact is that (in the weak-field approximation) the string tension and higher string coupling constants are the same for all large enough surfaces (C) . [The words "large enough" here mean the validity of the inequality $R \gg d$, i.e. the string length R should significantly exceed the exponentially large string thickness d . Indeed, the general n -th term of the derivative (or curvature) expansion of S_{str} is of the order of $m_D^2 R^4 (R=d)^{-2n}$.] It is, however, worth noting that some of the terms in the expansion (4.15) vanish at the surface of the minimal area corresponding to a given contour C (e.g. in the case of a flat surface considered in section 3). This concerns, for instance, the rigidity term ¹.

Let us finally demonstrate how to introduce the Kalb-Ramond field in a way alternative to its definition via the \tilde{B} -field and incorporating automatically the free-photonic contribution to the Wilson loop. It is based on the direct combination of Eqs. (3.1) and (3.7), that yields

$$\begin{aligned}
 W_a = & \sum_{N=0}^{\infty} \frac{1}{N!} \exp \left[-\frac{g_m^2}{2} \int d^3x d^3y \tilde{y}^N(x) D_0(x-y) \tilde{y}^N(y) \right] \\
 & + i \int_a \tilde{d}^3x \tilde{y}^N = \frac{1}{Z_{\text{mon}}^N} D_{\tilde{\kappa}} \\
 & \left(\int d^3x \left[\frac{1}{2} \tilde{\kappa} \tilde{a} \tilde{\kappa} + \frac{1}{2} \tilde{\kappa} \tilde{a} \tilde{\kappa} \right] \cos(g_m \tilde{\kappa} \tilde{a} \tilde{\kappa}) \right); \quad (4.18)
 \end{aligned}$$

where $\tilde{\kappa} = \tilde{y} + \frac{\tilde{a}}{g_m}$. One can further use the following formula, which shows explicitly how the Kalb-Ramond field unifies the monopole and the free-photonic contributions to the Wilson loop:

$$\begin{aligned}
 & \exp \left[\frac{1}{2} \int d^3x \tilde{\kappa} \tilde{a} \tilde{\kappa} + \frac{1}{4N} \int d^3x \tilde{\kappa} \tilde{a} \tilde{\kappa} \right] = \\
 & = \int d^3x \exp \left[\frac{1}{4} \tilde{\kappa}^2 + \frac{i}{2} \tilde{\kappa} \tilde{a} \tilde{\kappa} \right] \int d^3x \tilde{y}^N : \quad (4.19)
 \end{aligned}$$

¹Indeed, one can prove the equality $g^{ab}(\partial_a t)(\partial_b t) = (D^a D_a x)(D^b D_b x)$, where $D_a D_b x = \partial_a \partial_b x - \Gamma_{ab}^c \partial_c x$, and Γ_{ab}^c is the Christoffel symbol corresponding to the metric g^{ab} . On the other hand, the surface of the minimal area is defined by the equation $D^a D_a x(\cdot) = 0$ together with the respective boundary condition at the contour C .

In fact, one can prove that both sides of this formula are equal to^m

$$\exp \int_Z d^3x (\partial_\mu \tilde{\kappa} + g \tilde{\kappa} \tilde{a}_\mu)^2 :$$

Inserting further Eq. (4.19) into Eq. (4.18), we obtain

$$\begin{aligned} W_a \equiv W(C)_a^{\text{tot}} &= \frac{1}{Z^{\text{tot}}} \int_Z D\tilde{\kappa} D\tilde{a}_\mu \exp \int d^3x \left[\frac{1}{4} \tilde{\kappa}^2 + \right. \\ &+ \left. \frac{i}{2} \tilde{\kappa} \partial_\mu \tilde{a}_\mu + \sum_i \cos(g_m \alpha_i \tilde{\kappa}) + \frac{ig}{2} \tilde{a}_\mu \tilde{\kappa} \right] : \end{aligned} \quad (4.20)$$

Denoting $\tilde{\kappa} \equiv \tilde{\kappa}$, we arrive back to Eq. (4.1) with $Z_{\text{mon}}^N \equiv Z^{\text{tot}}$ and the constraint $\partial_\mu \tilde{\kappa} = 0$ abolished, as it should be.

4.2. Adjoint representation

Let us now extend the ideas of the previous subsection to the case of the Wilson loop in the adjoint representation. The charges of quarks in this representation are distributed along the roots, so that in the formulae for the Wilson loop, Eqs. (3.6) and (3.7), \tilde{a}_μ should be replaced by α_i . Noting that any root is a difference of two weights, we will henceforth in this subsection label roots by two indices running from 1 to N , e.g. $\alpha_{ab} = \tilde{a}_\mu - \tilde{a}_\mu^n$. Equation (3.14) then leads to the following formula:

$$\alpha_{ab} \alpha_{cd} = \frac{1}{2} (a_c + b_d - a_d - b_c); \quad (4.21)$$

according to which the product $\alpha_{ab} \alpha_{cd}$ may take the values $0; \frac{1}{2}; 1$. Owing to this fact, the ratio (3.8) is again equal to 1, i.e. for adjoint quarks, whose charge obeys the quantization condition $gg_m = 4$, the Wilson loop is as surface independent, as it is for fundamental quarks.

The adjoint-case version of Eq. (4.1) with the constraint $\partial_\mu \tilde{\kappa} = 0$ abolished [or of Eq. (4.20)] has the form:

$$\begin{aligned} W(C)_{ab}^{\text{tot}} &= \frac{1}{Z^{\text{tot}}} \int_Z D\tilde{\kappa} D\tilde{a}_\mu \exp \int d^3x \left[\frac{1}{4} \tilde{\kappa}^2 + \right. \\ &+ \left. \frac{i}{2} \tilde{\kappa} \partial_\mu \tilde{a}_\mu + \sum_{c;d=1}^N \cos(g_m \alpha_{cd} \tilde{\kappa}) + \frac{ig}{2} \alpha_{ab} \tilde{a}_\mu \tilde{\kappa} \right] : \end{aligned} \quad (4.22)$$

^m Obviously, in the noncompact case, when monopoles are disregarded and $\tilde{\kappa} = \partial_\mu \tilde{A} - \partial_\mu \tilde{A}$, the r.h.s. of Eq. (4.19) yields the free-photonic contribution to the Wilson loop, Eq. (4.12).

ⁿ In particular, this makes explicit the number of positive roots, $\frac{N^2 - N}{2}$.

28 D. Antonov and M. C. Diamantini

The saddle-point equation, emerging from the above formula in the course of integration over $\tilde{\nu}$,

$$\frac{i}{2} \tilde{\nu}'' @ \tilde{\nu} = g_m \sum_{c,d=1}^{X^N} \epsilon_{cd} \sin(g_m \epsilon_{cd} \tilde{\nu}); \tag{4.23}$$

is to be solved in a way similar to the one of the previous subsection, namely by using the Ansatz $\tilde{\nu} = \epsilon_{ab} h$, $\tilde{\nu} = \epsilon_{ab}$. Multiplying both sides of Eq. (4.23) by ϵ_{ab} , we obtain:

$$\frac{i}{2g_m} \tilde{\nu}'' @ h = \sum_{c,d=1}^{X^N} \epsilon_{ab} \epsilon_{cd} \sin(g_m \epsilon_{ab} \epsilon_{cd} h); \tag{4.24}$$

Let us now compute the sum on the r.h.s. of this equation by virtue of Eq. (4.21). To this end, imagine ourselves the antisymmetric $N \times N$ -matrix of roots ϵ_{cd} 's. Obviously, for a fixed root ϵ_{ab} , there is one root ϵ_{cd} (equal to ϵ_{ab}) and a negative symmetric to it, whose scalar products with ϵ_{ab} are equal to 1 and -1. Our aim is to calculate the number n of roots, whose scalar product with ϵ_{ab} is equal to $\frac{1}{2}$.^o According to Eq. (4.21), these roots belong either to the rows $c = a, c = b$, or to the columns $d = a, d = b$, which in total contain $4N - 4$ elements. Among these, two are diagonal and other two are those, which yield the scalar product -1. The remaining $n = 4(N - 2)$ roots are precisely those, which yield the scalar product $\frac{1}{2}$, so that this product equals $\frac{1}{2}$ for $2(N - 2)$ of them and $\frac{1}{2}$ for the other $2(N - 2)$. The sum on the r.h.s. of Eq. (4.24) thus takes the form

$$\begin{aligned} \sin(2h) &= \sin(h) + 2(N - 2) \frac{1}{2} \sin(h) - \frac{1}{2} \sin(h) = \\ &= 2[\sin(h) + (N - 2) \sin(h)]; \end{aligned}$$

where the field h has been defined after Eq. (3.15). Let us now consider the limit $N \rightarrow \infty$, in which Eq. (4.24) takes the form $\tilde{\nu}'' = iH_{ab}$, where [cf. Eq. (4.3)]

$$H_{ab} = \frac{1}{4N g_m} \epsilon_{ab} \tilde{\nu}'' @ \tilde{\nu};$$

The analogue of the potential (4.2) in the same limit then reads

$$V[\tilde{\nu}] = 4N \int d^3x \left[\frac{Z}{\alpha^3} H_{ab} \operatorname{arcsinh} H_{ab} + \frac{q}{1 + H_{ab}^2} \right];$$

^o For all other $N^2 - N - 2 - n$ roots, the scalar product vanishes, and so does the r.h.s. of Eq. (4.24).

The string representation of the adjoint Wilson loop in the large- N limit is therefore given by Eq. (4.22) with the substitution

$$Z = \int \prod_{ij} d^3x_{ij} \frac{1}{2} \left(\frac{1}{\Lambda} \right)^{2N} \int \prod_{cd=1}^3 \cos(g_m \mathcal{Q}_{cd} \sim) \int \prod_{i=1}^h V_{ij}$$

(and the symbol $\Lambda \sim$ removed).

We are naturally interested in the adjoint-case counterparts of Eqs. (4.16), (4.17), one can obtain in the weak-field limit $\mathcal{H}_{abj} \sim 1$.^P In this limit, the formula (4.11) recovers itself, with the substitution $\tilde{a} \sim \mathcal{Q}_{ab}$. As a consequence, the ratios of adjoint-case values of string couplings g_{adj}^2 , g_{adj}^2 , and \mathcal{H}_{adj} to the respective fundamental-case values, Eqs. (4.16) and (4.17), are equal to $\frac{2N}{N-1} \sim 2$. In particular, for the string tensions this ratio coincides with the known leading large- N QCD result (see e.g. Ref. [66]). However, unlike the fundamental string, the adjoint one is unstable at large distances due to the production of W^+W^- pairs. The string-breaking distance, R_c , can be estimated from the balance of the string free energy, $g_{adj} R_c$, and the mass of the produced pair, $2m_W$ [50]. By virtue of Eq. (4.16), one obtains $R_c / d \sim 1$. Therefore, the breaking length of the adjoint string is in the factor ~ 1 larger than its thickness. However, since the very existence of the string implies that its length is much larger than its thickness [cf. the discussion in the paragraph preceding Eq. (4.18)], the question of (non-)existence of the adjoint string becomes purely numerical.

4.3. k -strings

The large- N ideas discussed in the Introduction have recently found a novel realization in the studies of spectrum of k -strings in $SU(N)$ gauge theories. A k -string is defined as the conining flux tube between sources in higher representations, carrying a charge k with respect to the center of the gauge group Z_N , i.e. representations with nonvanishing N -ality. These sources can be seen as the superposition of k fundamental charges, and charge conjugation exchanges k - and $(N-k)$ -strings, so that non-trivial k -strings exist only for $N > 3$;^Q their string tensions σ_k can be – and should be – used to

^P Note that, using the formula $\mathcal{H}_{ab} = \frac{1}{2N} \mathcal{Q}_{ab} \sim$ and the Cauchy inequality, one gets the following analogue of Eq. (4.9): $\mathcal{H}_{abj} \sim \frac{1}{2N} \mathcal{Q}_{ab} \sim$. This clearly leads to the same definition of the weak-field limit in terms of the low-density approximation, as in the fundamental case. Namely, the weak-field limit corresponds to densities j which are of the order N times smaller than the mean one (4.10).

^Q We will not consider here high-dimensional representations that are screened by gluons and do not yield a genuine asymptotic string tension.

30 D. Antonov and M. C. Diamantini

constraint mechanisms of confinement [69, 70]. Results for the values of σ_k can be obtained by various approaches. Early results suggest the so-called "Casimir scaling" hypothesis for the ratio of string tensions [71]:

$$R(k; N) = \frac{\sigma_k}{\sigma_1} = \frac{k(N-k)}{N-1} C(k; N) \quad (4.25)$$

where σ_1 is the fundamental string tension. These are based on the arguments of the Parisi-Sourlas dimensional reduction of the 4D QCD to the 2D one, which is due to stochastic vacuum fields, and the further use of the fact that in 2D confinement is produced by the one-gluon exchange. Recent studies in supersymmetric Yang-Mills theories and M-theory suggest instead a "Sine scaling" formula:

$$R(k; N) = \frac{\sin(k\pi/N)}{\sin(\pi/N)} \quad (4.26)$$

Corrections are expected to both formulae, but the form of such corrections is unknown for the physically relevant case of a four dimensional, nonsupersymmetric, $SU(N)$ gauge theory.

In the large- N limit, where the interactions between flux tubes are suppressed by powers of $1/N$, the lowest-energy state of the system should be made of k fundamental flux tubes connecting the sources, hence:

$$R(k; N) \sim \frac{k!}{N!} \quad (4.27)$$

Both the Casimir and the Sine scaling formulae satisfy this constraint; they also remain invariant under the replacement $k \rightarrow (N-k)$, which corresponds to the exchange of quarks with antiquarks. However, it has been argued in Refs. [72, 73] that the correction to the large- N behavior should occur as a power series in $1/N^2$ rather than $1/N$. Clearly, such a behavior would exclude Casimir scaling as an exact description of the k -string spectrum.

Recent lattice calculations have provided new results for the spectrum of k -strings both in three and four dimensions [74-79]. They all confirm that Casimir scaling is a good approximation to the Yang-Mills results. To be more quantitative, one could say that all lattice results are within 10% of the Casimir scaling prediction, and that deviations from it are larger in four than they are in three dimensions, in agreement with strong-coupling predictions [78]. The taming of systematic errors is a crucial matter for such lattice calculations, and it can only be achieved by an intensive numerical

²One should study with some care whether the arguments presented in Refs. [72, 73] hold independently of the space-time dimensionality.

analysis. In four dimensions, the higher statistics simulations presented in Ref. [78] show that corrections to the Casimir scaling formula are statistically significant, and actually favor the Sine scaling. Finally, it has been pointed out in Ref. [79] that higher-dimensional representations with common N -ality do yield the same string tension, as expected because of gluon screening.

These numerical results trigger a few comments on Casimir scaling. The original argument [71] was based on the idea that a 4D gauge theory in a random magnetic field could be described by a 2D theory without such a field. Besides the numerical results, there is little support for such an argument in QCD; moreover it is not clear that the same hypothesis could explain the approximate Casimir scaling observed in three dimensions. On the other hand, Casimir scaling appears "naturally" as the lowest-order result, both at strong-coupling in the case of k -strings in the hamiltonian formulation of gauge theories, and in the case of the spectrum of bound states in chiral models. Corrections can be computed in the strong-coupling formulation and they turn out to be $\sim 1/(D-2)N$ - see e.g. Ref. [78] for a summary of results and references. While strong-coupling calculations are not directly relevant to describe the physics of the continuum theory, it is nonetheless instructive to have some quantitative analytic control within that framework. Last but not least, Casimir scaling also appears at the lowest order in the stochastic model of the QCD vacuum [80]. In view of these considerations, it is fair to say that approximate Casimir scaling should be a prerequisite for any model of confinement, that corrections should be expected, and that these corrections are liable to yield further information about the non-perturbative dynamics of strong interactions. Moreover, it would be very interesting to improve our understanding of some other aspects of the k -string spectrum, like e.g. the origin of the Sine scaling for non-supersymmetric theories, or the structure of the corrections to this scaling form.

Below in this subsection, we will prove that the Casimir scaling with a high accuracy takes place in the low-density approximation of the $SU(N)$ 3D GG model [81]. The k -string tension is defined by means of the k -th power of the fundamental Wilson loop. The surface-dependent part of the latter can be written, in terms of the dual-photon field, as follows:

$$W_k(C)_{\text{mon}} = \int_{a_1, \dots, a_k=1}^{N^*} \exp \left[ig \sum_{i=1}^k \int_{\text{mon}} d^3x \, \mathbf{x} \cdot \mathbf{A}(\mathbf{x}) \right] : \quad (4.28)$$

In this equation, $\int_{\text{mon}} d^3x \, \mathbf{x} \cdot \mathbf{A}(\mathbf{x}) \equiv \int_{(C)} \mathbf{x} \cdot \mathbf{A}(\mathbf{x})$, where again (C) is an arbitrary surface bounded by the contour C and parametrized by the vector $\mathbf{x}(\sigma)$.

32 D. Antonov and M. C. Diamantini

The independence of Eq. (4.28) of the choice of (C) can readily be seen in the same way as for the $(k = 1)$ -case. The ρ -dependence rather appears in the weak-field, or low-density, approximation, which is equivalent to keeping only the quadratic term in the expansion of the cosine in Eq. (3.3). As it has been demonstrated in subsection 4.1 [cf. the discussion around Eqs. (4.9), (4.10)], the notion "low-density" implies that the typical monopole density is related to the mean one, $\rho_{\text{typical}} = \rho_{\text{mean}} = \frac{1}{N(N-1)}$, by the following sequence of inequalities:

$$\rho_{\text{typical}} \leq \rho_{\text{mean}} = \frac{1}{N(N-1)} \leq \frac{1}{N^2} \quad (4.29)$$

Below [namely, in the paragraph following after Eq. (4.36)] we will discuss in some more details the correspondence between the low-density approximation and the large- N one.

Denoting for brevity a $\int_{(C)} d^3x \, \chi(\mathbf{x}) \sim \chi(\mathbf{x})$, we can rewrite Eq. (4.28) as

$$\langle \chi^k(C) \rangle_{\text{mon}} = \int_{\substack{a_1, \dots, a_k=1 \\ \text{with possible repetitions}}} d^3x \, e^{i\mathbf{a} \cdot (\sim_{a_1} + \dots + \sim_{a_k})} \quad (4.30)$$

where in the low-density approximation the average is defined with respect to the action

$$Z = \int d^3x \, \frac{1}{2} (\partial \cdot \sim)^2 + \frac{m_D^2}{2} \sim^2 \quad (4.31)$$

Similarly to the fundamental representation, in this approximation, the string tension for a given \mathbf{p} is the same for all surfaces (C) , which are large enough in the sense $S \gg \lambda$, where S is the area of (C) . In particular, the fundamental string tension, found above, reads $\sigma_1 = \frac{1}{2N} \frac{1}{g_m^2}$, where $\frac{1}{g_m^2} = \frac{1}{4} \frac{1}{N}$, and the factor $\frac{1}{2N}$ is the square of a weight vector.

To evaluate Eq. (4.30) for $k > 1$, we should calculate the expressions of the form

$$\int d^3x \, \chi^{(n)}(\mathbf{x}) \sim_{a_1} \dots \sim_{a_n} \quad (4.32)$$

where $(k = n)$ weight vectors \sim_{a_j} 's are mutually different and also different from the vector \sim_{a_1} . By virtue of the formula $\sim_a \cdot \sim_b = \frac{1}{2} \delta_{ab} \frac{1}{N}$, we obtain for Eq. (4.32):

$$\frac{1}{2N} \frac{1}{n^2 + k} \int d^3x \, \chi^{(n)}(\mathbf{x}) \sim_{a_1} \dots \sim_{a_n} = \frac{1}{2N} \frac{1}{2n(k-n) + 2} \int d^3x \, \chi^{(n)}(\mathbf{x}) \sim_{a_1} \dots \sim_{a_n} = \frac{k(N-k)}{2N} + \frac{1}{2} n^2 \quad (4.33)$$

We should further calculate the number of times a term with a given n appears in the sum (4.32). In what follows, we will consider the case $k < N$, although k may be of the order of N . Then, $C_k^n = \frac{k!}{n!(k-n)!}$ possibilities exist to choose out of k weight vectors n coinciding ones, whose index may acquire any values from 1 to N . The index of any weight vector out of other $(k-n)$ ones may then acquire only $(N-1)$ values, and so on. Finally, the index of the last weight vector may acquire $(N-k+n)$ values. Therefore, the desired number of times, a term with a given n appears in the sum (4.32), reads:

$$C_k^n N (k-n)(N-1) (k-n-1)(N-2) \dots 1(N-k+n) = \\ = C_k^n A_N^{k-n+1} (k-n)! = \frac{k N!}{n!(n+N-k-1)!}; \quad (4.34)$$

where $A_N^{k-n+1} = \frac{N!}{(N-k+n-1)!}$, Equations (4.33) and (4.34) together yield for the monopole contribution to the Wilson loop, Eq. (4.30):

$$hW_k(C)_{\text{mon}} = k N e^{-C S} \sum_{n=1}^{X^k} \frac{1}{n!(n+N-k-1)!} e^{\frac{n^2-n}{2} S}; \quad (4.35)$$

where $C = \frac{k(N-k)}{2N}$ is proportional to the Casimir of the rank- k antisymmetric representation of $SU(N)$. We have thus arrived at a Feynman-Kac type formula, where, in the asymptotic regime of interest, $S \gg 1$, only the first term in the sum is essential. The k -string tension therefore reads $\sigma_k = C$, that yields the Casimir-scaling law (4.25). It is interesting to note that the Casimir of the original unbroken $SU(N)$ group is recovered. This is a consequence of the Dirac quantization condition [61], which distributes the quark charges along the weights of the fundamental representation and the monopole ones along the roots. The orthonormality of the roots then yields the action (4.31), which is diagonal in the dual magnetic variables; the sum of the weights squared is responsible for the Casimir factor, since $C = (\sum_{a_1} \gamma + \sum_{a_k} \gamma)^2$ where all k weight vectors are different from each other. Therefore, terms where all k weight vectors are mutually different yield the dominant contribution to the sum (4.30). Their number in the sum is equal to $\frac{k N!}{(N-k)!}$, that corresponds to the $(n=1)$ -term in Eq. (4.35).

Let us further address the leading correction to the obtained Casimir scaling, which originates from the non-diluteness of plasma. Expanding the cosine up to the quartic term, we obtain the action [One should use the formula

$$\sum_{i=1}^{N(N-1)/2} q_i q_i q_i q_i = \frac{N}{2(N+1)} + \dots ;$$

34 D. Antonov and M. C. Diamantini

which stems from the orthonormality of roots.]

$$\int d^3x \left[\frac{1}{2} (\partial_\mu \tilde{\phi})^2 + \frac{m_D^2}{2} \tilde{\phi}^2 - \frac{g_m^2}{12(N+1)} \tilde{\phi}^4 \right] \quad (4.36)$$

By virtue of this formula, one can analyze the correspondence between the $1/N$ -expansion and corrections to the low-density approximation. The natural choice for defining the behavior of the electric coupling constant in the large- N limit is the QCD-inspired one, $g = O(N^{-1/2})$. To make some estimates, let us use an obvious argument that a $\tilde{\phi}$ -field configuration dominating in the partition function is the one, where every term in the action (4.36) is of the order of unity. When applied to the kinetic term, this demand tells us that the characteristic wavelength l of the field $\tilde{\phi}$ is related to the amplitude of this field as $l \sim j^{-2}$. Substituting further this estimate into the condition $\int d^3x m_D^2 \tilde{\phi}^2 \sim 1$, we get $l \sim j^{-2} m_D^{-1}$. The ratio of the quartic and mass terms, being of the order of $j^{-2} g_m^2 \sim N$, can then be estimated as $\frac{m_D g_m^2}{N} = \frac{g_m^3}{N} N^{p-}$. With the exponentially high accuracy, this ratio is small, provided $N \gg O(e^{S_0=2})$. Therefore, the non-diluteness corrections are suppressed only for N 's bounded from above by a certain parameter. However, due to the exponential largeness of this parameter, the derived constraint leaves enough space for N to be sufficiently large, in order to provide the validity of the inequalities (4.29).

To proceed with the study of the non-diluteness correction, one needs to solve iteratively the saddle-point equation, corresponding to the average (4.30) taken with respect to the approximate action (4.36). Since it has been demonstrated above that the string tension is defined by the averages $e^{ia(\tilde{a}_1 + \tilde{a}_k) \cdot \tilde{\phi}}$, where all k weight vectors are mutually different, let us restrict ourselves to such terms in the sum (4.30) only. Solving then the saddle-point equation with the Ansatz $\tilde{\phi} = \tilde{\phi}_0 + \tilde{\phi}_1$, where $\tilde{\phi}_1 \cdot \tilde{\phi}_0 = 0$, we obtain for such a term:

$$\ln \int d^3x e^{ia(\tilde{a}_1 + \tilde{a}_k) \cdot \tilde{\phi}} = \frac{g^2}{2} C \int d^3x d^3y (\tilde{x}) D_{m_D}(\tilde{x} - \tilde{y}) (\tilde{y}) + S \quad (4.37)$$

The first term on the r.h.s. of Eq. (4.37) yields the string tension $\sigma_k = C$,

^sThis constraint is similar to those which were derived at the end of sect. 2 as the necessary conditions for the stochasticity of the Higgs vacuum. We therefore conclude that the conditions of stochasticity of the Higgs vacuum and of the validity of the dilute-plasma approximation parallel each other.

while the second term yields the desired correction. This term reads

$$\begin{aligned}
 S &= \frac{2}{3} \frac{(gm_D C)^2}{N+1} \int_{\mathbb{R}^3} d^3x \int_{\mathbb{R}^4} d^4Y \int_{\mathbb{R}^Z} d^Z x_1 D_{m_D}(\mathbf{x} - \mathbf{x}_1) (\mathbf{x}_1) = \\
 &= \frac{2}{6} \frac{(gm_D C)^2}{N+1} \int_{\mathbb{R}^Z} d^Z (\mathbf{x}_1) \int_{\mathbb{R}^Z} d^Z (\mathbf{x}_2) \int_{\mathbb{R}^Z} d^Z (\mathbf{x}_3) \int_{\mathbb{R}^Z} d^Z (\mathbf{x}_4) e^{x_1} e^{x_2} e^{x_3} e^{x_4} I ; \tag{4.38}
 \end{aligned}$$

where $I = \int_{\mathbb{R}^3} d^3x \int_{\mathbb{R}^4} d^4Y \int_{\mathbb{R}^Z} d^Z x_1 D_{m_D}(\mathbf{x} - \mathbf{x}_1)$. The action (4.38) can be represented in the form

$$\begin{aligned}
 S &= \frac{(gC)^2}{(N+1)m_D^5} \int_{\mathbb{R}^Z} d^Z (\mathbf{x}_1) \int_{\mathbb{R}^Z} d^Z (\mathbf{x}_2) D(\mathbf{x}_1 - \mathbf{x}_2) \\
 &\quad \int_{\mathbb{R}^Z} d^Z (\mathbf{x}_3) \int_{\mathbb{R}^Z} d^Z (\mathbf{x}_4) D(\mathbf{x}_3 - \mathbf{x}_4) G(\mathbf{x}_1 - \mathbf{x}_3) ; \tag{4.39}
 \end{aligned}$$

Here, D and G are some positive functions, which depend on $m_D |\mathbf{x}_i - \mathbf{x}_j|$ and vanish exponentially at the distances $\gg d$. They can be represented as $D(\mathbf{x}) = m_D^4 D(m_D |\mathbf{x}|)$, $G(\mathbf{x}) = m_D^4 G(m_D |\mathbf{x}|)$, where the functions D and G are dimensionless.^t

The derivative expansion yields as leading terms the Nambu-Goto actions:

$$\int_{\mathbb{R}^Z} d^Z (\mathbf{x}_1) \int_{\mathbb{R}^Z} d^Z (\mathbf{x}_2) D(\mathbf{x}_1 - \mathbf{x}_2) = \int_{\mathbb{R}^Z} d^Z d^2 \int_{\mathbb{R}^P} \frac{1}{g(\mathbf{x}_1)} + O\left(\frac{D}{m_D^2}\right) ; \tag{4.40}$$

Here, $\int_{\mathbb{R}^Z} d^Z = \int_{\mathbb{R}^P} \frac{2m_D^2}{g(\mathbf{x})} d^2z D(z)$ (with z being dimensionless). Recalling that $\int_{\mathbb{R}^Z} d^Z (\mathbf{x}) = \int_{\mathbb{R}^Z} d^Z (\mathbf{x}) d^2$, we may further take into account that we are interested in the leading term of the derivative expansion of the action S , which corresponds to the so short distance $|\mathbf{x}_1 - \mathbf{x}_3|$ that $t(\mathbf{x}_1) t(\mathbf{x}_3) \approx 2$. (Higher terms of the derivative expansion contain derivatives of t and do not contribute to the string tension.) This yields for the integral in Eq. (4.39):

$$\begin{aligned}
 &\int_{\mathbb{R}^Z} d^Z \int_{\mathbb{R}^Z} d^Z \int_{\mathbb{R}^P} \frac{1}{g(\mathbf{x}_1)g(\mathbf{x}_3)} G(\mathbf{x}_1 - \mathbf{x}_3) ' \\
 &' \frac{2}{2} \int_{\mathbb{R}^Z} d^Z (\mathbf{x}_1) \int_{\mathbb{R}^Z} d^Z (\mathbf{x}_3) G(\mathbf{x}_1 - \mathbf{x}_3) = \frac{2}{2} \int_{\mathbb{R}^Z} d^Z \int_{\mathbb{R}^P} \frac{1}{g} + O\left(\frac{G}{m_D^2}\right) ;
 \end{aligned}$$

where $\int_{\mathbb{R}^Z} d^Z = \int_{\mathbb{R}^P} \frac{2m_D^2}{g} d^2z G(z)$. We finally obtain from Eq. (4.39):

^t Our investigations can readily be translated to the stochastic vacuum model of QCD [80] for the evaluation of a correction to the string tension, produced by the four-point irreducible average of field strengths. In that case, the functions D and G would be proportional to the gluonic condensate.

36 D. Antonov and M. C. Diamantini

$$k', \frac{(gC)^2 \frac{2}{D} G}{2(N+1)m_D^5} = \frac{(gC)^2 m_D}{4(N+1)} = \frac{C^2}{N+1};$$

where k' is some dimensionless positive constant. This yields

$$k + k = C \left(1 + \frac{C}{N+1} \right);$$

In the limit $k \ll N - 1$ of interest, the obtained correction is $O(1)$, while for $k = O(1)$ it is $O(1/N)$. The latter fact enables one to write down the following final result for the leading correction to Eq. (4.25) due to the non-diluteness of monopole plasma:

$$R(k;N) + R(k;N) = \frac{k + k}{1 + 1} = C(k;N) \left(1 + \frac{(k-1)(N-k-1)}{2N(N+1)} \right);$$

This expression is as invariant under the replacement $k \leftrightarrow (N - k)$ as the expression (4.25), which does not account for non-diluteness. The fact that, at $k \ll N - 1$, the obtained correction to the Casimir-scaling law is $O(1)$, indicates that non-diluteness effects can significantly distort the Casimir-scaling behavior.

5. Generalization to the $SU(N)$ -analogue of 4D compact QED with the θ -term

In this section, we will consider the 4D-case and introduce the field-theoretical θ -term⁴. As it has been first found for compact QED in Refs. [33,36] by means of the derivative expansion of the resulting non-local string effective action, this term generates the string θ -term. Being proportional to the number of self-intersections of the world sheet, the latter might be important for the solution of the problem of crumpling of large world sheets [3,31]. In this section, we will perform the respective analysis for the general $SU(N)$ -case under study, in the fundamental and in the adjoint representations, as well as for k -strings. It is worth noting that, in the lattice 4D compact QED, confinement holds only at strong coupling. On the opposite, the continuum counterpart [33,36] [of the $SU(N)$ -version] of this model, we are going to explore, possesses confinement at arbitrary values of coupling. However, we will see that the solution to the problem of crumpling

⁴ In the case when the ensemble of Abelian-projected monopoles is modeled by the magnetically-charged dual Higgs field, the effects, produced by this term to the respective string effective action, have been studied in Ref. [67].

due to the μ -term is only possible in the strong-coupling regime, implied in a certain sense. Note also that the continuum sine-Gordon theory of the dual-photon field possesses an ultraviolet cutoff, Λ , which appears in course of the path-integral average over the shapes of monopole loops. However, the μ -parameter is dimensionless, and consequently its values, at which one may expect the disappearance of crumpling, will be cutoff-independent.

The full partition function, including the μ -term and the average over free photons, reads [cf. Eq. (3.1)]

$$Z^N = \int_{\mathcal{D}\tilde{A}} e^{\frac{1}{4} \int_{\mathbb{R}^4} d^4x F^2} \sum_{N=0}^{\infty} \frac{1}{N!} \exp \left[\int_{\mathcal{S}} d^4x \tilde{j}^N; \tilde{A} \right] \int_{\mathcal{M}_{\text{on}}} \mathcal{H}^{i_0 E} ; \quad (5.1)$$

where

$$\mathcal{H}^{i_0 E} ; \tilde{A} = \int_{\mathbb{R}^4} d^4x d^4y \tilde{j}^N(x) D_0(x-y) \tilde{j}^N(y) + \frac{i g^2}{8} \int_{\mathbb{R}^4} d^4x \tilde{A} \tilde{j}^N ; \quad (5.2)$$

and $F^2 = \partial \tilde{A} \wedge \partial \tilde{A}$. For $N = 0$, the monopole current \tilde{j}^N is equal to zero, whereas for $N \geq 1$, it is defined as $\tilde{j}^N = g_m \int_{\mathbb{P}^1} \epsilon_{ik}^H dz^k(\sigma) \times x^k(\sigma)$. The couplings g and g_m are now dimensionless and obey the same condition $gg_m = 4$ as in the 3D-case. We have also parametrized the trajectory of the k -th monopole by the vector $x^k(\sigma) = y^k + z^k(\sigma)$, where $y^k = \int_0^R d\sigma x^k(\sigma)$ is the position of the trajectory, whereas the vector $z^k(\sigma)$ describes its shape, both of which should be averaged over. The fugacity of a single monopole loop, μ , entering Eq. (5.1), has the dimensionality $[\text{mass}]^4$, $\mu = e^{-S_{\text{mon}}}$, where the action of a single k -th loop, obeying the estimate $S_{\text{mon}} \sim \frac{1}{g^2} \int_0^R d\sigma (z^k)^2$, is assumed to be of the same order of magnitude for all loops. Finally, in Eq. (5.1), $D_0(x) = 1/(4\pi^2 x^2)$ is the 4D Coulomb propagator, and the average over monopole loops is defined similarly to Eq. (3.2) as follows:

$$\langle \mathcal{H}^{i_0 E} \rangle_{\mathcal{M}_{\text{on}}} = \int_{\mathcal{M}_{\text{on}}} \int_{\mathcal{D}z_n} d^4y_n \prod_{i_n=1}^N \int_{\mathcal{D}z_n} \mathcal{H}^{i_0 E} ; \quad (5.3)$$

The particular form of the path-integral average over the shapes of the loops, $z_n(\sigma)$'s, here is immaterial for the final (ultraviolet-cutoff dependent) expression for the partition function [see e.g. Ref. [68] for a similar situation in the plasma of closed dual strings in the Abelian-Higgs (type models)]. The only

38 D. Antonov and M. C. Diamantini

thing which matters is the normalization $\text{hli}_{z_n(\cdot)} = 1$, that will be implied henceforth. The analogue of the partition function (3.9), (3.10) then reads

$$Z_{\text{mon}}^N \text{h i} Z_n \text{h io} = \int D\tilde{\gamma} \exp S[\tilde{\gamma}; \tilde{A}] \quad \#)$$

$$D \sim \exp \int d^4x \frac{1}{2} X \cos \frac{g_1 \tilde{\gamma} \cdot \tilde{\gamma}}{i} + i \tilde{\gamma} \quad ; \quad (5.3)$$

with the action S given by Eq. (5.2), while the full partition function is

$$Z^N = \int D\tilde{A} e^{\frac{1}{4} \int d^4x F^2} Z_{\text{mon}}^N \text{h i} :$$

In Eq. (5.3),

$$\tilde{\gamma} \cdot \tilde{\gamma} = \frac{q}{1} ; \dots ; \frac{q}{N-1} ; \dots ; \frac{q}{N-1} ; \quad ; \quad (5.4)$$

and $y^k = z^{k-2} z^{k-1}$ is the ultraviolet cutoff. Clearly, unlike the 3D case without the $\tilde{\gamma}$ -term, the \tilde{A} -field is now coupled to $\tilde{\gamma}$, making Z_{mon}^N \tilde{A} -dependent. This eventually leads to the change of the mass of the dual photon $\tilde{\gamma}$, as well as to the appearance of the string-term.

Let us address the fundamental case first. After the saddle-point integration over $\tilde{\gamma}$,^v the analogue of Eqs. (4.1), (4.2) takes the form

$$W_a = \frac{1}{Z^N} \int D\tilde{\eta} \exp \left(\int d^4x \text{h i} S[\tilde{\eta}] + \frac{ig}{2} \int d^3x \tilde{\eta} \cdot \tilde{\eta} \right) ; \quad (5.5)$$

(c)

where the constraint $\partial \tilde{\eta} = 0$ was already abolished. Here, the Kalb-Ramond action reads

$$S[\tilde{\eta}] = \int d^4x \left[\frac{1}{4} \tilde{\eta}^2 - \frac{ig^2}{32} \tilde{\eta} \cdot \tilde{\eta} \right] + V[\tilde{\eta}] ; \quad (5.6)$$

The potential V in this equation is given by Eq. (4.2) with the symbol $\int d^3x$ replaced by $\int d^4x$, and

^v Unlike the superrenormalizable 3D case, where loop corrections to the saddle point are rapidly converging, this is not necessarily the case in four dimensions. This fact is, however, clearly unimportant in the weak-field limit [Eq. (5.7) below], where the cosine in Eq. (5.3) is approximated by the quadratic term only, and the saddle-point integration over $\tilde{\gamma}$ becomes Gaussian. As we have seen in the previous section, this limit is already enough for the discussion of the string representation. One of the authors (D.A.) is grateful to H. Gies and E. Vicari for pointing out his attention to this issue.

$$H_a = \frac{g}{(N-1)} \tilde{\gamma}_a \in \tilde{\mathfrak{h}} \quad ;$$

In these formulae, $\tilde{\gamma} = \frac{1}{2} \mathbf{0}$, and the absolute value is defined in the same way as in Eq. (5.4), i.e. again with respect to the Lorentz indices only. Note that the form, to which the $\tilde{\gamma}$ -term has been transformed, is quite natural, since the respective initial expression of Eq. (5.2) can be rewritten modulo full derivatives as

$$\frac{i g^2}{8} \int d^4 x \tilde{\gamma}^N = \frac{i g^2}{32} \int d^4 x (F^2 + F^N \cdot \tilde{F} + \tilde{F}^N) \quad ;$$

where [cf. Eq. (3.4)]

$$F^N(x) = \int d^4 y D_0(x-y) j^N(y); \quad \tilde{F}^N = j^N \quad ;$$

Next, the mass of the Kalb-Ramond field, equal to the Debye mass of the dual photon, which follows from the action (5.6), reads $m_D = \frac{g}{4} \sqrt{\frac{4}{g^2} + \frac{1}{2}}$, where $\frac{1}{N} = \frac{1}{2}$. In the extreme strong-coupling limit, $g \rightarrow 1$, this expression demonstrates the important difference of the case $\epsilon = 0$ from the case $\epsilon \neq 0$. Namely, since $(g)/e^{\text{const} \cdot g^2} \rightarrow 1$, $m_D \rightarrow 0$ at $\epsilon = 0$, whereas $m_D \rightarrow 1$ at $\epsilon \neq 0$. In other words, in the extreme strong-coupling limit, the correlation length of the vacuum, equal to d , goes large (small) at $\epsilon = 0$ ($\epsilon \neq 0$).

In the weak-field limit, $H_a \rightarrow 1$, Eq. (5.5) yields [cf. Eq. (4.11)]:

$$\begin{aligned} W(C; \gamma)_{\text{weak field}}^{\text{tot}; a} &= \frac{1}{Z^{\text{tot}}} \int D\tilde{\mathfrak{h}} \\ \exp \int d^4 x & \left[\frac{g^2}{12} H^2 + \frac{1}{4} \tilde{\mathfrak{h}}^2 - \frac{i g^2}{32} \tilde{\mathfrak{h}} \cdot \tilde{\mathfrak{h}} - \frac{i g}{2} \tilde{\gamma}_a \tilde{\mathfrak{h}} \right] \quad ; \end{aligned} \quad (5.7)$$

where again Z^{tot} is the same integral over $\tilde{\mathfrak{h}}$, but with $\tilde{\gamma}$ set to zero. Integrating over $\tilde{\mathfrak{h}}$, we then obtain:

$$\begin{aligned} W(C; \gamma)_{\text{weak field}}^{\text{tot}; a} &= \exp \left[\frac{N}{4N} \int_C \int_C g^2 dx^0 dx^0 D_{m_D}(x-x^0) + \right. \\ & \left. + \frac{1}{2} \int d^4 x d^4 x^0 D_{m_D}(x-x^0) \left((x) \cdot (x^0) + \frac{i g^2}{8} (x) \cdot (x^0) \right) \right] \quad ; \end{aligned} \quad (5.8)$$

40 D. Antonov and M. C. Diamantini

where $D_m(x) = m K_1(m|x|) = (4 - 2|x|)$ is the 4D Yukawa propagator with K_1 denoting the modified Bessel function. [Note that, at $\epsilon = 0$, Eq. (5.8) takes the form of Eq. (4.13).] Further curvature expansion of the $\tilde{\mathcal{L}}$ interaction [analogous to the expansion of the \mathcal{L} interaction in the action (4.14)] yields the string term equal to ic_{fund} . Here, $\frac{1}{2} \int d^2 \sigma \sqrt{-g} g^{ab} (\partial_a t) (\partial_b \tau)$ is the number of self-intersections of the world sheet, and the coupling constant c reads

$$c_{\text{fund}} = \frac{h(N-1)}{8N \left(\frac{4}{g^2} + \frac{1}{2} \right)} \quad (5.9)$$

As it has already been discussed, c_{fund} is ϵ -independent, since (similarly to the rigidity coupling constant) it is dimensionless. We therefore see that, at

$$g_{\text{fund}} = \frac{2}{4} \frac{N-1}{2N} \left(\frac{N-1}{2N} \right)^2 \frac{16}{g^2} \frac{3}{5}; \quad (5.10)$$

c_{fund} becomes equal to $\frac{1}{4}$, and self-intersections are weighted in the string partition function with the factor $(-1)^j$, that might cure the problem of crumpling. This is only possible at $g = g_{\text{fund}} = 4 \frac{2-N}{N-1}$,^w that parallels the strong-coupling regime, which should hold in the lattice version of the model (cf. the discussion in the beginning of this section). In the extreme strong-coupling limit, understood in the sense $g = g_{\text{fund}}$, only one critical value, g_{fund}^+ , survives, $g_{\text{fund}}^+ = \frac{N-1}{2N}$, whereas $g_{\text{fund}}^- = 0$, i.e. g_{fund} becomes a spurious solution, since $c_{\text{fund}}|_{g=0} = 0$.

With the use of the results of subsection 4.2, the adjoint-case large- N counterpart of Eq. (5.10) can readily be found. Indeed, Eq. (5.9) in that case becomes replaced by

$$c_{\text{adj}} = \frac{h}{4 \left(\frac{4}{g^2} + \frac{1}{2} \right)};$$

so that c_{adj} is equal to $\frac{2}{4} \frac{1}{1} \left(\frac{16}{g^2} \right) \frac{3}{5}$:

$$g_{\text{adj}} = \frac{2}{4} \frac{1}{1} \frac{16}{g^2} \frac{3}{5};$$

Note that the respective lower bound for the critical value of g , $g_{N-1}^{\text{adj}} = 4 \frac{1}{2}$,^x is only slightly different from the value $g_{N-1}^{\text{fund}} = 4 \frac{2-N}{N-1}$. Similarly

^w Clearly, at $g = g_{\text{fund}}$, $g_{\text{fund}}^+ = g_{\text{fund}} = \frac{N-1}{4N}$.

^x As in the previous footnote, at the particular value of g , $g = g_{N-1}^{\text{adj}}$, $g_{\text{adj}}^+ = g_{\text{adj}} = \frac{1}{2}$.

to the fundamental case, at $g = g_{N-1}^{\text{adj}}$, g^{adj} becomes a spurious solution, whereas g^{fund} is not. Thus, at $N = 1$ and in the strong-coupling limit, understood in the sense $g = g_{N-1}^{\text{fund}}$, the critical fundamental- and adjoint-case values of g , at which the problem of condensing might be solved, are given by the following simple formula: $g^{\text{fund}} = \frac{1}{2} g^{\text{adj}} = \frac{1}{2}$.

Finally, for k -strings, the ratio $\alpha_k = C_{\text{fund}}$ is the same as in 3D, $C(k; N)$, as the ratio of string tensions, as long as $N \geq 0$.^Y This enables one to readily find g_k as well. Namely,

$$g_k = \frac{1}{2} \sqrt{\frac{4}{2N C(k; N)} \left(\frac{N-1}{2N C(k; N)} \right)^2 + \frac{16}{g^2}}^{3/5};$$

that is only possible at $g > g_k = g^{\text{fund}P} \frac{1}{C(k; N)} = g^{\text{fund}}$.

6. Geometric aspects of condensing strings: the physics of negative stiffness

The role of antisymmetric tensor field theories for condensation has been studied in detail in Ref. [33]. By analyzing compact antisymmetric tensor field theories of rank $h = 1$ Quevedo and Tugueberger have shown that, starting from a "Coulomb" phase, the condensation of $(d-h-1)$ -branes (where $D = d+1$ is the space-time dimension) leads to a generalized condensation phase for $(h-1)$ -branes. Each phase in the model has two dual descriptions in terms of antisymmetric tensor of different ranks, massless for the Coulomb phase and massive for the condensation phase. Upon the integration over the massive antisymmetric tensor field (in the case of even number of dimensions when the θ -term is absent), one obtains string effective action (4.14). The derivative expansion of this nonlocal string effective action, up to the term next to the rigidity one, produces the following condensing-string action:

$$S = \int d^D x \sqrt{-g} g^{ab} D_a x^\mu D_b x^\nu \left(2T \sqrt{-D^2} + \frac{1}{M^2} D^4 D_b x^\nu \right); \quad (6.1)$$

Here, D_a is the covariant derivative with respect to the induced metric $g_{ab} = (g_{\mu\nu} x^\mu x^\nu)$ on the surface $x^\mu(0; 1)$.

In (6.1), the first term provides a bare surface tension $2T$, while the second accounts for rigidity with stiffness parameters. The last term can be written

^Y In 4D, the ratio of string tensions indeed equals $C(k; N)$ as in 3D, as long as the dual-photon field can be treated as a free massive field, since the rest of the derivation of Eq. (4.35) is based on the properties of fundamental weights only.

42 D. Antonov and M. C. Diamantini

(up to surface terms) as a combination of the fourth power and the square of the gradient of the extrinsic curvature matrices, with M being a new mass scale. It thus suppresses world-sheet configurations with rapidly changing extrinsic curvature; due to its presence, the stiffness may not necessarily be negative, as in Eq. (4.17) above, but also positive. We analyze the model (6.1) in the large- D approximation. To this end, we introduce a Lagrange multiplier matrix L^{ab} to impose the constraint $g_{ab} = (\partial_a x^\mu)(\partial_b x^\nu)g_{\mu\nu}$, extending the action (6.1) to

$$S + \int d^2 \sigma \bar{g} L^{ab} (\partial_a x^\mu \partial_b x^\nu g_{\mu\nu}) : \quad (6.2)$$

Then we parametrize the world-sheet in a Gauss map by $x^\mu(\sigma) = (x^0; x^i)$, ($i = 1, \dots, D-1$), where $x^0 = 2\sigma^0 = 2\tau$, $R = 2\sigma^1$, $R = 2$, and $x^i(\sigma)$ describe the $D-2$ transverse fluctuations. With the usual homogeneity and isotropy Ansatz $g_{ab} = \gamma_{ab}$, $L^{ab} = L g^{ab}$ of infinite surfaces ($R \gg 1$) at the saddle point, we obtain

$$S = 2 \int d^2 \sigma [T + L(1 - \gamma)] + \int d^2 \sigma \gamma_{ij} V(T; s; M; L; D^2) \gamma_a^i \gamma_b^j ; \quad (6.3)$$

where

$$V(T; s; M; L; D^2) = T + L \left(sD^2 + \frac{1}{M^2} D^4 \right) ; \quad (6.4)$$

Integrating over the transverse fluctuations, in the infinite-area limit, we get the effective action

$$S^e = 2A_{\text{ext}} [T + L(1 - \gamma)] + A_{\text{ext}} \frac{D^2}{8} \int d^2 \sigma \ln p^2 V(T; s; M; L; p) ; \quad (6.5)$$

where $A_{\text{ext}} = R^2$ is the extrinsic, physical, space-time area. For large D , the fluctuations of L and γ are suppressed and these variables take their "classical values", determined by the two saddle-point equations

$$0 = f(T; s; M; L) ; \quad \gamma = \frac{1}{f^0(T; s; M; L)} ; \quad (6.6)$$

where the prime denotes a derivative with respect to L and the "saddle-function" f is defined by

$$f(T; s; M; L) = L \frac{D^2}{8} \int dp p \ln p^2 V(T; s; M; L; p) : \quad (6.7)$$

Using (6.6) in (6.5) we get $S^e = 2(T + L)A_{\text{ext}}$ showing that $T = 2(T + L)$ is the physical string tension.

The stability condition for the Euclidean surfaces is that $V(T; s; M; L; p)$ be positive for all $p^2 > 0$. However, we will require the same condition also for $p^2 = 0$, so that strings propagating in Minkowski space-time are not affected by the propagating states of negative norm which plague rigid strings. The stability condition becomes thus $\frac{M^2 p^2}{T + L} - \frac{sM^2}{4} = 2j$, which allows us to introduce the real variables R and I defined by

$$R^2 = \frac{M^2 p^2}{2(T + L)} + \frac{sM^2}{4}; \quad I^2 = \frac{M^2 p^2}{2(T + L)} - \frac{sM^2}{4}; \quad (6.8)$$

In terms of these, the kernel V can be written as

$$M^2 V(T; s; M; L; p) = R^2 + I^2 + 2R^2 I^2 p^2 + p^4; \quad (6.9)$$

In order to analyze the geometric properties of the string model (6.1) we will study two correlation functions. First, we consider the orientational correlation function

$$g_{ab}(\sigma^0) = \langle h_a^i(\sigma) \mathcal{Q}_b^i(\sigma^0) \rangle$$

for the normal components of tangent vectors to the world-sheet.

Secondly, we compute the scaling law of the distance d_E in embedding space between two points on the world-sheet when changing its projection d on the reference plane. The exact relation between the two lengths is

$$d_E^2 = d^2 + \sum_i \langle h_j^i(\sigma) \mathcal{Q}_i^j(\sigma^0) \rangle; \quad (6.10)$$

When d_E^2 / d^2 this implies that the Hausdorff dimension of the surfaces, defined as $d_E^2 = (d^2)^{2/D_H}$, is equal to 2, and surface is smooth.

In order to establish the properties of our model, we analyze the saddle-point function $f(T; s; M; L)$ in (6.7). To this end, we must prescribe a regularization for the ultraviolet divergent integral. We use dimensional regularization, computing the integral in $(2 - \epsilon)$ dimensions. For small ϵ , this leads to

$$f(T; s; M; L) = L + \frac{1}{16^{3/2}} R^2 I^2 \ln \frac{R^2 + I^2}{2} - \frac{1}{8^{3/2}} R I \frac{1}{2} + \arctan \frac{I^2 - R^2}{2RI}; \quad (6.11)$$

where $\exp(2\epsilon)$ and μ is a reference scale which must be introduced for dimensional reasons. The scale μ plays the role of an ultraviolet cutoff, diverging for $\epsilon \rightarrow 0$.

The saddle-point function above is best studied by introducing the dimensionless couplings $t = T = \mu^2$, $m = M = \mu$ and $l = L = \mu^2$. We will study in detail the case $s = 0$, in which the saddle-point equations can be solved

44 D. Antonov and M. C. Diamantini

analytically since $R = I$. This choice is not too restrictive since, as we will show, $s = 0$ is the infrared fixed point. We get

$$1 = \frac{m^2 c^2}{2} \left(1 + \frac{4t}{m^2 c^2} \right)^{-1}; \quad (6.12)$$

$$= 1 - \frac{m c}{2} \frac{1}{t+1}; \quad (6.13)$$

where $c = 1/32$. This shows that the point $(t = 0, s = 0, m = 0)$ constitutes an infrared-stable fixed-point with vanishing physical string tension T . This point is characterized by long-range correlations $g(d) = a^{-d}$, with a constant a , and by the scaling law

$$d_E^2 = \frac{2}{a} d^2; \quad 1 - \frac{1}{2a} d; \quad (6.14)$$

which shows that the Hausdorff dimension of world sheets is $D_H = 2$. For $s = 0$, the constant a can be computed analytically:

$$a^2 = \lim_{\substack{t \rightarrow 0 \\ m \rightarrow 0}} \frac{1 + (2t - m^2 c^2) + \frac{1}{1 + 4t - m^2 c^2}}{2}; \quad (6.15)$$

from which we recognize that $1 = 2$.

At the infrared fixed point we can remove the cutoff. The renormalization of the model is easily obtained by noting that the effective action for transverse fluctuations to quadratic order decouples from other modes and is identical with the second term in (6.3) with $D^2 = \partial^2$ and taking its saddle-point value. From here we identify the physical tension, stiffness and mass as

$$T = \frac{1}{2} (t + 1); \quad S = \frac{S}{2}; \quad M = m; \quad (6.16)$$

For $s = 0$, we can compute analytically the corresponding β -functions:

$$\frac{d}{dt} \ln t = 2 + O\left(\frac{t}{m^2}\right); \quad (6.17)$$

$$\frac{d}{dm} \ln m = 1 + O\left(\frac{t^2}{m^4}\right); \quad (6.18)$$

The vicinity of the infrared fixed-point defines a new theory of smooth strings for which the range of the orientational correlations in embedding space is always of the same order or bigger than the length scale $l_p = \sqrt{T}$ associated with the tension. The naively irrelevant term $D^4 = M^2$ in (6.1) becomes relevant in the large- D approximation since it generates a string

tension proportional to M^2 which takes over the control of the fluctuations after the orientational correlations die off. Note moreover, that it is exactly this new quartic term which guarantees that the spectrum $p^2 V(\Gamma; s; M; L; p)$ has no other pole than $p = 0$, contrary to the rigid string, which necessarily has a ghost pole at $p^2 = -T/s$.

By studying the finite-size scaling [87] of the Euclidean model (6.1) on a cylinder of (spatial) circumference R it is possible to determine the universality class of confining strings. In the limit of large R , the effective action on the cylinder takes the form [41]

$$\lim_{R \rightarrow \infty} \frac{S^e}{R} = TR \left[\frac{c(D-2)}{6R} + \dots \right]; \quad (6.19)$$

for $(D-2)$ transverse degrees of freedom, the universality class being encoded in the pure number c . This suggests that the effective theory describing the infrared behavior is a conformal field theory (CFT) with central charge c . In this case the number c also fixes the Luscher term [88] in the quark-antiquark potential:

$$V(R) = TR \left[\frac{c(D-2)}{24R} + \dots \right]; \quad (6.20)$$

In [41] it has been shown that confining strings are characterized by $c = 1$. Although they share the same value of c , confining strings are clearly different $c = 1$ theories than Nambu-Goto or rigid strings. Indeed, the former are smooth strings on any scale, while the latter crumple and fill the ambient space, at least in the infrared region. Our result $c = 1$ is in agreement with recent precision numerical determinations of Luscher and Weisz [30] of this constant.

Having established that the model (6.1) describes smooth strings with $c = 1$, the question arises as to how much these results depend on the truncation of the original non-local action after the D^4 term. In [41] it has been proved that the answer to this question is no: the value of c and the smooth geometric properties are independent of an infinite set of truncations, provided that a solution for the polynomial "gap" equation exists. These properties are presumably common to a large class of non-local world-sheet interactions.

7. High temperature behavior of confining strings

The high-temperature behavior of large- N QCD has been studied by Polchinski in [46], where he shows that the deconfining transition in QCD is due to the condensation of Wilson lines, and the partition function of QCD

flux tubes can be continued above the deconfining transition; this high-temperature continuation can be evaluated perturbatively. So, any string theory that is equivalent to QCD must reproduce this behavior. However, the Nambu-Goto action has the wrong temperature dependence, while the rigid string parametrically has the correct high-temperature behavior but with a wrong sign and an imaginary part signaling a world-sheet instability [89].

The high temperature behavior of the confining string model proposed in [40] has been studied in [45], where it has been shown that this model has a high-temperature behavior that agrees in temperature dependence, sign and reality properties with the large- N QCD result [46]. The starting point will be again (6.2). In the Gaussian map, the value of the periodic coordinate ϕ_0 is $\phi_0 = 2\pi\alpha_0 = 2\pi\alpha_0$ with $\alpha_0 = 1/T$ and T the temperature. Note that, at high temperatures ($\alpha_0 \ll 1$), the scale M^2 can be temperature-dependent. This is not unusual in closed string theory as has been shown by Atick and Witten [90]. The value of α_1 is $\alpha_1 = R^2\alpha_0 = R^2\alpha_0$; $\alpha_1(\alpha_0)$ describe the $D-2$ transverse fluctuations. We look for a saddle-point solution with a diagonal metric $g_{ab} = \text{diag}(\alpha_0; \alpha_1)$, and a Lagrange multiplier of the form $\lambda^{ab} = \text{diag}(\lambda_0=0; \lambda_1=1)$.

After integration over transverse fluctuations we obtain, in the limit $R \rightarrow \infty$, four gap equations:

$$\frac{1}{\alpha_0} = 0; \tag{7.1}$$

$$\frac{1}{\alpha_1} = 1 - \frac{D-2}{2} \frac{1}{4} \frac{P}{2M^2 (\alpha_1 + t)^{3/4}}; \tag{7.2}$$

$$\frac{1}{2} (t - \alpha_1) + \frac{1}{2\alpha_1} (\alpha_1 + t) - t - \alpha_0 + \frac{D-2}{2} \frac{2}{2^2} = 0; \tag{7.3}$$

$$(t - \alpha_1) - \frac{1}{\alpha_1} (\alpha_1 + t) + \frac{D-2}{2} \frac{1}{2M^2 (\alpha_1 + t)^{1/4}} - \alpha_0 = 0; \tag{7.4}$$

and a simplified form of the effective action:

$$S^e = A_{\text{ext}} T \frac{r}{\alpha_1}; \tag{7.5}$$

with $D = 2(\alpha_1 + t)$ representing the physical string tension. By inserting (7.2) into (7.4), we obtain an equation for $(\alpha_1 + t)$ alone:

$$(\alpha_1 + t) - \frac{D-2}{2} \frac{5}{8} \frac{P}{2M^2 (\alpha_1 + t)^{1/4}} + \frac{D-2}{2} \frac{2}{2^2} - t = 0; \tag{7.6}$$

Without loss of generality we set

$$(\alpha_1 + t)^{1=4} = \frac{P}{2M} ; \tag{7.7}$$

where α_1 is a dimensionless parameter. It is possible to show that, at high temperatures, when

$$t^2 \ll \frac{D-2}{2} ; \tag{7.8}$$

we can completely neglect t in (7.6). Indeed, as we now show, α_1 is proportional to $(D-2)^{1/2}$. We can thus rewrite (7.6) as

$$\alpha_1 \frac{D-2}{2} \frac{5}{8} \alpha_1^{1=2} + \frac{D-2}{2} \frac{2}{2^2} = 0 ; \tag{7.9}$$

We now restrict ourselves to the regime

$$G(\alpha_1; D) = \frac{25}{64} \alpha_1^2 \frac{D-2}{2} \alpha_1^{1=2} - \frac{D-2}{2} > 0 ; \tag{7.10}$$

for which (7.9) admits two real solutions:

$$(\alpha_1^{1=2})^1 = \frac{5}{16} \frac{D-2}{2} + \frac{1}{2} \sqrt{\frac{P}{G(\alpha_1; D)}} ; \tag{7.11}$$

$$(\alpha_1^{1=2})^2 = \frac{5}{16} \frac{D-2}{2} - \frac{1}{2} \sqrt{\frac{P}{G(\alpha_1; D)}} ; \tag{7.12}$$

In both cases, α_1 is proportional to $(D-2)^{1/2}$, which justifies neglecting t in (7.6) and implies that the scale M^2 must be chosen proportional to $1=(D-2)$. Moreover, since the physical string tension is real we are guaranteed that $M^2 > 0$, as required by the stability of our model. Any complex solutions for D would have been incompatible with the stability of the truncation.

Let us start with analyzing the first solution (7.11). By inserting (7.11) into (7.3), we obtain the following equation for α_1 :

$$\frac{1}{\alpha_1} = 1 - \frac{4}{5 + \sqrt{25 - \frac{128}{2^{D-2}}}} ; \tag{7.13}$$

Owing to the condition (7.10), $1=(\alpha_1^{1=2})^1$ is positive and, since $\alpha_1^{1=2}$ is real, the squared free energy is also positive:

$$F^2(\alpha_1) = \frac{S_e^2}{R^2} = \frac{1}{2} \frac{5}{16} \frac{D-2}{2} \frac{1}{2} \sqrt{\frac{P}{G(\alpha_1; D)}}^4 \left(1 - \frac{4}{5 + \sqrt{25 - \frac{128}{2^{D-2}}}} \right)^2 A ; \tag{7.14}$$

In this case the high-temperature behavior is the same as in QCD, but the sign is wrong, exactly as for the rigid string. There is, however, a crucial difference: (7.14) is real, while the squared free energy for the rigid string is imaginary, signaling an instability in the model.

If we now look at the behavior of β_1 at low temperatures, below the deconfining transition [41], we see that β_1 is positive. The deconfining transition is indeed determined by the vanishing of β_1 at $T = T_{dec}$. In the case of (7.11) this means that β_1 is positive below the Hagedorn transition, touches zero at T_{dec} and remains positive above it. Exactly the same will happen also for F^2 , which is positive below T_{dec} , touches zero at T_{dec} and remains positive above it. This solution thus describes an unphysical "error" of the low-temperature behavior of the confining string, without a real deconfining Hagedorn transition. For this reason we discard it.

Let us now study the solution (7.12). Again, by inserting (7.12) into (7.3), we obtain for β_1 the equation:

$$\frac{1}{\beta_1} = 1 - \frac{5}{25} e^{-\frac{4}{25} \left(\frac{128}{2^{D/2}} \right)^2} ; \tag{7.15}$$

In this case, when

$$r > 4 \frac{D-2}{3} ; \tag{7.16}$$

β_1 becomes negative. The condition (7.16) is consistent with (7.10) and will be taken to fix the values of the range of parameter r that enters (7.7). We will restrict to those that satisfy (7.16). Since $\beta_0 = 1$ and β_1 is real and proportional to β_1^2 , we obtain the following form of the squared free energy:

$$F^2(\beta) = \frac{1}{2} \left(\frac{5}{16} \frac{D-2}{2} \frac{1}{2} \right)^{D/2} \frac{1}{G(\beta; D)} e^{-\frac{4}{25} \left(\frac{128}{2^{D/2}} \right)^2} \beta^A ; \tag{7.17}$$

In the range defined by (7.16) this is negative. For this solution, thus, both β_1 and F^2 pass from positive values at low temperatures to negative values at high temperatures, exactly as one would expect for a string model undergoing the Hagedorn transition at an intermediate temperature. In fact, this is also what happens in the rigid string case, but there, above the Hagedorn transition, there is a second transition above which, at high temperature, β_1 becomes large and essentially imaginary, giving a positive squared free energy. This second transition is absent in our model.

Let us now compare the result (7.17) with the corresponding one for large- N QCD [46]:

$$F^2(\mu)_{\text{QCD}} = \frac{2g^2(\mu)N}{2} ; \tag{7.18}$$

where $g^2(\mu)$ is the QCD coupling constant. First of all let us simplify our result by choosing large values of μ :

$$\frac{r}{25} \frac{128}{2} \frac{D}{2} \mu^{1=2} ;$$

In this case (7.17) reduces to

$$F^2(\mu) = \frac{1}{2} \frac{8^3 D}{125} \frac{\mu^2}{2} ; \tag{7.19}$$

This corresponds exactly to the QCD result (7.18) with the identifications

$$g^2 / \frac{1}{2} ;$$

$$N / D \mu^2 ;$$

The weak μ -dependence of the QCD coupling $g^2(\mu)$ can be accommodated in the parameter μ . Note that our result is valid at large values of μ , i.e. small values of g^2 , as it should be for QCD at high temperatures [91]. Note also the interesting identification between the order of the gauge group and the number of transverse space-time dimensions. Moreover, since the sign of μ_1 does not change at high temperatures, the field x is not unstable. The opposite happens in the rigid string case [89], where the change of sign of μ_1 gives rise to a world-sheet instability.

8. The influence of matter fields to the deconfinement phase transition in the 3D GG model at finite temperature

8.1. Introduction

The phase structure of the 3D GG model at finite temperature has for the first time been addressed in Ref. [51], where it has been shown that, in the absence of W -bosons, the weakly coupled monopole plasma undergoes the Berezinsky-Kosterlitz-Thouless (BKT) [82] phase transition into the molecular phase at the temperature $T_{\text{BKT}} = g^2/2$. Then, in Ref. [52], it has been shown that the true phase transition, which takes into account W -bosons, occurs at approximately twice smaller temperature. Let us first discuss in some more details the above-mentioned BKT phase transition, which takes

50 D. Antonov and M. C. Diamantini

place in the absence of W -bosons (i.e. in the continuum limit of the 3D lattice compact QED extended by the Higgs field).

At finite temperature $T = 1/\beta$, equations of motion of the fields \mathbf{A} and \mathbf{A}_m , entering the partition function (2.2),

$$Z_{\text{mon}} = \int \mathcal{D}\mathbf{A} \mathcal{D}\mathbf{A}_m \exp \left[-\int d^3x L[\mathbf{A}; \mathbf{A}_m] \right];$$

should be supplemented by the periodic boundary conditions in the temporal direction, with the period equal to β . Because of that, the lines of magnetic field emitted by a monopole cannot cross the boundary of the one-period region. Consequently, at the distances larger than β in the direction perpendicular to the temporal one, magnetic lines should go almost parallel to this boundary, approaching it. Therefore, monopoles separated by such distances interact via the 2D Coulomb potential, rather than the 3D one. Since the average distance between monopoles in the plasma is of the order of $\beta^{1/3}$, we see that at $T \sim \beta^{-1/3}$, the monopole ensemble becomes two-dimensional. Owing to the fact that β is exponentially small in the weak-coupling regime under discussion, the idea of dimensional reduction is perfectly applicable at the temperatures of the order of the above-mentioned critical temperature T_{BKT} .

Note that, due to the T -dependence of the strength of the monopole-antimonopole interaction, which is a consequence of the dimensional reduction, the BKT phase transition in the 3D Georgi-Glashow model is inverse with respect to the standard one of the 2D XY model. Namely, monopoles exist in the plasma phase at the temperatures below T_{BKT} and in the molecular phase above this temperature. As it has already been discussed, the analogy with the 2D XY model established in Ref. [62] is that spin waves of that model correspond to free photons of the 3D GG model at zero temperature, while vortices correspond to magnetic monopoles. At finite temperature, disorder is rather produced by the thermally excited W -bosons [52], whereas monopoles order the system, binding W -bosons into pairs. However, the analogy is still true in case of the continuum version of 3D compact QED (with the Higgs field) under discussion.

Let us therefore briefly discuss the BKT phase transition, occurring in the 2D XY model at a certain critical temperature $T = T_{\text{BKT}}^{\text{XY}}$. At $T < T_{\text{BKT}}^{\text{XY}}$, the spectrum of the model is dominated by massless spin waves, and the periodicity of the angular variable is unimportant in this phase. The spin waves are unable to disorder the spin-spin correlation functions, and those decrease at large distances by some power law. On the contrary, at

$T > T_{BKT}^{XY}$, the periodicity of the angular variable becomes important. This leads to the appearance of topological singularities (vortices) of the angular variable, which, contrary to spin waves, have nonvanishing winding numbers. Such vortices condense and disorder the spin-spin correlation functions, so that those start decreasing exponentially with the distance. Thus, the nature of the BKT phase transition is the condensation of vortices at $T > T_{BKT}^{XY}$. In other words, at $T > T_{BKT}^{XY}$, free vortices do exist and mix in the ground state (vortex condensate) of an infinite global vorticity. Contrary to that, at $T < T_{BKT}^{XY}$, free vortices cannot exist, and they rather mutually couple into bound states of vortex-antivortex pairs. Such vortex-antivortex molecules are small-sized and short-living (virtual) objects. Their dipole-type fields are short-ranged and therefore cannot disorder significantly the spin-spin correlation functions. However, when the temperature starts rising, the size of these molecules grows, until at $T = T_{BKT}^{XY}$ it diverges, that corresponds to the dissociation of the molecules into pairs. Therefore, one of the methods to determine the critical temperature of the BKT phase transition is to evaluate the mean squared separation in the molecule and to find the temperature at which it starts diverging.

Let us now return to the continuum limit of the finite-temperature 3D compact QED, extended by the Higgs field, and determine there the mean squared separation in the monopole-antimonopole molecule. One can then see that, up to exponentially small corrections, the respective critical temperature is unaffected by the finiteness of the Higgs-boson mass. Indeed, the mean squared separation reads²

$$L^2 = \frac{\int_{-m_w}^{m_w} d^2x \int_{-m_w}^{m_w} d^2y \exp\left[-\frac{4T}{g^2} K_0(m_H |x-y|)\right]}{\int_{-m_w}^{m_w} d^2x \int_{-m_w}^{m_w} d^2y \exp\left[-\frac{4T}{g^2} K_0(m_H |x-y|)\right]}; \quad (8.1)$$

where K_0 denotes the modified Bessel function. Disregarding the exponential factors in the numerator and denominator of this equation, we obtain $L^2 \sim \frac{4T}{2m_w^2(2T-g^2)}$, that yields the above-mentioned value of the BKT critical temperature $T_{BKT} = g^2/2$. Besides that, we see that, as long as T does not tend to T_{BKT} , and in the weak-coupling regime under study, the value of $\langle L^2 \rangle$ is exponentially smaller than the characteristic distance in the monopole plasma, λ^3 , i.e. molecules are very small-sized with respect to that distance.

² In this section, 3-vectors are denoted as \mathbf{a} , whereas 2-vectors are denoted as \mathbf{a} .

52 D. Antonov and M. C. Diamantini

The factor at the action of the dimensionally-reduced theory, $S_{d: r} := \int d^2x L[\dots]$, can be removed [and this action can be cast to the original form of Eq. (2.2) with the substitution $d^3x \rightarrow d^2x$] by the obvious rescaling:

$$S_{d: r} := \int d^2x L_{new}(\dots) \quad (8.2)$$

Here, $K_{new} = \dots$, and in what follows we will denote for brevity K_{new} and \dots simply as K and \dots , respectively. Averaging then over the field with the use of the cumulant expansion we arrive at the following action:

$$S_{d: r}' = \int d^2x \left[\frac{1}{2} (\dots)^2 - 2 \cos \frac{\dots}{K} \right] \int d^2x d^2y \cos \frac{\dots}{K} (x) K^{(2)}(x-y) \cos \frac{\dots}{K} (y) \quad (8.3)$$

In this expression, similarly to Eq. (2.6), we have disregarded all the cumulants higher than the quadratic one, and the limits of applicability of this bilocal approximation will be discussed below. Further, in Eq. (8.3), $K^{(2)}(x) = e^{K D_{m_H}^{(2)}(x)} - 1$, where $D_{m_H}^{(2)}(x) = K_0(m_H, x, x) = 2$ is the 2D Yukawa propagator, and $\exp \frac{K}{2} D_{m_H}^{(2)} m_W^{-1}$ denotes the monopole fugacity modified by the interaction of monopoles via the Higgs field. The exponential factor entering reads

$$\frac{4}{g^2} m_W + T \ln \frac{e}{2} c \quad :$$

Here, we have introduced the notation $c = m_H / m_W$, $c < 1$, and $\gamma = 0.577$ is the Euler constant, so that $\frac{e}{2} \gamma = 0.89 < 1$. We see that the modified fugacity remains exponentially small, provided that

$$T < \frac{m_W}{\ln \frac{e}{2} c} \quad (8.4)$$

This constraint should be updated by another one, which would provide the convergence of the cumulant expansion. Analogously to the zero-temperature case, the divergence of the cumulant expansion would indicate that the Higgs vacuum loses its normal stochastic property and becomes a coherent one. In order to get the new constraint, notice that the parameter of the cumulant expansion reads $I^{(2)}$, where $I^{(2)} = \int d^2x K^{(2)}(x)$. Evaluation of the integral $I^{(2)}$ yields [53]:

$$I^{(2)} = \frac{2}{m_H^2} \left(\frac{1}{2} c^2 + 1 + \frac{2}{e} \frac{1 - \frac{c^2}{g^2}}{1 - \frac{4T}{g^2}} \right) A + e^{a=ae} \left(1 + \frac{a}{e} \right)^5 : (8.5)$$

(Note that at $T = g^2/4$, $\frac{1 - \frac{c^2}{g^2}}{1 - \frac{4T}{g^2}} = 2 \ln c$, i.e. $I^{(2)}$ remains finite.)

In the derivation of this expression, the parameter $a = \frac{p}{2} \frac{1}{T=g^2}$ was assumed to be of the order of unity. That is because the temperatures we are working at are of the order of T_{BKT} . Due to the exponential term in Eq. (8.5), the violation of the cumulant expansion may occur at high enough temperatures [that parallels the above-obtained constraint (8.4)]. The most essential, exponential, part of the parameter of the cumulant expansion thus reads

$$I^{(2)} / \exp \left(\frac{4}{g^2} m_W + T \ln \frac{e}{2} c \right) = \frac{p}{e} : (##)$$

Therefore, the cumulant expansion converges at the temperatures obeying the inequality

$$T < \frac{m_W}{\frac{p}{e} \ln \frac{e}{2} c} ;$$

which strengthens the inequality (8.4). On the other hand, since we are working in the plasma phase, i.e. $T > T_{BKT}$, it is enough to impose the following upper bound on the parameter of the weak-coupling approximation,

$$\frac{p}{e} > \frac{2}{\ln \frac{e}{2} c} :$$

Note that, although this inequality is satisfied automatically at $\frac{e}{2} c = 1$ (or $c = 1$), since it then takes the form $\frac{p}{2e}$, this is not so automatic in the BPS limit, $c = 1$. Indeed, in this case, we have $\ln \frac{2}{e} = 2$. It is, however, quite feasible to obey this inequality, since the logarithm is a weak function.

8.2. Higgs-inspired corrections to the RG flow in the absence of W-bosons

Although we have seen in the previous subsection that the propagating Higgs field does not change the value of the critical temperature T_{BKT} , it is instructive to derive corrections it produces to the RG flow. Such a derivation can

54 D. Antonov and M. C. Diamantini

also be extended to the $SU(N)$ -case (2.5). In that case [83], it is qualitatively clear that T_{BKT} should remain the same, since it only differs from that of the $SU(2)$ -case in the factor ϵ_1^2 , which is equal to unity. However, some peculiarities of the $SU(N)$ -case at $N > 2$ become clear only upon a derivation of the RG equations, which will be quoted below.

Let us thus start with the $SU(2)$ -case (2.2), whose action after the dimensional reduction has the form (8.2). In what follows, we will use the usual RG strategy based on the integration over the high-frequency modes. Note that this procedure will be applied to all the fields, i.e. not only to ϕ , but also to ψ . Splitting the momenta into two ranges, $0 < p < \Lambda$ and $\Lambda < p < \infty$, one can define the high-frequency modes as $h = \int_{\Lambda < p < \infty} \frac{d^2 p}{(2\pi)^2} e^{ip \cdot x} O(p)$ and consequently, e.g.

$$h(x) = \int_{\Lambda < p < \infty} \frac{d^2 p}{(2\pi)^2} e^{ip \cdot x} O(p). \text{ The partition function,}$$

$$Z = \int_{0 < p < \infty} D(p) D(\psi) \exp \int d^2 x [\dots]$$

can be rewritten as follows:

$$Z = \int_{0 < p < \infty} D(p) D(\psi) \exp \frac{1}{2} \int d^2 x [\dots]$$

where

$$Z^0 = \int_{0 < p < \infty} D(p) D(\psi) \exp \int d^2 x [\dots]$$

and \dots . Owing to the exponential smallness of the fugacity, Z^0 can further be expanded as

$$Z^0 = 1 + 2 \int d^2 x e^{p \cdot (x)} \cos \frac{p \cdot (x) + h(x)}{K} + 2 \int d^2 x d^2 y \dots$$

where

$$\begin{aligned}
 \langle i_h \rangle &= \frac{\int_{0 < p < \infty} \int_{\mathbb{R}^D} \langle p \rangle \exp \left[\frac{1}{2} \int_{\mathbb{R}^D} d^2x h \langle \partial^2 h \rangle \right]}{\int_{0 < p < \infty} \int_{\mathbb{R}^D} \langle p \rangle \exp \left[\frac{1}{2} \int_{\mathbb{R}^D} d^2x h \langle \partial^2 h \rangle \right]} ; \\
 \langle i \rangle &= \frac{\int_{0 < p < \infty} \int_{\mathbb{R}^D} \langle p \rangle \exp \left[\frac{1}{2} \int_{\mathbb{R}^D} d^2x \langle \partial^2 \rangle - m_H^2 \right]}{\int_{0 < p < \infty} \int_{\mathbb{R}^D} \langle p \rangle \exp \left[\frac{1}{2} \int_{\mathbb{R}^D} d^2x \langle \partial^2 \rangle - m_H^2 \right]} :
 \end{aligned}$$

Carrying out the averages we arrive at the following expression for Z^0 :

$$\begin{aligned}
 Z^0 &= \int_{\mathbb{R}^D} d^2x e^{i p \cdot x} \cos \left[\frac{p \cdot x}{K} \right] + \left(\int_{\mathbb{R}^D} d^2x d^2y \right. \\
 &\quad \left. e^{i p \cdot (x+y)} \int_{k=1}^X A^{2k}(x-y) B^2(x-y) \right) \\
 &\quad \cos \left[\frac{p \cdot (x+k \cdot y)}{K} \right] ; \tag{8.6}
 \end{aligned}$$

where $A(x) = \int_{\mathbb{R}^D} d^2p \frac{e^{ipx}}{(2\pi)^2 p^2}$, $B(x) = \int_{\mathbb{R}^D} d^2p \frac{e^{ipx}}{(2\pi)^2 p^2 + m_H^2}$,

$$G_h(x) = \int_{0 < p < \infty} \frac{d^2p}{(2\pi)^2} \frac{e^{ipx}}{p^2} ; \quad G(x) = \int_{0 < p < \infty} \frac{d^2p}{(2\pi)^2} \frac{e^{ipx}}{p^2 + m_H^2} ;$$

Since in what follows we will take $\infty = d$, the factors $A^{2k}(x-y)B^2(x-y)$, $k = 1$, are small. Owing to this fact, it is convenient to introduce the coordinates $r = x-y$ and $R = \frac{1}{2}(x+y)$, and Taylor expand Eq. (8.6) in powers of r . Clearly, this expansion should be performed up to the induced-interaction term $\int_{\mathbb{R}^D} d^2R e^{i p \cdot R} \cos \left[\frac{p \cdot R}{K} \right]$, that itself should already be disregarded. As a result, we obtain the following expression for Z (For the sake of uniformity, we replace d^2R by d^2x):

$$\begin{aligned}
 Z &= \int_{0 < p < \infty} \int_{\mathbb{R}^D} \langle p \rangle \int_{\mathbb{R}^D} d^2x \exp \left[\frac{1}{2} \int_{\mathbb{R}^D} d^2x \langle \partial^2 \rangle - m_H^2 \right] + \\
 &+ a_2 \left(\int_{\mathbb{R}^D} d^2x d^2y \right) e^{i p \cdot (x+y)} \int_{k=1}^X A^{2k}(x-y) B^2(x-y) \\
 &\quad \cos \left[\frac{p \cdot (x+k \cdot y)}{K} \right] ; \tag{8.7}
 \end{aligned}$$

56 D. Antonov and M. C. Diamantini

where we have introduced the notations

$$a_1 = \int d^2r r^2 A^{-2}(r) B^2(r) \quad ; \quad a_2 = \int d^2r A^{-2}(r) B^2(r) \quad ; \quad (8.8)$$

Taking into account that $\phi = d$ it is straightforward to get

$$a_1 = \frac{1}{2} K \frac{d}{m_H^2} \left(1 + \frac{d^2}{m_H^2} \right) \quad ; \quad a_2 = \frac{1}{2} K \frac{d}{m_H^2} \left(1 + \frac{d^2}{m_H^2} \right) \quad ;$$

Here, $\alpha_{1,2}$ stand for some momentum-space-slicing dependent positive constants, whose concrete values will turn out to be unimportant for the final expressions describing the RG flow.

Next, since $a_{1,2}$ occur to be infinitesimal (because these are proportional to d), the terms containing these constants on the r.h.s. of Eq. (8.7) can be treated in the leading-order approximation of the cumulant expansion that we will apply for the average over ϕ . In fact, we have

$$\begin{aligned} & \int_{0 < p < \phi} d^2p \exp \left[\frac{1}{2} \int d^2x \phi^2 \left(\frac{p^2}{m_H^2} \right) \right] \\ & \exp \left[b \int d^2x e^{2i\mathbf{p} \cdot \mathbf{x}} f(\mathbf{x}) \right] \cdot \exp \left[b e^{2K G(0)} \int d^2x f(\mathbf{x}) \right] + \\ & + \frac{b^2}{2} e^{4K G(0)} \int d^2x d^2y e^{4K G(\mathbf{x} - \mathbf{y})} f(\mathbf{x}) f(\mathbf{y}) \quad ; \end{aligned}$$

where f is equal either to unity or to $(\phi - \phi_0)^2$, $b = \alpha_{1,2} (A(0)B(0))^2$, and

$$G(\mathbf{x}) = \int_{0 < p < \phi} \frac{d^2p}{(2\pi)^2} \frac{e^{i\mathbf{p} \cdot \mathbf{x}}}{p^2 + m_H^2} \quad ; \quad G(0) = \frac{1}{4} \ln \left(1 + \frac{\phi^2}{m_H^2} \right) \quad ;$$

In order to estimate the parameter of the cumulant expansion, $b e^{2K G(0)} \int d^2x e^{4K G(\mathbf{x})} \approx 1$, note that we are working in the phase where monopoles form the plasma, i.e. below T_{BKT} . Because of this fact, $4K |G(\mathbf{x})| \approx 32 \left(\frac{\phi_0}{m_H} \right)^2$, that, due to the factor $\left(\frac{\phi_0}{m_H} \right)^2$, is generally much smaller than unity. Owing to that, we get

$$b \approx 1 + \frac{\phi^2}{m_H^2} \int d^2x G(\mathbf{x}) \approx \frac{2bK}{m_H^2} \left(1 + \frac{K}{2} \frac{\phi^2}{m_H^2} \right) \quad ;$$

Choosing for concreteness $b = a_2 (A(0)B(0))^2$ and taking into account that

$$A(0) \approx 1 - \frac{K}{2} G_h(0) = 1 - \frac{K}{4} \frac{d}{m_H^2} \quad ; \quad (8.9)$$

$$B(0)' = 1 + \frac{K}{2}G(0)' = 1 + \frac{K}{4} \frac{d}{m_H^2}; \quad (8.10)$$

we obtain to the leading order: $\sim 2a_2K^{-2} = (m_H^2)^{-1}$. This quantity possesses the double smallness (firstly, because a_2 is infinitesimal and, secondly, due to the exponential smallness of \dots).

Such an extremely rapid convergence of the cumulant expansion enables one to replace $e^{2\frac{K}{C^2}}$ in the terms proportional to $a_{1,2}$ on the r.h.s. of Eq. (8.7) by the average value of this exponent equal to $1 + \frac{\langle \dots \rangle}{m_H^2}$. Comparing the so-obtained expression with the initial one, we arrive at the following renormalizations of fields and parameters of the Lagrangian:

$$\begin{aligned} C_0^{new} &= C_0; & C_1^{new} &= C_1; & K^{new} &= \frac{K}{C^2}; & m_H^{new} &= \frac{m_H}{C^2}; & A(0)B(0) &= A(0)B(0); \end{aligned} \quad (8.11)$$

where m_H^2 ,

$$\begin{aligned} C_1 &= 4 \left[1 + \frac{K a_1}{2} (A(0)B(0))^2 \right] \left[1 + \frac{\langle \dots \rangle}{m_H^2} \right]^{1-2}, \\ &= 1 + \frac{K a_1}{2} (A(0)B(0))^2 \left[1 + \frac{2K}{m_H^2} \right]^{1-2}; \end{aligned} \quad (8.12)$$

Besides that, we obtain the following shift of the free-energy density $F = \frac{\ln Z^0}{V}$:

$$\begin{aligned} F &= F^{new} - a_2 (A(0)B(0))^2 \left[1 + \frac{\langle \dots \rangle}{m_H^2} \right]^{1-2}, \\ &= F^{new} - a_2 (A(0)B(0))^2 \left[1 + \frac{K}{2} \frac{\langle \dots \rangle}{m_H^2} \right]; \end{aligned} \quad (8.13)$$

where V is the 2D volume (i.e. area) of the system.

By making use of the relations (8.9), (8.10), it is further straightforward to derive from Eqs. (8.11)–(8.13) the RG equations in the differential form. Those read

$$\begin{aligned} d &= \frac{K}{4} \left[1 - \frac{d}{m_H^2} \right]; & dK &= \frac{1}{2}K^3 \left[1 + \frac{K}{2} + 1 - \frac{d}{5} \right]; \\ d &= \frac{1}{2} (K)^2 \left[1 + \frac{K}{2} + 1 - \frac{d}{5} \right]; \end{aligned}$$

58 D. Antonov and M. C. Diamantini

$$dF = \frac{1}{2}K^2 \left(1 + \frac{K}{2} + 1 \right) \frac{d}{a^3} :$$

Let us further make the change of the momentum scale to the real-space one: $da = a^2 da$, that modifies the above equations as

$$d = \frac{K}{4} \frac{da}{a} \left(1 + \frac{1}{a^2} \right) ; \tag{8.14}$$

$$dK = \frac{1}{2}K^3 \frac{da}{a^3} \left(1 + \frac{K}{2} + 1 \right) \frac{1}{a^2} ; \tag{8.15}$$

$$d = \frac{1}{2} (K - y)^2 \frac{da}{a^3} \left(1 + \frac{K}{2} + 1 \right) \frac{1}{a^2} ; \tag{8.16}$$

$$dF = \frac{1}{2}K^2 \frac{da}{a} \left(1 + \frac{K}{2} + 1 \right) \frac{1}{a^2} ; \tag{8.17}$$

Our main aim below is to derive from Eqs. (8.14)-(8.16) the leading-order corrections in $(a^2)^{-1}$ to the BKT RG flow in the vicinity of the critical point, $K_c^{(0)} = 8$ (that clearly corresponds to T_{BKT}), $y_c^{(0)} = 0$, where $y = \frac{2}{a}$, and the superscript $^{(0)}$ denotes the zeroth order in the $(a^2)^{-1}$ expansion. These values of $K_c^{(0)}$ and $y_c^{(0)}$ will be derived below. Besides that, it will be demonstrated that, in the critical region, β is evolving very slowly. Owing to this fact, the initial assumption on the largeness of β (namely, that it is of the order of m_w^2), will be preserved by the RG flow, at least in that region. This enables one to consider β as almost a constant and seek corrections to the RG flow of $K^{(0)}$ in powers of $(a^2)^{-1}$. The zeroth-order equation stemming from Eq. (8.15) then reads

$$dK^{(0)} = \frac{1}{2}K^{(0)3} \frac{da}{a^3} ; \tag{8.18}$$

Respectively, the zeroth-order in $(a^2)^{-1}$ equation for y has the form

$$dy^{(0)2} = 2 \frac{da}{a} y^{(0)2} x ; \tag{8.19}$$

where $x = 2 \frac{K^{(0)}}{4}$. Equations (8.18) and (8.19) yield the above-mentioned leading critical value of K , $K_c^{(0)}$. Next, with this value of $K_c^{(0)}$, Eq. (8.18) can be rewritten in the vicinity of the critical point as

$$dx = (8 - K)^2 \frac{da}{a^3} ; \tag{8.20}$$

Introducing further instead of y a new variable $z = (8 - K)^2 y^2$ and performing the rescaling $a^{new} = a \frac{8 - K}{2}$ we get from Eqs. (8.17), (8.19),

and (8.20) the following system of equations:

$$dz^{(0)} = 2 \frac{da}{a} x z^{(0)}; dx = z^{(0)} \frac{da}{a}; dF^{(0)} = z^{(0)} \frac{da}{a^3}; \quad (8.21)$$

These equations yield the standard RG flow in the vicinity of the critical point, $x = z^{(0)} = 0$, which has the form [82, 84] $z^{(0)} \propto x^2 = \frac{1}{P} \frac{1}{z}$, where $\frac{1}{P} = (\mathbb{T}_{BKT} - T) = \mathbb{T}_{BKT}$ is some constant. In particular, $x \propto \frac{1}{z}$ at $T = \mathbb{T}_{BKT}$.

Owing to the first of Eqs. (8.21), this relation yields $z_{in}^{(0) \frac{1}{2}} = \ln(a = a_{in})$, where the subscript "in" means the initial value. Taking into account that $z_{in}^{(0)}$ is exponentially small, while $z^{(0)} = 1$ (the value at which the growth of $z^{(0)}$ stops), we obtain in the case $x_{in} = \frac{1}{P}$: $\ln(a = a_{in}) = z_{in}^{(0) \frac{1}{2}} = 1 = 2$. According to this relation, at $T = \mathbb{T}_{BKT} = 0$, the correlation length diverges with an essential singularity as $a \rightarrow 0$ ($\exp(\text{const} \frac{1}{P})$). (At $T < \mathbb{T}_{BKT}$, $a \rightarrow 0$, while at $T > \mathbb{T}_{BKT}$, the correlation length becomes finite due to the short-rangeness of molecular fields.) As far as the leading part of the free-energy density is concerned, it scales as $F^{(0)} \propto a^{-2}$ and therefore remains continuous in the critical region. Moreover, the correction to this behavior stemming from the finiteness of the Higgs-boson mass [the last term on the r.h.s. of Eq. (8.17)] is clearly of the same functional form, $\exp(\text{const} \frac{1}{P})$, i.e. it is also continuous.

We are now in the position to address the leading-order [$\ln(a^{-2})^{-1}$] corrections to the above-discussed BKT RG flow of $K^{(0)}$ and $z^{(0)}$. To this end, let us represent K and z as $K = K^{(0)} + K^{(1)} = (a^{-2})$, $z = z^{(0)} + z^{(1)} = (a^{-2})$ that, by virtue of Eqs. (8.14) and (8.15), leads to the following novel equations:

$$dK^{(1)} = 2K^{(1)} \frac{da}{a} = 4 \frac{da}{a} z^{(0)} + \frac{z^{(1)}}{a^2} \left(1 + \frac{K^{(0)}}{2} + \frac{3K^{(1)}}{K^{(0)}} \right); \quad (8.22)$$

$$dz^{(1)} = 2z^{(1)} \frac{da}{a} = 2 \frac{da}{a} z^{(1)} \frac{K^{(0)}}{4} + \frac{z^{(0)}}{4} K^{(1)} - K^{(0)}; \quad (8.23)$$

In the vicinity of the critical point, we can insert into Eq. (8.23) the above-obtained critical values of $K^{(0)}$ and $z^{(0)}$, that yields:

$$dz^{(1)} = 2z^{(1)} \frac{da}{a}; \quad (8.24)$$

Therefore, $z^{(1)} = C_1 a^2$, where C_1 is the integration constant of dimensionality (mass)², $C_1 = \dots$. Inserting further this solution into Eq. (8.22), consid-

60 D. Antonov and M. C. Diamantini

ered in the vicinity of the critical point, one obtains the following equation:

$$dK^{(1)} - 2K^{(1)} \frac{da}{a} = \frac{4 C_1 da}{a} - \frac{3K^{(1)}}{8} + 5 : \quad (8.25)$$

Its integration is straightforward and yields

$$K^{(1)} = C_2 a^{2-1 \frac{3C_1}{4}} + \frac{40 C_1}{4 - 3C_1} ; \quad (8.26)$$

where the dimensionless integration constant C_2 should be much smaller than $(a^2)^{\frac{3C_1}{4}}$. [Note that the last addendum in Eq. (8.26) is positive.] Therefore, the total correction, $K^{(1)} = (a^2)$, approximately scales with a in the critical region as $\frac{40 C_1}{(4 - 3C_1) a^2}$. We see that, at the critical point, this expression vanishes due to the divergence of the correlation length. Note also that Eq. (8.26) can obviously be rewritten as the following dependence of $K^{(1)} = (a^2)$ on $z^{(1)} = (z^{(1)} a^2)$:

$$\frac{K^{(1)}}{a^2} = C_2 a^{2 - \frac{3z^{(1)}}{4 a^2}} + \frac{40 z^{(1)}}{4 a^2 - 3z^{(1)} a^2} :$$

With the above-discussed critical behavior of the correlation length, $a(\epsilon)$, this relation determines the correction to the BKT RG flow, $z^{(0)}$

$$2 \frac{K^{(0)}}{4} = \dots$$

Finally, in order to justify the above-adopted approximation, within which ϵ was considered as a constant, we should check that, under the RG flow, ϵ indeed evolves slowly. To this end, let us pass in Eq. (8.16), considered in the critical region, from the variable ϵ to the above-introduced variable z and perform again the rescaling $a \rightarrow a^{new}$. This yields $\frac{d\epsilon}{\epsilon} = \frac{z da}{2 a}$ or $d\epsilon = \frac{C_1 da}{2 a}$. Since $C_1 \ll 1$, we conclude that $\frac{d\epsilon}{\epsilon} \ll \frac{da}{a}$. The obtained inequality means that, in the vicinity of the BKT critical point, ϵ is really evolving slowly. This fact justifies our treatment of ϵ as a large (with respect to ϵ^2) constant quantity, approximately equal to its initial value, of the order of m_W^2 .

Let us now proceed with a similar RG analysis of the $SU(N)$ -version of the theory (2.2) given by Eq. (2.5). Applying to the respective dimensionally-reduced theory the above-described RG procedure, we arrive at the following analogue of Eq. (8.7):

$$Z = \int_{0 < p < \infty} D \sim (p) D(p) \exp \int_{\mathbb{R}^d} d^2 x \left[\frac{1}{2} \phi^T \partial^2 \phi - m_H^2 \phi^T \phi + \dots \right]$$

$$\begin{aligned}
 & + (A(0)B(0))^2 e^{2\frac{p}{K} \int_0^X \cos \frac{p}{K} \varphi_i \sim 0} a_2^{ij} \cos \frac{h_p}{K} (\varphi_i - \varphi_j) \sim 0 + 2 A(0)B(0) \\
 & e^{\frac{p}{K} \int_0^X \cos \frac{p}{K} \varphi_i \sim 0} \frac{1}{2} \sum_{ab} + (A(0)B(0))^2 \frac{K}{8} e^{2\frac{p}{K} \int_0^X \cos \frac{p}{K} \varphi_i \sim 0} \\
 & \sum_{ij} a_1^{ij} (\varphi_i + \varphi_j) (\varphi_i + \varphi_j) \cos \frac{h_p}{K} (\varphi_i - \varphi_j) \sim 0 \quad \text{; ; ;} : \\
 & \hspace{15em} (8.27)
 \end{aligned}$$

Here, $i, j = 1; \dots; (N - 1)$, and we have introduced the notations similar to (8.8),

$$a_1^{ij} = \int d^2 r r^2 B^2(r) e^{K \varphi_i \varphi_j G_h(r)} \quad ; \quad a_2^{ij} = \int d^2 r B^2(r) e^{K \varphi_i \varphi_j G_h(r)} \quad ;$$

so that, at $\theta = 0$, $d = d$,

$$a_1^{ij} = \frac{1}{K} \frac{d}{5} \varphi_i \varphi_j + \frac{2}{m_H^2} \quad ; \quad a_2^{ij} = \frac{2}{K} \frac{d}{3} \varphi_i \varphi_j + \frac{2}{m_H^2} \quad ;$$

The main difference of Eq. (8.27) from Eq. (8.7) is due to the terms containing $\cos \frac{h_p}{K} (\varphi_i - \varphi_j) \sim 0$, which violate the RG invariance. Nevertheless, approximately this invariance does hold, since the respective sums are dominated by the terms with $i = j$. Working within this approximation and making use of the identity $\sum_i \varphi_i^2 = \frac{N}{2}$, we obtain

$$\begin{aligned}
 & \int d^2 x \exp \left[-a_2 \frac{N(N-1)}{2} (A(0)B(0))^2 \left(1 + \frac{\theta^2}{m_H^2} \right) \right] \\
 & \int d^2 x \exp \left[-\frac{1}{2} \int_0^X \cos \frac{p}{K} \varphi_i \sim 0 + 2 A(0)B(0) e^{\frac{p}{K} \int_0^X \cos \frac{p}{K} \varphi_i \sim 0} \right] \\
 & \frac{1}{2} \left(1 + \frac{\theta^2}{m_H^2} \right) \frac{N K a_1}{4} (A(0)B(0))^2 (\langle \varphi_i \sim 0 \rangle)^2 \quad ; \quad (8.28)
 \end{aligned}$$

This expression has again been derived in the leading order of the cumulant expansion applied in course of the average over φ_i .

The shift of the free-energy density stemming from Eq. (8.28) [cf. Eq. (8.13)] reads

$$F^{new} = F + \frac{N(N-1)}{2} (A(0)B(0))^2 \left(1 + \frac{K}{2} \frac{\theta^2}{m_H^2} \right) \quad ; \quad (8.29)$$

62 D. Antonov and M. C. Diamantini

As far as the renormalization of fields and coupling constants is concerned, it is given by Eq. (8.11), where the first equation should be modified as $\tilde{g}_0^{\text{new}} = C \tilde{g}_0$, and the parameter C from Eq. (8.12) now reads

$$C = 4 \left[1 + \frac{K N a_1}{4} (A(0)B(0))^2 \right]^{-1} \left[1 + \frac{g^2}{m_H^2} \left(\frac{N}{2} \right)^{3/2} \right]^{-1},$$

$$\tilde{g}_0^{\text{new}} = \tilde{g}_0 \left[1 + \frac{K N a_1}{4} (A(0)B(0))^2 \right]^{-1} \left[1 + \frac{g^2}{m_H^2} \left(\frac{N}{2} \right)^{3/2} \right]^{-1} : \quad (8.30)$$

From Eqs. (8.29) and (8.30) we deduce that, in the $SU(N)$ -case at $N > 2$, the RG flow of couplings and of the free-energy density is identical to that of the $SU(2)$ -case. Indeed, all the N -dependence can be absorbed into the constants $a_{1,2}$ by rescaling them as $a_1 = N^{-1/2}$, $a_2 = N(N-1)^{-1/2}$ and further redefining [cf. the notations introduced after Eq. (8.20)] $z = (8/3)^{1/2} a_1^2$ and $a^{\text{new}} = a_2^2 a_1^{-1/2}$. In particular, the critical temperature T_{BKT} remains the same as in the $SU(2)$ -case. (As it has already been mentioned, this follows also from the estimate of the mean squared separation in the monopole-antimonopole molecule, if one takes into account that the square of any root vector is equal to unity.) Thus, the principal difference of the $SU(N)$ -case, $N > 2$, from the $SU(2)$ -one is that, while in the $SU(2)$ -case the RG invariance is exact (modulo the negligibly small higher-order terms of the cumulant expansion applied to the average over \mathcal{U}), in the $SU(N)$ -case it is only approximate, even in the limit $m_H \rightarrow 1$.

8.3. Finite-temperature 3D compact QED with massless fundamental fermions

Let us consider the extension of the model (2.1) by fundamental dynamical quarks [55] (for the sake of simplicity, we omit the summation over the flavor indices, although consider the general case of an arbitrary number of flavors):

$$S = \int_{\mathcal{R}} d^3x \left[\bar{\psi} \not{D} \psi + h \frac{a}{2} \bar{\psi} \psi \right].$$

In this formula, the Yukawa coupling h has the dimensionality $(\text{mass})^{1-2}$, $D = \partial + ig \frac{a}{2} A^a$, $\gamma = \gamma^y$, where the Euclidean Dirac matrices are defined as $\gamma = \gamma^i$ with $\gamma^0 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$,

$\gamma^i = \begin{pmatrix} 0 & \sigma^i \\ \sigma^i & 0 \end{pmatrix}$, and σ^i denote the Pauli matrices. As it will be demonstrated, at $T > T_{\text{BKT}}$, quark zero modes in the monopole field lead to the additional attraction of a monopole and an antimonopole (M and \bar{M} for shortness) in the molecule. In particular, when the number of these modes (equal to

the number of massless flavors) is sufficiently large, the molecule shrinks so strongly that its size becomes of the order of the inverse W -boson mass. Another factor which defines the size of the molecule is the characteristic range of localization of zero modes. Namely, it can be shown that the stronger zero modes are localized in the vicinity of the monopole center, the smaller molecular size is. We will consider the case when the Yukawa coupling h of quarks with the Higgs field vanishes, and originally massless quarks do not acquire any mass. This means that zero modes are maximally delocalized. Such a weakness of the quark-mediated interaction of monopoles opens a possibility for molecules to undergo eventually the phase transition into the plasma phase at the temperatures of the order of T_{BKT} . However, this will be shown to occur only provided that the number of flavors is equal to one, whereas at any larger number of flavors, the respective critical temperature becomes exponentially small. This means that the interaction mediated by such a number of zero modes is already strong enough to maintain the molecular phase at any temperature larger than that one.

For a short while, let us consider the general case $h \neq 0$. One can then see that the Dirac equation in the field of the third isotopic component of the 't Hooft-Polyakov monopole [27] decomposes into two equations for the components of the $SU(2)$ -doublet ψ . The masses of these components stemming from such equations are equal to each other and read $m_q = h = 2$. Next, the Dirac equation in the full monopole potential has been shown [85] to possess the zero mode, whose asymptotic behavior at $r \rightarrow \infty$ $\psi \sim m_q^{-1}$ has the following form:

$$\psi_n^+ = N \frac{e^{-m_q r}}{r} s^+ s_n \quad s^- s_n^+ ; \quad \psi_n = 0: \quad (8.31)$$

Here, ψ_n are the upper and the lower components of the mode, i.e. $\psi = \begin{pmatrix} \psi_n^+ \\ \psi_n^- \end{pmatrix}$. Next, $n = 1; 2$ is the isotopic index, $\mu = 1; 2$ is the Dirac index, $s^+ = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$, $s^- = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$, and N is the normalization constant.

It is a well known fact that, in 3D, the 't Hooft-Polyakov monopole is actually an instanton [3,26]. Therefore, we can use the results of Ref. [86] on the quark contribution to the effective action of the instanton-antiinstanton molecule in QCD. Let us thus recapitulate the analysis of Ref. [86] adopting it to our model. To this end, we fix the gauge $A^a = \delta^{a3}$ and define the analogue of the free propagator S_0 by the relation $S_0^{-1} = -i \not{\partial} + m_q \gamma^3$. Next, we define the propagator S_M in the field of a monopole located at the origin, $A^{aM} = [A_i^{aM} \delta^{ij} - \epsilon^{aij} x^j] = gr^2$ at $r = m_W^{-1}$, by the formula $S_M^{-1} = S_0^{-1} - g \frac{a}{2} A^{aM}$. Obviously, the propagator S_M in the field of an antimonopole

64 D. Antonov and M. C. Diamantini

located at a certain point R , $A^{aM}(\mathbf{x}) = A^{aM}(\mathbf{x} - R)$, is defined by the equation for S_M^{-1} with the replacement $A^{aM} \rightarrow A^{aM} + A^a$. Finally, one can consider the molecule made out of these monopole and antimonopole and define the total propagator S in the field of such a molecule, $A^a = A^{aM} + A^{aM}$, by means of the equation for S_M^{-1} with A^{aM} replaced by A^a .

One can further introduce the notation j_n , $n = 0; 1; 2; \dots$, for the eigenfunctions of the operator $-iD$ defined at the field of the molecule, namely $-iD j_n = \lambda_n j_n$, where $\lambda_0 = 0$. This yields the following form of spectral representation for the total propagator S :

$$S(\mathbf{x}; \mathbf{y}) = \sum_{n=0}^{\infty} \frac{j_n(\mathbf{x}) i h_n(\mathbf{y})}{i m_q^3}.$$

Next, it is worth recalling the mean-field approximation, discussed in section 3, within which zero modes dominate in the quark propagator, i.e.

$$S(\mathbf{x}; \mathbf{y}) \approx \frac{j_0(\mathbf{x}) i h_0(\mathbf{y})}{i m_q^3} + S_0(\mathbf{x}; \mathbf{y}). \quad (8.32)$$

The approximation (8.32) remains valid for the molecular phase near the phase transition (i.e. when the temperature approaches $T_{BK T}$ from above), we will be interested in. That is merely due to the fact that, in this regime, molecules become very much inflated being about to dissociate.

Within the notations adopted, one now has $S = S_M^{-1} + S_M^{-1} S_0^{-1} S_M^{-1} = S_M^{-1} S^{-1} S_M$, where

$$S = S_0^{-1} (S_M^{-1} S_0^{-1} S_M^{-1} S_0^{-1}) = S_0^{-1} \frac{M_0}{i m_q^3} S_0^{-1} \frac{M_0}{i m_q^3};$$

and M_0 , M_0 are the zero modes of the operator $-iD$ defined at the field of a monopole and an antimonopole, respectively. Denoting further $a = \frac{M_0}{0} \sim \frac{a}{2} A^{aM} \frac{M_0}{0}$, it is straightforward to see, by the definition of the zero mode, that $a = \frac{M_0}{0} i \not{\partial} \frac{M_0}{0} = \frac{M_0}{0} S_0^{-1} \frac{M_0}{0}$. This yields $S = S_0 + (a \not{m}_q^2) \frac{M_0}{0} \frac{M_0}{0}$, where the star stands for the complex conjugation, and therefore $\det S = 1 + (\not{a} \not{m}_q)^2 \det S_0$. Finally, defining the desired effective action as $\Gamma = \ln[\det S^{-1} = \det S_0^{-1}]$, we obtain for it, in the general case with N_f flavors, the following expression: $\Gamma = \text{const} + N_f \ln m_q^2 + \not{a} \not{a}$. The constant in this formula, standing for the sum of effective actions defined at the monopole and at the antimonopole, cancels out in the normalized expression for the mean squared separation in the $M M$ molecule.

Let us further set h equal to zero, and so m_q is equal to zero as well. Notice first of all that, although in this case the direct Yukawa interaction of the Higgs boson with quarks is absent, they keep interacting with each other via the gauge field. Owing to this fact, the problem of finding a quark zero mode in the monopole field is still legitimate. Note that, according to Eq. (8.31), this mode will be non-normalizable in the sense of a discrete spectrum. However, it is clear that, in the gapless case $m_q = 0$ under discussion, the zero mode, which lies exactly on the border of the two contiguous Dirac seas, should be treated not as an isolated state of a discrete spectrum, but rather as a state of a continuum spectrum. This means that it should be understood as follows: $j_0(\mathbf{x}) \sim \lim_{p \rightarrow 0} e^{i\mathbf{p}\cdot\mathbf{r}} = r$, where $\mathbf{p} = |\mathbf{p}|\hat{\mathbf{j}}$. Once being considered in this way, zero modes are normalizable by the standard condition of normalization of the radial parts of spherical waves, R_{p1} , which reads [92] $\int_0^R dr r^2 R_{p0} R_{p1} = 2 \int_0^R dr r^2 R_{p1}^2$. The dependence of the absolute value of the matrix element a on the distance R between a monopole and an antimonopole can now be readily found. Indeed, we have $\int_{\mathbb{R}^3} d^3r = r^2 \int_0^\infty R^{-1} dr \int_0^{2\pi} d\phi \int_0^\pi \sin\theta d\theta = 4 \ln(R)$, where Λ stands for the infrared cutoff.

We can now switch on the temperature and explore a possible modification of the BKT critical temperature, T_{BKT} , due to the zero-mode mediated interaction. As it has been discussed, this can be done upon the evaluation of the mean squared separation in the MM molecule and further finding the temperature below which it starts diverging. In this way we should take into account that, in the dimensionally-reduced theory, the usual Coulomb interaction of monopoles (without the loss of generality, we consider the molecule with the temporal component of \mathbf{R} equal to zero.)

$R^{-1} = \sum_{n=1}^{\infty} R^{-2 + (n)^2} \ln R$ goes over into $2T \ln(R)$, where R denotes the absolute value of the 2-vector \mathbf{R} . This statement can be checked e.g. by virtue of the Euler-MacLaurin formula. As far as the novel logarithmic interaction, $\ln(R) = \sum_{n=1}^{\infty} \ln \left(R^2 + (n)^2 \right)^{-1/2}$, is concerned, it transforms into

$$TR + \ln [1 - \exp(-2TR)] \ln 2 : \tag{8.33}$$

Let us prove this statement. To this end, we make use of the following formula [93]:

$$\sum_{n=1}^{\infty} \frac{1}{n^2 + x^2} = \frac{1}{2x} \coth(x) + \frac{1}{x} :$$

66 D. Antonov and M. C. Diamantini

This yields

$$\sum_{n=1}^{\infty} \frac{x^{2n-1}}{x^2 + (2n-a)^2} = \frac{1}{x} + \frac{xa^2}{2} \sum_{n=1}^{\infty} \frac{x^{2n-1}}{n^2 + (xa-2)^2} = \frac{a}{2} \coth \frac{ax}{2} :$$

On the other hand, the l.h.s. of this expression can be written as

$$\frac{1}{2} \frac{d}{dx} \sum_{n=1}^{\infty} \ln \left(x^2 + \frac{2n^2}{a} \right) :$$

Integrating over x , we get

$$\sum_{n=1}^{\infty} \ln \left(x^2 + \frac{2n^2}{a} \right) = a \int dx \coth \frac{ax}{2} = 2 \ln \sinh \frac{ax}{2} = ax + 2 \ln \left(1 - e^{-ax} \right) - 2 \ln 2 :$$

Setting $\frac{2}{a} = R$ and $x = R$ we arrive at Eq. (8.33).

Thus, the statistical weight of the quark-mediated MM -interaction in the molecule at high temperatures has the form $\exp(-2N_f \ln \frac{R}{a}) / [TR + \ln [1 - \exp(-2TR)]] \ln 2^{2N_f}$. Accounting for both, (former) logarithmic and Coulomb, interactions we eventually arrive at the following expression for the mean squared separation L^2 in the molecule as a function of T , g , and N_f :

$$L^2 = \frac{\int_0^R dR R^3 \frac{8T}{g^2} [TR + \ln [1 - \exp(-2TR)]] \ln 2^{2N_f}}{\int_0^R dR R^1 \frac{8T}{g^2} [TR + \ln [1 - \exp(-2TR)]] \ln 2^{2N_f}} : \quad (8.34)$$

At large R , $\ln 2 - TR$ and $\ln [1 - \exp(-2TR)] \sim \exp(-2TR)$

TR . Consequently, we see that L^2 is finite at $T > T_{BKT}^{N_f} = (2 - N_f)g^2/4$, that reproduces T_{BKT} at $N_f = 0$. For $N_f = 1$, the plasma phase is still present at $T < g^2/4$. Instead, when $N_f \geq 2$, at whatever temperatures, which are parametrically larger than the temperature of dimensional reduction, $O(1=3)$, the monopole ensemble exists only in the molecular phase. At such temperatures, massless dynamical quarks produce their own deconfinement. They also destroy the monopole-based confinement of any other fundamental matter. Clearly, the obtained temperature $T_{BKT}^{(N_f=1)}$ is exponentially larger than $1=3$ as T_{BKT} , that fully validates the idea of dimensional reduction. Note also that, at $N_f = \max\{1; 4 - T/g^2\}$, $\ln 2 \sim 1/m_w$, which means that such a large number of zero modes shrinks the molecule to the

minimal admissible size. In conclusion, notice that, in the presence of W -bosons, the influence of heavy fundamental scalar matter to the dynamics of the deconfining phase transition has been studied in Ref. [56].

8.4. 3D Georgi-Glashow model and its supersymmetric generalization at finite temperature

The crucial difference in the finite-temperature behavior of the 3D GG model [52] from that of 3D compact QED [51] stems from the presence in the former of the heavy charged W -bosons. Although being practically irrelevant at zero temperature due to their heaviness, at finite temperature these bosons, due to their thermal excitation, form a plasma whose density is compatible to that of monopoles. The density of the W -plasma can be evaluated as follows (see e.g. Ref. [94]):

$$\begin{aligned}
 n_W &= 6T \int \frac{d^3p}{(2\pi)^3} \ln \int_0^\infty e^{-\beta(\epsilon(p) - \mu)} \epsilon(p) d\epsilon(p) \\
 &= 6 \int \frac{d^3p}{(2\pi)^3} \frac{1}{e^{\beta(\epsilon(p) - \mu)} - 1} \\
 &= \frac{3m_W^2}{1} \int_0^\infty \frac{dz}{e^{m_W z} - 1}, \quad \frac{3m_W^2}{1} \int_0^\infty dz e^{-m_W z} = \frac{3m_W T}{1 + \frac{T}{m_W}} e^{-m_W} :
 \end{aligned}$$

Here, μ stands for the chemical potential, $\epsilon(p) = \sqrt{p^2 + m_W^2}$, and the factor $\int d\epsilon(p)$ represents the total number of spin states of W^+ - and W^- -bosons. We have also denoted $z = \epsilon(p) - m_W$ and taken into account that, since we want to stay in the plasma phase of monopoles and the temperatures under consideration should not exceed T_{BKT} , $T \ll m_W$. Therefore, up to small corrections, $n_W = \frac{3m_W T}{1 + \frac{T}{m_W}} e^{-m_W}$.

It has been found in [52], further elaborated in [53, 56], etc., and reviewed in [95] that, when the finite-temperature 3D compact QED is promoted, upon the incorporation of W -bosons, to the finite-temperature 3D GG model, the critical temperature T_{BKT} and the $U(1)$ universality class of the deconfining phase transition are changed by $T_c = \frac{g^2}{4}$ and the 2D-Ising universality class, respectively. In what follows, we will survey the main ideas of Ref. [52], proceeding further to the analysis of the SUSY 3D GG model at finite temperature [54].

The Lagrangian of 3D compact QED, being rewritten in terms of the vortex operator $V = e^{i\mathcal{G}} = 2$,

$$\mathcal{L}_{3D} = \frac{1}{2} (\partial_\mu \mathcal{G})^2 - 2 \cos(\mathcal{G}) = \frac{2}{g_m^2} \mathcal{J}^\mu V \mathcal{J}_\mu - V^2 + (V')^2 ;$$

explicitly has the magnetic Z_2 symmetry [50]. At zero temperature, this symmetry is spontaneously broken, since $\langle V(\mathbf{x})V(\mathbf{0}) \rangle \sim |\mathbf{x}|^{-1}$. It can be shown [50] that the breakdown of the magnetic symmetry implies confinement. Since this symmetry is inherited in the 3D compact QED from the initial 3D GG model, the deconfinement phase transition, occurring in the latter at a certain temperature (equal to T_c), should be associated with the restoration of this very symmetry. Already this fact alone indicates that the phase transition in the finite-temperature 3D GG model should be of the same kind as that of the (Z_2 -invariant) 2D Ising model, rather than the BKT phase transition of compact QED. It has quantitatively been shown in Ref. [52] that this is indeed the case if W -bosons are additionally included in the compact-QED Lagrangian. At finite temperature, this can be done by noting that W -bosons are nothing but vortices of the \mathcal{A} -eld and they can be incorporated into the dimensionally-reduced Lagrangian, $L_{2D} = \frac{1}{2}(\mathcal{A})^2 - 2 \cos(\mathcal{K})$, by adding to it the term $-2 \cos \sim$ (from now on, \mathcal{A} will denote the dual-photon \mathcal{A} -eld in the dimensionally-reduced theory). The \mathcal{A} -eld \sim , dual to the \mathcal{A} -eld \mathcal{A} , is defined through the relation $i\mathcal{A} \sim = g \mathcal{A}$. The fugacity of W -bosons, \mathcal{A} , is proportional to their density n_W , therefore $\mathcal{A} \sim n_W / m_W T e^{-m_W/T}$. Note also that an alternative way to introduce W -bosons into L_{2D} has been proposed in Ref. [96]. Within that approach, the above definition of \sim through \mathcal{A} is abolished. Instead, an extra interaction of these, now independent, \mathcal{A} -elds of the type $i(\mathcal{A}_{x_1}) (\mathcal{A}_{x_2} \sim)$ appears.]

Owing to the novel cosine term, even in the absence of the monopole plasma, the dual photon never becomes massless [52]. Its mass m_D rather increases, and the vacuum correlation length d decreases, with the increase of the temperature. Consequently, contrary to what we had in the case of the inverse BKT phase transition, where the correlation length was becoming infinite at $T > T_{BKT}$, now it never becomes infinite. This result parallels the general expectation that, with the increase of the temperature in any local \mathcal{A} -eld theory, the degree of disorder becomes higher, and the correlation length decreases. Indeed, in the absence of monopoles (i.e. at $\mathcal{A} = 0$),

$$L_{2D} = \frac{1}{2}(\mathcal{A})^2 - 2 \cos g \mathcal{A} \quad ; \quad (8.35)$$

where $\mathcal{A} \sim = g \mathcal{A}$. Therefore, $m_D^2(\mathcal{A}) = 2 g^2 / m_W g^2 e^{-m_W/T}$. We see that m_D grows with the decrease of \mathcal{A} , i.e. with the increase of T , that proves the above statement.

Evaluating further the mean squared separation in the W^+W^- molecule

in the same way as we have done it earlier for the MM molecule [cf. Eqs. (8.1) and (8.34)], we have $L^2 / \int d^2R R^2 \frac{g^2}{2T}$. The convergence of this integral at $T < T_{XY} = \frac{g^2}{8}$ means that, in the absence of monopoles, with the increase of temperature W -bosons pass from the molecular phase into the plasma one at $T = T_{XY}$. In the same case of absence of dynamical monopoles, a static MM -pair becomes linearly confined at $T > T_{XY}$. Indeed, the MM -potential, V , is related to the correlation function of two vortex operators as $\langle V(\mathbf{x})V(\mathbf{y}) \rangle = e^{-V(\mathbf{x}-\mathbf{y})}$. In the phase where static monopoles are confined, one has

$$\langle V(\mathbf{x})V(\mathbf{y}) \rangle \sim e^{-\mu|\mathbf{x}-\mathbf{y}|}, \tag{8.36}$$

and the magnetic Z_2 -symmetry is restored. Let us clarify the origin of Eq. (8.36) and get an idea of a mass parameter which appears in it. To this end, notice that we are dealing with the situation dual to that of finite-temperature 3D compact QED below T_{BKT} [51]. In the latter case, external static quarks, represented by Polyakov loops, are linearly confined. Therefore, similarly to the correlation function of two Polyakov loops in that theory, which decreases with the distance at the inverse soliton mass, the same happens in our case with the correlation function of two vortex operators. In fact, it can be shown that vortex operators create solitons in the theory (8.35). Solitons in this theory carry a unit topological charge, corresponding to the $U(1)$ symmetry generated by the topological current $j = \frac{g}{2} \epsilon^{\mu\nu\lambda} \partial_\mu \mathbf{A}_\nu \times \mathbf{A}_\lambda$. In the dimensionally-reduced theory, the 3D vortex operator, $V(\mathbf{x}) = e^{i\int_{\mathbb{S}^1} \mathbf{A} \cdot d\mathbf{l}(\mathbf{x})}$, becomes replaced by $V(\mathbf{x}) = e^{i\int_{\mathbb{S}^1} \mathbf{K} \cdot d\mathbf{l}(\mathbf{x})}$, or equivalently,

$$V(\mathbf{x}) = \exp \left[i \int_{\mathbb{S}^1} \frac{p}{2} \mathbf{K} \cdot d\mathbf{l}(\mathbf{x}) \right] \tag{8.37}$$

The integration contour here goes either to infinity or to a point where the vortex operator of the opposite magnetic charge,

$$V(\mathbf{y}) = \exp \left[i \int_{\mathbb{S}^1} \frac{p}{2} \mathbf{K} \cdot d\mathbf{l}(\mathbf{y}) \right] \tag{8.38}$$

is defined. One can further demonstrate that the operators (8.37) and (8.38) have topological charges 1 and -1 and are nothing, but the creation operators of a soliton and an antisoliton, respectively. This fact stems from the relations $\langle V(\mathbf{0}) \rangle = \langle V(\mathbf{0}) \rangle$ and $\langle V(\mathbf{0}) \rangle = \langle V(\mathbf{0}) \rangle$, where

70 D. Antonov and M. C. Diamantini

$Q = \frac{1}{2\pi} \int_C dx j(x)$ is the topological-charge operator with the contour C encircling anticlockwise the origin. These relations stem from another ones

$$\langle j(x) V(0) \rangle \approx \frac{x}{2x^2} \langle V(0) \rangle; \quad \langle j(x) V(0) \rangle \approx \frac{x}{2x^2} \langle V(0) \rangle$$

(where \approx means "neglecting correlation functions of topological currents higher than the two-point one"), that themselves can be obtained from the two-point correlation function of topological currents,

$$\langle j(x) j(0) \rangle = \frac{g^2}{(2\pi)^3} \frac{x \cdot x}{x^2} = \frac{1}{x^2} :$$

Owing to the fact that the operator (8.37) [(8.38)] creates a soliton [antisoliton], the large-distance behavior of the correlation function (8.36) is determined by the lightest state with the unit topological charge, i.e. by the soliton. One can therefore identify $\langle j(x) j(0) \rangle$ on the r.h.s. of Eq. (8.36) with the soliton mass m_{sol} , therefore $m_{sol} = T m_{sol}$.

We have considered above the two idealistic situations, namely when either dynamical monopoles or W -bosons are absent. In reality, when both are present and interact with each other, the deconfining phase transition temperature is neither T_{BKT} nor T_{XY} , but is rather T_c , which in fact lies between the two. The deconfining phase transition takes place when the densities of monopoles and W -bosons become equal, i.e. $2 = n_W$. Up to inessential preexponential factors, this happens when the monopole action S_0 is equal to m_W , i.e. at $T = T_c$. At this temperature, the thickness of the string connecting a W^+ and a W^- boson is of the order of its length. The thickness of the string has been discussed [cf. e.g. after Eq. (3.17)] to be the inverse Debye mass of the dual photon, $d / \epsilon^{1/2}$. The length of the string is clearly of the order of the average distance between W 's, that is $\epsilon^{1/2}$. Therefore, again up to preexponential factors, the thickness and the length of the string are equal at $T = T_c$. Note that these qualitative arguments can be formalized by the RG procedure [52]. In the presence of W -bosons, the RG equations possess three fixed points. The first two are the zero- and infinite-temperature ones. The third fixed point of the RG flow is a nontrivial infrared unstable one, $T = T_c$, $\epsilon = \epsilon_c$, which corresponds to the phase transition. [In this point, however, both fugacities become infinite. It has been demonstrated in Ref. [96] that a correcting factor $(\epsilon + 2) = (2 + \epsilon)$, by which T_c should be multiplied, appears, if one demands that the RG flow of the fugacities should be stopped when at least one of them becomes of the order of unity.]

Note that an independent condition of the deconfining phase transi-

tion stems from the coincidence of scaling dimensions of the operators $:\cos \frac{K}{4} \phi^-:$ and $:\cos g \phi^-:$ at the critical temperature. The scaling dimensions are equal $\frac{K}{4}$ and $\frac{g^2}{4}$ respectively, so that the monopole cosine term is relevant at $T < T_{BKT}$, whereas the cosine term of W -bosons is relevant at $T > T_{XY}$. The two scaling dimensions become equal when $g_m^2 T = g^2$, that yields the value of the critical temperature $\frac{g^2}{4}$. As we see, this value is indeed equal to T_c , up to the factor $\frac{1}{4}$, which is of the order of unity and is generated by the loop corrections. At the temperature $\frac{g^2}{4}$, both scaling dimensions are equal to unity, therefore both cosine terms in L_{2D} are relevant at this temperature. In fact, in the whole region of temperatures $T_{XY} < T < T_{BKT}$, both terms are relevant.

The supersymmetric version of the 3D GG model [54] contains, in addition to the dual photon, its superpartner { an adjoint fermion, which we will call photino. This model possesses a discrete parity symmetry, which should lead to the masslessness of photinos. At zero temperature, this is however not the case, since the parity is spontaneously broken via a nonvanishing photino condensate. Thus, at finite temperature, one can anticipate two phase transitions { one related to the vanishing of the photino condensate and the other one due to deconfinement. These two transitions could either be distinct and happen at different temperatures, or could coincide. In this respect, the model is similar to QCD with adjoint quarks, where a similar question can be asked about the (non-)coincidence of deconfinement and restoration of discrete chiral symmetry.

The Lagrangian of the model contains the bosonic fields of the non-supersymmetric GG model, that are the massless photon, the heavy W vector bosons and the massive Higgs field, as well as their superpartners { photino, winos and Higgsino. It has been shown in [97] that, just like in the GG model, the monopole effects render the dual photon massive, although the mass in this case is parametrically smaller, since it is due to the contribution from a two-monopole sector, rather than a single-monopole sector as in the GG model. The low-energy sector of the theory is described by the supersymmetric sine-Gordon model. Its Euclidean action in the superfield notation reads [We adopt here the notations of Ref. [98], in particular $d^2 = 1$.]

$$S = \int d^3x d^2 \left[\frac{1}{2} D D + \cos(g\phi) \right] ; \tag{8.39}$$

In this equation, the scalar supermultiplet and supercovariant derivatives have the form

72 D. Antonov and M. C. Diamantini

$$(x; \sigma) = \sigma + \frac{1}{2} F; D = \frac{g}{g} (\hat{\sigma}); D = \frac{g}{g} (\hat{\sigma}) :$$

Here, σ again denotes the dual-photon field (real scalar), ψ is the photino field, which is the two-component Majorana spinor ($\psi = \psi^T$), F is an auxiliary scalar field, $\hat{\sigma} = \sigma_1 \sigma_2$, and $\sim = \sim$. The monopole fugacity has dimensionality $[m_{ass}]^2$ and is exponentially small. In terms of the disorder operator, the action (8.39) in the component notations can be rewritten, up to an inessential constant, as

$$S = \int d^3x \left[\frac{2}{g_m^2} \psi^\dagger V \psi + \frac{1}{2} \hat{\sigma} \left(\frac{g_m^2}{2} V^2 + V^{-2} - \frac{(g_m)^2}{2} V^4 + V^{-4} \right) \right];$$

where $\psi = \psi = 4$. Besides the magnetic Z_2 symmetry, this action has an additional discrete parity symmetry inherited from the full supersymmetric GG action,

$$V(x_1; x_2; x_3) \rightarrow iV(x_1; x_2; x_3); \psi(x_1; x_2; x_3) \rightarrow \psi^3(x_1; x_2; x_3) :$$

The photino mass term is odd under the parity transformation. Thus, the photino can acquire a mass only if the parity is spontaneously broken, that is indeed the case. The breaking of parity results in the non-vanishing photino condensate $\langle \psi \rangle = m_W$ and leads to the non-vanishing photino mass $m = 2g_m^2$ (equal to the Debye mass of the dual photon). It can be proved [54] that the equality of the dual-photon and the photino masses is preserved on the quantum level as well.

At finite temperatures, one can integrate photinos out in the dimensionally-reduced theory. Including again W -bosons, we arrive at a theory similar to the one we had in the non-supersymmetric case:

$$S_{d: r} = \int d^2x \left[\frac{1}{2} (\hat{\sigma})^2 + 2 \cos \frac{p}{K} - 2 \cos g \right] ; \quad (8.40)$$

where β / Λ^2 is a certain positive and exponentially small fugacity. The crucial difference from the non-supersymmetric GG model is due to the factor $\sqrt{2}$ in the term $\cos(2/K)$, which makes the model (8.40) Z_4 -invariant. Similarly to the non-supersymmetric case, up to higher-loop corrections, the condition for the determination of the deconfinement phase transition temperature reads $4g_m^2 T = g^2$, that yields $T_c = \frac{g^2}{8}$. The universality class of the phase transition is therefore Z_4 rather than Z_2 , as it was in the non-supersymmetric case. Further, it has been argued in Ref. [54] that the parity order parameter, $V^2 + V^{-2}$, vanishes at $T > T_c$. Therefore, the parity

restoring phase transition takes place at the same temperature T_c as the deconining transition. While this is an interesting phenomenon, it seems to be somewhat non-generic. In particular, in 4D gauge theory with adjoint fermions there is no reason to expect the deconining and chiral-symmetry restoring phase transitions to coincide. The physical order parameter for deconement is the 't Hooft loop V [99], while for chiral symmetry it is the fermionic bilinear form χ . In four dimensions, the two have very different nature. While χ is a local field, V is a string-like object. It is thus difficult to imagine these two order parameters combining into a single one as it is the case in the 3D theory discussed above. The lattice results indeed indicate that, at least in the $SU(3)$ -theory, in four dimensions the two transitions are distinct [100].

9. Summary

Below we would like to mention once again some specific issues discussed in this review.

3D GG model at $T = 0, N \geq 2$

- { $m_H < 1$) N must be smaller than $\exp[(\text{certain constant}) \frac{m_H}{m_W} = g^2]$, otherwise the Higgs vacuum is not stochastic;
- { $m_H = 1$) string tension of the flat Wilson loop in the fundamental representation is obtained; it possesses an ambiguity in the numerical factor due to the exponentially large thickness of the string;
- { the Kalb-Ramond field, which incorporates both monopoles and free photons) numerical factor at all the string coupling constants (in particular, at the string tension) is fixed for an arbitrarily shaped surface, in the weak-field (low-density) approximation; a generalization of the so-obtained theory of conining strings to the adjoint case and to k -strings: $\frac{adj}{fund} \sim 2$ at $N = 1$; $\frac{k}{fund} = \frac{k(N-k)}{N-1}$ (Casimir scaling) in the low-density approximation; the leading non-diluteness correction is derived, such that, at $k = N - 1$, it can significantly distort the Casimir scaling;
- { 4D-case with the field-theoretical term in the strong-coupling regime) string term; fundamental, adjoint, and k -case critical values of β , at which crumpling might disappear; a modification of the vacuum structure due to the term.

3D GG model at $T = T_{cr}^{1=3}$, effects of W -bosons are neglected

- { $N = 2, m_H < 1$) $\frac{g^2}{m_W}$ may not be larger than a certain function of $\frac{m_H}{m_W}$, otherwise the Higgs vacuum is not stochastic;

74 D. Antonov and M. C. Diamantini

{ Higgs-induced corrections to the BKT RG flow in the leading order in m_H^{-1} ; m_H itself evolves very slowly in the vicinity of the BKT critical point; $N > 2$) unlike the $SU_P(2)$ -case, the RG invariance holds only modulo the approximation $a_{ij} \cos[(\phi_i - \phi_j)P] \approx a_{ii}$, even at $m_H \neq 1$;

{ in the presence of N_f dynamical fundamental quark flavors, at $N_f = 1$ and $m_q = 0$, $T_{BKT} = \frac{g^2}{4}$; at $N_f \geq 2$ and/or $m_q \neq 0$, any fundamental matter (including quarks themselves) is deconfined at $T = T_{dtr}$;

When bosons are taken into account, supersymmetric generalization) the deconfining and the discrete-parity restoring phase transitions occur at the same temperature, $\frac{g^2}{8}$; the universality class of the deconfining phase transition is the same as in Z_4 -invariant spin models.

confining string

{ the presence of the negative-stiffness term forces the introduction of a term into the string action that suppresses the formation of spikes and, thus, prevents the crumpling of the world-sheet. This action, in the large- D limit, has an infrared stable fixed point at zero stiffness, that corresponds to a tensionless smooth string with Hausdorff dimension 2;

{ the effective theory describing the infrared behavior of the confining string is a conformal field theory with central charge $c = 1$;

{ at high temperature, the free energy of the confining string goes like $F^2(\beta) \sim (1-\beta^2)$ with $\beta = 1/T$ and it agrees in sign, temperature behavior, and the reality property with the large- N QCD result obtained by Polchinski.

Acknowledgments

D. A. is grateful to Profs. A. Di Giacomo and D. Ebert, as well as to the staffs of the Physics Departments of the University of Pisa and of the Humboldt University of Berlin, where his part of this work has been respectively initiated and finalized, for cordial hospitality. He also acknowledges the financial support of INFN and of the Alexander von Humboldt foundation at the initial and final stages of this work, respectively. M. C. D. thanks CERN, where part of this review was written, for kind hospitality.

References

1. Yu. A. Simonov, *Phys. Usp.* 39, 313 (1996); A. Di Giacomo, H. G. Dosch, V. I. Shevchenko and Yu. A. Simonov, *Phys. Rept.* 372, 319 (2002).
2. A. Di Giacomo, *Topics in non-perturbative QCD*, hep-lat/0012013; *QCD vacuum and confinement*, in: *Campos do Jordao 2002, New states of matter in hadronic interactions*, p. 168 [hep-lat/0204001].
3. A. M. Polyakov, *Gauge fields and strings* (Harwood Academic Publishers, 1987).
4. K. Nishijima, *Int. J. Mod. Phys. A* 9, 3799 (1994); *ibid.* A 10, 3155 (1995); M. Chaichian and K. Nishijima, *Eur. Phys. J. C* 22, 463 (2001).
5. G. S. Bali, *Phys. Rept.* 343, 1 (2001).
6. *Proceedings of "Lattice 2003"*: *Nucl. Phys. Proc. Suppl.* 129 (2004).
7. M. G. Alford, K. Rajagopal and F. Wilczek, *Phys. Lett. B* 422, 247 (1998); M. G. Alford, K. Rajagopal and F. Wilczek, *Nucl. Phys. B* 537, 443 (1999); D. T. Son, *Phys. Rev. D* 59, 094019 (1999); for a review see: M. G. Alford, *Annu. Rev. Nucl. Part. Sci.* 51, 131 (2001).
8. J. Berges, N. Tetradis and C. Wetterich, *Phys. Rept.* 363, 223 (2002).
9. G. 't Hooft, *Nucl. Phys. B* 72, 461 (1974).
10. Yu. M. Makeenko and A. A. Migdal, *Phys. Lett. B* 88, 135 (1979) [Erratum -*ibid.* B 89, 437 (1980)]; Yu. M. Makeenko and A. A. Migdal, *Nucl. Phys. B* 188, 269 (1981).
11. *The large-N expansion in quantum field theory and statistical physics: From spin systems to 2-dimensional gravity*, Eds. E. Brezin and S. R. Wadia (World Scientific, Singapore, 1993); Yu. M. Makeenko, *Methods of contemporary gauge theory* (Cambridge University Press, Cambridge, 2002).
12. K. K. Onishi and K. Takenaga, *Phys. Lett. B* 508, 392 (2001).
13. L. D. Faddeev and A. J. Niemi, *Phys. Rev. Lett.* 82, 1624 (1999).
14. G. 't Hooft, *Nucl. Phys. B* 190, 455 (1981).
15. A. Di Giacomo, B. Lucini, L. Montesi and G. Pauti, *Phys. Rev. D* 61, 034503 (2000); *ibid.* D 61, 034504 (2000); J. M. Camona, M. D'Elia, A. Di Giacomo, B. Lucini and G. Pauti, *Phys. Rev. D* 64, 114507 (2001); L. Del Debbio, A. Di Giacomo, B. Lucini and G. Pauti, *Abelian projection in SU(N) gauge theories*, hep-lat/0203023; J. M. Camona, M. D'Elia, L. Del Debbio, A. Di Giacomo, B. Lucini and G. Pauti, *Phys. Rev. D* 66, 011503 (2002).
16. A. Di Giacomo, *Color confinement and dual superconductivity: An update*, hep-lat/0204032; J. M. Camona, M. D'Elia, L. Del Debbio, A. Di Giacomo, B. Lucini, G. Pauti and C. Pica, *Nucl. Phys. A* 715, 883 (2003); A. Di Giacomo, *Confinement of color: A review*, hep-lat/0310023; A. Di Giacomo, *Confinement of color: Recent progress*, hep-lat/0310021.
17. Y. Nambu and G. Jona-Lasinio, *Phys. Rev.* 122, 345 (1961); *ibid.* 124, 246 (1961); V. G. Vaks and A. I. Larkin, *ZhETF* 40, 282 (1961).
18. S. Weinberg, *Phys. Rev. Lett.* 18, 188 (1967); S. R. Coleman, J. Wess and B. Zumino, *Phys. Rev.* 177, 2239 (1969); C. G. Callan, S. R. Coleman, J. Wess and B. Zumino, *ibid.* 177, 2247 (1969).
19. J. Gasser and H. Leutwyler, *Nucl. Phys. B* 250, 517 (1985); *ibid.* B 250, 465 (1985); D. Ebert and M. K. Volkov, *Z. Phys. C* 16, 205 (1983); D. Ebert and H. Reinhardt, *Nucl. Phys. B* 271, 188 (1986).
20. D. Ebert, H. Reinhardt and M. K. Volkov, *Prog. Part. Nucl. Phys.* 33, 1 (1994).
21. A. A. Belavin, A. M. Polyakov, A. S. Shvarts and Y. S. Tyupkin, *Phys. Lett. B* 59, 85 (1975); for a recent review see e.g. T. Schafer and E. V. Shuryak, *Rev. Mod. Phys.* 70, 323 (1998).

76 D. Antonov and M. C. Diamantini

22. R. Jackiw, *Rev. Mod. Phys.* **49**, 681 (1977); B. J. Harrington and H. K. Shepard, *Phys. Rev. D* **17**, 2122 (1978).
23. E. M. Ilgenfritz and M. Müller-Preussker, *Nucl. Phys. B* **184**, 443 (1981); D. Diakonov and V. Yu. Petrov, *Nucl. Phys. B* **272**, 457 (1986).
24. A. Gonzalez-Arroyo and Yu. A. Simonov, *Nucl. Phys. B* **460**, 429 (1996).
25. M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, *Nucl. Phys. B* **147**, 385 (1979); *ibid.* **B 147**, 448 (1979).
26. A. M. Polyakov, *Nucl. Phys. B* **120**, 429 (1977).
27. G. 't Hooft, *Nucl. Phys. B* **79**, 276 (1974); A. M. Polyakov, *JETP Lett.* **20**, 194 (1974) [reprinted in: *Magnetic monopoles*, Eds. A. S. Goldhaber and W. P. Trower (American Assoc. Phys. Teachers, 1990) p. 103, and in: *Solitons and particles*, Eds. C. Rebbi and G. Soliani (World Scientific, 1984) p. 522].
28. For a review, see: M. B. Green, J. H. Schwarz and E. Witten, *Superstring theory* (Cambridge University Press, Cambridge, 1987); J. Polchinski, *String theory* (Cambridge University Press, Cambridge, 1998).
29. A. M. Polyakov, *Phys. Scripta T* **15**, 191 (1987).
30. M. Luscher and P. Weisz, *JHEP* **07**, 049 (2002).
31. A. M. Polyakov, *Nucl. Phys. B* **268**, 406 (1986);
32. H. Kleinert, *Phys. Lett. B* **174**, 335 (1986).
33. F. Quevedo and C. A. Trugenberger, *Nucl. Phys. B* **501**, 143 (1997).
34. A. M. Polyakov, *Nucl. Phys. B* **486**, 23 (1997).
35. E. Alvarez, C. Gomez and T. Ortin, *Nucl. Phys. B* **545**, 217 (1999).
36. M. C. Diamantini, F. Quevedo and C. A. Trugenberger, *Phys. Lett. B* **396**, 115 (1997).
37. A. M. Polyakov, *Int. J. Mod. Phys. A* **14**, 645 (1999).
38. E. Alvarez and C. Gomez, *Nucl. Phys. B* **550**, 169 (1999); *JHEP* **05**, 012 (2000).
39. S. S. Gubser, I. R. Klebanov and A. M. Polyakov, *Phys. Lett. B* **428**, 105 (1998).
40. M. C. Diamantini, H. Kleinert and C. A. Trugenberger, *Phys. Rev. Lett.* **82**, 267 (1999).
41. M. C. Diamantini, H. Kleinert and C. A. Trugenberger, *Phys. Lett. B* **457**, 87 (1999).
42. M. C. Diamantini and C. A. Trugenberger, *Phys. Lett. B* **421**, 196 (1998).
43. M. C. Diamantini and C. A. Trugenberger, *Nucl. Phys. B* **531**, 151 (1998).
44. M. N. Chernodub, M. I. Polikarpov, A. I. Veselov and M. A. Zubkov, *Phys. Lett. B* **432**, 182 (1998).
45. M. C. Diamantini and C. A. Trugenberger, *Phys. Rev. Lett.* **88**, 251601 (2002); M. C. Diamantini and C. A. Trugenberger, *JHEP* **04**, 032 (2002).
46. J. Polchinski, *Phys. Rev. Lett.* **68**, 1267 (1992).
47. M. C. Diamantini, H. Kleinert and C. A. Trugenberger, *Phys. Lett. A* **269**, 1 (2000).
48. R. S. Lakes, *Phys. Rev. Lett.* **86**, 2897 (2001); *Nat.* **410**, 565 (2001).
49. E. T. Akhmedov, M. N. Chernodub, M. I. Polikarpov, and M. A. Zubkov, *Phys. Rev. D* **53**, 2087 (1996).
50. A. Kovner, *Int. J. Mod. Phys. A* **17**, 2113 (2002); Confinement, magnetic Z_N symmetry and low-energy effective theory of gluodynamics, in: *At the frontier of particle physics* Ed. M. Shifman (World Scientific, 2002) Vol. 3, p. 1777 [hep-ph/0009138]; A. Kovner and B. Rosenstein, *Int. J. Mod. Phys. A* **7**, 7419 (1992).
51. N. O. Agasian and K. Zarembo, *Phys. Rev. D* **57**, 2475 (1998).
52. G. V. Dunne, I. I. Kogan, A. Kovner and B. Tekin, *JHEP* **01**, 032 (2001).
53. D. Antonov, *Phys. Lett. B* **535**, 236 (2002).
54. D. Antonov and A. Kovner, *Phys. Lett. B* **563**, 203 (2003).

3D Georgi-Glashow model and condensing strings 77

55. N. O. Agasian and D. Antonov, *Phys. Lett. B* 530, 153 (2002).
56. G. V. Dunne, A. Kovner and S. M. Nishigaki, *Phys. Lett. B* 544, 215 (2002).
57. M. K. Prasad and C. M. Sommerfeld, *Phys. Rev. Lett.* 35, 760 (1975) [Reprinted in: *Solitons and particles* Eds. C. Rebbi and G. Soliani (World Scientific, 1984) p. 530]; E. B. Bogomolnyi, *Sov. J. Nucl. Phys.* 24, 449 (1976) [Reprinted in: *Solitons and particles* Eds. C. Rebbi and G. Soliani (World Scientific, 1984) p. 389].
58. T. W. Kirkman and C. K. Zachos, *Phys. Rev. D* 24, 999 (1981).
59. K. Dietz and Th. Filk, *Nucl. Phys. B* 164, 536 (1980).
60. V. G. Kiselev and K. G. Selivanov, *Phys. Lett. B* 213, 165 (1988).
61. S. R. Wadia and S. R. Das, *Phys. Lett. B* 106, 386 (1981) [Erratum *ibid.* B 108, 435 (1982)]; N. J. Snyderman, *Nucl. Phys. B* 218, 381 (1983).
62. T. Banks, R. Myerson and J. B. Kogut, *Nucl. Phys. B* 129, 493 (1977).
63. R. Gilmore, *Lie groups, Lie algebras, and some of their applications* (John Wiley & Sons, 1974).
64. D. Antonov, *Mod. Phys. Lett.* 17, 279 (2002).
65. P. Orland, *Nucl. Phys. B* 428, 221 (1994); D. V. Antonov, D. Ebert, and Yu. A. Simonov, *Mod. Phys. Lett.* 11, 1905 (1996).
66. S. B. Khokhlov and Yu. M. Makeenko, *Phys. Lett. B* 101, 403 (1981).
67. E. T. Akhmedov, *JETP Lett.* 64, 82 (1996); E. T. Akhmedov, M. N. Chernodub and M. I. Polikarpov, *JETP Lett.* 67, 389 (1998); D. Antonov, *Phys. Lett. B* 475, 81 (2000); *Phys. Lett. B* 543, 53 (2002).
68. D. Antonov, *JHEP* 07, 055 (2000).
69. L. Del Debbio and D. Diakonov, *Phys. Lett. B* 544, 202 (2002).
70. J. Greensite and S. Olejnik, *JHEP* 09, 039 (2002).
71. J. Ambjorn, P. Olesen and C. Peterson, *Nucl. Phys. B* 240, 189 (1984).
72. A. Ammon and M. Shifman, *Nucl. Phys. B* 664, 233 (2003).
73. A. Ammon and M. Shifman, *Nucl. Phys. B* 671, 67 (2003).
74. B. Lucini and M. Teper, *Phys. Lett. B* 501, 128 (2001).
75. B. Lucini and M. Teper, *JHEP* 06, 050 (2001).
76. B. Lucini and M. Teper, *Phys. Rev. D* 64, 105019 (2001).
77. L. Del Debbio, H. Panagopoulos, P. Rossi and E. Vicari, *Phys. Rev. D* 65, 021501 (2002).
78. L. Del Debbio, H. Panagopoulos, P. Rossi and E. Vicari, *JHEP* 01, 009 (2002).
79. L. Del Debbio, H. Panagopoulos and E. Vicari, *JHEP* 09, 034 (2003).
80. V. I. Shevchenko and Yu. A. Simonov, *Phys. Rev. Lett.* 85, 1811 (2000); On Casimir scaling in QCD, hep-ph/0104135; *Int. J. Mod. Phys. A* 18, 127 (2003); A. I. Shoshi, F. D. Steffen, H. G. Dosch and H. J. Pimer, *Phys. Rev. D* 68, 074004 (2003).
81. D. Antonov and L. Del Debbio, *JHEP* 12, 060 (2003).
82. V. L. Berezinsky, *JETP* 32, 493 (1971); J. M. Kosterlitz and D. J. Thouless, *J. Phys. C* 6, 1181 (1973); J. M. Kosterlitz, *J. Phys. C* 7, 1046 (1974); for a review see e.g. J. Zinn-Justin, *Quantum field theory and critical phenomena* (Oxford Univ. Press, New York, 2nd edition, 1993).
83. D. Antonov, *Mod. Phys. Lett. A* 17, 851 (2002).
84. J. B. Kogut, *Rev. Mod. Phys.* 51, 659 (1979); B. Svetitsky and L. G. Yafe, *Nucl. Phys. B* 210 [FS6], 423 (1982).
85. R. Jackiw and C. Rebbi, *Phys. Rev. D* 13, 3398 (1976) [Reprinted in: *Solitons and particles*, Eds. C. Rebbi and G. Soliani (World Scientific, 1984) p. 331].
86. C. Lee and W. A. Bardeen, *Nucl. Phys. B* 153, 210 (1979).
87. For a review see: J. Cardy, *Scaling and renormalization in statistical physics* (Cambridge University Press, Cambridge, 1996).

78 D. Antonov and M. C. Diamantini

88. M. Luscher, K. Symanzik and P. Weisz, Nucl. Phys. B 173, 365 (1980).
89. J. Polchinski and Z. Yang, Phys. Rev. D 46, 3667 (1992).
90. J. J. Atick and E. Witten, Nucl. Phys. B 310, 291 (1988).
91. F. Wilczek, QCD in extreme conditions, in: Ban 1999, Theoretical physics at the end of the twentieth century, p. 567 [hep-ph/0003183].
92. V. B. Berestetsky, E. M. Lifshits, and L. P. Pitaevsky, Quantum electrodynamics (Pergamon, Oxford, 1982).
93. I. S. Gradshteyn and I. M. Ryzhik, Table of integrals, series, and products (Academic Press, Orlando, 1980).
94. R. P. Feynman, Statistical mechanics. A set of lectures (Addison-Wesley, Reading, MA, 1972).
95. I. I. Kogan and A. Kovner, in: At the frontier of particle physics, Ed. M. Shifman (World Scientific, 2002) Vol. 4, p. 2335 [hep-th/0205026].
96. Yu. V. Kovchegov and D. T. Son, JHEP 01, 050 (2003).
97. I. A. Heck, J. Harvey, E. Witten, Nucl. Phys. B 206, 413 (1982).
98. M. Moshe and J. Zinn-Justin, Nucl. Phys. B 648, 131 (2003).
99. C. Korshals-Aliès and A. Kovner, Phys. Rev. D 62, 096008 (2000).
100. F. Karsch and M. Lutgemeier, Nucl. Phys. B 550, 449 (1999).