

# Cardy-Verlinde Formula and Holographic Dark Energy

Ke-Ke<sup>1</sup> and Miao Li<sup>1,2</sup>

<sup>1</sup> Institute of Theoretical Physics  
Academia Sinica, P.O. Box 2735  
Beijing 100080, China

and

<sup>2</sup> Interdisciplinary Center of Theoretical Studies  
Academia Sinica, Beijing 100080, China  
kek@itp.ac.cn, mli@itp.ac.cn

If we separate energy in a holographic theory into an extensive part and an intrinsic part, where the extensive part is given by the cosmological constant, and assume entropy be given by the Gibbon-Hawking formula, the Cardy-Verlinde formula then implies an intrinsic part which agrees with a term recently proposed by Hsu and Zee. Moreover, the cosmological constant so derived is in the form of the holographic dark energy, and the coefficient is just the one proposed recently by Li. If we replace the entropy by the so-called Hubble bound, we show that the Cardy-Verlinde formula is the same as the Friedmann equation in which the intrinsic energy is always dark energy. We work in an arbitrary dimension.

July, 2004

Recently there has been considerable interest in explaining the observed dark energy [1,2] by the holographic dark energy model [3]. (See also [4].) A. Cohen and collaborators suggested some time ago that the zero-point energy in quantum field theory is affected by an infrared cut-off  $L$ , thus, the cosmological constant is given by a formula  $\Lambda = M_p^2 L^{-2}$ . If one takes  $L^{-1}$  be the Hubble constant  $H$ , then dark energy is close to the critical energy density observed. However, Hu pointed out that in a universe with this dark energy and matter, this formula yields a wrong equation of state [5]. Li subsequently proposed that the infrared cut-off  $L$  should be given by the size of event horizon [6], namely  $L = a \int_{t^0}^{R_1} dt = a(t^0)$ , then the observed data can be nicely fitted. This model was further studied in [7]

Hu and Zee recently made the following interesting observation [8]: Suppose there be an infrared cut-off  $L$  in an effective action describing our universe, the cosmological constant term contributes an term  $\Lambda L^4$  to this action, suppose further for some reason (maybe due to quantum corrections, as suggested by these authors) there be an additional term  $M_p^4$  in this effective action, then minimizing the action yields  $\Lambda = M_p^2 L^{-2}$ . Hu and Zee propose the following picture, let the energy scale associated with the cosmological constant be  $m$ , the energy scale associated with the infrared cut-off be  $m_U$ , thus  $m = \frac{M_p}{m_U}$ , a formula reminiscent of the seesaw mechanism.

In Cohen's paper [3], the holographic form of dark energy is argued for by setting the UV and IR cutoffs to saturate the bound set by formation of a black hole. In this note, we assume that the dark energy has a holographic origin. By relating it to the Cardy-Verlinde formula, we thus argue for the holographic form of dark energy from another viewpoint.

Verlinde suggested, motivated both by holography and by Cardy's formula for a 2 dimensional conformal field theory, a generalization of Cardy's formula to a  $n$  dimensional conformal field theory [9]

$$S = \frac{4\pi R}{n-1} \frac{E_c E_e}{E_c E_e}; \quad (1)$$

where  $S$  is entropy,  $E_c$  is a part of energy similar to Casimir energy, and  $E_e$  is the extensive part of energy (our convention for  $E_c$  differs from that of Verlinde by a factor of  $\frac{1}{2}$ ). The CFT is supposed to live in spacetime with topology  $R \times S^{n-1}$ ,  $n-1$  is the spatial dimension.

The effective action proposed in [8] can be written in a form of effective energy

$$E = \frac{M_p^4}{L^3} + \frac{M_p^4}{L}; \quad (2)$$

Minimizing the effective energy also results in the correct scaling for  $\Lambda$ .  $\tilde{E}$  is of course the extensive energy in bulk, it is reasonable to assume that it is also the extensive energy

in the field theory dual to the cosmology theory in question, since by dimensional analysis,  $\rho = \frac{E_e}{L}$  can be viewed as the 2d energy density, then  $E_e = \rho L^2$  scales correctly in terms of the 2d volume.  $M_p^4 = (\frac{1}{L})$  is the intrinsic (Casimir) energy, then  $\frac{1}{L} \overline{E_c E_e} L = M_p^2 L^2$ , a quantity resembling the Gibbons-Hawking entropy if we identify  $L$  with  $R$ , the size of the cosmic horizon. This remarkable coincidence strongly suggests to us that the new term  $M_p^4 = (\frac{1}{L})$  has a holographic origin: it arises as the Casimir energy in a dual theory, if so, its nature is indeed quantum mechanical.

Note that while Verlinde originally works with a CFT with spacetime dimension the same as that of the whole universe, in our problem we prefer to interpret his formula as the one in a dual theory with one fewer dimension. There are many papers applying the Cardy-Verlinde formula to dS space, in the spirit of dS/CFT correspondence. For example, in [10], the author suggested that there is a CFT theory dual to dS and checked the Cardy-Verlinde formula. Here, we also assume the dark energy in the bulk dual to a CFT theory on the boundary. There is a correspondence between the extensive holographic dark energy in the bulk and extensive energy in the CFT theory. So we can use the Cardy-Verlinde formula (1). Of course, our purpose as well as the definition of energy are completely different from those in [10]. Other discussions on relation between the Cardy-Verlinde formula and the Friedmann equation with a cosmological constant can be found for instance in [11].

In Li's paper [6], there is an undetermined parameter  $c$ . The author suggested  $c = 1$  through an argument compared to blackhole. Here, we will use the Cardy-Verlinde formula to determine this parameter. Let us work instead in an arbitrary dimension and be more accurate numerically. Suppose in our universe there be a cosmic horizon of size  $R$ , or more generally, an infrared cut-off  $R$ . Let spacetime be  $n+1$  dimensional. The Gibbons-Hawking entropy is

$$S_{GH} = \frac{\frac{1}{2} \Omega_{n-1} R^{n-1}}{4G} = \frac{1}{2} \Omega_{n-1} M_p^{n-1} R^{n-1}; \quad (3)$$

where  $\Omega_{n-1}$  is the volume of the unit sphere  $S^{n-1}$ ,  $M_p^{n-1} = 1/(8G)$  the reduced Planck mass in  $n+1$  dimension. Note that we need to use the Gibbons-Hawking entropy for our purpose, this is why we interpret  $R$  as the size of the cosmic horizon.

Next, with the presence of dark energy density  $\rho$ , we propose that the bulk energy be  $E_e = \rho V$ , where  $V = \frac{1}{n} \Omega_n R^n$  is the volume enclosed by the cosmic horizon, and interpret this as the extensive energy in a dual theory. It is certainly difficult to regard  $E_e$  as energy

in the flat patch  $ds^2 = -dt^2 + a(t)^2 (dx^i)^2$ , since there is no notion of conserved energy in this system of coordinates. If we work in the static coordinates

$$ds^2 = -\left(1 - \frac{r^2}{R^2}\right)dt^2 + \left(1 - \frac{r^2}{R^2}\right)^{-1}dr^2 + r^2 d\Omega_{n-1}^2; \quad (4)$$

then  $E = R^p \frac{1}{G_{00}G_{rr}} = V$  is the conserved energy conjugate to  $t$ , the proper time of the comoving observer sitting at  $r = 0$ . However, as we shall explain later, it still makes sense to interpret  $E$  calculated in the flat patch as energy in the dual theory.

Let  $E_c$  be unknown, applying the Cardy-Verlinde formula (1) to  $E_e$ ,  $S_{GH}$  and  $E_c$  we solve  $E_c$  in terms of  $E_e$  and  $S_{GH}$  and find

$$E_c = \frac{n(n-1)^2}{4} M_p^{2n-2} R^{n-4}; \quad (5)$$

This is a  $n+1$  dimensional generalization of proposal in [8]. Note that we assume that the Cardy-Verlinde formula is used in a  $n$  dimensional dual theory. In the dual theory, the total energy is then

$$E = E_e + E_c = \frac{1}{n} M_p^{n-1} R^n + \frac{n(n-1)^2}{4} M_p^{2n-2} R^{n-4}; \quad (6)$$

Minimizing  $E$ , we obtain

$$= \frac{n(n-1)}{2} M_p^{n-1} R^{-2}; \quad (7)$$

the holographic dark energy in  $n+1$  dimension. Take  $n = 3$ , this result agrees with the one proposed in [6] with parameter  $c = 1$  advocated there.

It is easy to check that the above result is consistent with a  $n+1$  dimensional de Sitter space. The Friedmann equation in  $n+1$  dimensions is

$$\frac{n(n-1)}{2} M_p^{n-1} H^2 = \rho; \quad (8)$$

Replacing  $\rho$  by  $E/V$  and plugging (7) into the above equation, we have  $H^2 = R^{-2}$ , this is exactly the correct relation between the Hubble constant  $H$  and the cosmic horizon size  $R$  for a de Sitter space.

Having succeeded in deriving the dark energy formula in a holographic dual, we now come to a universe with both dark energy and other form of energy. If we insist on the Cardy-Verlinde formula, since at least the extensive energy is larger than the one corresponding to dark energy, entropy should be greater than the Gibbons-Hawking entropy.

This entropy, just like the one in a closed universe discussed in [9], must be the Hubble bound  $S_H$ . We propose the following formula for the Hubble bound

$$S_H = \frac{nH V}{4G} = 2^{-n-1} M_p^{n-1} H R^n; \quad (9)$$

where we work in the flat patch, only in this case  $H$  is defined. This formula is similar to the one proposed in [9], but with a slight difference in coefficient since we have a flat universe with boundary. We may or may not interpret  $R$  as the size of event horizon, but it must be the same infrared cut-off appearing in the dark energy formula. In a pure de Sitter space, the Hubble bound agrees with the Gibbons-Hawking entropy only when  $R$  is taken to be the horizon size.

Solving  $H$  in terms of  $S_H$

$$H^2 = \frac{S_H^2}{(2^{-n-1} M_p^{n-1} R^n)^2}; \quad (10)$$

On the L.H.S. we use the Friedmann equation

$$M_p^{n-1} H^2 = \frac{2}{n(n-1)} (\rho + \rho_m); \quad (11)$$

where  $\rho$  is the holographic dark energy (7), and  $\rho_m$  is whatever energy density in question, we deduce

$$\begin{aligned} S_H^2 &= \frac{8^{-2} M_p^{n-1} R^{2n-2}}{n(n-1)} (\rho + \rho_m) \\ &= \left( \frac{4 R}{n-1} \right)^2 E_c (E_c + E_m); \end{aligned} \quad (12)$$

where  $E_c = \rho V = E_e$  is the "on-shell" Casimir energy associated with  $\rho$ , and  $E_c + E_m = E_e + \rho_m V$  is the total extensive energy. We thus find that the Cardy-Verlinde formula is equivalent to Friedmann equation provided we always use the same formula for the Casimir energy without any contribution from the matter part. This result is similar to Verlinde's: When the Casimir energy is replaced by Bekenstein-Hawking energy (in our case, Bekenstein-Hawking energy is similar to the on-shell Casimir energy), the Cardy-Verlinde formula is equals to the Friedmann equation.

Note that the presumed holographic dual itself hides the time evolution in the bulk, the Friedmann equation governs time evolution in the bulk, but its interpretation in the dual theory is a static relation. Similarly, one can not ask the question as to why the total energy is not conserved in the bulk. By the same token,  $S_H$  is interpreted in the dual

theory as total entropy, while in the bulk it is interpreted as a upper bound only. One can not take  $S_H$  as the real entropy in the bulk, since in general  $S_H$  viewed as a function of time does not observe the second law of thermodynamics. To see this, let us compute

$$\frac{d}{dt}(H R^n) = R^{n-2} \left[ 1 + \frac{p}{1+w} - \frac{n}{2} \frac{1}{(1+w)} + n \left( \frac{1}{1+w} - \frac{1}{2} \right) \right]; \quad (13)$$

where  $\frac{p}{1+w}$  is the fraction of the dark energy,  $w$  is the equation of state index of matter. One can check numerically that in later time when  $\frac{p}{1+w}$  approaches 1, the above quantity can be negative.

Our understanding of this issue is the following. Just as the total energy  $E_e + E_m$ ,  $S_H$  can not be interpreted as a physical quantity in the bulk, its role is just a bound on entropy. We propose that the Gibbons-Hawking entropy is real entropy in the bulk, and it is certainly bounded by the Hubble bound. The Gibbons-Hawking entropy always increases with time.

To summarize, we used the Cardy-Verlinde formula twice. In the first usage, we apply the Gibbons-Hawking entropy and obtain the Casimir energy, by minimizing the total energy we obtain the dark energy formula. In the second usage, we apply the Hubble entropy and obtain the Friedmann equation. The first usage is valid both for an "empty" de Sitter space as well as a spacetime with other form of energy, if this application of the Cardy-Verlinde formula is correct, we effectively "derived" the formula proposed in [6]. The second usage can be regarded as a definition of the Hubble entropy, thus less conjectural and less consequential.

Acknowledgments.

We are grateful to R.G.Cai and A.Zee for useful discussions. This work was supported by a grant of CAS and a grant of NSFC.

## References

- [1] A . G . Riess et al., *Astron. J.* 116 (1998) 1009; S. Perlmutter et al., *APJ* 517 (1999) 565.
- [2] C . L . Bennett et al., astro-ph/0302207, *Astrophys.J.Suppl.* 148 (2003) 1; D . N . Spergel et al., astro-ph/0302209, *Astrophys.J.Suppl.* 148 (2003) 175; H . V . P . Peiris et al., astro-ph/0302225, *Astrophys.J.Suppl.* 148 (2003) 213.
- [3] A . Cohen, D . Kaplan and A . Nelson, hep-th/9803132, *Phys.Rev.Lett.* 82 (1999) 4971.
- [4] S . Thomas, *Phys. Rev. Lett.* 89 (2002) 081301.
- [5] S . D . H . Hsu, *Phys.Lett. B* 594 (2004) 13-16.
- [6] M . Li, hep-th/0403127, *Phys. Lett. B* 603 (2004) 1.
- [7] Q . G . Huang and Y . Gong, astro-ph/0403590, *JCAP* 0408 (2004) 006; Y . Gong, hep-th/0404030, *Phys. Rev. D* 70 (2004) 064029; Q . G . Huang and M . Li, astro-ph/0404229, *JCAP* 0408 (2004) 013; M . Ito, hep-th/0405281; K . Enqvist and M . S . Sloth, hep-th/0406019.
- [8] S . Hsu and A . Zee, hep-th/0406142.
- [9] E . Verlinde, hep-th/0008140.
- [10] R . G . Cai, hep-th/0111093, *Phys. Lett. B* 525 (2002) 331.
- [11] B . Wang, E . Abdalla, R . K . Su, hep-th/0101073, *Phys.Lett. B* 503 (2001) 394; S Nojiri, S D Odintsov and S Ogushi, hep-th/0205187, *Int. Journ. Mod. Phys. A* 17 (2002) 4809.