

A Model of Perfect Fluid with Spin and Non-Abelian Color Charge

O. V. Babourova,

Department of Theoretical Physics, Faculty of Physics,
Moscow State University, Leninskie Gory, d.1, st.2, Moscow 119992, Russia,

A. S. Vshivtsev,^y

Academician V. P. Maslennikov,
Keldysh Institute of Applied Mathematics, Russian Academy of Sciences,
Musskaya pl. 4, Moscow 125045, Russia,

B. N. Frolov,^z

Department of Mathematics, Moscow State Pedagogical University,
Krasnoprudnaya 14, Moscow 107140, Russia

At present, there is a tendency for mutual penetration of the methods of the continuous medium mechanics and the methods of classical and quantum field theory. On the one hand, the use of the Yang-Mills formalism in perfect fluid hydrodynamics allows one to construct new gauge-invariant models [1]. On the other hand, in field theory, the hydrodynamic approach (according to which the quantum system of interacting quarks and gluons is approximated classically by the perfect fluid with spin and non-Abelian color charge) is being successfully developed [2],[3]. In this work, a relativistic variational theory of the perfect spin fluid with color charge in an external non-Abelian Yang-Mills field will be derived in the Riemann-Cartan space U_4 with curvature and torsion, which will take into account the spin polarization chromomagnetic effects.

As the dynamic variables describing the fluid, we take the null-form fields and their conjugates, which are transformed by the representations of the direct product of the Lorentz group and $SU(3)$ color group. An element of the fluid has a vector of four-dimensional velocity $u = u^a e_a$, whose corresponding three- and one-forms of velocity may be taken, respectively, as $u := u^c = u^a e_a$ and $u = u_a e^a = g(u; \cdot)$, with $u \wedge u = 0$, which is the conventional condition for the squared velocity: $g(u; u) = c^2$. Here e^a are the basis one-forms, ϵ is the volume four-form. $\cdot$ denotes the inner product, and $\wedge$ denotes the Hodge dualization. We also introduce the three- and two-form fields [4] $\omega_b := e_b c = \omega_b$ and $\omega_{ab} := e_b c_a = (e_a \wedge e_b)$, respectively. The basis e_a is supposed to be nonholonomic orthogonal with $g(e_a, e_b) =: g_{ab} = \text{diag}(1; 1; 1; 1)$.

The internal energy density ϵ of the fluid depends on the extensive (additive) thermodynamic parameters $\{$ the number of particles per unit volume (concentration) n and the entropy per particle s , which are calculated in the intrinsic frame of reference $\}$ and on the quantities describing the internal degrees of freedom of the fluid particle, namely, the spin

^{E-mail:} babourova@orc.ru

^yDeceased

^zE-mail: frolovbn@orc.ru

tensor $S_{ab} = M_{ab}$ and the color charge $J_m = I_m$, where M_{ab} and I_m are the generators of the corresponding representations of the Lorentz and SU (3) color groups, respectively. The quantities characterizing the uid (including n and s) are considered to be averaged over a certain elementary three-volume that contains what is referred to as a uid element.

Furthermore, the uid element in itself represents a statistical subsystem with a sufficiently large number of particles to consider it as a quasi-closed system whose properties coincide with the statistical properties of the uid in a macroscopic state. For the uid element, the first law of thermodynamics has the following form :

$$d(n; s; S_{ab}; J_m) = \frac{n + p}{n} dn + n T ds + \frac{1}{2} n!^{ab} dS_{ab} + n!^m dJ_m ; \quad (1)$$

where p is the hydrodynamic pressure in the uid. In (1), the coefficient multiplying dn has the meaning of the chemical potential, and $!^{ab}$ and $!^m$ describe, respectively, the possible spin and color-charge exchanges between uid elements.

We suppose that the laws of conservation of the number of particles and entropy [which can be expressed by the equalities $d(nu) = 0$ and $d(nsu) = 0$, where d is the operator of exterior differentiation] are valid. The first of these equations ensures the continuity of flow lines of the uid particles. The second equation expresses the constancy of entropy along the flow lines of the uid and thus constitutes the second law of thermodynamics for adiabatic flow of the uid.

The fact that the spin tensor meets the Frenkel condition $S_{ab}u^b = 0$ expresses the fundamental physical property of the tensor, its spacelike character. In terms of the exterior forms, this condition can be written by $(e_a \lrcorner S) \wedge u = 0$, where the spin two-form $S = (1/2)S_{ab} e^a \wedge e^b$ has been introduced.

The four-form of Lagrangian density, whose integral determines (the action of the theory, is chosen by us in the form

$$L_{fluid} = L_{fluid} = (n; s; ;) + n D \wedge u - n J_m F^m \wedge S + n_1 (u \wedge u \lrcorner c^2) + n u \wedge d_2 + n_3 u \wedge ds + n^a (e_a \lrcorner S) \wedge u : \quad (2)$$

In (2), $n, s, ,$ and u are regarded as independent variables describing the perfect uid dynamics. In doing so, one should take into account that, in view of (1), the dependence of the internal energy on n and s is realized only through its dependence on the spin tensor and color charge: $= (n; s; M_{ab}; I_m)$. The constraints imposed on the independent variables are taken into account by using the undetermined Lagrange multipliers $_1, _2, _3$ and a . The second term in (2) is a kinetic term in which D designates the operation of exterior covariant differentiation with respect to both gauge groups [the Lorentz and SU (3) color groups]. The term with coupling constant $_1$ in (2), in which $F^m = (1/2)F^m_{ab} e^a \wedge e^b$ designates the two-form of strength of the non-Abelian gauge color field, describes the possible spin polarization chromomagnetic effects.

On variation of (2) over the undetermined Lagrange multipliers, we obtain the corresponding constraints, and on variation over the dynamic variables describing the perfect uid, we obtain the variational equations of motion of the perfect uid with non-Abelian color charge. Using these equations, together with the values of the undetermined Lagrange multipliers determined by means of these equations, it is possible to show that the four-form of Lagrangian density (2) is proportional to the hydrodynamic pressure in the uid $L_{fluid} = p$. This indicates that the variational theory of perfect spin uid with internal color charge transforms into the variational theory of a conventional perfect uid as a result of correct passage to the limit.

We use the equations of motion and the identity $D M_{ab} = 0$ to deduce the law that governs changes of the spin tensor of the fluid particle:

$$u^\alpha \wedge D S_{ab} + \frac{2}{c^2} S_{[a}^c u_{b]} u_c = 2 S_{[a}^f c_{b]}^d (F^m_{cd} J_m + !_{cd}) : \quad (3)$$

Here, the projection tensor is designated by $c_b^c = \delta_b^c + c^2 u^c u_b$ and the Newtonian dot signifies the differentiation that is defined for an arbitrary object by the equality $\dot{a}_b := (u^\alpha \wedge D a_b)$. With the zero right-hand side, the law governing changes of the spin tensor (3) takes the form of the corresponding Weyssenhof-Raabe law [5] for the perfect spin fluid.

To derive the equations for the non-Abelian gauge color field, we should add the four-form of the color-field Lagrangian density to the four-form of Lagrangian density (2):

$$L_{matter} = L_{fluid} + L_{field}; \quad L_{field} = -\frac{1}{2} F^m \wedge F_m : \quad (4)$$

Here, g is the coupling constant. Lifting and lowering the indices like m at the gauge field, which are transformed by an adjoint representation of the $SU(3)$ color group, are effected with the metric tensor $g_{mn} = -(1/2) c_m^p c_n^q$, where c_m^p are the structure constants of the $SU(3)$ group.

The equations for the field are obtained by variation of (4) over the one-form A^m of potential of the gauge color field:

$$D (F_m + g J_m S) = g J_m u : \quad (5)$$

This equation must be supplemented with the Bianchi identity for a non-Abelian gauge field $D F^m = 0$.

Using the equation of motion for the fluid and identity $D I_m = 0$, we derive the law that governs changes of the non-Abelian color charge:

$$u^\alpha \wedge D J_m = c_m^p c_n^q (F^n \wedge S + !^n) : \quad (6)$$

The derived classical equations of motion for the non-Abelian color field, which take into account spin and color degrees of freedom, form a self-consistent set. Equation (6) shows that, in the case of a non-Abelian field, the conservation of color charge along a fluid flow line may be violated because of the spin-chromomagnetic interaction. It is notable that, in the case of an Abelian field, the conservation of electric charge along a fluid flow line always takes place, since in that case $c_m^p c_n^q = 0$.

According to the general principles of the field theory for the Riemann-Cartan space [4], [6], [7], geometric properties of space-time are defined by the canonical energy-momentum tensor and the spin momentum tensor of the matter. In the exterior form formalism, the role of the latter tensor is played by the three-form of spin momentum τ_a^b , which is defined as a variational derivative of (4) with respect to the connection one-form ω_a^b of the Riemann-Cartan space U_4 . The role of the canonical energy-momentum tensor is played by the energy-momentum three-form τ_a , which is defined as a variational derivative of the Lagrangian density four-form (4) with respect to the basis one-form θ^a . On calculating τ_a for the Lagrangian density (4) with the use of the equations of motion of the fluid and the expressions for the undetermined Lagrangian multipliers, we arrive at

$$\tau_a = \frac{L_{matter}}{\theta^a} = \tau_a^{fluid} + \tau_a^{field}; \quad (7)$$

$$\begin{aligned}
\frac{f_{\text{fluid}}}{a} &= p_a + n_a + \frac{p}{nc^2} u_a u + n (e_a c F^m J_m) \wedge S; \\
\frac{f_{\text{field}}}{a} &= (F^m_{ac} F^{bc}_m - \frac{1}{4} F^m_{ac} F^{cd}_m F^{ab}_m) \wedge b = \\
&= \frac{1}{2} ((e_a c F^m) \wedge F_m - F^m \wedge (e_a c F_m)) :
\end{aligned} \tag{8}$$

In these expressions, the dynamic momentum per particle in a fluid element is introduced:

$$u_a = \frac{1}{nc^2} u \wedge \frac{f_{\text{fluid}}}{a}; \quad a = \frac{1}{c^2} u_a - \frac{1}{c^2} S_a^c u_c + u^b (J_m F^{bc}_m + !_{bc});$$

Here u denotes the effective energy per fluid particle, $u = u_0 + J_m F^m \wedge S$, where $u_0 = u/n$. This expression generalizes the corresponding result obtained for the case of an electromagnetic field [8] to the non-Abelian case. We see that the role of the fluid pressure p is twofold: first, it prevents the compression of the fluid [the first term in (8)], and, second, it effectively increases the mass of the fluid element [the second term in (8)].

It is well known in the theory of electromagnetic fields that the equations of motion of charged particles follow from the energy-momentum conservation law of the system together with the field equations, and, in turn, the energy-momentum conservation law is a consequence of the invariant properties of the system. In the field theory in U_4 space, a similar statement is valid, but the laws of conservation have a more complicated form [4]:

$$D_a = (e_a c T^b) \wedge b - (e_a c R^c_b) \wedge b^b_c; \quad D_{ab} = !_{ab} \wedge b; \tag{9}$$

Here, T^b is the torsion two-form and R^c_b is the curvature two-form of U_4 space. The first of these equations is a consequence of invariance of the system's Lagrangian under diffeomorphisms of the energy-momentum, and the second equation is a consequence of local Lorentz invariance and a generalization of the law of conservation of the spin momentum. Both of these equations become identities if and only if the equations of motion of the system in U_4 space hold. Thus, these equations are in fact just another representation of the laws of motion.

We use the first of equations (9) to derive the equation of motion of the perfect spin fluid with color charge in the form generalizing the Euler hydrodynamic equation. Substituting the energy-momentum tensor (7) in (9) and taking into account the gauge color field equation (5), we obtain

$$\begin{aligned}
u \wedge D \left(a + \frac{p}{nc^2} u_a \right) &= \frac{1}{n} r_a p - (e_a c T^b) \wedge \left(b + \frac{p}{nc^2} u_b \right) u \\
(e_a c R^{bc}) \wedge \left(\frac{1}{2} S_{bc} u \right) &- (e_a c F^m) \wedge (J_m u) + (r_a \wedge F^m) \wedge (J_m S);
\end{aligned} \tag{10}$$

where r_a designates the covariant differentiation along the direction of the basis field e_a and $r_a \wedge$ in the last term denotes the covariant differentiation of the tensor-like forms. The third term in the right-hand side of equation (10) represents the Mathisson force arising from the interaction of the fluid-particle spin with the space curvature; the second term is characteristic of the U_4 space and reflects the interaction between the torsion of the space and the dynamic momentum of the particle. The fourth term in the right-hand of this equation generalizes the Lorentz force to the case of a non-Abelian gauge field.

Now let us consider a zero fluid-pressure limiting case. In this case, equation (10) describes the motion of a single fluid particle that has a spin and color charge. Equations

(10), (3) and (6), together constitute the classical equations of motion of the color particle, which generalize the Wong equations [9] to the case of the $SU(3)$ color group and allow for the spin of the particles.

The constructed theory may form the basis of a new model that hydrodynamically describes the quark-gluon plasma and differs from the Landau model in that it allows for the structure of quantum chromodynamics vacuum and color charge of quarks in an explicit form [10]. Moreover, on the basis of the theory of perfect spin-dilaton fluid developed in [11], this model can be generalized to the case of a fluid possessing the dilaton charge, apart from the spin and the color charge, and thus provide the foundation for studies of the non-perturbed gluon condensate up to the phase transition from hadron matter to the quark-gluon plasma.

References

- [1] Guzev M. A. and Myasnikov V. P., *Izv. Akad. Nauk, Mekh. Zhidk. Gaza*, 1993, no. 4, pp. 25{29.
- [2] Holm D. D., Kupersmidt B. A., *Phys. Rev. D : Part. Fields*, 1984, vol. 30, pp. 2557{2560.
- [3] Amorim R., *Phys. Rev. D : Part. Fields*, 1986, vol. 33, pp. 2796{2800.
- [4] Trautman A., *Symposia Mathematica*, vol. 12, Bologna: Monograph, 1973, pp. 139{162.
- [5] Ray J. R., Smalley L. L., *Phys. Rev. D : Part Fields*, 1983, vol. 27, pp. 1383{1385.
- [6] Kibble T. W. B., *J. Math. Phys.*, 1961, vol. 2, pp. 212{221.
- [7] Frolov B. N., *Vestn. Mosk. Univ., Ser. 3: Fiz.-Astron.*, 1963, no. 6, pp. 48{58.
- [8] Corben H. C., *Classical and Quantum Theory of Spinning Particles*, San Francisco: Holden-Day, 1968.
- [9] Wong S. K., *Nuovo Cimento B*, 1970, ser. 10, vol. 65, pp. 689{694.
- [10] Vshivtsev A. S. and Peregudov D. V., *Yad. Fiz.*, 1997, no. 7, pp. 825{834.
- [11] Babourova O. V., Frolov B. N., *The Variational Theory of Perfect Hypermomentum Fluid*, Preprint of Los Alamos Archive 1996, Los Alamos, no. gr-qc/9612055.