# N on-perturbative A spects of K aluza {K lein M odes in F ive-dim ensional Supersymmetric Q C D on S<sup>1</sup>

Yukiko Ohtake

Department of Physics, Faculty of Science, Ochanomizu University Institute of Humanities and Sciences, Ochanomizu University 2-1-1 Otsuka, Bunkyo-ku, Tokyo 112-8610, Japan

#### A bstract

We study the stability of Kaluza {K lein (KK) modes in ve-dimensional N = 1 supersymmetric QCD compactied on S<sup>1</sup> using the D3-brane probe realization. We nd a phenomenon in which the quark KK mode with the KK number n = 1 decays while other KK modes are stable. This is contrary to the ordinary assumption that the state with n = 1 is the most stable in quark KK modes. In addition, we show that a massive gauge singlet state carrying the KK number exists stably. This provides a proper candidate for dark matter.

### 1 Introduction

Encouraged by great success of the brane world scenario [1], m any scientists have studied various m odels in which our four-dimensional world is embedded in a higher dimensional spacetime. In the research, the KK m ode plays a central role since it enables us to observe extra-dimensions from the four-dimensional world. Despite of the importance, however, the stability of KK m odes has been studied only perturbatively [2]. Thus we would like to investigate non-perturbative aspects of the stability in this paper.

For simplicity, we study ve-dimensional SU (2) Yang{M ills theory with  $N_f$  fundamental m atters compacti ed on S<sup>1</sup>. In addition we introduce N = 1 supersymmetry in

ve-dimension, which provides N = 2 supersymmetry in the four-dimensional elective theory. Then we analyze the model by using Seibreg{W itten theory  $\beta$ ], which tells us the exact elective theory and the stability of the states in it. A lthough determining the stability is complicated in eld theory, we can easily determine it in IIB string theory, where the elective theory appears on the D 3-brane in a 7-brane background[4].

In the following we review how the theory appears on the D3-brane and how the stability is determined. A fler that, we apply the method to study the stability of phenom enologically important KK states. In section 3, we determ ine the stability of quark KK modes. As a result, we nd a phenom enon in which the KK mode with the KK number n = 1 decays while other KK modes are stable. In section 4, we show that there exists a massive gauge singlet, which gives a proper candidate for dark matter.

### 2 Brane Construction of Five-dim ensional SQCD

Five-dimensional N = 1 SU (2) QCD appears in IIA string theory as the world volume theory of a D4-brane in the 8-brane background consisting of an O8-plane and N<sub>f</sub> D8-branes parallel to the D4-brane[5]. In this picture, a string connecting a D8-brane and the D4-brane provides a quark, while a string with the both ends on the D4-brane provides a gauge eld. As we set the origin of the space transverse to the 8-branes at the position of the O8-plane, the positions of D8-branes correspond to the quark bare m ass parameters and that of D4-brane corresponds to the moduli parameter which comes from the vacuum expectation value of an adjoint Higgs eld.

To compactify the eld theory on  $S^1$ , we compactify the IIA string theory on  $S^1$  parallel to the D 4-brane. Moreover we perform T-duality along  $S^1$  to obtain the string picture which describes the elective four-dimensional theory well. The T-duality transforms IIA theory into IIB theory, the N<sub>f</sub> D 8-brane into N<sub>f</sub> D 7-branes, the O 8-plane into two O 7-planes, and the D 4-brane into a D 3-brane, on which the elective theory appears. As the

ve-dimensional case, the positions of branes correspond to the parameters of the eld theory and strings ending on the D 3-brane provide quarks and gauge elds. In addition, the distance between the O 7-planes is related to 1=R where R is the radius of  $S^1$ , and the winding number of strings around the O 7-planes corresponds to the KK number in the

eld theory.

M oreover, there appear various strings providing solitons on the D 3-brane[6]. To see this, we introduce the notation used in [7]: we call a fundamental string (1;0), a D 7-brane [1;0], the SL (2;Z) dual strings (p;q) and the dual 7-branes [p;q], where p and q are relatively prime integers. As (1;0) emanates from [1;0], (p;q) emanates from [p;q]. In this notation the background of N<sub>f</sub> D 7-branes and two O 7-planes is written as  $[1;0]^{N_f}$  ([2; 1][0;1])<sup>2</sup> because an O 7-plane is identi ed with [2; 1][0;1][8]. Thus there appear not only (1;0) but also (2; 1), (0;1) and their composites. O ne of these strings, (p;q), is observed as a particle with the electric charge of unbroken U (1) gauge symmetry of SU (2) p and the magnetic charge q. W hat composites exist stably depends on details of the 7-brane background. In the following we shall review a gravitational description of the background.

#### 2.1 7-brane backgrounds

A 7-brane background is expressed as a solution of the ten-dimensional supergravity which is the low energy e ective theory of IIB theory [9]. The solution consists with an eightdimensional at spacetime parallel to the 7-branes and a two-dimensional curved space denoted by z-plane. The z-plane is described by an elliptic curve,

$$y^2 = x^3 + f(z)x + g(z);$$
 (1)

where x and y are complex parameters, and f(z) and g(z) are polynomials of z. The equation (1) gives a torus for a xed value of z. The periods of the torus are

$$\frac{\mathrm{da}(z)}{\mathrm{dz}} = \begin{bmatrix} 1 & \frac{\mathrm{dx}}{\mathrm{y}}; & \frac{\mathrm{da}_{\mathrm{D}}(z)}{\mathrm{dz}} = \begin{bmatrix} 1 & \frac{\mathrm{dx}}{\mathrm{y}}; & (2) \end{bmatrix}$$

where f; g is an orthogonal basis of two-cycles of the torus. U sing a (z) and  $a_D$  (z), we obtain the vacuum expectation value of the dilaton and the R am ond R am ond zero-form (z), and the metric on the z-plane ds<sup>2</sup> as follow s;

$$(z) = ie^{-(z)} (z) = \frac{da_{D}(z)}{da(z)};$$
 (3)

$$ds^{2} = Im \quad (z) \frac{da(z)}{dz} dz^{2} : \qquad (4)$$

The positions of 7-branes are zeros of the discrim inant of (1),  $(z) = 4f (z)^3 + 27g (z)^2$ . The number of 7-branes at a zero is the order of the zero. Types of 7-branes at a zero of (z) are determined from the monodrom y matrix acting on  $t(\frac{da_D}{dz}; \frac{da}{dz})$  when we go around the zero anti-clockwisely [7]. When the matrix at an n-th order zero is expressed as  $p_{1,2} p_{1,2} M_{2,2}$ , where

$$M_{[p,q]} = \begin{array}{c} 1 \quad pq \quad p^2 \\ q^2 \quad 1 + pq \end{array};$$
(5)

we decide that the 7-branes at the zero are  $[p_1;q_1][p_2;q_2]$   $_n;[p_n]$ .

#### 2.2 string junctions

In this subsection we introduce strings in a 7-brane background. If a string is stable, it stretches along a certain curve on the z-plane to m inim ize its mass. We call such curve geodesic in this paper. If a geodesic passes z = u, the corresponding string can end on the D 3-brane at z = u and provides a stable particle in the world volum e theory with the moduli parameter u. Therefore the region on the z-plane where the geodesic of a string (p;q) passes corresponds to the region on the moduli u-plane where the particle (p;q) exists stably. Hence we determ ine geodesics in (4) to investigate the stability of particles.

W e start with a simplest background, a 7-brane [p;q] at  $z = z_1$ . In this case, the string em anating from the background is (p;q). The mass of the string connecting the 7-brane and the D 3-brane at z = u along a curve C is given by

$$M_{(p,q)}(u) = \int_{C}^{Z} T_{p,q} ds; \qquad (6)$$

where  $T_{p,q} = \dot{p} + \dot{q} = \dot{p}$  is the tension of (p;q) [6]. Substituting (3) and (4) for (6), we obtain the lower bound of the mass as follows:

$$M_{(p;q)}(u) = \int_{c}^{2} p \frac{da(z)}{dz} q \frac{da_{D}(z)}{dz} j dz j$$

$$jpa(u) qa_{D}(u) pa(z_{1}) + qa_{D}(z_{1}) j; \qquad (7)$$

The minimum is realized if and only if all the points on C satisfy

Arg (pa (z) 
$$qa_D$$
 (z)  $pa(z_1) + qa_D(z_1)) = ;$  (8)

where is a constant between 0 and 2.

The equation (8) for a xed value of gives a geodesic of (p;q). In this background we can use pa  $qa_D$   $c(z z_1)[\beta]$ , so that (8) gives a half line from  $z = z_1$  at an angle of

A rgc to the positive realaxis. Therefore, as varies from 0 to 2, the geodesic sweeps all the z-plane. Thism eans that the state corresponding to the string (p;q) appears stably in the theory with any value of the m oduli parameter u.

Next we consider the background of two 7-branes,  $[p_1;q_1]$  at  $z = z_1$  and  $[p_2;q_2]$  at  $z = z_2$ . We assume  $z_1 \notin z_2$  because two 7-branes never coincide unless  $[p_1;q_1][p_2;q_2] = [1;0][1;1], [1;0][1;0]$  or their SL (2;Z) dual[10], and geodesics for the exceptions are derived from the limit  $z_1 ! z_2$  in the case of  $z_1 \notin z_2$ . In this background there exists a string  $(p_i;q_i)$  (i = 1;2) connecting  $[p_i;q_i]$  at  $z = z_i$  and the D3-brane at z = u. The geodesic is described by (8) where we replace (p;q) with  $(p_i;q_i)$  and  $z_1$  with  $z_i$ . The equation is di cult to solve, how ever, qualitatively the geodesic should be a line bent by gravity of the other 7-brane. Thus the geodesic sweeps all the z-plane as in the background of a 7-brane. Then the particle  $(p_i;q_i)$  appears stably in the theory with any value of the moduli parameter.

In addition to the strings, there exists a string junction [11]. A string junction is constructed from  $n_1(p_1;q_1)$  and  $n_2(p_2;q_2)$  emanating from the 7-branes and merge into



Figure 1: A con guration of string junction (a) and the related CMS (b).

a string  $(p;q) = \prod_{i=1,2}^{P} n_i (p_i;q_i)$  as in Fig.1a. The mass of a junction is the sum of the component string masses. These are derived similar to (7) and the geodesics are descrived by

Arg 
$$(p_i a(z) q_i a_D(z) p_i a(z_i) + q_i a_D(z_i)) = _i;$$
 (9)

Arg (pa (z) 
$$qa_D(z)$$
 pa ( $z_J$ ) +  $qa_D(z_J)$ ) = ; (10)

where  $z = z_J$  is the point where the strings m erge. Thus the m ass of the string junction is given by

$$M_{(p;q)}(u) = \frac{ja_{a}(u)}{X} qa_{b}(u) pa(z_{J}) + qa_{b}(z_{J})j + n_{i}jp_{i}a(z_{J}) q_{i}a_{b}(z_{J}) p_{i}a(z_{i}) + q_{i}a_{b}(z_{i})j i = 1;2 pa(u) qa_{b}(u) X n_{i}(p_{i}a(z_{i}) q_{i}a_{b}(z_{i})); (11) i = 1;2$$

where the lower bound is realized when  $_1 = _2 =$ . Therefore the geodesics (9) and (10) are rewritten as follows;

- -

$$A \operatorname{rg} (p_i a (z) \quad q_i a_D (z) \quad p_i a (z_i) + q_i a_D (z_i)) = ;$$
 (12)

Arg@pa(z) 
$$qa_{D}(z)$$
  $n_{i}(p_{i}a(z_{i}) q_{i}a_{D}(z_{i}))^{A} = :$  (13)  
 $\sum_{i=1,2}^{X}$ 

The geodesics (12) and (13) for a proper value of are estimated as in Fig.1a. If we change continuously, the geodesic of  $(p_i;q_i)$  rotates around  $z = z_i$ . Then the merging point  $z = z_J$  draws a circle as depicted in Fig.1b, and the string (p;q) sweeps only the outside of the circle. Hence the particle (p;q) exists when the moduli parameter u takes a value outside the circle and disappears when u takes a value inside it. When u takes a value on the circle, (p;q) becomes marginally stable: in this case, the D 3-brane is located at  $z = z_J$  in Fig.1a, so that (p;q) can decay to  $n_1$   $(p_1;q_1)$  and  $n_2$   $(p_2;q_2)$ . Then the curve drawn by  $z = z_J$  is called a curve of marginal stability (CMS). Since a point on CMS

satis es (12) and (13), CMS obeys

$$\operatorname{Im} \frac{p_{i}a(z) \quad q_{i}a_{D}}{pa(z) \quad q_{a}(z) \quad P} \frac{p_{i}a(z_{i}) + q_{i}a_{D}(z_{i})}{p_{j-1,2}n_{j}(p_{j}a(z_{j}) \quad q_{j}a_{D}(z_{j}))} = 0:$$
(14)

Finally, we consider a background of N 7-branes,  $[p_i;q_i]$  at  $z = z_i$  (i = 1;2;:::;N (3)). As discussed before, a string emanating from each 7-branes,  $(p_i;q_i)$ , sweeps all the z-plane. Then the corresponding state exists in the eld theory with any value of u. Moreover, there exist string junctions. We call  $(p;q)_n$  a string junction constructed from  $n_i(p_i;q_i)$ , where  $(p;q) = \prod_{i=1}^{P} n_i(p_i;q_i)$  and  $n = (n_1;n_2;:::;n_N)$ . The con guration is complicated because  $(p_i;q_i)$  merge one after another. To x the order in which the strings merge, we have to solve quantitatively the geodesics of  $(p_i;q_i)$  which are described by (12). On the other hand, the mass and the geodesics of the component strings are the same form independent of the order;

$$M_{(p;q)_{n}}(u) = pa(u) \quad qa_{D}(u) \qquad \prod_{i=1}^{X^{n}} n_{i}(p_{i}a(z_{i}) - q_{i}a_{D}(z_{i})); \qquad (15)$$

Arg 
$$p^{0}a(z) q^{0}a_{D}(z) = ;$$
 (16)

where  $(p^0;q^0)_{n^0}$  runs all the component strings. From (16) we see that CMS is given by

$$\operatorname{Im} \frac{p^{0}a(z) \quad q^{0}a_{D}(z)}{pa(z) \quad q_{D}(z) \quad p_{j=1}^{P} n_{j}^{0}(p_{j}a(z_{j}) \quad q_{j}a_{D}(z_{j}))} = 0; \qquad (17)$$

where  $(p^0;q^0)_{n^0}$  is a string merging the outgoing string  $(p;q)_n$ . To specify  $(p^0;q^0)_{n^0}$ , we have to x the order in which strings merge. In the following section, we do this in some backgrounds.

### 3 Stability of quark Kaluza {K lein m odes

In the previous section, we review a method of determ ining the stability of particles appearing on the D 3-brane in a 7-brane background. Using the method, we shall derive the spectrum of quark KK modes in ve-dimensional N = 1 SU (2) QCD compactified on S<sup>1</sup>, which is realized on the D 3-brane in the 7-brane background  $[1;0]^{N_f}$  ([2; 1][0;1])<sup>2</sup>. For simplicity, we concentrate on the cases in which N<sub>f</sub> = 7;6;5, all the quarks are massless, and the gauge coupling constant g ! 1. In these cases the global symmetry is enhanced to  $E_{N_f+1}$ , and all the 7-branes but one coincide to produce the symmetry [12].

The explicit form s of a and  $a_D$  for the backgrounds are derived in [13] as follows;

$$\begin{array}{c} a_{\rm D}(z) \\ a(z) \end{array} = \frac{\sin}{2 (1 2) R} \qquad \begin{array}{c} 2_2 & 2_1 & V_1(z) \\ \frac{!}{\sin} & 2 \frac{!}{\sin} & 1 \end{array} , \qquad \begin{array}{c} V_2(z) \end{array} ; \qquad (18)$$

where  $_{1} = (2) = {}^{2}()$ ,  $_{2} = (2 2) = {}^{2}(1)$ ,  $! = e^{i(\frac{1}{2})}$  and = 1=6;1=4;1=3 for N  $_{f} = 7;6;5$ , respectively. A lso V<sub>i</sub>(z) (i = 1;2) are given by

$$V_1(z) = \frac{1}{z} {}_{3}F_2(;;;2;1+;z)$$
 (19)

$$V_{2}(z) = \frac{1}{1} z^{1} {}_{3}F_{2}(1 ; 1 ; 1 ; 2(1 ); 2 ; z):$$
(20)

In these backgrounds 7-branes are located at z = 0 and  $z = 1^{1}$ . The m onodrom y m atrices around these points are

.

$$M_{0} = \begin{array}{ccc} N_{f} & 7 & N_{f} & 8 \\ 1 & 1 & 7 \end{array}; M_{1} = \begin{array}{ccc} 1 & 0 \\ 1 & 1 \end{array}; (21)$$

These are rewritten as  $M_0 = M_{[1;0]}^{N_f} M_{[2;1]} M_{[2;1]} M_{[2;1]}$  and  $M_1 = M_{[0;1]}$ , where  $M_{[p;q]}$  is given in (5). Thus 7-branes  $[1;0]^{N_f} [2; 1][0;1][2; 1]$  are located at z = 0 and a 7-brane [0;1] at z = 1. For the later use we also provide the monodrom y matrices acting on  $t_{(a_D;a;1=R)}$ ;

In this background there exists a string junction constructed from (p;q n) emanating from z = 0 and n(0;1) from z = 1. The mass form ula is obtained from (15),

$$M_{(p,q)_{n}} = jpa(u) \quad qa_{D}(u) \quad pa(0) + (q \quad n)a_{D}(0) + na_{D}(1)j$$
  
=  $pa(u) \quad qa_{D}(u) + \frac{n}{R};$  (23)

where we have used  $a(0) = a_D(0) = 0$  and  $a_D(1) = 1=R$ . The geodesics of the outgoing string of the junction is derived from (16),

Arg pa(z) 
$$qa_{D}(z) + \frac{n}{R} = :$$
 (24)

W e call  $(p;q)_n$  a string whose con guration satisfying (24) throughout this paper. In this notation a string (1;0) from z = 0 which corresponds to a quark is represented by  $(1;0)_0$ . In addition, a string corresponding to a quark KK mode with the KK number n is  $(1;0)_n$ , since the number of strings from z = 1 is identi ed with the KK number from (23).

Here we introduce branch cuts on the z-plane to make the functions a(z) and  $a_D(z)$  single-valued [14]. We take the cuts running from each 7-brane to z = 1 along the real

 $<sup>^1</sup>$  T he distance between the 7-branes actually depends on the radius R. Here we elim inate it by rescaling the original z to  $\frac{2}{27}R^6z$ ,  $\frac{1}{8}R^4z$ ,  $\frac{4i}{27}R^3z$  for N  $_{\rm f}$  = 7;6;5, respectively.



Figure 2:  $(1;0)_1$  and the related CMS.

axis. If we go across the cut en anating from z = i (i = 0;1) from the lower half-plane, <sup>t</sup>( $a_D$ ;a;1=R) changes to  $M_i^{1t}(a_D;a;1=R)$ . Simultaneously, we should change <sup>t</sup>(p;q;n) into  $K_i^{t}(p;q;n)$  where

to keep the geodesic (24) unchanged. We assume in the following that the cut from z = 0 is in nitesimally lower than the cut from z = 1. Thus<sup>t</sup> (p;q;n) changes to K<sub>1</sub>K<sub>0</sub><sup>t</sup> (p;q;n) when we go across the cuts on Rez > 1 from the lower half-plane.

W ith these branch cuts, we can determ ine a and  $a_D$  numerically using M athematica and solve the geodesics (24) approximately. To do this, we take a lattice of the spacing d on the z-plane for the region Rezj s and Jim zj s. Then we generate data of the numerical values of a and  $a_D$  at the lattice points, and m ake data of signs of Im (pa  $qa_D$  n)= $e^i$ . If the signs at both ends of a link are dimerent, at least a solution of (24) exists on the link. Thus we look for such links and plot the middle of each as an approximate solution. These plots tell us an approximate conguration of (p;q) at . We made plots for d = 0.01 and s = 5 for N<sub>f</sub> = 7;6;5 independently, and found that the geodesics in N<sub>f</sub> = 7;6;5 are similar. Then we shall give a gure to show the results of the three cases at once.

Now we are ready to consider our main problem, the conguration of  $(1;0)_h$ . First we derive the geodesics (24) for  $(1;0)_0$ . Taking an appropriate value of , we me that the conguration of  $(1;0)_h$  is as depicted in Fig.2a. The geodesic crosses the cuts on Rez > 1, then the set of charges t(1;0;0) changes to  $K_1K_0^t(1;0;0) = (1;0;1)$ . Thus we join  $(1;0)_0$  to  $(1;0)_1$  on the cuts. When we change gradually, we me that the geodesic  $(1;0)_h$  sweeps the whole z-plane. Therefore the quark  $(1;0)_0$  exists in the theory with any value of the moduli parameter.

Second we consider how  $(1;0)_1$  appears. Recall that it already appears in Fig2a as  $(1;0)_0$  crosses the cuts on Rez > 1. Therefore, as we change in Fig2a,  $(1;0)_1$  of the string sweeps the right-hand side of the upper-half z-plane until the string collides with the 7-brane at z = 1. If we change further after the collision, the string changes to a string junction with the outgoing string  $(1;0)_1$  as depicted in Fig2b [7, 15]. This is

(p,q) $(p_1,q_1)$  $(p_2,q_2)$ 



Figure 3:  $(1;0)_2$  and the related CMS.

understood in the following way. A fler the collision,  $(1;0)_0$  gets over the cut on Rez < 1 and the set of charges (1;0;0) changes to K $_0^t(1;0;0) = (1; 1;0)$ . Thus  $(1;0)_0$  changes to  $(1; 1)_0$  as depicted in Fig.2b while  $(1;0)_1$  preserves its charges. Consequently, to ll the di erence of these charges,  $(0;1)_1$  should be created at z = 1. A swe change in Fig.2b, a point where strings m erge, say  $z = z_1$ , draw s a half circle C<sub>1</sub> as in Fig.2c. Sim ultaneously  $(1;0)_1$  sweeps all the z-plane except for the region S<sub>1</sub> surrounded by C<sub>1</sub> and the real axis. A ll the lower-half plane is swept because  $(1;0)_1$  is hardly bent by 7-branes at z = 0, which act as a large gravitational source. From these facts, we conclude that the quark KK m ode  $(1;0)_1$  exists outside S<sub>1</sub>, decays to  $(1; 1)_0 + (0;1)_1$  on C<sub>1</sub>, and disappears inside S<sub>1</sub>.

Third we consider  $(1;0)_2$ . The string appears in Fig.3a as  $(1;0)_1$  crosses the cuts on Rez > 1. As we change in Fig.3a, the outgoing string collides the 7-brane at z = 1 and changes to another string junction as depicted in Fig.3b. During the transition, an additional  $(0;1)_1$  is created. The geodesics of two  $(0;1)_1$  are the same, but we separate them in Fig.3b to show the structure of the junction. One of  $(0;1)_1$  merges with  $(1; 1)_1$  which appears as  $(1;0)_1$  crosses the cut while the other merges with  $(1; 1)_0$ . Changing

further, we nd that the merging point of (1; 1) and  $(0;1)_1$ , say  $z = z_2$ , draws a half circle  $C_2$  as depicted in Fig.3c. Also we nd that the outgoing string (1;0) sweeps all the z-plane except for the region  $S_2$  surrounded by  $C_2$  and the real axis. Thus we conclude that  $(1;0)_2$  exists outside  $S_2$ , decays to  $(1; 1)_1 + (0;1)_1$  on  $C_2$ , and disappear inside  $S_2$ .

Repeating the operation, we can construct  $(1;0)_n$  for n 3. The results are similar to the cases of n = 1;2. A geodesic  $(1;0)_n$  appears as  $(1;0)_{n-1}$  crosses the cuts on Rez > 1. As we change , the outgoing string becomes a junction in which  $(0;1)_1$  and a string junction  $(1; 1)_{n-1}$  merge into  $(1;0)_n$  at a point. The merging point draws a half circle  $C_n$  connecting z = 0 and z = 1. Simultaneously,  $(1;0)_n$  sweeps all the z-plane except for the region  $S_n$  surrounded by  $C_n$  and the realaxis. Thus the quark KK mode  $(1;0)_n$  exists outside  $S_n$ , decays to  $(1; 1)_{n-1} + (0;1)_1$  on  $C_n$ , and disappears inside  $S_n$ . We checked the result for n = 3;4 num erically and for other n by using the approximation introduced in [14], where we neglect the e ects of the 7-brane at z = 1 on the metric.



Figure 4: CM S of quark KK m odes.

Here we compare the sizes of  $C_1$  and  $C_2$ . For this purpose, let us notice the geodesic  $(0;1)_1$  in Fig.3b and two points on it,  $z = z_1$  and  $z = z_2$ . If we change ,  $(0;1)_1$  rotates around z = 1 and each point  $z = z_i$  (i = 1;2) draws a half circle  $C_i$ . We see that  $C_1$  is larger than  $C_2$  since  $z = z_1$  is farther from z = 1 than  $z = z_2$ . Sim ilar consideration is valid for other pairs of positive KK numbers,  $n_1$  and  $n_2$ . Then we conclude that  $C_{n_1}$  is larger than  $C_{n_2}$  when  $n_1 < n_2$  (Fig.4). Accordingly,  $(1;0)_1$  disappears is and other states  $(1;0)_n$  (n 2) disappear in numeric order of n when we change the moduli parameter from a value above  $C_1$  to the segment [0;1] on the real axis. This result is contrary to the ordinary assumption that  $(1;0)_1$  is the most stable in quark KK modes.

Before concluding this section, we comment on the quark KK modes with the negative KK numbers,  $(1;0)_n$  (n 1). The outgoing string  $(1;0)_n$  appears when we rotate  $(1;0)_0$  clockwisely n times. The resulting conguration of the junction is the mirror image of the conguration of  $(1;0)_n$  with respect to the realaxis. A ctually, the replacement  $z \mid \overline{z}$  in (24) corresponds to  $(p;q)_n \mid (p + (8 N_f)q; q)_n$  [16]. Thus  $(1;0)_n$  exists outside the mirror image of  $S_n$ , decays to  $(N_f - 7;1)_{n+1} + (8 N_f; 1)_1$  on the mirror image of  $C_n$ , and disappears inside the mirror image of  $S_n$ .

## 4 Stability of a gauge singlet state

In this section we study a bop string encircling all the 7-branes. The string has a nite length so that it should have nite mass. In addition, when we locate the D 3-brane on the loop, the total string charges absorbed by the brane are apparently zero. Then the string corresponds to a massive gauge singlet in eld theory, and provides a proper candidate for dark matter. This motivate us to study the string.

First we consider a loop string in the simple backgrounds given by (18). The mass of a string stretching along a loop C surrounding all the 7-branes is derived as follows;

$$M (u) = \prod_{\substack{c \\ c \\ lim, ja(z i) \\ e \\ lim, a(z + i) + \frac{1}{R}} T_{1;0} ds$$

$$= \lim_{\substack{c \\ lim, ja(z i) \\ e \\ lim, a(z + i) + \frac{1}{R}} a(z + i) = \frac{1}{R}; \qquad (26)$$

where z = z is the intersection of C and the realaxis on Rez 1, and we have used (22)



Figure 5: Con gurations of (0;0).

to derive the third line. The lower bound is realized when all the points on C satisfy

$$Arg(a(z) = a(z + i)) = 0;$$
 (27)

where is a positive in nitesimal number. Note that the mass formula is identical with (23) when  $(p;q)_n = (0;0)_1$ , and the charges of a particle corresponding to the string is  $(0;0)_1$ . Thus we call the string and the particle  $(0;0)_1$  in the following.

Now we solve the geodesics (27) numerically. The resulting congurations are closed bops labeled by z 0. As we decrease z, the loop gets smaller and nally collapses to a point at z = 0. Note here that a loop of z > 1 is closed not only in shape but also in the string charges: when the string crosses the cuts on Rez > 1, the charges (1;0) remain unchanged. On the other hand, a loop of 0 z 1 is not closed in charges because (1;0) changes to (1; 1) as it crosses the cut on Rez 1 from the lower half plane. Thus a string (0;1)<sub>1</sub> should be added to 11 the di erence of the charges. From these facts we conclude that the conguration of (0;0) is as depicted in Fig.5a when z > 1, Fig.5b when 0 < z 1 and Fig.5c when z = 0. One can show that the strings in the gures have the same mass, 1=R.

From the gures we determ ine how (0;0) is detected by the D 3-brane probe and nd out the stability of the corresponding state. First we focus on a part of  $(0;0)_1$ , a string (1;0) which makes a bop. As we change z from 0 to 1, the string sweeps all the z-plane but S, the segment [0;1] on the real axis. Hence  $(0;0)_1$  is detected through the string if the D 3-brane is not on S. The D 3-brane on the bop does not divide  $(0;0)_1$  in parts so that the string cannot decay. Therefore the corresponding state is stable in this case. Next we consider the rest of  $(0;0)_1$ , the string  $(0;1)_1$ . It appears only on S. W hen we bocate the D 3-brane on it, the brane separates  $(0;0)_1$  into two parts,  $(0;1)_1$  and  $(0; 1)_0$ . Thus  $(0;0)_1$  can decay to  $(0;1)_1 + (0; 1)_0$ , and the corresponding state is marginally stable. In summary, the state  $(0;0)_1$  exists for any value of the moduli parameter u, and it is stable if u is not on S and marginally stable if u is on S.

Next we shall generalize the results to the eld theory with other parameters. Recall that the parameters correspond to the positions of 7-branes. Thus we deform the background (18) by moving 7-branes, and determ ine how the deform ation a ects the geodesics



Figure 6: Con gurations of loop strings.

of bop strings. Let us start with a large bop string in the background (18) as depicted in Fig.5a. In general, a geodesic of (p;q) changes continuously unless it hits a 7-brane [r;s] (for [p;q]), while it changes to a string junction when it hits the 7-brane. Therefore, if we move 7-branes nitely in the bop, the string remains a bop as depicted in Fig.6a. A loo if we drive a 7-brane [1;0] to in nity, which corresponds to reducing  $N_f$ , the loop of (1;0) remains a loop. On the other hand, if we drive [2; 1] or [0;1] to in nity, the loop string disappears: the loop changes to a junction as depicted in Fig.6b and the string connecting the loop and the 7-brane becomes in nitely long. Thus the corresponding state has in nite mass, and disappears from the spectrum. This is natural because the deform ations corresponds to the lim it R ! 0. In conclusion, a loop string large enough to encircle all the 7-branes exists in any background with nite R. Next we x the 7-branes and consider the region where the loop string sweeps. As we change the size of the loop from in nity to zero, the string hits 7-branes with di erent charges and changes to a junction as in Fig.6bc. Then the singlet sweeps all the z-plane and nally it becomes S, the string junction with the vanishing loop as depicted in Fig.6d. From the con gurations we see that the singlet is stable when u is not on S and marginally stable when u is on s.

A lthough we do not discuss in detail, it is interesting to study the decays occurring on the D 3-brane at the singularities. As an example, let us consider the case where the D 3-brane is located at z = 0 in the background (18). Junctions in this background are constructed from  $(0;1)_1$  and various  $(p;q)_0$ , and the length of  $(p;q)_0$  becomes zero when the D 3-brane is located at z = 0. Thus the state  $(P;Q)_n$  can decay to  $(P;Q n)_0 + n (0;1)_1$ . Then  $(0;1)_1$  can carry all the KK number of the theory. Recall here that the conguration of  $(0;0)_1$  observed at z = 0 is the same with  $(0;1)_1$  as depicted in Fig.5c. The dimension between  $(0;1)_1$  and  $(0;0)_1$  is that a massless string  $(0;1)_0$  is added or not. Thus the decay  $(0;1)_1 \\$   $(0;0)_1 + (0;1)_0$  would occur and then all the KK number could be carried by the singlet.

### A cknow ledgem ents

I would like to thank J.K am oshita, A. Sugam oto, T. Suzuki, T.U esugi and H.Yakazu for useful discussions. This work was partly supported by the Grant-in-Aid for Scienti c Research on Priority A reas (No.14039204) from the M inistry of Education, Culture, Sports, Science and Technology, Japan.

### References

- N.ArkaniHamed, S.D in opoulos and G.D vali, Phys. Lett. B 429 (1998) 263, hep-ph/9803315; Phys.Rev.D 59 (1999) 086004, hep-ph/9807344; L.R andalland R.Sundrum, Phys.Rev.Lett. 83 (1999) 3370, hep-ph/9905221; Phys.Rev.Lett. 83 (1999) 4690, hep-th/9906064.
- [2] H.-C. Cheng, K.T. M atchev and M. Schmaltz, Phys. Rev. D 66 (2002) 036005, hepph/0204342.
- [3] N. Seiberg and E. W itten, Nucl. Phys. B 426 (1994) 19; Erratum -ibid. B 430 (1994) 485, hep-th/9407087; Nucl. Phys. B 431 (1994) 484, hep-th/9408099.
- [4] O. Bergm an and A. Fayyazuddin, Nucl. Phys. B 531 (1998) 108, hep-th/9802033;
   A. Mikhailov, N. Nekrasov and S. Sethi, Nucl. Phys. B 531 (1998) 345, hep-th/9803142.
- [5] N. Seiberg, Phys. Lett. B 388 (1996) 753, hep-th/9608111
- [6] J.H. Schwarz, Phys. Lett. B 360 (1995) 13; Erratum -ibid. B 364 (1995) 252, hepth/9508143.
- [7] M.R.Gaberdieland B.Zwiebach, Nucl. Phys. B 518 (1998) 151, hep-th/9709013.
- [8] A.Sen, Nucl. Phys. B 475 (1996) 562, hep-th/9605150.
- [9] B.R.Greene, A. Shapere, C. Vafa and S.-T. Yau, Nucl. Phys. B 337 (1990) 1.
- [10] K.Kodaira, Annals of Math. 71 (1961) 111; 77 (1963) 563; 78 (1963) 1.
- [11] J.H. Schwarz, Nucl. Phys. Proc. Suppl. 55B (1997) 1, hep-th/9607201.
- [12] Yam ada and S.-Kil. Yang, Nucl. Phys. B 566 (2000) 642, hep-th/9907134; A. Sen and B. Zwiebach, JHEP 0003 (2000) 036, hep-th/9907164.
- [13] K.Mohri, Y.Ohtake and S.-Kil. Yang, Nucl. Phys. B 595 (2001) 138, hep-th/0007243.
- [14] M.R.Gaberdiel, T.Hauer and B.Zwiebach, Nucl. Phys. B 525 (1998) 117, hepth/9801205.

[15] O.Bergm an and A.Fayyazuddin, Nucl. Phys. B 535 (1998) 139, hep-th/9806011.
[16] Y.Ohtake, Prog. Theor. Phys. 102 (1999) 671, hep-th/9812227.