

TRACES ON THE SKLYANIN ALGEBRA AND CORRELATION FUNCTIONS OF THE EIGHT-VERTEX MODEL

H. BOOS, M. JIMBO, T. MIWA, F. SMIRNOV AND Y. TAKEYAMA

Abstract. We propose a conjectural formula for correlation functions of the \mathbb{Z} -invariant (inhomogeneous) eight-vertex model. We refer to this conjecture as Ansatz. It states that correlation functions are linear combinations of products of three transcendental functions, with theta functions and derivatives as coefficients. The transcendental functions are essentially logarithmic derivatives of the partition function per site. The coefficients are given in terms of a linear functional Tr on the Sklyanin algebra, which interpolates the usual trace on finite dimensional representations. We establish the existence of Tr and discuss the connection to the geometry of the classical limit. We also conjecture that the Ansatz satisfies the reduced qKZ equation. As a non-trivial example of the Ansatz, we present a new formula for the next-nearest neighbor correlation functions.

1. Introduction

Exact description of correlation functions and their analysis is one of the central problems of integrable lattice models. Significant progress has been made over the last decade toward this goal. In the study of correlation functions, a basic role is played by a multiple integral representation, first found for the archetypical example of the spin $1/2$ -XXZ chain [17, 15, 20]. Subsequently it has been generalized in several directions, to incorporate an external field [20], unequal time [19], non-zero temperature [13] and finite chains [18]. Earlier in the literature, extension to elliptic models has also been pursued. The free field construction used in the XXZ model was extended in [24] to the SOS models, resulting in an integral formula for correlation functions of the ABF model. In [25] an integral formula was obtained for the eight-vertex model by mapping the problem to the SOS counterpart. A novel free field representation of the eight-vertex model is being developed in [33, 34].

Recent studies have revealed another aspect of these integrals. Through examples at short distance, it has been observed in the case of the homogeneous XXX chain that the relevant integrals can be evaluated in terms of the Riemann zeta function at odd integers with rational coefficients [5]. Similar calculations have been performed for the XXZ chain [23, 37]. This phenomenon was explained later through a duality between the qKZ equations of level 0 and level -4 [6, 7]. Motivated by these works, we have established in our previous papers [3, 4] an algebraic representation (in the

sense no integrals are involved) for general correlation functions of the inhomogeneous six-vertex model and its degeneration¹. The aim of the present paper is to continue our study and examine the eight-vertex model.

We formulate a conjectural formula for correlation functions (the Ansatz) along the same line with the six-vertex case. Consider the eight-vertex model where each column i (resp. row j) carries an independent spectral parameter t_i (resp. 0). The object of our interest is the matrix

$$h_n(t_1; \dots, t_n; t) = \frac{1}{2^n} \sum_{i_1, \dots, i_n \neq 0} X^{a_{i_1}} \dots X^{a_{i_n}} W_{1, \dots, n}^{-1} (t_1, \dots, t_n)^T \text{End}(\mathbb{C}^2)^n$$

where h_i denotes the ground state average in the thermodynamic limit $\beta \rightarrow \infty$, $a = 1, 2, 3$ are the Pauli matrices, and T stands for the matrix transpose. Regard h_n as a vector via the identification $\text{End}(\mathbb{C}^2)^n \rightarrow (\mathbb{C}^2)^{2n}$, and let s_n denote the vector corresponding to the identity. Our Ansatz is that h_n can be represented in the form

$$h_n(t_1; \dots, t_n; t) = 2^{-n} \exp \left(\sum_{i < j} X^{a_{ij}} \right) \prod_{a=1}^3 !_a(t_{ij}) \mathcal{X}_{a,n}^{(i,j)}(t_1; \dots, t_n; t s_n)$$

Here $!_a(t)$ are scalar functions given explicitly in terms of the partition function per site (see (2.32) below). The matrices $\mathcal{X}_{a,n}^{(i,j)}$ are expressible by theta functions and derivatives. Leaving the details to Section 2.4, let us comment on the latter.

In the six-vertex case, $\mathcal{X}_{a,n}^{(i,j)}$ are defined in terms of a ‘trace’ of a monodromy matrix. Here ‘trace’ means the unique linear functional

$$\text{Tr} : U_q(\mathfrak{sl}_2) \rightarrow \mathbb{C}[q, q^{-1}] \rightarrow \mathbb{C}[q, q^{-1}];$$

which for $q \rightarrow 1$ reduces to the usual trace on the $(-1)^n$ -dimensional irreducible representation of $U_q(\mathfrak{sl}_2)$. In the eight-vertex case, we need an analogous functional Tr , defined on the Sklyanin algebra and taking values in the space of entire functions involving q , theta functions and derivatives. Compared with the trigonometric case, the existence of Tr is more difficult to establish. We do that by considering the classical limit and showing that, for generic values of the structure constants, the computation of the trace of an arbitrary monomial can be reduced to that of seven basic monomials. We have also compared our formula for Tr with the results by K. Fabricius and B. McCoy [8] for $n = 3, 4, 5$. In the classical limit, the Sklyanin algebra becomes the algebra of regular functions on an algebraic surface in \mathbb{C}^4 , which turns out to be a smoothing of a simple-elliptic singularity of K. Saito [29]. The reduction of the trace is closely connected with the de Rham cohomology of this surface (see Appendix A). Although we do not use Saito’s results for our immediate purposes, we find this connection intriguing.

¹Correlation functions of the XXZ and XXX chains are given in the limit where all the inhomogeneity parameters are chosen to be the same. However we have not succeeded in performing this homogeneous limit.

In the trigonometric case, it was shown [4] that the functions given by the above Ansatz satisfies the reduced qKZ equation. The steps of the proof carry over straightforwardly to the elliptic case, except for one property (the Cancellation identity). Unfortunately we have not succeeded in proving this last relation. It remains an open question to show that our Ansatz in the elliptic case satisfies the reduced qKZ equation.

To check the validity of the Ansatz, we examine the simplest case $n = 2$. In this case an exact answer for the homogeneous chain is obtained as derivatives of the ground state energy of the spin-chain Hamiltonian. Our formula matches with it. We also present an explicit formula for the correlators with $n = 3$. It agrees well numerically with the known integral formulas of [25, 26].

The plan of the paper is as follows. In Section 2, we introduce our notation and formulate the Ansatz for correlation functions. In Section 3, we discuss the validity of the Cancellation Identity and give arguments in its favor. Section 4 is devoted to the examples for correlators of the nearest and the next nearest neighbor spins. We also discuss briefly the trigonometric limit. In Appendix A we prove the existence of the trace functional. As was mentioned above, the classical limit of the Sklyanin algebra is related to an affine algebraic surface, and the trace functional tends to an integral over a certain cycle on it. We explain the connection between this picture and K. Saito's theory. In Appendix B, we give an explicit description of the integration cycles. Appendix C contains technical Lemmas about the trace. Finally in Appendix D we discuss the transformation properties of the matrices $X_{a,n}^{(i,j)}$.

2. Ansatz for correlation functions

In this section we introduce our notation and formulate the Ansatz for correlation functions of an inhomogeneous eight-vertex model, following the scheme developed in [4].

2.1. R matrix. We consider an elliptic R matrix depending on three complex parameters $t; q; \tau$. We assume $\text{Im } \tau > 0$ and $q \in \mathbb{C} \setminus \{0\}$. We will normally regard $q; \tau$ as fixed constants and suppress them from the notation. Let $\theta(t) = \theta(t; \tau)$ ($\theta_4(t) = \theta(t; 4\tau)$) denote the Jacobi elliptic theta functions associated with the lattice $\mathbb{Z} + \mathbb{Z}\tau$ [14]. We set

$$[t] := \frac{\theta_1(2t)}{\theta_1(2\tau)};$$

The R matrix is given by

$$(2.1) \quad R(t) := [t] \frac{r(t)}{[t+]} \text{End}(V \otimes V);$$

$$(2.2) \quad r(t) := \frac{1}{2} \sum_{n=0}^3 \frac{X^{3-n} \theta_{+1}(2t + n\tau)}{\theta_{+1}(n\tau)};$$

where $V = \mathbb{C}v_+ \oplus \mathbb{C}v_-$.

²Later on we also assume that τ is generic

The matrix $r(t) = r_{12}(t)$ is the unique entire function satisfying

$$\begin{aligned} r_{12}(0) &= P_{12}; \\ {}_1^a {}_2^a r_{12}(t) &= r_{12}(t) {}_1^a {}_2^a \quad (a = 1; 2; 3); \\ r_{12}(t + \frac{1}{2}) &= {}_1^1 r_{12}(t) {}_1^1; \\ r_{12}(t + \frac{1}{2}) &= {}_1^3 r_{12}(t) {}_1^3 e^{2i(2t + \frac{1}{2})} : \end{aligned}$$

Here $P \in \text{End}(V \otimes V)$ signifies the transposition $Pu \otimes v = v \otimes u$. As is customary, the suffix of a matrix indicates the tensor component on which it acts non-trivially, eg. ${}_1^1 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$, ${}_2^2 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$.

The normalizing factor $\gamma(t)$ is chosen to ensure that the partition function per site of the corresponding eight-vertex model equals to 1. Its explicit formula depends on the regime under consideration, and will be given later in (2.28), (2.29). In each case it satisfies

$$\gamma(t) \gamma(t) = 1; \quad \gamma(t) \gamma(t) = \frac{[t]}{[t]};$$

We will often write $t_{ij} = t_i \otimes t_j$. The basic properties of $R(t)$ are the Yang-Baxter equation

$$(2.3) \quad R_{12}(t_{12})R_{13}(t_{13})R_{23}(t_{23}) = R_{23}(t_{23})R_{13}(t_{13})R_{12}(t_{12});$$

and

$$(2.4) \quad R(t) = P R(t) P;$$

$$(2.5) \quad R(t) = 2P;$$

$$(2.6) \quad R_{12}(t)R_{21}(t) = 1;$$

$$(2.7) \quad R_{12}(t)P_{23} = R_{13}(t)P_{23}:$$

In (2.5), $P = (1 \otimes P) \in \text{End}(V \otimes V)$ denotes the projection onto the one-dimensional subspace spanned by

$$s = v_+ \otimes v_- - v_- \otimes v_+ \in V \otimes V:$$

We will use also

$$R(t) = P R(t):$$

2.2. Sklyanin algebra. Along with the R matrix, we will need the L operator whose entries are generators of the Sklyanin algebra [35, 36].

Recall that the Sklyanin algebra A is an associative unital \mathbb{C} -algebra defined through four generators S_i ($i = 0; 1; 2; 3$) and quadratic relations

$$(2.8) \quad [S_0; S_a] = iJ_{bc}(S_b S_c + S_c S_b);$$

$$(2.9) \quad [S_b; S_c] = i(S_0 S_a + S_a S_0);$$

where $(a;b;c)$ runs over cyclic permutations of $(1;2;3)$. The J_{bc} are the structure constants given by

$$(2.10) \quad J_{bc} = \frac{J_b J_c}{J_a} = \frac{1}{a} \frac{(b+1)^2 (c+1)^2}{(a+1)^2};$$

$$(2.11) \quad J_a = \frac{(a+1)(2)}{(a+1)^2};$$

where

$$J_2 = 1; \quad J_1 = 1 \quad (a \in \mathbb{Z}):$$

Here and after, theta functions without arguments stand for the theta zero values, $\theta_a = \theta_a(0)$ and $\theta_1 = \theta_1(0)$.

Since the defining relations are homogeneous, A is a \mathbb{Z}_0 -graded algebra, $A = \sum_{n \in \mathbb{Z}_0} A_n$, where the generators S_a all belong to A_1 . We have also a $\mathbb{Z}_2 \times \mathbb{Z}_2$ -grading, $A = \sum_{(m,n) \in \mathbb{Z}_2 \times \mathbb{Z}_2} A^{(m,n)}$, defined by the assignment $S \in A^{(1,0)}$, where

$$(2.12) \quad 0 = (0;0); \quad 1 = (1;0); \quad 2 = (1;1); \quad 3 = (0;1) \in \mathbb{Z}_2 \times \mathbb{Z}_2:$$

To make distinction, the \mathbb{Z}_0 -grading and the $\mathbb{Z}_2 \times \mathbb{Z}_2$ -grading will be referred to as 'degree' and 'color', respectively. Thus S has degree 1 and color 1.

There are two central elements of degree 2 and color 0,

$$(2.13) \quad K_0 := \sum_{a=0}^3 S^a; \quad K_2 := \sum_{a=1}^3 J_a S_a^2:$$

We call them Casimir elements.

Introduce the generating function (L operator)

$$L(t) := \sum_{a=0}^3 \frac{1}{2} \frac{X^a (2t + \theta_a)}{\theta_a} S^a \in \text{End}(V):$$

The defining relations (2.8), (2.9) are equivalent to

$$(2.14) \quad R_{12}(t-s) L_1(t) L_2(s) = L_2(s) L_1(t) R_{12}(t-s):$$

From (2.13) we have

$$(2.15) \quad L_1 \frac{t}{2} L_2 \frac{t}{2} \quad P_{12} = \frac{1}{4} \frac{(1-t)(1+t)}{(1)^2} K_0 + \frac{(t)^2}{(1)^2} K_2 \quad P_{12}:$$

We will be concerned with representations in series (a) of [36], which are analogs of finite-dimensional irreducible representations of sl_2 . For each non-negative integer k , let $V^{(k)}$ denote the vector space of entire functions $f(u)$ with the properties

$$f(u+1) = f(u) = f(-u); \quad f(u+\frac{1}{2}) = e^{2ik(2u+\frac{1}{2})} f(u):$$

We have $\dim V^{(k)} = k + 1$. The following formula defines a representation $\rho^{(k)} : A \rightarrow \text{End}(V^{(k)})$ [36]:³

$$(2.16) \quad \begin{aligned} \rho^{(k)}(S) f(u) &= \frac{P_{k+1}(u)}{P_1(2u)} f(u) + (2u - k) e^{\partial_u} P_{k+1}(2u - k) e^{\partial_u} f(u). \end{aligned}$$

Here $P_2 = 1$, $P_1 = 1(u - 2)$, and $(e^{\partial_u} f)(u) = f(u - 1)$. In particular, if $k = 1$, then in an appropriate basis we have

$$(1) \quad \rho(S) = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}; \quad (1) \quad \text{id } L(t) = r(t):$$

On $V^{(k)}$, the Casimir elements K_0, K_2 act as scalars $K_0(k+1), K_2(k+1)$ respectively, where

$$(2.17) \quad K_0(u) = 4 \frac{1}{P_1(2u)^2}; \quad K_2(u) = 4 \frac{1}{P_1(2u)^2} + \frac{1}{P_1(2u)}.$$

2.3. The functional Tr . In order to formulate the Ansatz, we need to consider the trace $\text{tr}_{V^{(k)}} \rho^{(k)}(A)$ of an element $A \in A$ as a function of the dimension $k + 1$. The precise meaning is as follows.

For each $A \in A$ one can assign a unique entire function $\text{Tr } A$ in \mathbb{C}^n with the following properties:

- (i) $\text{Tr } A|_{k+1} = \text{tr}_{V^{(k)}} \rho^{(k)}(A)$ holds for all $k \in \mathbb{Z}_{\geq 0}$,
- (ii) If $A \in A_n$, $\text{Tr } A$ has the functional form

$$(2.18) \quad \text{Tr } A = \frac{1}{n!} \left(g_{A,0}(t) + g_{A,1}(t) + g_{A,2}(t) + \dots \right) \quad (n: \text{odd});$$

$$(2.18) \quad \text{Tr } A = \frac{1}{n!} \left(g_{A,0}(t) + g_{A,1}(t) + g_{A,2}(t) + \dots \right) \quad (n: \text{even});$$

where $g_{A,0}(t)$, $g_{A,2}(t)$ and $g_{A,3}(t) = g_{A,1}(t+1) - g_{A,1}(t)$ are elliptic functions with periods 1; $g_{A,1}(t+1) = g_{A,1}(t)$.

For example,

$$(2.19) \quad \text{Tr } 1 = 1;$$

$$(2.20) \quad \text{Tr } S = 2 \frac{1}{P_1(2)};$$

$$(2.21) \quad \text{Tr } S^2 = \frac{2}{P_1(2)^3} (F_1(2) - F_2(2));$$

where

$$(2.22) \quad F_1(t) = \sum_{n=1}^{\infty} \frac{1}{n} \frac{\partial}{\partial t} \left(g_{A,1}(t+n) - g_{A,1}(t) \right);$$

$$(2.23) \quad F_2(t) = \sum_{n=1}^{\infty} \frac{1}{n} \frac{\partial}{\partial t} \left(g_{A,1}(t+n) - g_{A,1}(t) \right):$$

For reference we set

$$(2.24) \quad F_3(t) = \sum_{n=1}^{\infty} \frac{1}{n} \frac{\partial}{\partial t} \left(g_{A,1}(t+n) - g_{A,1}(t) \right):$$

³We have modified eq.(6) of [36] by a factor $1/P_1(2)$.

Tr satisfies also

$$(2.25) \quad \text{Tr}(AB) = \text{Tr}(BA);$$

$$(2.26) \quad \text{Tr}(K_i A) = K_i(\lambda) \text{Tr}(A) \quad (i = 0, 2);$$

$$(2.27) \quad \text{Tr} A = 0 \quad (A \in A^{(m,n)}; (m, n) \in (0, 0));$$

The derivation of (2.27) as well as (2.20), (2.21) is sketched in Appendix C. In Appendix A we show that, for generic λ , any element $A \in A^{(3)} = \bigoplus_{i=0}^3 [S_i; A]$ can be written as a $\mathbb{C}[K_0, K_2]$ -linear combination of seven monomials: $1; S_0; S_1; S_2; S_3; S_0^2; S_3^2$. Hence $\text{Tr} A$ is completely determined by the property (2.25)–(2.27) along with (2.19)–(2.21). However an effective algorithm for the reduction is not known to us. The situation is in sharp contrast to the trigonometric case, where a simple recursive procedure for calculating Tr is available (see [4]). It would be useful if one could find a more direct expression for the trace using Q -operators, as is done for the XXZ model in [22].

In this connection, notice that by the above rule, $\text{Tr} A$ has a simpler structure for elements of odd degree than those of even degree, since in the former case it is a polynomial of $\lambda(t + c)$ ($c \in \mathbb{C}$) without involving derivatives. The subtle difference between finite spin chains with odd length and those with even length has been noticed in the context of Q -operators [27, 9, 21].

2.4. The Ansatz. Consider an inhomogeneous eight vertex model, where each column i (resp. row j) carries a spectral parameter t_i (resp. 0). The Boltzmann weights are given by the entries $R_{1,1}^{0,0}; R_{1,2}^{0,0}(t_i)$ of the R -matrix (2.1). We choose the normalizing factor $\chi(t)$ in (2.1) in accordance with the two regimes

- (i) $t_i \in i\mathbb{R}, \quad \lambda > 0$ (disordered regime)
- (ii) $t_i \in \mathbb{R}, \quad \lambda < 0$ (ordered regime)

In the disordered regime, $\chi(t) = \chi^{\text{dis}}(t)$ is given by [1]

$$(2.28) \quad \chi^{\text{dis}}(t) := e^{-2i\lambda t} \frac{(2 - 2t)(4 + 2t)}{(2 + 2t)(4 - 2t)};$$

$$(u) = (u; 4; \lambda);$$

where

$$(u; \lambda; \lambda) := \prod_{j,k=0}^Y \frac{1 - e^{2i((j+1) + (k+1) - u)}}{1 - e^{2i(j+k+u)}}$$

is the elliptic gamma function [28]. In the ordered regime, the formula for $\chi(t) = \chi^{\text{ord}}(t)$ is changed to

$$(2.29) \quad \chi^{\text{ord}}(t) = e^{-4i\lambda t} \chi^0(t^0);$$

where $\chi^0(t^0)$ is given by the right hand side of (2.28) with $t; \lambda$ being replaced by

$$(2.30) \quad t^0 = \frac{t}{\lambda}; \quad \lambda^0 = -; \quad \lambda^0 = \frac{1}{\lambda};$$

respectively.

By correlation functions we mean the ground state averages $\langle h_{1^1 \dots n^1} \rangle$ of a product of spin operators on consecutive columns $1, \dots, n$ on a same row of the lattice. The thermodynamic limit is assumed. We arrange them into a matrix

$$h_n(t_1; \dots; t_n; t) = \frac{1}{2^n} \sum_{i_1, \dots, i_n \neq 0} X^{i_1 \dots i_n} h_{1^1 \dots n^1} (i_1 \dots i_n)^T \quad n)^T \in \text{End}(V^{\otimes n});$$

where $(i_1 \dots i_n)^T = (i_n \dots i_1)$ stands for the transposed matrix. Because of the \mathbb{Z} -invariance [2], it does not depend on t_i with $i < 1$ or $i > n$. When $t_1 = \dots = t_n = \bar{t}$, each $\langle h_{1^1 \dots n^1} \rangle$ is a correlation function of the infinite XYZ spin chain

$$H_{XYZ} = \sum_{j=1}^X I_j^{11} I_{j+1}^{11} + I_j^{22} I_{j+1}^{22} + I_j^{33} I_{j+1}^{33}$$

at zero temperature (the coefficients I^a will be given below in (4.5)). The h_n may be viewed as the density matrix of a finite sub-system of length n , regarding the rest of the spins as an environment.

From now on we fix n , and write $j = 2n - j + 1$. We regard h_n as a 2^{2n} -dimensional vector through the identification $\text{End}(V^{\otimes n}) \cong V^{\otimes 2n}$ given by

$$(2.31) \quad E_{i_1, i_1} \dots E_{i_n, i_n} \prod_{j=1}^n E_{i_j, i_j} \prod_{j=1}^n E_{i_j, i_j} \dots E_{i_n, i_n} \prod_{j=1}^n E_{i_j, i_j}$$

where $E_{i, i} = (E_{i, i}^0, E_{i, i}^1, E_{i, i}^2, E_{i, i}^3)$.

Let us explain the constituents which enter the Ansatz.

First, we define three functions in terms of the factor $\phi(t)$ (given in (2.28) or (2.29)) by

$$(2.32) \quad \phi_1(t) := \frac{\partial}{\partial t} \log \phi(t); \quad \phi_2(t) := \frac{\partial}{\partial t} \log \phi'(t); \quad \phi_3(t) := \frac{\partial}{\partial t} \log \phi''(t);$$

where we have set

$$\phi'(t) := \phi(t)^4 \frac{1(2 - 2t)}{1(2 + 2t)};$$

They are a meromorphic solution of the system of difference equations

$$\begin{aligned} \phi_1(t) + \phi_1(t) &= q_1(t); \\ \phi_2(t) + \phi_2(t) - \phi_1(t) &= q_2(t); \\ \phi_3(t) + \phi_3(t) &= q_3(t); \end{aligned}$$

where

$$(2.33) \quad q_1(t) := \frac{\partial}{\partial t} \log \phi(t); \quad q_2(t) := \frac{\partial}{\partial t} \log \phi'(t); \quad q_3(t) := \frac{\partial}{\partial t} \log \phi''(t);$$

and

$$\phi(t) := \frac{1(2t)^3 - 1(2t - 4)}{1(2t - 2)^3 - 1(2t + 2)};$$

The next ingredient are the matrices $\mathcal{X}_{a,n}^{(i,j)}$ ($a = 1, 2, 3, 1 \leq i \leq j \leq n$). Consider a 'transfer matrix'

$$(2.34) \quad \mathcal{X}_n(t_1; \dots; t_n); t = \frac{1}{[t_{1,2}] \prod_{p=3}^n [t_{1,p}][t_{2,p}]} \text{Tr}_{t_{1,2}} = T_n^{[1]} \frac{t_1 + t_2}{2}; t_1; \dots; t_n; t P_{12} P_{11} P_{22} :$$

We used the functional Tr introduced in the previous section, and

$$T_n^{[1]}(t; t_1; \dots; t_n) = L_2(t - t_2) \dots L_n(t - t_n) L_n(t - t_n) \dots L_2(t - t_2) :$$

Notice the presence of the permutation P_{12} and the projectors $P_{11} P_{22}$ in (2.34).

For $i < j$, we define

$$(2.35) \quad \mathcal{X}_n^{(i,j)}(t_1; \dots; t_n); t = \mathcal{X}_n^{(j,i)}(t_1; \dots; t_n); t = R_n^{(i,j)}(t_1; \dots; t_n) \mathcal{X}_n(t_1; t_j; t_1; \dots; t_i; \dots; t_j; \dots; t_n); t R_n^{(i,j)}(t_1; \dots; t_n); t^{-1} :$$

Here $R_n^{(i,j)}$ stands for the product of R matrices

$$(2.36) \quad R_n^{(i,j)}(t_1; \dots; t_n); t = R_{i,i-1}(t_{i,i-1}) R_{2,1}(t_{1,1}) R_{j,j-1}(t_{j,j-1}) R_{i+2,i+1}(t_{j,i+1}) R_{i+1,i}(t_{j,i-1}) R_{3,2}(t_{j,1}) R_{i-1,i}(t_{i-1,i}) R_{12}(t_{1,i}) R_{j-1,j}(t_{j-1,j}) R_{i+1,i+2}(t_{i+1,i,j}) R_{i,i+1}(t_{i-1,j}) R_{2,3}(t_{1,j}) :$$

Finally, for all $i \leq j$, $\mathcal{X}_{a,n}^{(i,j)}$ are defined by

$$(2.37) \quad c \mathcal{X}_{1,n}^{(i,j)}(t_1; \dots; t_n); t = \mathcal{X}_n^{(i,j)}(t_1; \dots; t_n); t t_{i,j-1} \mathcal{X}_n^{(i,j)}(t_1; \dots; t_n); t$$

$$(2.38) \quad c \mathcal{X}_{2,n}^{(i,j)}(t_1; \dots; t_n); t = \mathcal{X}_n^{(i,j)}(t_1; \dots; t_n); t$$

$$(2.39) \quad c \mathcal{X}_{3,n}^{(i,j)}(t_1; \dots; t_n); t = \mathcal{X}_n^{(i,j)}(t_1; \dots; t_n); t \mathcal{X}_n^{(i,j)}(t_1; \dots; t_n); t$$

where $c = 2 \frac{0}{1} = 1$ (2) and

$$\mathcal{X}_a^{(i)} f(t_i; t) = f(t_i + a t) f(t_i; t) :$$

As we show in Appendix D, the $\mathcal{X}_{a,n}^{(i,j)}(t_1; \dots; t_n); t$ are doubly periodic in t_k with periods $1; \dots$. The only exception is the case $a = 1$, $k = i$ or j and with respect to the shift by \dots , where the transformation law becomes

$$\mathcal{X}_{1,n}^{(i,j)}(t_1; \dots; t_n); t = \mathcal{X}_{1,n}^{(j,i)}(t_1; \dots; t_n); t \mathcal{X}_{3,n}^{(i,j)}(t_1; \dots; t_n); t$$

Conversely we have

$$\mathcal{X}_n^{(i,j)}(t_1; \dots; t_n); t = c \mathcal{X}_{1,n}^{(i,j)}(t_1; \dots; t_n); t \frac{t_{i,j}}{2} \mathcal{X}_{2,n}^{(i,j)}(t_1; \dots; t_n); t :$$

We compute the trace in the formula (2.35) by using the formulas (2.17) and (2.21). The separation of $\mathcal{X}_n^{(i,j)}$ into two parts $\mathcal{X}_{1,n}^{(i,j)}$ and $\mathcal{X}_{2,n}^{(i,j)}$ comes from that of $\text{Tr } S^2$ into $F_1(\cdot)$ and $F_2(\cdot)$. Note that $\mathcal{X}_{1,n}^{(j,i)} = \mathcal{X}_{1,n}^{(i,j)}$ and $\mathcal{X}_{a,n}^{(j,i)} = \mathcal{X}_{a,n}^{(i,j)}$ for $a = 2, 3$.

We are now in a position to state our conjecture. Let

$$S_n = \sum_{p=1}^n S_{pp}$$

be the vector corresponding to the identity by the map (2.31).

Conjecture. Correlation functions of the inhomogeneous eight-vertex model are given by the formula

$$(2.40) \quad h_n(t_1; \dots, t_n) = 2^{-n} \exp \left(\sum_{1 \leq j < a=1}^n X^3 \right) !_a(t_{ij}) \mathcal{X}_{a,n}^{(i,j)}(t_1; \dots, t_n) ; t S_n ;$$

where $!_a(t)$ and $\mathcal{X}_{a,n}^{(i,j)}$ are defined respectively by (2.32) and (2.34)–(2.39). \square

3. Reduced qKZ equation

The h_n is known to satisfy the following set of equations [16]:

$$(3.1) \quad h_n(t_1; \dots, t_{j+1}; \dots, t_n) = R_{j,j+1}(t_{j,j+1}) R_{j+1,j}(t_{j+1,j}) h_n(t_1; \dots, t_j; t_{j+1}; \dots, t_n) (1 - t_{j,j+1}^{-1});$$

$$(3.2) \quad h_n(t_1; \dots, t_j; t; \dots, t_n) = \frac{(j)}{n} A(t; \dots, t_n) h_n(t_1; \dots, t_j; t; \dots, t_n);$$

$$(3.3) \quad P_{1,l} h(t_1; \dots, t_n) ; t_{1,\dots,n,m,\dots,l} = \frac{1}{2} s_{11} h_{n-1}(t_2; \dots, t_n) ; t_{2,\dots,n,m,\dots,l} ;$$

Here

$$(3.4) \quad A_n^{(j)}(t_1; \dots, t_n) ; t = (1)^n R_{j,j-1}(t_{j,j-1}) \dots R_{j,l}(t_{j,l}) R_{j,j+1}(t_{j,j+1}) \dots R_{j,n}(t_{j,n}) P_{j,j} R_{j,n}(t_{j,n}) \dots R_{j,j+1}(t_{j,j+1}) R_{j,l}(t_{j,l}) \dots R_{j,j-1}(t_{j,j-1}) ;$$

In this section, assuming a conjectural identity, we explain that these relations are valid also for the Ansatz.

3.1. Properties of $b_n^{(i,j)}$. Consider the expression

$$b_n^{(i,j)}(t_1; \dots, t_n) = \sum_{a=1}^n X^3 !_a(t_{ij}) \mathcal{X}_{a,n}^{(i,j)}(t_1; \dots, t_n) ; t$$

which enters the Ansatz (2.40). In [4] for the XXZ model, the following relations are derived.

Exchange relation:

$$(3.5) \quad R_{k,k+1}(t_{k,k+1}) R_{k+1,k}(t_{k+1,k}) b_n^{(i,j)}(t_k; t_{k+1}; \dots, t_n) = b_n^{(i,j)}(t_k; t_{k+1}; \dots, t_n) R_{k,k+1}(t_{k,k+1}) R_{k+1,k}(t_{k+1,k});$$

Here t_k signifies the transposition $(k; k+1)$.

Besides the obvious translation invariance, $Q_n = Q_n^{(1)}$ has the following properties.

$$(3.12) \quad Q_n \left(\prod_{j=2}^n (2t_{1,j}) \right) Q_n(t_1; \dots; t_n); \text{ is entire};$$

$$(3.13) \quad Q_n \left(\prod_{j=1}^n t_j^{-\frac{1}{2}}; \dots \right) \frac{1}{j} Q_n \left(\prod_{j=1}^n t_j^{\frac{1}{2}}; \dots \right) (1 - j - n);$$

$$(3.14) \quad Q_n \left(\prod_{j=1}^n t_j^{-\frac{3}{2}}; \dots \right) \frac{3}{j} Q_n \left(\prod_{j=1}^n t_j^{\frac{3}{2}}; \dots \right) (1 - j - n);$$

$$(3.15) \quad R_{j;j+1}(t_{j;j+1}) Q_n \left(\prod_{j=1}^n t_j^{\frac{1}{2}}; \dots \right) \\ = Q_n \left(\prod_{j=1}^n t_j^{\frac{1}{2}}; \dots \right) R_{j;j+1}(t_{j;j+1}); \quad (2 - j - n - 1);$$

$$(3.16) \quad Q_n(t_1; \dots; t_n; t_n) P_{n-1, n} \big|_{t_{n-1}=t_n} = Q_{n-2}(t_1; \dots; t_n) P_{n-1, n};$$

$$(3.17) \quad t_{n-1} Q_n(t_1; \dots; t_n) \neq 0;$$

$$(3.18) \quad t_n Q_n(t_1; \dots; t_n) \neq Q_{n-1}(t_1; \dots; t_n);$$

These relations are verified in a way similar to those in [4]. The derivation of (3.13)–(3.14) rests on the transformation laws of the $X_{a,m}^{(i,j)}$, which we discuss in Appendix D.

From the properties (3.12), (3.13), (3.14), Q_n can be written as

$$(3.19) \quad \sum_{j=2}^n \prod_{j=2}^n (2t_{1,j}) Q_n(t_1; \dots; t_n); t \\ = \sum_{\substack{j=1 \\ 1; \dots; n \neq 0}}^n X^3 \prod_{j=1}^n \frac{Y^n}{j+1} \frac{(2t_{1,j})}{(j+1)^{-1}} \prod_{j=1}^n t_j^{-1};$$

with some $c_{n-1,2} \in \mathbb{C}$. Terms with $j=0$ are actually absent in the sum, in accordance with (3.17). For convenience we set $c_{n-2,0} = 0$. Note that (3.13), (3.14) and the translation invariance imply

$$(3.20) \quad c_{n-1,1} = 0 \quad \text{unless} \quad \prod_{j=1}^n P_{j-1,j} = (0;0);$$

By induction, assume $c_m = 0$ for $m < n$. We are going to argue that Q_n is then determined up to a multiplicative constant (see Lemma 3.1 below).

By (3.16), the induction hypothesis and (3.15), we have

$$(3.21) \quad Q_n \left(\prod_{j=1}^n t_j^{\frac{1}{2}}; \dots \right) P_{j;j+1} \big|_{t_j=t_{j+1}} = 0 \quad (2 - j - n - 1);$$

By (3.18) we may also assume

$$(3.22) \quad c_{n-1,1} = 0 \quad \text{unless} \quad c_n \neq 0;$$

Quite generally, consider a matrix of the form

$$U_{1,2}(u;v) = \sum_{j=0}^X \frac{+1}{+1} \frac{(2u)}{(j)} \frac{+1}{+1} \frac{(2v)}{(j)} \prod_{j=1}^2;$$

Then the relations

$$(3.23) \quad R_{12}(u-v)U_{1,2}(u;v) = U_{1,2}(v;u)R_{12}(u-v);$$

$$(3.24) \quad U_{1,2}(u+;u)P_{1,2} = 0;$$

are equivalent to the following relations for the coefficients α_a :

$$\begin{aligned} \alpha_a;0 - \alpha_0;a &= iJ_{bc}(\alpha_c;b + \alpha_b;c); \\ \alpha_b;a - \alpha_a;b &= i(\alpha_c;0 + \alpha_0;c); \\ X^3 &= 0; \\ J_a - \alpha_a;a &= 0: \end{aligned}$$

Here a,b,c are cyclic permutations of $1,2,3$, and J_a, J_{bc} are as in (2.10), (2.11). The above relations have the same form as those derived from the quadratic relations (2.8), (2.9) and from the Casimir elements (2.13), respectively. Consider the quotient A of the Sklyanin algebra modulo the relations that the Casimir elements are zero. This is a graded algebra,

$$A = \bigoplus_{n=0}^M A_n:$$

From the above considerations one easily concludes that there exist three linear functionals α_a ($a = 1,2,3$) on A_{n-1} such that

$$\alpha_a(S_n) = \alpha_a(S_{n-2})$$

which satisfy the additional condition

$$\alpha_a(S_0A) = 0:$$

In the Sklyanin algebra with generic parameter α , any monomial $S_2 \dots S_n$ can be reduced to a linear combination of ordered monomials $S_0^0 S_3^3 S_1^2 S_2^1$ with $\alpha_1; \alpha_2 \in \mathbb{C}$, by using the quadratic relations and Casimir elements (PBW basis) [10, 11]. Together with (3.20) this means that each functional α_a is defined by one constant, that is,

$$(3.25) \quad \begin{aligned} \alpha_1(S_3^{n-2}S_1); \quad \alpha_2(S_3^{n-2}S_2); \quad \alpha_3(S_3^{n-1}) & \quad \text{for } n \text{ even;} \\ \alpha_1(S_3^{n-2}S_2); \quad \alpha_2(S_3^{n-2}S_1); \quad \alpha_3(S_3^{n-3}S_1S_2) & \quad \text{for } n \text{ odd:} \end{aligned}$$

There remain three coefficients. In order to finish the proof of the Cancellation Identity, it remains to show that these coefficients vanish.

In addition to (3.15), we have also the relation

$$R_{12}(\alpha)Q_n(t_1; t_2; \alpha) = Q_n^{(2)}(t_2; t_1; \alpha):$$

The poles of the R matrix in the left hand side are not the poles of $Q_n^{(2)}$. This entails the relation

$$P_{12}Q_n(t_1; t_2; \alpha) = 0;$$

which can be rewritten in terms of functionals φ_a as follows:

$$(3.26) \quad 2\varphi_a(AS_0) - i(1 + J_{bc})\varphi_c(AS_b) + i(1 - J_{bc})\varphi_b(AS_c) = 0 \quad \forall a \in A_{n-2}:$$

These equations can be viewed as a system of linear equations for three constants (3.25). Certainly, these equations are not explicit since for every A we have to perform the procedure of reducing to PBW form. This huge system of homogeneous linear equations does not allow us to prove that the constants in question vanish; rather they reduce them to one constant. Let us explain this point. First, it is clear that the equations (3.26) correspond to the following relation in A_n

$$2S_0S_a - i(1 + J_{bc})S_bS_c + i(1 - J_{bc})S_cS_b = 0;$$

obtained by solving (2.8), (2.9) for S_0S_a . So, it is easy to see that all our equations including (3.26) are satisfied by the following construction. Consider a linear functional φ on A_n such that $\varphi(S_1 \dots S_n) = 0$ unless $\sum_{j=1}^n j = (0; 0)$, and

$$\varphi(AS_0) = 0:$$

Then all the requirements are satisfied by

$$\varphi_a(A) = \varphi(AS_a):$$

On the other hand the number of solutions to the system of linear equations (3.26) for three constants cannot be bigger for arbitrary n than it is for $n = 0$. In the latter case the algebra is commutative (see Appendix A, notably (A.4)), and the equations (3.26) become

$$\varphi_a^{cl}(S_bA) = \varphi_b^{cl}(S_aA)$$

with additional condition $\varphi_a^{cl}(S_0A) = 0$. It is easy to see that this gives a system of three equations for three constants whose rank equals 2.

Thus we come to the conclusion:

Lemma 3.1. Under the induction hypothesis, we have

$$\varphi_n(S_1 \dots S_n) = \varphi(S_n \dots S_1)$$

where φ is a linear functional on A_n satisfying

$$\varphi(S_n \dots S_1) = 0 \quad \text{unless} \quad \sum_{j=1}^n j = (0; 0)$$

$$\varphi(AS_0) = \varphi(S_0A) = 0$$

and as such is defined by one constant:

$$\begin{aligned} & \varphi(S_3^n) && \text{for } n \text{ even;} \\ & \varphi(S_3^{n-2}S_1S_2) && \text{for } n \text{ odd;} \end{aligned}$$

Unfortunately, we were not able to show that this remaining constant equals zero. The problem is still open.

4. Examples

In this section we write down the Ansatz in the simplest cases $n = 2; 3$. We also consider the trigonometric limit.

4.1. The case $n = 2$. In the case $n = 2$, ${}^{(1,2)}_2(t_1; t_2)$ can be readily found from (2.21). The function $h_2(t_1; t_2)$ is given as follows:

$$h_2(t_1; t_2) = \frac{1}{4} \sum_{a=1}^3 \frac{X^a}{4[t_{12}]} H_{a+1}(2t_{12})^{-a};$$

where

$$H_{a+1}(2t) = \frac{v_{a+1}^2(2t)}{4(v_1^2(2t))^{a+1}} v_{a+1}^0(2t) v_1(t) + \frac{v_{a+1}^0(2t)}{v_{a+1}^2(2t)} v_2(t) v_3(t);$$

This gives the formula for the nearest neighbor correlators of the inhomogeneous chain:

$$(4.1) \quad h_{12}^a = \frac{v_1^2(2t)}{v_1^2(2t)} H_{a+1}(2t);$$

where $a = 1; 2; 3$ and $t = t_{12}$. Noting that $H_{a+1}(2t)$ is odd in t , we obtain $h_{12}^a = \frac{v_1^0(2t)}{v_{a+1}^0(2t)} v_1^0(0) = \frac{v_1^0(0)}{v_{a+1}^0(0)}$ in the homogeneous limit $t \rightarrow 0$, or more explicitly we have

$$(4.2) \quad h_{12}^a = \frac{v_{a+1}^2}{8 v_1^3(2t)} \left(v_{a+1}^0(0) v_{a+1}^2(2t) + v_{a+1}^0(2t) v_{a+1}^0(0) \right) \frac{v_1^0(0)}{v_{a+1}^0(0)};$$

Let us check the formula (4.2) against known results. As is well known, the XYZ Hamiltonian is obtained by differentiating the transfer matrix of the eight-vertex model

$$T_L(t) = \text{tr} R_{0L}(t) R_{L0}(t)$$

as

$$(4.3) \quad T_L(0)^{-1} T_L^0(0) = \sum_{j=1}^L \sum_{a=1}^3 v_a^0(0) \frac{v_a^2}{v_{j+1}^2} + L v_0^0(0);$$

where L is the length of the chain, and we have set $R(t) = \prod_{j=0}^{L-1} v_j(t)$. As it was mentioned already, the R matrix (2.1) is so normalized that in the thermodynamic limit $L \rightarrow \infty$ the free energy per site of the eight-vertex model is 0. Therefore, taking the ground state average of (4.3), we obtain

$$(4.4) \quad \lim_{L \rightarrow \infty} \frac{1}{L} \ln T_L^0(0) = 0;$$

with $I = v^0(0)_{-1}(2) = \frac{0}{1}$. Explicitly we have

$$(4.5) \quad I^a = \frac{a+1}{a+1} (2) \quad (a = 1; 2; 3);$$

$$I^0 = \frac{1}{0} (2) \frac{1}{1} = \frac{1}{4} !_{-1}(0):$$

The average over the normalized ground state has the property $\langle h_{XYZ} \rangle = \langle h_{XYZ} \rangle$, where $\langle \cdot \rangle$ stands for the variation of the coefficients I^a . Hence we have in addition

$$(4.6) \quad \sum_{a=1}^3 \frac{\partial I^a}{\partial} h_{-1}^a \frac{\partial}{\partial} = \frac{\partial I^0}{\partial};$$

$$(4.7) \quad \sum_{a=1}^3 \frac{\partial I^a}{\partial} h_{-1}^a \frac{\partial}{\partial} = \frac{\partial I^0}{\partial}:$$

The nearest neighbor correlators $\langle h_{-1}^a \frac{\partial}{\partial} \rangle$ are completely determined by the linear equations (4.4), (4.6), (4.7). Using Riemann's identity and the heat equation $4 \frac{\partial}{\partial} (t_j) = \frac{\partial}{\partial} = \frac{\partial}{\partial} (t_j) = \frac{\partial}{\partial} t_j^2$, one can verify that our formula (4.2) indeed gives the unique solution.

4.2. The case $n = 3$. Let us proceed to the next case $n = 3$. Written in full, $h_3(t_1; t_2; t_3)$ reads

$$h_3(t_1; t_2; t_3) = \frac{1}{8} \frac{1}{16} \frac{1}{[t_{12}][t_{13}][t_{23}]} \sum_{\substack{(i,j,k) \in (0;0;0) \\ i+j+k=0}} X \quad \sum_{1 \leq j < k \leq 3} I_{i,j,k}^{(j,k)}(t_1; t_2; t_3):$$

The coefficients $I_{i,j,k}^{(j,k)}$ are given as follows:

$$I_{0,1,1}^{(1,2)} = 0;$$

$$I_{1,0,1}^{(1,2)} = \frac{2}{2(2)} \frac{4(2t_{13}) - 3(2t_{23})}{4(2)^3} H_3(2t_{12}) + \frac{3(2t_{13}) - 4(2t_{23})}{3(2)^4} H_4(2t_{12});$$

$$I_{1,1,0}^{(1,2)} = 2[t_{13}][t_{23}]H_2(2t_{12});$$

$$I_{1,2,3}^{(1,2)} = (-i) [t_{13}] \frac{4(2t_{23})}{4(2)} H_2(2t_{12}) - \frac{2-3(2)}{2(2)^3} \frac{4(2t_{13})}{4(2)} [t_{23}] H_3(2t_{12});$$

$$I_{1,3,2}^{(1,2)} = (-i) \frac{2-4(2)}{2(2)^4} \frac{3(2t_{13})}{3(2)} [t_{23}] H_4(2t_{12}) - [t_{13}] \frac{3(2t_{23})}{3(2)} H_2(2t_{12});$$

$$\begin{aligned}
I_{0;1;1}^{(1;3)} &= I_{1;1;0}^{(1;3)} = 0; \\
I_{1;0;1}^{(1;3)} &= \frac{\frac{2}{3} \binom{2}{4} \frac{2}{4} + \frac{2}{4} \binom{2}{4} \frac{2}{3}}{\binom{2}{3} \binom{2}{4} \binom{2}{4} \binom{2}{4}} [t_{12}] [t_{23}] H_2(2t_{13}) \\
&\quad - \frac{2}{2 \binom{2}{2}} \frac{4 \binom{2}{4} \binom{2}{4} \binom{2}{4}}{4 \binom{2}{4} \binom{2}{4}} H_3(2t_{13}) + \frac{3 \binom{2}{4} \binom{2}{4} \binom{2}{4}}{3 \binom{2}{4} \binom{2}{4}} H_4(2t_{13}) ; \\
I_{1;2;3}^{(1;3)} &= (i) \frac{2}{3} \frac{3 \binom{2}{4} \binom{2}{4}}{2 \binom{2}{4}} [t_{23}] H_4(2t_{13}) - [t_{12}] \frac{4}{3} \frac{3 \binom{2}{4} \binom{2}{4}}{4 \binom{2}{4} \binom{2}{4}} H_2(2t_{13}) ; \\
I_{1;3;2}^{(1;3)} &= (i) [t_{12}] \frac{3}{4} \frac{4 \binom{2}{4} \binom{2}{4}}{3 \binom{2}{4} \binom{2}{4}} H_2(2t_{13}) - \frac{2}{4} \frac{4 \binom{2}{4} \binom{2}{4}}{2 \binom{2}{4} \binom{2}{4}} [t_{23}] H_3(2t_{13}) ; \\
I_{1;1;0}^{(2;3)} &= 0; \\
I_{1;0;1}^{(2;3)} &= \frac{2}{2 \binom{2}{2}} \frac{3 \binom{2}{4} \binom{2}{4} \binom{2}{4}}{3 \binom{2}{4} \binom{2}{4}} H_3(2t_{23}) + \frac{4 \binom{2}{4} \binom{2}{4} \binom{2}{4}}{3 \binom{2}{4} \binom{2}{4}} H_4(2t_{23}) ; \\
I_{0;1;1}^{(2;3)} &= 2 [t_{12}] [t_{13}] H_2(2t_{23}); \\
I_{1;2;3}^{(2;3)} &= (i) \frac{3 \binom{2}{4} \binom{2}{4} \binom{2}{4}}{3 \binom{2}{4} \binom{2}{4} \binom{2}{4}} [t_{12}] H_3(2t_{23}) - [t_{13}] \frac{2 \binom{2}{4} \binom{2}{4}}{2 \binom{2}{4} \binom{2}{4}} H_4(2t_{23}) ; \\
I_{1;3;2}^{(2;3)} &= (i) [t_{13}] \frac{2 \binom{2}{4} \binom{2}{4}}{2 \binom{2}{4} \binom{2}{4}} H_3(2t_{23}) - \frac{3 \binom{2}{4} \binom{2}{4} \binom{2}{4}}{3 \binom{2}{4} \binom{2}{4} \binom{2}{4}} \frac{2 \binom{2}{4} \binom{2}{4}}{2 \binom{2}{4} \binom{2}{4}} [t_{12}] H_4(2t_{23}) :
\end{aligned}$$

The rest are given by the cyclic change $1 \rightarrow 2 \rightarrow 3 \rightarrow 1$ of the indices i, j, k in $I_{i;j;k}^{(j;k)}$ with the change $2 \rightarrow 3 \rightarrow 4 \rightarrow 2$ of the indices in a and H_a . The correlators of the inhomogeneous chain are

$$h_{1 \ 2 \ 3}^{(j;k)} = \frac{1}{2 [t_{12}] [t_{13}] [t_{23}]} \sum_{j < k \leq 3} I_{i;j;k}^{(j;k)}(t_1; t_2; t_3);$$

With the abbreviation $H_a^0 = H_a^0(0)$ and $H_a^{\infty} = H_a^{\infty}(0)$, we obtain a new formula for the next nearest neighbor correlators for the homogeneous chain

$$\begin{aligned}
h_{1 \ 3}^{a \ a} &= \frac{1}{4} \frac{\binom{2}{1}^n}{\binom{0}{1}} \frac{2 \binom{2}{b+1} \binom{2}{c+1} + \binom{2}{c+1} \binom{2}{b+1}}{\binom{2}{b+1} \binom{2}{c+1} \binom{2}{b+1} \binom{2}{c+1}} H_{a+1}^0 \\
&\quad + \frac{1}{\binom{0}{1}} \frac{\binom{2}{a+1}}{\binom{2}{a+1} \binom{2}{c+1}} \frac{n}{\binom{2}{c+1}} \left(\frac{\binom{0}{b+1}}{\binom{2}{b+1}} H_{b+1}^0 + \frac{2 \binom{0}{c+1}}{\binom{2}{c+1}} H_{b+1}^0 - H_{b+1}^{\infty} \right) \\
&\quad + \frac{\binom{0}{b+1}}{\binom{2}{b+1} \binom{2}{c+1}} \left(\frac{\binom{0}{c+1}}{\binom{2}{c+1}} H_{c+1}^0 + \frac{2 \binom{0}{b+1}}{\binom{2}{b+1}} H_{c+1}^0 - H_{c+1}^{\infty} \right) :
\end{aligned}$$

We have in addition

$$h_{1 \ 2 \ 3}^{a \ b \ c} = 0; \quad h_{1 \ 2 \ 3}^{c \ b \ a} = 0;$$

In both formulas, $(a; b; c) = (1; 2; 3); (2; 3; 1); (3; 1; 2)$.

M. Lashkevich communicated to us a program for numerically calculating correlation functions from the integral formula of [25, 26]. For $n = 2$ and $n = 3$, we found agreement between their results and ours to within the precision 10^{-4} .

4.3. Trigonometric limit. Finally we briefly touch upon the trigonometric limit, and discuss how various quantities which appear in (4.1) are related to the trigonometric counterpart.

First we consider the limit to the massive regime. For this purpose, it is convenient to rewrite the R matrix in terms of the parameters $t; q; \hbar$ in (2.30) as

$$R(t) = \frac{t^0}{t^0 + \hbar^0} (U - U)r^0(t^0) (U - U)^{-1};$$

where $t^0 = \frac{1}{2}(2t^0)^0 = \frac{1}{2}(2^0 j^0)$, $r^0(t^0)$ is obtained from (2.2) by replacing $t; q; \hbar$ by $t^0; q^0; \hbar^0$, and $U = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$. In the limit $\hbar \rightarrow +i\hbar$ while keeping $t = t^0$ and $q = q^0$ fixed, the R matrix tends to

$$(4.8) \quad R_{XXZ}(\hbar) = r_{XXZ}(\hbar) \frac{r_{XXZ}(\hbar)}{[\hbar + 1]_{XXZ}};$$

In the above, $[\hbar]_{XXZ} = \sin \frac{\hbar}{2} = \sin \frac{\hbar}{2}$, and

$$(4.9) \quad r_{XXZ}(\hbar) = \frac{1}{2} \frac{\sin(\hbar + 1/2)}{\sin \hbar/2} \hbar^0 \hbar^0 + \frac{1}{2} \hbar^0 \hbar^0 + \frac{1}{2} \hbar^0 \hbar^0 + \frac{\cos(\hbar + 1/2)}{\cos \hbar/2} \hbar^3 \hbar^3;$$

$$(4.10) \quad r_{XXZ}(\hbar) = \frac{(q^2 - 2)_1 (\hbar^2)_1}{(q^2 - 2)_1 (\hbar^2)_1};$$

where $\hbar = e^i$, $q = e^i$, $(x)_1 = \prod_{j=0}^{\infty} (1 - q^{4j} x)$. It is easy to see that

$$\hbar_1(t) = \frac{8}{\sin \hbar} \hbar(\hbar); \quad \hbar_2(t) = \frac{8}{\sin \hbar} \hbar(\hbar); \quad \hbar_3(t) = 0;$$

where the functions $\hbar(\hbar); \hbar(\hbar)$ are given by [4], eqs.(13.2){(13.5) for the massive regime. Hence the limit of (4.1) becomes

$$(4.11) \quad \lim_{\hbar \rightarrow i\hbar} h_{1/2}^{3/2} i = \frac{4}{(q - q^{-1})^2} \hbar(\hbar) + \frac{\hbar + 1}{(q - q^{-1})(\hbar - 1)} \hbar(\hbar);$$

$$\lim_{\hbar \rightarrow i\hbar} h_{1/2}^{1/2} i = \lim_{\hbar \rightarrow i\hbar} h_{1/2}^{2/2} i = 2 \frac{\hbar + 1}{(q - q^{-1})^2} \hbar(\hbar) + \frac{q + q^{-1}}{(q - q^{-1})(\hbar - 1)} \hbar(\hbar);$$

which reproduces the formulas in the massive regime (see [4], Example in Section 3).

Second let us consider the limit to the massless regime. We set

$$\hbar = \frac{1}{i} r; \quad \hbar = \frac{1}{2i} r; \quad t = \frac{1}{2i} r$$

for a constant $(0 < r < 1)$ and take the limit $r \neq 0$ with \hbar and t fixed.

The limit of the R matrix is given by the same formula (4.8){(4.9), with $r_{XXZ}(\hbar)$ being replaced by

$$r_{XXZ}(\hbar) = \frac{S_2(\hbar) S_2(1 + \hbar)}{S_2(\hbar) S_2(1 - \hbar)};$$

Here $S_2(x) = S_2(x; 1)$ signifies the double sine function. In the limit we have

$$r!_1(t) \sim \frac{8^{-2i}}{\sin} !(\cdot); \quad r!_2(t) \sim \frac{8^{-2i}}{\sin} \ddagger(\cdot);$$

where now $!(\cdot)$ and $\ddagger(\cdot)$ stand for the functions given by [4], eqs. (13.2)–(13.5) for the massless regime. Moreover we have

$$re^{\frac{2}{r}} \left(t \frac{\partial}{\partial t} + \frac{\partial}{\partial} + \frac{\partial}{\partial} \log' \right) = 0.$$

From the formulas above, we see that in the massless limit the function $h_2(t_1; t_2)$ tends to the solution $h_2(\cdot; \cdot)$ of the reduced qKZ equation given in [4].

Appendix A. Existence of Tr

For every finite-dimensional representation of the Sklyanin algebra A , we can define the trace, which is a functional on A whose main property is cyclicity. In order to formulate our Ansatz for correlation functions, we need an analytic continuation of this functional with respect to the dimension. We denote this analytic continuation by $\text{Tr } A$, where $A \in A$ and $\dim A = k + 1$ for $(k) \in A$. In Section 2.3, we presented the formulas for $\text{Tr } S$, $\text{Tr } S^2$ ($\dim A = 0; 1; 2; 3$). In this appendix, we discuss the general case. In fact, we prove that for generic parameters J_1, J_2, J_3 , the definition of $\text{Tr } A$ for general $A \in A$ can be reduced to these known cases.

Consider the polynomial ring $F = K[K_0; K_2]$ with $K = \mathbb{C}(J_1; J_2; J_3)$. Here, we consider $K_0; K_2; J_1; J_2; J_3$ as variables, whereas they are parameterized by θ, ϕ and ψ in Section 2 and Appendix B. We use the parameterization in order to define finite-dimensional representations. The discussion in this appendix is mainly concerned with the algebraic relations in the Sklyanin algebra only.

We denote by A the Sklyanin algebra defined over the field K . It is a graded vector space,

$$A = \bigoplus_{n=0}^{\infty} A_n; \quad \dim A_n < \infty.$$

Multiplication by the central elements (2.13) endows A with an F -algebra structure. Suppose we try to define some F -linear functional Tr on A which satisfies cyclicity $\text{Tr}(AB) = \text{Tr}(BA)$. Then the question is, for how many independent elements of A this functional should be defined. In other words, describe the F -module

$$H = A = A^0$$

where

$$A^0 = \bigoplus_{n=0}^{\infty} A_n = \bigoplus_{n=0}^{\infty} [S; A_n]:$$

Note that $H = \bigoplus_{n=0}^{\infty} H_n$ where $H_n = A_n = A_n^0$, $A_n^0 = \bigoplus_{i=0}^n [S; A_{n-i}]$.

We prove

Theorem A.1. The F -module H is a rank 7 free module generated by the monomials

$$(A.1) \quad (m_i)_{1 \leq i \leq 7} = (1; S_0; S_1; S_2; S_3; S_0^2; S_3^2):$$

The F -linear independence of these elements follows from (2.20)-(2.23). Indeed, suppose there is a relation $\sum_{i=1}^7 c_i m_i = 0$ with $c_i \in F$. The sum over elements of even degree and of odd degree must vanish separately. Specialize J_i to the value (2.11) with $\theta \in \mathbb{Q} + \mathbb{Q}\sqrt{3}$, $\text{Im } \theta > 0$, and take the trace of both sides on the representation $V^{(k)}$ for $k \in \mathbb{Z}_{\geq 0}$. By Lemma C.2, it follows that $c_i = 0$ except for $i = 3, 4, 5$. To see that the latter vanish, it is enough to apply the automorphism s^{-1} (see (D.1), (D.2) and two lines above) and take the trace.

Let us prove the spanning property.

Consider the tensor algebra T over the field K generated by four independent variables S_0, S_1, S_2 and S_3 . Set $R = T[[K_0; K_2]]$. It is a graded algebra: $R = \sum_{n=0}^{\infty} R_n$, where we have $\dim_K R_n < \infty$. We have the isomorphism of K -vector spaces

$$(A.2) \quad \begin{aligned} H_n \subset R_n = & \sum_{a=0}^n R_{n-2a} [S_0; S_a] iJ_{b;c} (S_b S_c + S_c S_b) \\ & + \sum_{a=0}^n R_{n-2a} [S_b; S_c] i(S_0 S_a + S_a S_0) + \sum_{a=1}^n [S_a; R_{n-1}] \\ & + \sum_{a=0}^n R_{n-2a} S^2 K_0 + \sum_{a=1}^n R_{n-2a} J_a S_a^2 K_2 : \end{aligned}$$

The K -vector space R_n is spanned by the monomials of the form $K_0^{m_0} K_2^{m_2} S_1^{m_1} S_3^{m_3}$ where $2m_0 + 2m_2 + m_1 + m_3 = n$. The relations which define H_n in (A.2) are linear relations for these monomials. For each n the coefficients of these linear relations form a matrix M_n with entries in K .

The spanning property is clear for $n = 0, 1$. Suppose that $n \geq 2$. Divide the set of monomials of degree n into two groups: the first group is the monomials such that the part $S_1^{m_1} S_3^{m_3}$ is equal to one of m_i ($1 \leq i \leq 7$) and the second group is the rest. The matrix M_n is divided into two blocks $M_n = (M_n^{(0)}; M_n^{(1)})$ where $M_n^{(0)}$ (resp., $M_n^{(1)}$) corresponds to the first (resp., the second) group of monomials. It suffices to show that the rank of $M_n^{(1)}$ is equal to the cardinality of the second group.

The proof of this statement exploits the classical limit:

$$(A.3) \quad J_a = 1 - \hbar^2 j_a; \quad \hbar \rightarrow 0:$$

We introduce new variables s_i ($i = 0, 1, 2, 3$) and k_0, k_1 by

$$(A.4) \quad S_0 = \hbar s_0; \quad S_a = s_a \quad (a = 1, 2, 3);$$

$$(A.5) \quad K_0 = k_0; \quad K_2 = k_0 - \hbar^2 k_1;$$

In Appendix B, the classical limit is taken as $\hbar \rightarrow 0$ instead of $\hbar \rightarrow 0$. In this appendix, we avoid the parametrization by θ , ϕ and ψ in order to simplify the argument.

In the limit $\hbar \rightarrow 0$, we have

$$(A.6) \quad [s_i; s_j] = i\hbar f_{ij} s_i s_j + O(\hbar^2);$$

where the Poisson bracket is defined by

$$(A.7) \quad f s_0; s_a g = 2 j_{b;c} s_b s_c i$$

$$(A.8) \quad f s_b; s_c g = 2 s_0 s_a i$$

Here $j_{ab} = j_{ba} = j_a j_b$ and $(a;b;c)$ runs over cyclic permutations of $(1;2;3)$. In the classical limit, the variables s_a become commutative. Let $K^{cl} = C(j_1; j_2; j_3)$ denote the field of rational functions in j_a , and let $F^{cl} = K^{cl}[k_0; k_1]$, $A^{cl} = K^{cl}[s_0; s_1; s_2; s_3]$ denote the polynomial ring in indeterminates k_0, k_1 and s_0, \dots, s_3 , respectively. The Casimir relations (2.13) become the algebraic relations

$$(A.9) \quad s_1^2 + s_2^2 + s_3^2 = k_0;$$

$$(A.10) \quad s_0^2 + j_1 s_1^2 + j_2 s_2^2 + j_3 s_3^2 = k_1$$

in $K^{cl}[s_0; s_1; s_2; s_3; k_0; k_1]$:

$$A^{cl} \subset K^{cl}[s_0; s_1; s_2; s_3; k_0; k_1] \models (s_1^2 + s_2^2 + s_3^2 - k_0; s_0^2 + j_1 s_1^2 + j_2 s_2^2 + j_3 s_3^2 - k_1):$$

This makes A^{cl} an F^{cl} -algebra. The algebra A^{cl} is graded as well: $A^{cl} = \sum_{n=0}^1 A_n^{cl}$. Let $(A_n^{cl})_n^0 = A_n^{cl}$ be the limit (in the appropriate Grassmannian, such that we consider s_a as commutative variables) of A_n^0 as $\hbar \rightarrow 0$. Then, we see $\sum_{n=0}^3 f s_n; A_{n-1}^{cl} g$ $(A_n^{cl})_n^0$ from (A.6). In order to show the spanning property of $(m_i)_{1 \leq i \leq 7}$ in H , it is therefore sufficient to show the spanning property for

$$(A.11) \quad (m_i)_{1 \leq i \leq 7} = (1; s_0; s_1; s_2; s_3; s_0^2; s_3^2):$$

in $H^{cl} = \sum_{n=0}^1 H_n^{cl}$, where

$$(A.12) \quad H_n^{cl} = A_n^{cl} = \sum_{X^3=0} f s_n; A_{n-1}^{cl} g:$$

In conclusion, Theorem A.1 follows from

Proposition A.2. The F^{cl} -module H^{cl} is a rank 7 free module generated by the monomials (A.11).

Proof. The F^{cl} -linear independence of m_i^{cl} follows from the same argument as in the quantum case. In place of trace on $V^{(k)}$ we use the non-degenerate pairing between cycles and cocycles given in Appendix B.

We prove the spanning property. Define

$$r_0 = j_{2,3} s_2 s_3 \frac{\partial}{\partial s_1} + j_{3,1} s_3 s_1 \frac{\partial}{\partial s_2} + j_{1,2} s_1 s_2 \frac{\partial}{\partial s_3};$$

$$r_a = j_{b,c} s_b s_c \frac{\partial}{\partial s_0} + s_0 s_c \frac{\partial}{\partial s_b} - s_0 s_b \frac{\partial}{\partial s_c};$$

where $(a;b;c) = (1;2;3), (2;3;1), (3;1;2)$. We have $r \cdot P = \frac{1}{2} f s; P g$. These are F^{cl} -linear.

We want to show that modulo $\sum_{P=0}^3 r \cdot A^{cl}$ any monomial $s_0^{m_0} s_1^{m_1} s_2^{m_2} s_3^{m_3}$ can be reduced to an element in $F^{cl} \cdot 1 + \sum_{P=0}^3 F^{cl} \cdot s + \sum_{P=0}^3 F^{cl} \cdot s^2$. Set

$$A^{cl(i-1)} = \sum_{0 \leq X \leq 3} s \cdot s \cdot A^{cl};$$

$$H^{cl(i-1)} = A^{cl(i-1)} = \sum_{X^3=0} r \cdot A^{cl};$$

We also denote $H^{\text{cl}[0]} = H^{\text{cl}}$ and $A^{\text{cl}[0]} = A^{\text{cl}}$.

Since

$$A^{\text{cl}} = \sum_{n=0}^{\infty} \frac{X^3}{n!} K^{\text{cl}} S^n A^{\text{cl}[1]};$$

by using (A.9) and (A.10) the above statement follows from the following:

$$(A.13) \quad H^{\text{cl}[1]} = F^{\text{cl}} k_0 s_0 + \sum_{a=1}^3 F^{\text{cl}} (k_1 - j_a k_0) s_a + \sum_{a=1}^3 F^{\text{cl}} (k_1 - j_a k_0) s_a^2;$$

Note that

$$k_1 - j_a k_0 = s_0^2 + j_{b;a} s_b^2 + j_{c;a} s_c^2;$$

and therefore we have $k_0 s_0, (k_1 - j_a k_0) s_a, (k_1 - j_a k_0) s_a^2 \in A^{\text{cl}[1]}$.

Let us prove (A.13). Suppose that a monomial $m = s_0^{n_0} s_1^{n_1} s_2^{n_2} s_3^{n_3} \in H^{\text{cl}[1]}$ is such that $j_a n_a \in 2\mathbb{Z} + 1$: e.g., $n_0, n_1 \in 2\mathbb{Z} + 1$. By using the Casimir relations (A.9) and (A.10) we can replace s_0^2 and s_1^2 with s_2^2 and s_3^2 . Therefore, we have

$$m \in s_0 s_1 F^{\text{cl}}[s_2; s_3];$$

Since

$$r_2(s_2^j s_3^k) = k s_0 s_1 s_2^j s_3^{k-1};$$

we have

$$m = 0;$$

Next consider the case $j_a n_a \in 2\mathbb{Z} + 1$, $g = 1$. Suppose $n_0 \in 2\mathbb{Z} + 1$. Then, we have

$$m \in s_0 s_2^2 F^{\text{cl}}[s_2^2; s_3^2] + s_0 s_3^2 F^{\text{cl}}[s_2^2; s_3^2];$$

Since

$$r_1(s_1^i s_2^j s_3^k) = l s_0 s_1^i s_2^{j-1} s_3^{k+1} - k s_0 s_1^i s_2^{j+1} s_3^{k-1};$$

the monomial m belongs to $K^{\text{cl}} s_0 s_2^{2j} \in H^{\text{cl}[1]}$ where $2j + 1 = n_0 + n_1 + n_2 + n_3$. Similarly, we see that

$$k_0^j s_0 \in K^{\text{cl}} s_0 s_2^{2j};$$

Since $k_0^j s_0$ is a non-zero element in $H^{\text{cl}[1]}$, we have

$$m \in K^{\text{cl}} k_0^j s_0;$$

The case $n_a \in 2\mathbb{Z} + 1$ ($a = 1, 2, 3$) is similar.

The remaining case is $n_0, n_1, n_2, n_3 \in 2\mathbb{Z}$ and $j_a n_a > 0$. We have $\deg m = 4$. Note that

$$H^{\text{cl}[4]} = \sum_{0 \leq i < j \leq 3} K^{\text{cl}} s_i^2 s_j^2;$$

We have the following relations in $H_4^{\text{cl}[-1]}$:

$$\begin{aligned} r_1(s_0 s_2 s_3) &= j_{2,3} s_2^2 s_3^2 + s_0^2 s_3^2 - s_0^2 s_2^2; \\ r_2(s_0 s_3 s_1) &= j_{3,1} s_3^2 s_1^2 + s_0^2 s_1^2 - s_0^2 s_3^2; \\ r_3(s_0 s_1 s_2) &= j_{1,2} s_1^2 s_2^2 + s_0^2 s_2^2 - s_0^2 s_1^2. \end{aligned}$$

Therefore, we have $\dim_K \text{cl} H_4^{\text{cl}[-1]} = 3$. On the other hand $(k_1 - j_a k_0) s_a^2 \in H_4^{\text{cl}[-1]}$ for $a = 1, 2, 3$, and they are K^{cl} -linearly independent. Therefore, we have

$$H_4^{\text{cl}[-1]} = \sum_{a=1}^3 K^{\text{cl}} (k_1 - j_a k_0) s_a^2.$$

Observe that for $(a; b; c) = (1; 2; 3), (2; 3; 1), (3; 1; 2)$ we have

$$\begin{aligned} r_a(s_0^{n_0+1} s_1^{n_1} s_2^{n_2} s_3^{n_3}) &= (n_0 + 1) j_{b,c} s_0^{n_0} s_a^{n_a} s_b^{n_b+1} s_c^{n_c+1} \\ &\quad + n_b s_0^{n_0+2} s_a^{n_a} s_b^{n_b-1} s_c^{n_c+1} - n_c s_0^{n_0+2} s_a^{n_a} s_b^{n_b+1} s_c^{n_c-1}. \end{aligned}$$

We can increase the power in s_0 by rewriting $s_0^{n_0} s_a^{n_a} s_b^{n_b+1} s_c^{n_c+1}$ in terms of $s_0^{n_0+2} s_a^{n_a} s_b^{n_b-1} s_c^{n_c+1}$ and $s_0^{n_0+2} s_a^{n_a} s_b^{n_b+1} s_c^{n_c-1}$. Thus, we see that the K^{cl} vector space $H_{2j}^{\text{cl}[-1]} (j \geq 3)$ is spanned by

$$s_0^{2d} s_a^{2(j-d)} \quad (a = 1; 2; 3; 1 - d - j - 1):$$

On the other hand, the space $H_{2j}^{\text{cl}[-1]}$ contains K^{cl} -linearly independent elements

$$(A.14) \quad k_0^d k_1^{j-d-2} (k_1 - j_a k_0) s_a^2 \quad (a = 1; 2; 3; 0 - d - j - 2):$$

Therefore, the elements (A.14) span $H_{2j}^{\text{cl}[-1]}$. \square

In the rest of this appendix, we explain a mathematical background of Proposition A.2, which is the de Rham cohomology of the affine algebraic variety defined by the two quadrics (A.9) and (A.10). Although our proof is independent, the statement of Proposition A.2 is closely related to a result of K. Saito.

We set

$$M = f(j_1; j_2; j_3) \in C^3 \text{ for } a \in \mathbb{Z}:$$

Let

$$(A.15) \quad \gamma : X = C^4 \times M \rightarrow Y = C^2 \times M$$

be the mapping such that $\gamma = (\gamma_1; \dots; \gamma_5)$ and for $(s; j) = (s_0; s_1; s_2; s_3; j_1; j_2; j_3) \in C^4 \times M$ we have

$$(A.16) \quad \gamma_1(s; j) = s_1^2 + s_2^2 + s_3^2;$$

$$(A.17) \quad \gamma_2(s; j) = s_0^2 + j_1 s_1^2 + j_2 s_2^2 + j_3 s_3^2;$$

$$(A.18) \quad \gamma_3(s; j) = j_1;$$

$$(A.19) \quad \gamma_4(s; j) = j_2;$$

$$(A.20) \quad \gamma_5(s; j) = j_3;$$

The critical set $C \subset C^4 \times M$ of this mapping is given by the equation

$$d\gamma_1 \wedge \dots \wedge d\gamma_5 \neq 0:$$

We have

$$C = \sum_{j=0}^3 f(s; j) j^s = 0 \text{ for } s \notin g;$$

We have the commutative diagram ,

$$\begin{array}{ccc} X & & C \\ \downarrow & & \downarrow \\ Y & & Y \\ \downarrow & & \downarrow \\ Y & & D; \end{array}$$

where the discriminant set D is given by

$$(A.21) \quad D = \{f(k_0; k_1; j_1; j_2; j_3) \neq 0 \mid f(k_0; k_1; j_1; j_2; j_3) = 0\};$$

$$(A.22) \quad = \sum_{a=1}^3 k_0 (k_1 - j_a k_0):$$

The inverse image $\pi^{-1}(0)$ is called the simple elliptic singularity of type D_5 [29]. If $y \in Y$ does not belong to D , the inverse image $X_y = \pi^{-1}(y)$ is a non-singular affine complex surface and is called a smoothing of the singularity. The mapping

$$(A.23) \quad \pi: X \rightarrow Y$$

is a locally topologically trivial fiber space, and the homology group is of rank 7: $H_2(X_y; \mathbb{Z}) = \mathbb{Z}^7$. In Appendix B, we construct cycles in $H_2(X_y; \mathbb{Z})$. (In [31], K. Saito defined the extended affine root systems. The homology group $H_2(X_y; \mathbb{Z})$ is isomorphic to $D_5^{(1;1)}$ in his classification. We have not identified our cycles in $D_5^{(1;1)}$.)

Let \mathcal{O}_X^p be the sheaf of \mathcal{O}_X modules consisting of germs of holomorphic p -forms on X , and $\mathcal{O}_{X=Y}^p$ the quotient sheaf

$$(A.24) \quad \mathcal{O}_{X=Y}^p = \mathcal{O}_X^p / \sum_{i=1}^5 d_i^p \wedge \mathcal{O}_X^{p-1};$$

The relative de Rham complex $(\mathcal{O}_{X=Y}^p; d_{X=Y})$ is defined by the commutative diagram

$$\begin{array}{ccc} d_{X=Y} : \mathcal{O}_{X=Y}^p & \rightarrow & \mathcal{O}_{X=Y}^{p+1} \\ & \searrow & \searrow \\ d : \mathcal{O}_X^p & \rightarrow & \mathcal{O}_X^{p+1} \end{array}$$

This is an exact sequence of \mathcal{O}_X modules. The following is K. Saito's result [32].

Theorem A.3. The cohomology group

$$H^2(\pi^{-1}(\mathcal{O}_{X=Y}^p)) = \text{Ker}(\pi^{-1}(\mathcal{O}_{X=Y}^2) \xrightarrow{d_{X=Y}} \pi^{-1}(\mathcal{O}_{X=Y}^3)) = \text{Im}(\pi^{-1}(\mathcal{O}_{X=Y}^1) \xrightarrow{d_{X=Y}} \pi^{-1}(\mathcal{O}_{X=Y}^2))$$

is an \mathcal{O}_Y -locally free module of rank 7.

We connect the above algebro-geometric setting to ours. The following proposition is a corollary to Theorem A.3. Here we give a proof in the line of this appendix without using Saito's result.

Proposition A.4. Consider a complex of F^{cl} -modules:

$$\begin{aligned} Z^P &= \bigoplus_{0 \leq i_1 < i_2 < i_3}^M A^{\text{cl}} ds_{i_1} \wedge \dots \wedge ds_{i_3} \\ Z^P &= Z^P = \bigoplus_{j=1,2}^X d'_{i_j} \wedge Z^{P-1}: \end{aligned}$$

The cohomology group $H^{\text{cl}[2]} \stackrel{\text{def}}{=} \text{Ker}(Z^2 \xrightarrow{d} Z^3) = \text{Im}(Z^1 \xrightarrow{d} Z^2)$ is a F^{cl} -free module of rank 7, where the action of k_0 (resp., k_1) is given by the multiplication of $'_1$ (resp., $'_2$).

Proof. The key idea of the proof is to identify $A^{\text{cl}[1]}$ with Z^2 .

Let us introduce a holomorphic section $!$ of the sheaf $\Omega_{X/Y}^2$.

$$\begin{aligned} ! &= \frac{ds_1 \wedge ds_2}{s_0 s_3} = \frac{ds_2 \wedge ds_3}{s_0 s_1} = \frac{ds_3 \wedge ds_1}{s_0 s_2} \\ &= \frac{ds_0 \wedge ds_1}{j_{2,3} s_2 s_3} = \frac{ds_0 \wedge ds_2}{j_{3,1} s_3 s_1} = \frac{ds_0 \wedge ds_3}{j_{1,2} s_1 s_2} \end{aligned}$$

Consider the F^{cl} -module

$$\tilde{A}^{\text{cl}} \stackrel{\text{def}}{=} K^{\text{cl}}_{C[j_1, j_2, j_3]} C[s_0; s_1; s_2; s_3; j_1; j_2; j_3]!$$

We have a canonical isomorphism of F^{cl} -modules

$$A^{\text{cl}} \rightarrow \tilde{A}^{\text{cl}};$$

sending $P \in A^{\text{cl}}$ to $P! \in \tilde{A}^{\text{cl}}$. It is easy to see that

$$A^{\text{cl}[1]}! = Z^2;$$

We have

$$(A.25) \quad d(P ds) = r(P)!$$

for $P \in A^{\text{cl}}$. From (A.25) we have

$$\text{Im}(Z^1 \xrightarrow{d} Z^2) = \bigoplus_{i=0}^3 r(A^{\text{cl}})!$$

We have already constructed the F^{cl} -bases of the modules $A^{\text{cl}} = \bigoplus_{i=0}^3 r(A^{\text{cl}})$ and $A^{\text{cl}[1]} = \bigoplus_{i=0}^3 r(A^{\text{cl}})$. We will construct a basis of $\text{Ker}(Z^2 \xrightarrow{d} Z^3) = \text{Im}(Z^1 \xrightarrow{d} Z^2)$. First, observe that

$$Z^3 = \bigoplus_{n=0}^M K^{\text{cl}} s_0^n ds_1 \wedge ds_2 \wedge ds_3 + \bigoplus_{a=1}^3 \bigoplus_{n=0}^M K^{\text{cl}} s_a^n ds_0 \wedge ds_b \wedge ds_c$$

where $(a; b; c) = (1; 2; 3), (2; 3; 1), (3; 1; 2)$. From this we see that

$$\begin{aligned} \text{Ker}(\mathbb{Z}^2 \xrightarrow{d} \mathbb{Z}^3) = & \sum_{0 \leq a \leq b \leq c} A^{cl} s_a s_b s_c! \sum_{\substack{m, n \geq 0 \\ m+n=2}} K^{cl} s_a^m s_b^n! \\ & + \sum_{\substack{n \geq 1 \\ n \neq 2}} \sum_{1 \leq a \leq b \leq 3} K^{cl} (s_0^2 + j_{ab} s_b^2) s_a^n! + \sum_{1 \leq a \leq 2} K^{cl} (s_a^2 + s_{a+1}^2) s_0^n! \\ & + \sum_{1 \leq a < b \leq 3} K^{cl} (j_{ab} s_a^2 s_b^2 + s_0^2 s_a^2 + s_0^2 s_b^2)! : \end{aligned}$$

We know that the F^{cl} -module $\mathbb{Z}^2 = \text{Im}(\mathbb{Z}^1 \xrightarrow{d} \mathbb{Z}^2)$ has the free generators $s_0 = k_0 s_0!$, $s_a = (k_1 - j_a k_0) s_a!$ ($a = 1; 2; 3$) and $s_a = (k_1 - j_a k_0) s_a^2!$ ($a = 1; 2; 3$). For each degree $e \geq 3$ and color $c = 0; 1; 2; 3$, we want to construct elements of degree e and color c in $\text{Ker}(\mathbb{Z}^2 \xrightarrow{d} \mathbb{Z}^3)$ by taking K^{cl} -linear combinations of $k_0^m k_1^n$ ($m = 0; 1; 2; 3$) and $k_0^m k_1^n s_a$ ($a = 1; 2; 3$). We set $s_0 = k_0 s_0^2!$. We have the relation $s_0 = s_1 + s_2 + s_3$. A straightforward calculation shows that for $e = 3; 4$ we have none; for $e = 5$ we find $s_0 = k_0 s_0$ for color 0 and $s_a = (k_1 - j_a k_0) s_a$ for color $a = 1; 2; 3$; for $e = 6$ we find two color 0 elements:

$$\begin{aligned} s_4 &= (k_1 - j_1 k_0) s_1 - (k_1 - j_2 k_0) s_2 - j_{1,2} k_0 s_0; \\ s_5 &= (k_1 - j_2 k_0) s_2 - (k_1 - j_3 k_0) s_3 - j_{2,3} k_0 s_0; \end{aligned}$$

For $e = 5; 6$ the above elements span the degree e cohomology classes. For $e = 8$, we have 4 obvious elements k_{i-j} ($i = 0; 1; j = 4; 5$), and in addition, we find

$$s_6 = \frac{1}{4} k_0^2 \sum_{a=1, b \neq a} j_{a,b} s_a + k_0 \sum_{a=1} (k_1 - j_a k_0) s_a :$$

These 5 elements span the degree 8 cohomology classes.

Finally, we show that s_a ($0 \leq a \leq 6$) are the generators of the F^{cl} -module $\text{Ker}(\mathbb{Z}^2 \xrightarrow{d} \mathbb{Z}^3) = \text{Im}(\mathbb{Z}^1 \xrightarrow{d} \mathbb{Z}^2)$. For odd $e = 2n + 3 \geq 7$, let us consider the color 0 case. The cases of other colors are similar. The degree $2n + 3$ and color 0 space in $\mathbb{Z}^2 = \text{Im}(\mathbb{Z}^1 \xrightarrow{d} \mathbb{Z}^2)$ has the K^{cl} -basis consisting of $n + 1$ elements $k_0^j k_1^{(n-j)}$ ($0 \leq j \leq n$). We have

$$d(k_1^n s_0) = 3 s_0^{2n} ds_1 \wedge ds_2 \wedge ds_3 \in \mathbb{Z}^3 :$$

Therefore, the K^{cl} -dimension of the degree $2n + 3$ and color 0 subspace of $\text{Ker}(\mathbb{Z}^2 \xrightarrow{d} \mathbb{Z}^3) = \text{Im}(\mathbb{Z}^1 \xrightarrow{d} \mathbb{Z}^2)$ is n . Since the K^{cl} -linearly independent elements $k_0^j k_1^{(n-j)}$ ($0 \leq j \leq n - 1$) belongs to this subspace, they span the subspace.

For even $e = 2n + 4 \geq 10$, the degree $2n + 4$ subspace of $\mathbb{Z}^2 = \text{Im}(\mathbb{Z}^1 \xrightarrow{d} \mathbb{Z}^2)$ has the K^{cl} -basis consisting of $3(n + 1)$ elements $k_0^j k_1^{(n-j)} s_a$ ($0 \leq j \leq n; a = 1; 2; 3$). A simple calculation shows that the elements $d(k_1^n s_0)$, $d(k_0^n s_a)$ ($a = 1; 2; 3$) are K^{cl} -linearly independent in \mathbb{Z}^3 . Therefore, the K^{cl} -dimension of the degree $2n + 4$ subspace of $\text{Ker}(\mathbb{Z}^2 \xrightarrow{d} \mathbb{Z}^3) = \text{Im}(\mathbb{Z}^1 \xrightarrow{d} \mathbb{Z}^2)$ is $3n + 1$. On the other hand, we obtain $3n + 1$

independent elements of the subspace from the degree 6 elements e_4, e_5 and the degree 8 element e_6 . We conclude that they span the subspace. \square

We have explicitly constructed free bases for the spaces $H^{cl[0]}, H^{cl[1]}, H^{cl[2]}$. In [32], similar spaces are studied for hypersurface singularities E_6, E_7, E_8 . The results in this appendix supplement some part of his construction in D_5 , where the elliptic singularity is given by a complete intersection of two quadrics.

Appendix B. Cycles and integrals in the classical limit

In the classical limit, the defining relations of the Sklyanin algebra turns into (A.9), (A.10) which define an algebraic surface $S \subset \mathbb{C}^4$. In this section, we study the classical limit of the functional Tr and identify it with an integral over a cycle on S .

Sklyanin's formulas for the representation (2.16) gives an explicit uniformization of S . Consider the limit

$$! 0; \quad ! 1; \quad \text{nite:}$$

In the right hand side of (2.16), we replace q_i by $2iv$ where v is the variable canonically conjugate to u ,

$$2fv; ug = 1:$$

In the limit, the formulas for $s_0 = S_0, s_a = S_a$ ($a = 1, 2, 3$) tend to

$$(B.1) \quad s(u; v) = c \frac{c_{+1}(2u) e^{2iv} + c_{-1}(2u) e^{2iv}}{c_1(2u)};$$

where $(u; v) \in \mathbb{C}^2$ and

$$c_0 = \frac{1}{2}; \quad c_a = p_a \pi_a \frac{a+1}{2} \frac{(0)}{1} \quad \text{for } a = 1, 2, 3:$$

Eq. (B.1) provides a parametrization of the surface (A.9), (A.10) with

$$(B.2) \quad k_0 = \frac{1}{0} \frac{()^2}{1}; \quad k_1 = \frac{1}{0} \frac{()^2}{1} \log_1():$$

For definiteness, we consider the case $2iR_{>0}, 0 < < 1=2$. The functions s have common periods

$$e_1 = (1; 0); \quad e_2 = (0; 1); \quad e_3 = (; 2);$$

so $(u; v)$ should be regarded as variables on $\mathbb{C}^2 = \left(\prod_{i=1}^3 \mathbb{Z} e_i \right)$. We shall consider the fundamental domain:

$$0 \leq \text{Re}(u) < 1; \quad 0 \leq \text{Im}(u) < \frac{1}{i}; \quad 0 \leq \text{Re}(v) < 1:$$

There are also pole divisors of s at $u = p_i$, where


$$(B.3) \quad p_0 = 0; \quad p_1 = \frac{1}{2}; \quad p_2 = \frac{1+}{2}; \quad p_3 = \frac{2}{2}:$$

If we neglect these divisors, there are, obviously, three non-trivial 2-cycles which are tori with generators $(e_1; e_3), (e_1; e_2), (e_2; e_3)$. We denote them by $\gamma_0, \gamma_1, \gamma_2$.

One can choose the tori τ_1, τ_2 so that they do not intersect with the pole divisors. As for τ_0 , the first impression is that it hits the divisors. But actually one has to be very careful at this point. From the formulae (B.1) it follows that there are no singularities at (B.3) if $u = p_0; p_1$ and $v = 0; 1=2$, or $u = p_2; p_3$ and $v = \infty; +1=2$. Hence the actual divisors are $D = D^0 \cup D^\infty$, with


$$(B.4) \quad \begin{aligned} D^0 &= \{ \sum_{i=0,1} f(p_i; v) \mid 0 \leq \operatorname{Re}(v) < 1; v \neq 0; 1=2g \} \\ D^\infty &= \{ \sum_{i=2,3} f(p_i; v) \mid 0 \leq \operatorname{Re}(v) < 1; v \neq \infty; +1=2g \} \end{aligned}$$

This means, first of all, that we can modify τ_0 at $v = 0$ and $v = \infty$ into a well-defined cycle without intersection with D , as depicted in Fig. 1 below.



g. 1

Now we have another possibility. We can draw spheres $\sigma_0; \sigma_1$ which have as south (resp. north) poles the points $(p_0; 0), (p_1; 0)$ (resp. $(p_0; 1=2), (p_1; 1=2)$). In the vicinity of these points they are parallel to the u -plane, and every section of it by the plane $\operatorname{Re}(v) = a$ for $0 < a < 1=2$ is a cycle around $p_0; p_1$ in the u -plane.



g. 2

These spheres do not intersect D . Similarly, we construct spheres $\sigma_2; \sigma_3$ which have as south (resp. north) poles the points $(p_2; \infty), (p_3; \infty)$ (resp. $(p_2; +1=2), (p_3; +1=2)$).

The homology group $H_2(S; \mathbb{Z}) \cong \mathbb{Z}^7$ is known. We will see below that our cycles $\sigma_0; \sigma_1; \sigma_2; \sigma_3; \tau_0; \tau_1; \tau_2 \in H_2(S; \mathbb{Z})$ give a linearly independent basis in $H_2(S; \mathbb{C})$.

Using our seven monomials we construct 2-forms

$$\begin{aligned} \omega &= \frac{1}{4} \frac{ds_1 \wedge ds_2}{s_0 s_3}; \quad \omega_0 = s_0^2 \omega; \quad \omega_3 = s_3^2 \omega; \\ &= s \omega; \quad \omega = 0; 1; 2; 3; \end{aligned}$$

By direct computation we check that in Sklyanin's parameterization (B.1)

$$(B.5) \quad \omega = du \wedge dv;$$

This formula allows us to calculate the integrals explicitly.

Let us start with the cycles Z . We obtain

$$\omega = \omega = 0; \quad \omega = 2 \quad c \frac{+1 \left(\begin{smallmatrix} 0 \\ 1 \end{smallmatrix} \right)}{1};$$

where ω are elements of the matrix:

$$\begin{aligned} &\begin{pmatrix} 0 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 \end{pmatrix} \\ &= \begin{pmatrix} B & B & B & B & B \\ C & C & C & C & C \\ A & A & A & A & A \end{pmatrix} : \end{aligned}$$

Note, in particular, that

$$(B.6) \quad \begin{aligned} &\omega \\ &P_3 = 0 \end{aligned}$$

for $\omega = 1; 2; 3$.

For the integrals over Z_k ($k = 1; 2$), we find

$$\begin{aligned} \omega &= 0; \quad \omega = 1; \quad \omega = ; \\ \int_{Z_k} \omega &= 2 \quad c \frac{+1 \left(\begin{smallmatrix} 0 \\ 1 \end{smallmatrix} \right)}{1} \quad \log_{+1} \left(\begin{smallmatrix} 0 \\ 1 \end{smallmatrix} \right); \\ \int_{Z_1} \omega &= 2 \quad c \frac{+1 \left(\begin{smallmatrix} 0 \\ 1 \end{smallmatrix} \right)}{1} \quad \log_{+1} \left(\begin{smallmatrix} 0 \\ 1 \end{smallmatrix} \right) + 2 \quad i; \\ \int_{Z_2} \omega &= 2 \quad c \frac{+1 \left(\begin{smallmatrix} 0 \\ 1 \end{smallmatrix} \right)}{1} \quad \log_{+1} \left(\begin{smallmatrix} 0 \\ 1 \end{smallmatrix} \right) + 2 \quad i; \end{aligned}$$

For ω_0 , we introduce

$$\omega_0 = \omega_0 \quad 2 \quad 1 + \frac{1}{2} X^3 : \quad \omega_0 = 0$$

Then, we have

$$\begin{aligned} \int_{Z_0} \omega &= \int_{Z_0} \omega = 0; \quad \int_{Z_0} \omega = 2 \quad c \frac{+1 \left(\begin{smallmatrix} 0 \\ 1 \end{smallmatrix} \right)}{1} \quad \log_{+1}^2 \left(\begin{smallmatrix} 0 \\ 1 \end{smallmatrix} \right); \\ \int_{\omega_0} \omega &= \int_{\omega_0} \omega = 0; \quad \int_{\omega_0} \omega = 2 \quad c \frac{+1 \left(\begin{smallmatrix} 0 \\ 1 \end{smallmatrix} \right)}{1} \quad \log_{+1}^2 \left(\begin{smallmatrix} 0 \\ 1 \end{smallmatrix} \right); \end{aligned}$$

Using the above formulae we can calculate the determinant of the period matrix as

$$\det(P) = \text{Const} \cdot (2\pi)^{-(1/2)} \left(\frac{1}{2} \right)^{1/2} \left(\frac{1}{2} \right)^{1/2} \frac{\partial}{\partial} \log \frac{1}{2} \left(\frac{1}{2} \right);$$

whence we conclude that for generic the period matrix is non-degenerate.

Comparison with the quantum formulae (2.20), (2.21) shows the exceptional role of the cycle γ_0 as $\gamma_0 \neq 0$,

$$\begin{aligned} \text{Tr}(S) &= \frac{1}{2} \sum_{s=0}^Z s! \\ \text{Tr}(S^2) &= \frac{1}{2} \sum_{s=0}^Z s^2! \quad \left(\begin{array}{l} 1 \quad (s=0); \\ \frac{1}{2} \quad (s=1;2;3); \end{array} \right. \end{aligned}$$

Actually, classical limits of all the elements of F in (2.22)–(2.24) can be found among integrals over the cycles $\gamma_0; \gamma_1; \gamma_2$. This fact played an important heuristic role in the calculation of traces.

Appendix C. Technical Lemmas

In this Appendix, we collect some technical matters related to Tr .

The first is the color conservation for Tr .

Lemma C.1. If $A \in A^{(m,n)}$, then

$$\text{Tr} A = 0 \quad \text{unless } (m;n) = (0;0):$$

Proof. Set

$$'_1(f)(u) := f(u + \frac{1}{2})e^{2iku}; \quad '_2(f)(u) := f(u + \frac{1}{2}):$$

One verifies easily that $'_1; '_2 \in \text{End}(V^{(k)})$ and

$$\begin{aligned} '_1^{(k)}(S) &= '_1^{-1} = (1)^{-1} \cdot (k) (S); \\ '_2^{(k)}(S) &= '_2^{-1} = (1)^{-2} \cdot (k) (S); \end{aligned}$$

where $\gamma = (\gamma_1; \gamma_2) \in \mathbb{Z}_2 \times \mathbb{Z}_2$. It follows that

$$\text{tr}_{V^{(k)}}^{(k)}(A) = (1)^m \text{tr}_{V^{(k)}}^{(k)}(A) = (1)^n \text{tr}_{V^{(k)}}^{(k)}(A);$$

whence the lemma. \square

The next Lemma is concerned about the uniqueness of the representation of Tr .

Lemma C.2. Assume $\text{Im} \gamma; \text{Im} \delta > 0$, $\gamma \in \mathbb{Q} + \mathbb{Q}i$. Let $g_i (i=1;2;3)$ be elliptic functions with periods 1 and γ , and set $f(u) = \gamma(u)g(u) + u g_2(u) + g_3(u)$, where $\gamma(u) = (1-2i) \frac{0}{1}(u) = \frac{0}{1}(u)$. If there exists an $N > 0$ such that $f(k\gamma) = 0$ holds for all integers $k > N$, then we have $g_i(u) = 0 (i=1;2;3)$.

Proof. We divide into three cases, (i) $g_1(u) = g_2(u) = 0$, (ii) $g_1(u) \neq 0$ and $g_2(u) = 0$, (iii) $g_2(u) \neq 0$.

Since $K = \{k \mid j_k > N\}$ has accumulation points, the assertion is evident in case (i). Let us show that (ii), (iii) lead to contradictions.

In case (ii), considering $f(u) = g_1(u)$ we may assume $g_1(u) = 1$. Then $f(u + \omega) = f(u) + 1$. Choose a point $u_0 \in \mathbb{C} \setminus L$ which is not a pole of $f(u)$. One can find a sequence of integers $k_1 < k_2 < \dots$, $k \geq 1$, such that $k_n \omega$ tends to u_0 in $\mathbb{C} = L$. Since $\text{Im } \omega > 0$, if we write $k_n \omega = a_n + b_n i$ ($a_n, b_n \in \mathbb{R}$), then the integer part of b_n diverges. Therefore $f(k_n \omega)$ diverges as $n \rightarrow \infty$, which contradicts to the assumption $f(k_n \omega) = 0$.

In case (iii), we may assume $g_2(u) = 1$. Set $F(u) = f(u + \omega) - f(u)$. We have $F(k \omega) = 0$ ($k > N$), $F(u + 1) = F(u)$, and $F(u + \omega) - F(u) = g(u + \omega) - g_1(u)$. Hence $F(u) = G(u) + \varphi(u)(g(u + \omega) - g_1(u))$ with some elliptic function $G(u)$. From cases (i), (ii) we conclude that $F(u) = 0$. In particular g_1 is a constant. Considering $f^0(u + \omega) = f^0(u)$, we find that $f(u)$ is a linear function. Clearly this is impossible. \square

Let us sketch the derivation of the formulas for $\text{Tr } S$, $\text{Tr } S^2$. We have the standard functional relation

$$t^{(1)} - t - \frac{k}{2} - t^{(k)} - t + \frac{1}{2} = (t + \omega)^{(k+1)}(t) + (t)t^{(k-1)}(t + \omega)$$

for the transfer matrices

$$t^{(k)}(t) := \text{tr}_{V^{(k)}} r_{a, N}^{(k;1)}(t - t_N) = \sum_{a,1}^{(k;1)} r(t - t_1) \in \text{End}(V^{\otimes N});$$

where $r^{(k;1)}(t) := (\text{id} \otimes L(t))$ and $(t) = \sum_{j=1}^N [t - t_j]$ ($k=2$). Choosing $N = 1$; $t_1 = 0$ and applying (C.1), we easily find (2.20). The same method is applicable for (2.21). We find it slightly simpler to use the difference equation for the matrices $\mathcal{X}_{a;2}^{(1;2)}$.

Appendix D. Transformation properties of $\mathcal{X}_n^{(i;j)}$

Let us study the transformation properties of $\mathcal{X}_n^{(i;j)}$ with respect to the shift of variables by half periods. For that purpose we exploit the order 4 automorphism s_1, s_3 of the Sklyanin algebra A given by

$$\begin{aligned} s_1(S_0) &= \frac{1(\omega)}{2(\omega)} S_1; \quad s_1(S_1) = \frac{2(\omega)}{1(\omega)} S_0; \quad s_1(S_2) = i \frac{3(\omega)}{0(\omega)} S_3; \quad s_1(S_3) = i \frac{0(\omega)}{3(\omega)} S_2; \\ s_3(S_0) &= \frac{1(\omega)}{0(\omega)} S_3; \quad s_3(S_1) = \frac{2(\omega)}{3(\omega)} S_2; \quad s_3(S_2) = \frac{3(\omega)}{2(\omega)} S_1; \quad s_3(S_3) = \frac{0(\omega)}{1(\omega)} S_0. \end{aligned}$$

In terms of the L-operator they can be written as

$$(D.1) \quad s_1(L(t)) = L\left(t + \frac{1}{4}\right) s_1;$$

$$(D.2) \quad s_3(L(t)) = L\left(t + \frac{1}{4}\right) s_3 = (i) e^{i(2t + \frac{1}{4})} :$$

Lemma D.1. Let $A \in A_n$ be an element of the Sklyanin algebra of even degree n , and let $\text{Tr}_{t=1}^n A = g_{A,1}(t) = g_{A,2}(t)$ where $g_{A,1}, g_{A,2}$ are as in (2.18). Then

$$\begin{aligned} \text{Tr}_{t=1+\frac{1}{2}}^n A &= g_{A,1}(t) \left(-t + \frac{1}{2} \right) g_{A,2}(t); \\ \text{Tr}_{t=1+\frac{3}{2}}^n A &= g_{A,1}(t) + \frac{1}{2} g_{A,3}(t) \left(-t + \frac{1}{2} \right) g_{A,2}(t) e^{i(-2+2t)\frac{n-2}{2}}; \end{aligned}$$

where $g_{A,3}(t) = g_{A,1}(t+1) - g_{A,1}(t)$ is an elliptic function.

Proof. In view of Theorem A.1, it is enough to consider elements A of the form $m \frac{S}{S}$, where m is a polynomial in $K_0; K_2$ of degree $(n-2)/2$. For $n=2$, the assertion can be verified from the explicit formula (2.21).

Let I_t denote the two-sided ideal of A generated by $K_0 - 4_1(t)^2, K_2 - 4_1(t+1)4_1(t-1) = 4_1(2)^2$, and let $\$ _t : A \rightarrow A/I_t$ be the projection. From (2.15) and (2.17) we have

$$\$ _t L_1 \frac{S}{2} L_2 \frac{S}{2} P_{12} = \frac{4_1(t-s)4_1(t+s)}{4_1(2)^2} P_{12};$$

Along with (D.1) and (D.2) it follows that for $i=0,2$

$$\begin{aligned} \$ _{t+1=2}^{-1}(K_i) &= \$ _t(K_i); \\ \$ _{t+2}^{-3}(K_i) &= \$ _t(K_i) - (1)e^{i(-2+2t)\frac{n-2}{2}}; \end{aligned}$$

The Lemma follows from these relations. \square

Proposition D.2. The $X_{a,n}^{(i,j)}$ obey the following transformation laws.

(D.3)

$$\begin{aligned} & \frac{1}{k} \frac{1}{k} X_{a,n}^{(i,j)} \left(-k + t \frac{1}{2}; \right) \\ &= X_{a,n}^{(i,j)} \left(-k; t \right) \begin{pmatrix} 1 & 1 \\ k & k \end{pmatrix} Q_{\substack{p \notin \{i,j\} \\ p \neq 1}} \frac{1}{p} \frac{1}{p} \quad (k \neq i,j); \\ & \quad (1)^{n-1} \quad (k = i,j); \end{aligned}$$

(D.4)

$$\begin{aligned} & \frac{3}{k} \frac{3}{k} X_{a,n}^{(i,j)} \left(-k + t \frac{1}{2}; \right) \\ &= \begin{pmatrix} 3 & 3 \\ k & k \end{pmatrix} X_{a,n}^{(i,j)} \left(-k; t \right) \begin{pmatrix} 3 & 3 \\ k & k \end{pmatrix} Q_{\substack{p \notin \{i,j\} \\ p \neq 1}} \frac{3}{p} \frac{3}{p} \quad (k \neq i,j); \\ & \quad \frac{1}{2} \frac{1}{2} X_{3,n}^{(i,j)} \left(-k; t \right) (1)^{n-1} \quad (k = i,j); \end{aligned}$$

In the last line the upper (resp. lower) sign is chosen for $k=i$ (resp. $k=j$). In particular, $X_{2,n}^{(i,j)}$; $X_{3,n}^{(i,j)}$ are elliptic functions of t ; n with periods 1, $\frac{1}{2}$.

Proof. It is enough to prove the case $(i; j) = (1; 2)$. If $k \notin 1; 2$, this is a simple consequence of the transformation law of the L-operator

$$\begin{aligned} L(t + \frac{1}{2}) &= {}^1L(t)^{-1}; \\ L(t + \frac{1}{2}) &= {}^3L(t)^{-3} e^{2i(2t + \frac{1}{2})}; \end{aligned}$$

Consider the case $k = 1$. Using the automorphism 1 we have

$$\begin{aligned} & \chi_n^{(1;2)}(t_1 + \frac{1}{2}; \dots)_{12} \prod_{p=3}^n [t_{1p} + 1][t_{2p}] \\ &= \text{Tr}_{t_{12} + \frac{1}{2}} T^{[1]} \left(\frac{t_1 + t_2}{2} + \frac{1}{4}; t_1 + \frac{1}{2}; \dots; t_{12} P_{12} P_{11} P_{22} \right) \\ &= \text{Tr}_{t_{12} + \frac{1}{2}} {}^1 T^{[1]} \left(\frac{t_1 + t_2}{2}; t_1; \dots; t_{12} \prod_{p=2}^n \frac{1}{p} P_{12} P_{11} P_{22} \right) \\ &= \frac{1}{1} \frac{1}{1} \text{Tr}_{t_{12} + \frac{1}{2}} {}^1 T^{[1]} \left(\frac{t_1 + t_2}{2}; t_1; \dots; t_{12} P_{12} P_{11} P_{22} \prod_{p=3}^n \frac{1}{p} \right); \end{aligned}$$

Applying Lemma D.1 we obtain (D.3) with $k = 1$. Eq. (D.4) is shown similarly. The case $k = 2$ can be obtained by using the translation invariance. \square

Acknowledgments. Research of HB is supported by the RFFI grant # 04-01-00352. Research of MJ is supported by the Grant-in-Aid for Scientific Research B2{16340033 and A2{14204012. Research of TM is supported by the Grant-in-Aid for Scientific Research A1{13304010. Research of FS is supported by INTAS grant # 03-51-3350 and by EC networks "EUCLID", contract number HPRN-CT-2002-00325 and "ENIGMA", contract number MRTN-CT-2004-5652. Research of YT is supported by University of Tsukuba Research Project. This work was also supported by the grant of 21st Century COE Program at Graduate School of Mathematical Sciences, the University of Tokyo, and at RIMS, Kyoto University.

MJ, TM and YT are grateful to K. Saito for explaining his theory of simple-elliptic singularities. MJ thanks V. Mangazeev, A. Zabrodin, A. Mironov, C. Kor and R. Weston for discussions. HB thanks M. Lashkevich for tight cooperation in numerical checking. He also thanks F. Gohmann, A. Klumper and K. Fabricius for discussions. Last but not least, HB and FS wish to thank the University of Tokyo for the invitation and hospitality where the present work was started, and HB, MJ, TM and YT to Universite Paris VI for hospitality, where it was completed.

References

- [1] R. Baxter, Exactly solved models in statistical mechanics, Academic Press, London 1982.
- [2] R. Baxter, Solvable eight-vertex model on an arbitrary planar lattice, Phil. Trans. Royal Soc. London, 289 (1978) 315{346.

- [3] H. Boos, M. Jimbo, T. Miwa, F. Smirnov and Y. Takeyama, A recursion formula for the correlation functions of an inhomogeneous XXX model, hep-th/0405044, to appear in Algebra and Analysis.
- [4] H. Boos, M. Jimbo, T. Miwa, F. Smirnov and Y. Takeyama, Reduced qKZ equation and correlation functions of the XXZ model, hep-th/0412191, to appear in Commun. Math. Phys.
- [5] H. Boos and V. Korepin, Quantum spin chains and Riemann zeta functions with odd arguments, hep-th/0104008, J. Phys. A 34 (2001) 5311{5316.
- [6] H. Boos, V. Korepin and F. Smirnov, Emptiness formation probability and quantum Knizhnik-Zamolodchikov equation, hep-th/0209246, Nucl. Phys. B Vol. 658/3 (2003) 417 {439.
- [7] H. Boos, V. Korepin and F. Smirnov, New formulae for solutions of quantum Knizhnik-Zamolodchikov equations on level 4, hep-th/0304077, J. Phys. A 37 (2004) 323{336.
- [8] K. Fabricius and B. McCoy, Functional equations and fusion matrices for the eight vertex model, Publ. RIMS, Kyoto Univ. 40 (2004) 905{932.
- [9] K. Fabricius and B. McCoy, New developments in the eight-vertex model II. Chains of odd length, cond-mat/0410113.
- [10] A. Odesskii and B. Feigin, Sklyanin elliptic algebra, Funct. Anal. and its Appl. 23 (1990) 207{214.
- [11] A. Odesskii, Elliptic algebras, Russian Math. Surveys 57 (2002) 1{50.
- [12] P. A. Grieth and Harris, Principles of algebraic geometry, A. Wiley Interscience, 1978
- [13] F. Gohmann, A. Klumper and A. Seel, Integral representation of the density matrix of the XXZ chain at finite temperatures, J. Phys. A 38 (2005) 1833-1842 ; cond-mat/0412062.
- [14] A. Erdelyi et al., Higher transcendental functions, McGraw-Hill, 1953.
- [15] M. Jimbo and T. Miwa, Quantum Knizhnik-Zamolodchikov equation at $\hbar=1$ and correlation functions of the XXZ model in the gapless regime, J. Phys. A 29 (1996) 2923{2958
- [16] M. Jimbo, T. Miwa and A. Nakayashiki, Difference equations for the correlation functions of the eight vertex model, J. Phys. A 26 (1993), 2199{2209.
- [17] M. Jimbo, T. Miwa, K. Miki and A. Nakayashiki, Correlation functions of the XXZ model for $\Delta < 1$, Phys. Lett. A 168 (1992), 256{263.
- [18] N. Kitanine, J.-M. Maillet, N. A. Slavnov and V. Terras, Master equation for spin-spin correlation functions of the XXZ chain, hep-th/0406190.
- [19] N. Kitanine, J.-M. Maillet, N. A. Slavnov and V. Terras, Dynamical correlation functions of the XXZ spin-1/2 chain, hep-th/0407223.
- [20] N. Kitanine, J.-M. Maillet and V. Terras, Correlation functions of the XXZ Heisenberg spin- $\frac{1}{2}$ -chain in a magnetic field, Nucl. Phys. B 567 (2000), 554{582.
- [21] C. Korner, Auxiliary matrices on both sides of the equator, J. Phys. A 38 (2005), 47{66.
- [22] C. Korner, A Q-operator identity for the correlation functions of the infinite XXZ spin-chain, hep-th/0503130
- [23] G. Kato, M. Shiroishi, M. Takahashi, K. Sakai, Third-neighbor and other four-point correlation functions of spin-1/2 XXZ chain, J. Phys. A: Math. Gen. 37 (2004) 5097.
- [24] S. Lukyanov and Y. Pugai, Multipoint local height probabilities in the integrable RSOS model, Nucl. Phys. B 473 [FS] (1996) 631-658.
- [25] M. Lashkevich and Y. Pugai, Free field construction for correlation functions of the eight-vertex model, Nucl. Phys. B 516 (1998) 623-651.
- [26] M. Lashkevich and Y. Pugai, Nearest neighbor two-point correlation function of the Z-invariant eight-vertex model, JETP Lett. 68 (1998) 257-262.
- [27] A. Pronko and Yu. Stroganov, Bethe equations \on the wrong side of equator, J. Phys. A 32 (1999), 2333-2340

- [28] S.N.M. Ruijsenaars, First order analytic difference equations and integrable quantum systems, *J. Math. Phys.* 38 (1997) 1069{1146
- [29] K. Saito, Einfach-elliptische Singularitäten, *Inventiones math.* 23 (1974), 289{325.
- [30] K. Saito, Period mapping associated to a primitive form, *Publ. RIMS, Kyoto Univ.* 19 (1983), 1231{1264.
- [31] K. Saito, Extended affine root systems I (Coxeter transformations), *Publ. RIMS, Kyoto Univ.* 21 (1985), 75{179.
- [32] K. Saito, On the periods of primitive integrals I, *RIMS preprint* 412 (1982).
- [33] J. Shiraishi, Free field constructions for the elliptic algebra $A_{qp}(\hat{\mathfrak{sl}}_2)$ and Baxter's eight-vertex model, *Int. J. Mod. Phys. A* 19 (2004) 363–380; *math.QA/0302097*.
- [34] J. Shiraishi, A commutative family of integral transformations and basic hypergeometric series I,II, *math.QA/0501251*, 0502228.
- [35] E.K. Sklyanin, Some algebraic structures connected with the Yang-Baxter equation, *Func. Anal. and Appl.* 16 (1982) 27{34.
- [36] E.K. Sklyanin, Some algebraic structures connected with the Yang-Baxter equation. Representations of quantum algebras, *Func. Anal. and Appl.* 17 (1983) 34{48.
- [37] M. Takahashi, G. Kato, M. Shiroishi, Next nearest-neighbor correlation functions of the spin-1/2 XXZ chain at massive region, *J. Phys. Soc. Jpn.* 73 (2004) 245.

HB: Physics Department, University of Wuppertal, D-42097, Wuppertal, Germany⁴

E-mail address: boos@physik.uni-wuppertal.de

MJ: Graduate School of Mathematical Sciences, The University of Tokyo, Tokyo 153-8914, Japan

E-mail address: jimbomic@ms.u-tokyo.ac.jp

TM: Department of Mathematics, Graduate School of Science, Kyoto University, Kyoto 606-8502, Japan

E-mail address: tetsuji@math.kyoto-u.ac.jp

FS⁵: Laboratoire de Physique Théorique et Hautes Energies, Université Pierre et Marie Curie, Tour 16 1^{er} étage, 4 Place Jussieu 75252 Paris Cedex 05, France

E-mail address: smirnov@lpthe.jussieu.fr

YT: Graduate School of Pure and Applied Sciences, Tsukuba University, Tsukuba, Ibaraki 305-8571, Japan

E-mail address: takeyama@math.tsukuba.ac.jp

⁴ on leave of absence from the Institute for High Energy Physics, Protvino, 142281, Russia

⁵ Membre du CNRS